



UNIVERSITI PUTRA MALAYSIA

***ISOLATION AND CHARACTERIZATION OF LACTIC ACID
BACTERIA FROM GASTROINTESTINAL TRACT OF SNAKEHEAD
(Channa striatus, Bloch) AS PROBIOTIC FOR FRESHWATER FISH***

SEYED KAMALEDDIN ALLAMEH

IB 2012 5

**ISOLATION AND CHARACTERIZATION OF LACTIC ACID BACTERIA
FROM GASTROINTESTINAL TRACT OF SNAKEHEAD
(*Channa striatus*, Bloch) AS PROBIOTIC
FOR FRESHWATER FISH**


BY

SEYED KAMALEDDIN ALLAMEH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirement for the Degree of
Doctor of Philosophy**

June 2012

In the name of God



“From water everything is alive”

(Quran, Sourah: Al-Anbiya, Vers: 30)

DEDICATION

To all teachers and researchers



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

ISOLATION AND CHARACTERIZATION OF LACTIC ACID BACTERIA FROM GASTROINTESTINAL TRACT OF SNAKEHEAD (*Channa striatus*, Bloch) AS PROBIOTIC FOR FRESHWATER FISH

By

SEYED KAMALEDDIN ALLAMEH

June 2012

Chairman: Associate Professor Hassan bin Hj. Mohd. Daud, PhD

Faculty/Institute: Institute of Bioscience

The intensive application of antibiotics to prevent and control the bacterial diseases in aquaculture has resulted in development of antibiotic-resistant bacteria. Therefore, probiotics as an alternative strategy have been suggested to be used as replacement for antimicrobial drugs and growth promoters. Lactic acid bacteria (LAB) constitute a group of G+ve bacteria with a high ability to produce antibacterial compounds and improve fish performance which makes them excellent probiotics. In the present study, snakehead (*Channa striatus*), an indigenous commercial freshwater fish in Southeast Asia countries was chosen for the isolation of LAB. Isolates obtained on de Man Rogosa and Sharp (MRS) agar and broth showed 27 and 9 pure isolates from the fingerling and adults fishes, respectively. According to the antagonistic plate test against *Aeromonas hydrophila* as a main freshwater fish pathogen, five LABs i.e. three isolates from the fingerlings and two from the adults that showed the greatest inhibition zone were selected. The selected LABs were identified as *Aerococcus*

viridans, *Lactobacillus delbreuckii* sp. *delbreuckii* and *Enterococcus faecalis* from fingerlings and *Lactobacillus fermentum* and *Leuconostoc mesenteroides* sp. *mesenteroides* from adult fish. Probiotic property evaluation of LAB candidates showed that they could survive and grow at pH 3 up to 8 ($P < 0.05$). They could also tolerate bile salt concentrations from 0.0 to 0.3% ($P < 0.05$). The LABs were active at different levels of NaCl (0.0 to 4%) and also, at various temperatures ranging from 15 °C to 45 °C, but showed no growth at 10 °C and 50 °C. Antagonistic effects against three tested fish pathogens i.e. *A. hydrophila*, *Pseudomonas aeruginosa* and *Shewanella putrefaciens* indicated that *En. faecalis*, *L. fermentum* and *Leu. mesenteroides* sp. *mesenteroides* had the highest inhibition activities. The antibiotic sensitivity test showed that *En. faecalis* had more antibiotic's resistance property against some antibiotics as compared to other LAB ($P < 0.05$). According to probiotic characterization as screening, *En. faecalis*, *L. fermentum* and *Leu. mesenteroides* were selected for *in vivo* experiments. The viability of LABs in feed stored at 4 °C was higher than 25 °C during four weeks storage. Significant increased of the LAB proportion was observed in the fish intestine fed LAB-fortified diets as compared to the control group ($P < 0.05$). In addition, the lower G-ve bacteria population in the fish intestine was observed for experimental groups as compared to the control group. Effects of mono and multi-species of LAB candidates on the body composition of Javanese carp (*Puntius gonionotus*) indicated that LAB-fortified diets could not seriously affect the chemical composition of experimental fish carcasses. The survivability was the same for all experimental groups. The concentration of 10^7 cfu/g diets was overall more effective than 10^5 and 10^9 cfu/g diets. The use of *En.*

faecalis as probiotic had more positive effects than *L. fermentum* and *Leu. mesenteroides* on growth, feed conversion ratio, specific growth rate, and protein efficiency ratio ($P < 0.05$). Moreover, the use of *En. faecalis* resulted in an increase in the production of protease and lipase enzymes in the digestive tract of Javanese carp. Results obtained from the short chain fatty acid determination which included acetate, propionate and butyrate showed that treated group with *En. faecalis* could significantly ($P < 0.05$) enhance the propionate and butyrate production as compared to the control. An *in vivo* challenge test of *A. hydrophila* as a fish pathogen with *En. faecalis* as a probiotic on Javanese carp indicated that this probiotic could protect the fish against *A. hydrophila* and showed a higher survivability compared to the control. The presence of *En. faecalis* in the diet could affect immune response to enhance the antibody level as a humoral response. The fish treated with a LAB-fortified diet (*En. faecalis*) and added pathogen (*A. hydrophila*) into the water showed the highest antibody level as compared to the control group ($P > 0.05$). In conclusion, LABs can be normal microbiota in the gastrointestinal tract of the snakehead. In addition, the use of *En. faecalis* as a probiotic had positive effects on overall fish performance, digestive enzymes and short chain fatty acid production, survivability in challenge with pathogen and increased humoral immune response in Javanese carp. This LAB as an environmental friendly agent can be suggested as an alternative to antibiotics in freshwater fish culture.

Abstrak tesis untuk dikemukakan kepada Senat, Universiti Putra Malaysia bagi memenuhi keperluan ijazah Doktor Falsafah

**PEMENCILAN DAN PENCIRIAN BAKTERIA ASID LAKTIK DARI
SALURAN GASTROUSUS IKAN HARUAN (*Channa striatus*, Bloch 1793)
SEBAGAI PROBIOTIK UNTUK IKAN AIR TAWAR**

Oleh

SEYED KAMALEDDIN ALLAMEH

Jun 2012

Pengerusi: Prof. Madya Hassan bin Hj. Mohd. Daud, PhD

Fakulti/Institut: Institut Biosains

Penggunaan antibiotik secara intensif bagi mengelak dan merawat penyakit bakteria dalam akuakultur telah menyebabkan terjadinya bakteria yang rentang antibiotik. Oleh itu probiotik adalah strategi alternatif yang telah disyorkan sebagai pengganti kepada dadah antimikrobial, penggalak pertumbuhan dan peransang imuniti. Bacteria asid laktik (BAL) terdiri dari kumpulan bakteria G+ve yang mempunyai ciri morfologikal, metabolik dan fisiologi yang menjadikan mereka probiotik yang amat baik. Di dalam kajian ini, ikan haruan (*Channa striatus*), sejenis ikan tempatan berkomersial di negara Asia Tenggara telah dipilih untuk pemencilan BAL. Percambahan koloni di atas agar *de Man Rogosa and Sharp* (MRS) dan kaldu memperolehi 27 isolat tulin dari ikan bersaiz jari dan sembilan dari ikan dewasa. Keputusan menunjukkan beberapa BAL yang lazim dalam perut dan saluran usus kedua-dua saiz ikan. Mengikut ujian plat antagonis melawan *Aeromonas hydrophila*

sebagai patogen ikan, lima BAL yakni tiga isolat dari ikan bersaiz jari dan dua dari ikan dewasa menunjukkan zon kesekatlakuan terbesar. Isolat tersebut dikenalpasti sebagai *Aerococcus viridans*, *Lactobacillus delbreuckii* sp. *delbreuckii* dan *Enterococcus faecalis* dari ikan bersaiz jari, dan *Lactobacillus fermentum* dan *Leuconostoc mesenteroides* sp. *mesenteroides* dari ikan dewasa. Penilaian sifat-sifat calon probiotik BAL menunjukkan bahawa mereka boleh hidup dan tumbuh pada pH 3 sehingga pH 8 ($P < 0.05$). Mereka juga boleh tahan dalam garam hempedu dari kepekatan 0.0 ke 0.3% ($P < 0.05$). BAL aktif pada beberapa paras NaCl berlainan (0.0 to 4%) dan juga pada suhu berjulat dari 15 °C to 45 °C, tetapi tiada pertumbuhan pada 10 and 50 °C. Kesan antagonis terhadap tiga patogen ikan iaitu *A. hydrophila*, *Pseudomonas aeruginosa* and *Shewanella putrefaciens* menunjukkan *En. faecalis*, *L. fermentum* dan *Leu. mesenteroides* sp. *mesenteroides* mempunyai aktiviti kesekatlakuan tertinggi. Ujian kepekaan antibiotic menunjukkan *En. faecalis* mempunyai lebih banyak kerentangan terhadap beberapa antibiotik dibandingkan dengan LAB lain (keertian pada $P < 0.05$). Berdasarkan pencirian probiotik, *En. faecalis*, *L. fermentum* and *Leu. mesenteroides* telah dipilih untuk eksperimen *in vivo*. BAL terpilih dengan kesan keertian ($P < 0.05$) menunjukkan kebolehidupan tinggi dalam diet disimpan pada 4 °C and 25 °C. Tambahan lagi, BAL juga dapat merendah percambahan populasi bakteria G-ve dalam saluran gastrousus. Pertambahan yang bererti BAL dapat dilihat dalam usus ($P < 0.05$). Kesan calon BAL secara tunggal dan pelbagai terhadap komposisi badan ikan lampam jawa (*Puntius gonionotus*) menunjukkan bahawa diet yang ditambah dengan BAL tidak memberi kesan jelas terhadap ke atas komposisi kimia karkas ikan ujian. Selain itu tiada kematian yang

dilihat. Oleh itu BAL yang dipilih adalah tidak berbahaya dan tidak memberi kesan sampingan terhadap ikan ujian. Diet berkepekatan 10^7 cfu/g pada keseluruhannya adalah lebih efektif dari kepekatan 10^5 dan 10^9 cfu/g. Walaubagaimana pun, data yang diperolehi menunjukkan penggunaan *En. faecalis* sebagai probiotik memberi lebih kesan positif dari *L. fermentum* and *Leu. mesenteroides* terhadap pertumbuhan, nisbah pertukaran makanan, kadar pertumbuhan tentu dan nisbah kecekapan protein (keertian pada $P < 0.05$). Tambahan lagi, penggunaan *En. faecalis* sebagai probiotik menyebabkan peningkatan pengeluaran enzim protease dan lipase dalam saluran penghadaman ikan lampam jawa (*P. gonionotus*). Juga bakteria ini lebih efektif merembeskan enzim protease dari lipase. Keputusan dari penentuan asid lemak berantai pendek termasuk asetat, propionat dan butirat menunjukkan kumpulan yang dirawat dengan *En. faecalis* boleh, secara bererti ($P < 0.05$) meningkatkan pengeluaran propionat and butirat berbanding dengan kawalan. Ujian cabaran *in vivo* *A. hydrophila* sebagai patogen ikan dengan *En. faecalis* sebagai probiotik ke atas lampam jawa menunjukkan probiotik ini boleh melindungi dari jangkitan dan menunjukkan kesan penghalang yang efektif terhadap patogen ini. Keputusan dari gerak balas sistem imun menunjukkan kehadiran *En. faecalis* dalam diet boleh merangsang gerak balas peningkatan paras antibodi sebagai keimunan humoral. Kumpulan ikan dengan diet yang ditambah dengan BAL dan bersama patogen (*A. hydrophila*) dalam air menunjukkan paras antibodi tertinggi berbanding dengan kawalan ($P > 0.05$). Pada kesimpulannya, BAL boleh menjadi mikrobiota normal dalam saluran gastrousus ikan haruan. Tambahan lagi, penggunaan *En. faecalis* sebagai probiotik memberi kesan positif ke atas keseluruhan perkembangan ikan,

enzim pencernaan dan pengeluaran asid lemak berantai pendek, kemandirian dalam cabaran dengan patogen dan peningkatan ransangan imun humoral di lampam jawa. Bakteria asid laktik ini sebagai satu agen yang mesra alam boleh disyorkan sebagai alternatif kepada penggunaan antibiotik dalam kultur ikan air tawar.



ACKNOWLEDGMENTS

First of all, we really must be grateful to God, who provide everything and opportunities to mankind and also someone's that deliver beneficences to man.

I would like to thank Prof. Datin Paduka Dr. Aini Ideris, the Deputy Vice Chancellor of UPM who honestly and modestly helped me to start my education in UPM and introduced me to Prof. Dr. Fatimah Md. Yusoff of IBS, who is one of the best lecturers in UPM.

I would like to record my gracious thank to my very pleasant supervisor, Assoc. Prof. Dr. Hassan bin Hj. Mohd. Daud, who is a practical scientific man and provided laboratory equipments, materials and guidance to perform the high quality research.

I should appreciate Prof. Dr. Fatimah Md. Yusoff, who as a supervisor and co-supervisor completely supported me to solve the problems during my education and conducted the successful research.

I would like to thank Assoc. Prof. Dr. Che Roos Saad, whose guidance helped me to do the research easily.

I really should appreciate one of the best friends of mine, Dr. Hassan Moeini, who honestly and strongly helped me; without his support and guidance I would not be able to perform all those experiments.

I would like to show gratitude to Dr. Sanjoy Banerjee and Dr. Helena Khatoon, who helped me to carry out the experiments in the lab.


I would like to thank Mr. Perumal Kuppan, who is hard working and always tries to solve student's problems in the Institute of Bioscience.

I should appreciate Ali Baradaran and Mahdi Ebrahimi, who helped me for laboratory work.

There are many people who influenced my research, during these past few years. It is not possible to mention all of them here, but I am grateful for all for their help and untiring support in Aquatic Animal Health Unit, Faculty of Veterinary Medicine, Institute of Bioscience, and Animal Science Department of Faculty of Agriculture.

Finally, much appreciation to my family for their supports and patience.

Lastly, even though there are many hardships on our way to scientific career, it feels great to Learn, Know and Understand.



I certify that a Thesis Examination Committee has met on 20 June 2012 to conduct the final examination of Seyed Kamaleddin Allameh on his thesis entitled “Isolation and Characterization of Lactic Acid Bacteria from Gastrointestinal Tract of Snakehead (*Channa striatus*, Bloch) as Probiotic for Freshwater Fish” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Abdul Rahman bin Omar, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Mohamed Ali bin Rajion, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Abdul Razak bin Alimon, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Einar Ringø, PhD

Professor
University of Tromsø, Norway
(External Examiner)

SEOW HENG FONG, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of **Doctor of Philosophy**. The members of the Supervisory Committee were as follows:

Hassan Mohd. Daud, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Fatimah Md. Yusoff, PhD

Professor
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Datin Paduka Aini Ideris, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Che Roos Saad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or other institutions.



SEYED KAMALEDDIN LLAMEH

Date: 20 June 2012

TABLE OF CONTENTS

	Page
DEDICATION	iii
ABSTRACT	iv
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF TABLES	xxi
LIST OF FIGURES	xxiii
LIST OF ABBREVIATIONS	xxv
CHAPTER	
1 INTRODUCTION	1
1.1 Background of study	1
1.2 Problem statement	2
1.3 Hypothesis	4
1.4 Objectives	4
2 LITERATURE REVIEW	5
2.1 Introduction	5
2.2 Snakehead fish (<i>Channa striatus</i>)	7
2.3 Development of the fish gastrointestinal tract	9
2.4 Microorganisms in fish	11
2.5 Development of probiotic definition	12
2.6 Mechanisms of action of probiotics	13
2.7 Lactic acid bacteria (LAB)	13
2.8 Isolation of LAB from aquatic animals	15
2.9 Probiotic criteria	17
2.10 Characterization of lactic acid bacteria	18
2.10.1 Importance of acid and bile salt tolerance	19
2.10.2 Growth of LAB at different NaCl and temperatures	20
2.10.3 Antagonistic effect against fish pathogen and antibiotic sensitivity	21
2.11 Colonization of bacteria in intestine	23
2.12 Effects of probiotics on fish performance	24
2.12.1 Growth	24
2.12.2 Feed conversion ratio (FCR)	26
2.12.3 Protein efficiency ratio (PER)	26
2.12.4 Digestibility	27
2.12.5 Body composition	28
2.13 Immune system response	29

2.14	Pathogenicity and challenge	31
3	ISOLATION AND IDENTIFICATION OF LACTIC ACID BACTERIA FROM THE STOMACH AND INTESTINE OF FINGERLING AND ADULT SNAKEHEADS (<i>Channa striatus</i>)	33
3.1	Introduction	33
3.2	Materials and Methods	34
3.2.1	Sampling	34
3.2.2	Isolation of LAB from fingerlings	35
3.2.3	Isolation of LAB from adult fish	36
3.2.4	Making stock from the isolated LABs	38
3.2.5	Selection of the isolates by antagonistic test	38
3.2.6	Identification of selected LABs	39
3.2.7	Statistical analysis	44
3.3	Results	44
3.3.1	Total colony count of bacteria in stomach and intestine	44
3.3.2	Isolation of lactic acid bacteria from GI tract	45
3.3.3	Selection of the isolates by antagonistic test	46
3.3.4	Identification of selected LABs	46
3.4	Discussion	50
3.4.1	Total colony count of bacteria from the stomach and intestine of fingerling and adult snakeheads	50
3.4.2	Isolation of LABs from GI tract of fingerling and adult snakeheads	51
3.4.3	Identification of the selected LAB	53
3.5	Conclusion	55
4	PROBIOTIC PROPERTIES OF LACTIC ACID BACTERIA ISOLATED FROM SNAKEHEAD (<i>Channa striatus</i>)	56
4.1	Introduction	56
4.2	Materials and Methods	58
4.2.1	pH tolerance test	58
4.2.2	Bile salt tolerance test	59
4.2.3	Growth at different NaCl concentrations	59
4.2.4	Growth at different temperatures	60
4.2.5	Antagonistic effect test against fish pathogens	60
4.2.6	Antibiotic sensitivity test	62
4.2.7	Statistical analysis	62
4.3	Results	63
4.3.1	pH tolerance test	63
4.3.2	Bile salt tolerance test	65
4.3.3	Growth at different NaCl concentrations	67
4.3.4	Growth at different temperatures	70
4.3.5	Antagonistic effect against fish pathogens	73

4.3.6	Antibiotic sensitivity test	76
4.4	Discussion	78
4.4.1	pH and bile salt tolerance	78
4.4.2	Growth at different NaCl concentrations and temperature	79
4.4.3	Antagonistic effect against fish pathogens and antibiotic sensitivity tests	80
4.5	Conclusion	82
5	EFFECT OF MONO AND MULTI-SPECIES OF SELECTED LACTIC ACID BACTERIA AS PROBIOTICS ON THE GROWTH PARAMETERS AND BODY COMPOSITION OF JAVANESE CARP (<i>Puntius gonionotus</i>)	84
5.1	Introduction	84
5.2	Materials and Methods	87
5.2.1	Experimental design	87
5.2.2	Preparation of experimental diets	88
5.2.3	Viability of LAB in diet	90
5.2.4	Survival of the LAB in the GI tract and their controlling effects on the population of Gram-negative bacteria in the experimental fishes	91
5.2.5	Body composition of fishes	92
5.2.6	Calculated parameters	92
5.2.7	Statistical analysis	93
5.3	Results	93
5.3.1	Viability of LAB in diet	93
5.3.2	Survival of the LAB in the GI tract and their controlling effects on the population of Gram-negative bacteria in the experimental fishes	96
5.3.3	Body composition of fish	99
5.3.4	Growth parameters	100
5.4	Discussion	103
5.4.1	Viability of LAB in diet	103
5.4.2	Survival of the LABs in the GI tract and their controlling effects on the population of Gram-negative bacteria in the experimental fishes	103
5.4.3	Growth parameters	104
5.5	Conclusion	107
6	DETERMINATION OF DIGESTIVE ENZYMES AND SHORT CHAIN FATTY ACID PRODUCTION FROM THE INTESTINE OF JAVANESE CARP (<i>Puntius gonionotus</i>) TREATED WITH <i>Enterococcus faecalis</i> AS PROBIOTIC	109
6.1	Introduction	109
6.2	Materials and Methods	111
6.2.1	Experimental design	111

6.2.2	Detection of digestive enzymes in the fish intestine	112
6.2.3	Short chain fatty acid determination in the fish intestine	116
6.2.4	Statistical analysis	117
6.3	Results	118
6.3.1	Protease and lipase activities in the fish intestine	118
6.3.2	Short chain fatty acid production in the fish intestine	118
6.4	Discussion	119
6.4.1	Protease and lipase enzymes determination	119
6.4.2	Short chain fatty acid assays	121
6.5	Conclusion	122
7	CHALLENGE TEST WITH FISH PATHOGEN AND EVALUATION OF <i>Enterococcus faecalis</i> ON IMMUNE SYSTEM RESPONSE OF JAVANESE CARP (<i>Puntius gonionotus</i>)	123
7.1	Introduction	123
7.2	Materials and Methods	125
7.2.1	Challenge test with fish pathogen	125
7.2.2	Evaluation of the immune system response	127
7.2.3	Statistical analysis	130
7.3	Results	130
7.3.1	Effects of challenge test with fish pathogen on experimental fishes	130
7.3.2	Humoral immune response evaluation	132
7.4	Discussion	133
7.4.1	Challenge test	133
7.4.2	Humoral immune response evaluation	135
7.5	Conclusion	136
8	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	138
8.1	Summary	138
8.2	General conclusion	139
8.2.1	Isolation, selection and identification of LAB	139
8.2.2	Probiotic properties of selected LABs	140
8.2.3	Effects of mono and multi species of selected LABs on rearing parameters	140
8.2.4	Effects of candidate probiotic on digestive enzyme and short chain fatty acid	141
8.2.5	Effect of candidate probiotic on survival in the challenge test and immune system response	142
8.3	Recommendation for future research	143
	REFERENCES	144
	APPENDIX	155



© COPYRIGHT UPM