



UNIVERSITI PUTRA MALAYSIA

**RESPONSE TO SALT STRESS AND STRATEGIES TO IMPROVE SALT
TOLERANCE IN CHINESE KALE (*Brassica oleracea* var. *alboglabra*)**

AMIN TAYEBI-MEIGOONI

FP 2012 43

**RESPONSE TO SALT STRESS AND STRATEGIES TO IMPROVE SALT
TOLERANCE IN CHINESE KALE (*Brassica oleracea* var. *alboglabra*)**



By
AMIN TAYEBI-MEIGOONI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirement for the Degree of Doctor of Philosophy**

June 2012

DEDICATION

I lovingly dedicate this thesis to

My motherland IRAN,

My beloved Parents,

My lovely Wife,

My dear daughterViana,

My beloved Brother & lovely Grandma,

Whose supported me each step of the way.



Abstract of thesis presented to senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

RESPONSE TO SALT STRESS AND STRATEGIES TO IMPROVE SALT TOLERANCE IN CHINESE KALE (*Brassica oleracea* var. *alboglabra*)

By

AMIN TAYEBI-MEIGOONI

June 2012

Chairman: Assoc. Prof. Yahya Awang, PhD

Faculty: Agriculture

Despite the large body of literature on salinity stress, our knowledge about the effects of salt stress on Chinese kale (*Brassica oleracea* var. *alboglabra*) is still little. Therefore, there is a pressing need to know in more detail how an important vegetable plant as Chinese kale responds and adapts to such conditions. With regard to the importance of Chinese kale in term of their benefits and high consumption in human diets, understanding the possible effects of salt stress on Chinese kale have never been exposed to salt stress can be significant. Therefore attempts to provide some practical, safe and environmentally sound techniques to retain productivity under salt stress is critical. The purposes of this study were to unveil the mechanism contributing to salinity tolerance during vegetative stage and to test for possible contribution of exogenous ascorbic acid and pre-treatment with low

concentration of hydrogen peroxide in association with the tolerance mechanisms during vegetative development of the crop.

Preliminary study involving four cultivars of Chinese kale ['Standard kailan', 'Hong Kong kailan', 'Kale Curly Leaf' and 'Hong Kong Stem Flower'] and four salinity levels (0, 25, 50 and 75) showed that final growth parameters, relative water content, concentration of photosynthetic pigments and maximum quantum yield of PSII (F_v/F_m) were significantly reduced by salinity. The treatment elevated plant's proline, hydrogen peroxide and lipid peroxidation. Among cultivars tested, cv. 'Standard kailan' showed minimum reduction of biomass. Cv. 'Standard kailan' also showed relatively higher adaptability to salinity due to its ability to regulate hydrogen peroxide (H_2O_2) generation and moderate oxidative damage to cell membranes as shown by malondialdehyde (MDA) content in leaf tissues at all salinity. Based on the results, cv. 'Standard kailan' was more salt-tolerant than other cultivated varieties and has been used for further experimentation.

Biochemical analysis of the leaves revealed that salinity also produced significant negative impacts on protein synthesis and leaf pigments (chlorophylls and carotenoid). The results may imply the increase in the energy dissipation *via* chlorophyll fluorescence along with reduction of chlorophylls maybe is a manifestation to preserve a balance between

harvesting and utilizing energy in order to reduce threat of oxidative stress. Consistent with results in the first experiment, detrimental effects of salinity were further illustrated in the generation and accumulation of hydrogen peroxide and MDA. However, salt stressed plants contained relatively higher concentrations of proline and activity of antioxidant enzymes especially ascorbate peroxidase (APX) and peroxidase (POX). Chinese kale was shown to be not affected by low salinity up to 25 mM and can be tolerant up to 75 mM NaCl.

Subsequent study further confirmed the results of the first experiment where the overall reduction in plant growth was coupled with reduced leaf expansion and net assimilation rate. Negative effects of salinity are clearly shown as reduction in net assimilation rate (NAR) and relative growth rate (RGR) and this was coupled with alteration of ionic balance in the plant tissue. Salinity reduced N, K⁺, Ca²⁺ and Mg²⁺ in leaves and roots. P was also decreased by the treatment in leaves, while it was increased in root. The concentrations of Na⁺ and Cl⁻ in leaves were markedly enhanced in plants tissues as the NaCl in the root zone increases. Reduction of N and Mg²⁺ availability for various processes such as chlorophyll biosynthesis, lack of P of bioenergetics pathway, besides imbalance of K⁺ and Ca²⁺ and negative effects of Na⁺ and Cl⁻ to different bioactive processes were also considered to be responsible for growth reduction in this plant.

Interactions of foliar application of ascorbic acid (AsA) and salt stress on the plants were also evaluated in this study. Application of AsA reduced portion of antioxidant enzyme activity catalase (CAT) and peroxidase (POX) may be due to its own talent for scavenging of reactive oxygen species. APX (acts by utilizing AsA) was stimulated by exogenous AsA, which assist the plants for higher resistance under salinity stress. Application exogenous H₂O₂ as a stress priming factor has resulted in a noticeable reduction of endogenous H₂O₂ and lipid peroxidation. Stimulation of POX and APX activities by exogenous H₂O₂ were greater than CAT activity, suggesting that POX and APX have had a higher affinity to scavenge ROS in these circumstances, and this could increase the plant's tolerance level to salinity.

Overall, results of the study showed that Chinese kale was sensitive to NaCl salinity stress but the plant can tolerate up to 75 mM NaCl. However, the deleterious effects of salinity can be partially moderated *via* application of exogenous ascorbic acid and H₂O₂ priming.

Abstrak tesis untuk dikemukaan kepada senat, Universiti Putra Malaysia
bagi memenuhi keperluan ijazah Doktor Falsafah

**TINDAK BALAS KEPADA STRES KEMASINAN DAN STRATEGI
UNTUK MEMPERBAIKI TAHAP KEMASINAN DALAM KAILAN
(*Brassica oleracea* var. *alboglabra*)**

Oleh

AMIN TAYEBI-MEIGOONI

Jun 2012

Pengerusi: Profesor Madya Yahya Awang, PhD

Fakulti: Pertanian

Kajian dan literasi berkaitan tegasan kemasinan adalah sangat meluas, namun pengetahuan mengenai kesan tegasan kemasinan terhadap kailan (*Brassica oleracea* var. *alboglabra*) sukar diperolehi. Oleh itu, pengetahuan tentang keupayaan dan cara bagaimana cara tanaman sayuran seperti kailan beradaptasi terhadap tekanan kemasinan sangat diperlukan. Berdasarkan kepada kepentingan kalian khususnya dari segi diet pemakanan dan permintaan, adalah sangat penting dan signifikan bagi menjalankan sebuah kajian untuk memahami kesan tegasan kemasinan terhadap tanaman ini. Selain daripada mengekalkan produktiviti di bawah tegasan kemasinan, salah satu perkara yang perlu dititikberatkan ialah untuk menyediakan suatu teknik pengeluaran yang praktikal, selamat dan mesra alam. Tujuan kajian ini dijalankan ialah untuk mendedahkan mekanisma yang terlibat

dalam toleransi terhadap tegasan kemasinan serta mengkaji peranan asid askorbik eksogenous dan pra-rawatan hidrogen peroksida pada kepekatan yang rendah terhadap mekanisma toleransi kailan pada peringkat perkembangan vegetatif.

Kajian permulaan yang melibatkan empat kultivar kailan menunjukkan bahawa parameter tumbesaran, kandungan relatif air, kepekatan pigmen fotosintesis dan hasil kuantum maksimum PSII (Fv/Fm) menurun secara signifikan oleh kemasinan. Rawatan yang diberikan telah meningkatkan kandungan proline, hidrogen peroksida dan lipid peroksida (sebagai malondialdehyd, MDA) dalam kailan. Kultivar ‘Standard kailan’ menunjukkan pengurangan biomass yang minimum berbanding kultivar-kultivar yang lain. Secara perbandingannya, ‘Standard kailan’ juga menunjukkan adaptasi kemasinan yang tertinggi pada semua tahap kemasinan berdasarkan kemampuannya mengawal pengeluaran hidrogen peroksida dan kesan pengoksidaan yang sederhana pada membran sel seperti yang ditunjukkan oleh kandungan MDA di dalam tisu daun.

Analisis biokimia daun menunjukkan kemasinan memberikan kesan negatif yang signifikan ke atas sintesis protein dan pigmen daun (klorofil dan karotenoid). Keputusan ini mungkin menunjukkan peningkatan dalam pelesapan tenaga menerusi klorofil floresens dengan pengurangan klorofil sebagai suatu manifestasi untuk mengekalkan keseimbangan di antara

penuaan dan penggunaan tenaga demi mengurangkan tegasan oksidatif. Konsisten dengan keputusan eksperimen yang pertama, kesan kemasinan dihasilkan berdasarkan kepada penghasilan dan pengumpulan hidrogen peroksida dan MDA. Walaubagaimanapun, kailan yang mengalami tegasan kemasinan mengandungi prolin dan aktiviti enzim antioksida khususnya aksorbat peroksida dan peroksida pada kepekatan yang tinggi. Kailan menunjukkan ketahanan terhadap kemasinan sehingga kepada 75 mM.

Kajian yang berikutnya mengesahkan keputusan eksperimen yang pertama iaitu penurunan pertumbuhan tumbuhan ditambah pula dengan penurunan pengembangan daun dan kadar fotosintesis bersih. Kesan negatif kemasinan ditunjukkan secara jelas oleh penurunan kadar asimilasi bersih (NAR) dan kadar pertumbuhan relatif (RGR) dan ini ditambah lagi dengan gangguan pada keseimbangan ion pada tisu tumbuhan. Kemasinan menurunkan N, K⁺, Ca²⁺ dan Mg²⁺ di dalam daun dan akar. Kepekatan P di dalam daun menurun dengan rawatan NaCl. Peningkatan kepekatan Na⁺ dan Cl⁻ di dalam tisu daun jelas dengan peningkatan NaCl di kawasan akar. Penurunan N dan Mg²⁺ yang berfungsi dalam pelbagai proses seperti biosintesis klorofil, kekurangan P untuk tindakan bioenergetik, disamping ketidakseimbangan K⁺ dan Ca²⁺, dan kesan negatif Na⁺ dan Cl⁻ terhadap pelbagai proses bioaktif adalah merupakan antara faktor-faktor yang bertanggungjawab terhadap penurunan pada pertumbuhan kailan.

Kesan interaksi di antara rawatan asid askorbik (AsA) dan kemasinan terhadap tumbuhan juga diperhatikan di dalam kajian ini. Aplikasi AsA telah menyebabkan penurunan aktiviti enzim antioksidan (CAT dan POX) dan ini mungkin disebabkan oleh sifat semulajadinya yang mengikat oksigen yang reaktif. APX (menggunakan AsA untuk berfungsi) telah dirangsang oleh AsA untuk membantu tumbuhan untuk lebih tahan kepada tegasan kemasinan. Aplikasi H₂O₂ yang merupakan faktor utama penyebab tegasan telah menyebabkan penurunan H₂O₂ dalam tisu pokok dan lipid peroksid. Rangsangan terhadap aktiviti POX dan APX oleh H₂O₂ adalah sangat besar berbanding aktiviti CAT, di dalam situasi begini POX dan APX mempunyai afiniti yang tinggi untuk mengikat ROS dan seterusnya meningkatkan ketahanan tumbuhan terhadap kemasinan.

Secara keseluruhannya, keputusan kajian menunjukkan bahawa kailan sangat sensitif kepada tekanan kemasinan NaCl tetapi kailan mempunyai ketahanan sehingga 75 mM NaCl. Walaubagaimanapun, kesan kerosakan oleh tegasan kemasinan boleh dikurangkan melalui aplikasi AsA dan H₂O₂.

ACKNOWLEDGEMENTS

Thanks to my almighty God for helping me achieving this work. Without his guidance, this work would never have been accomplished.

I would like to express my deepest gratitude and sincere thanks to Assoc. Prof. Dr. Yahya Awang who provided me with this opportunity for successful completion of my Ph.D. program under his scientific supervision, continuous support and invaluable help in Malaysia.

I would be honoured to convey my heartfelt thanks and true gratitude to my co-supervisors Prof. Dr. Maziah Mahmoud, Dr. Ahmad Selamat and Prof. Dr. Zakaria Wahab for their kind encouragement, permanent advice and the moral support they always gave.

I am particularly grateful to Prof. Dr. Rosli Mohamad for his valuable suggestions and kind encouragement. I am equally thankful to all the staff of UPM, especially those at the Faculty of Agriculture who contributed to my learning process, especially Mr. Abdul Rahman for his help during using lab and field equipments.

I am also thankful to many friends that my family and I met and who contributed to make a pleasant and memorable part of our lives, although different language and culture. Last but not least, I wish to thank my wife Khatereh and my family for their nonstop support, willing help, permanent patience and continuous love.

APPROVAL

I certify that a Thesis Examination Committee has met on June 11, 2012 to conduct the final examination of Amin Tayebi-Meigooni on his thesis entitled "**Response to salt stress and strategies to improve salt tolerance in Chinese kale (*Brassica oleracea* var. *alboglabra*)**" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee are as follows:

Uma Rani Sinniah, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairperson)

Mohd Ridzwan A. Halim, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

Siti Aishah Hassan, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

Muhammad Ashraf, PhD

Professor

Faculty of Science

University of Agriculture Faisalabad

Pakistan

(External Examiner)

SEOW HENG FONG, PhD
Professor and Deputy Dean School
of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the Degree of **Doctor of philosophy**. The members of the Supervisory Committee were as follows:

Yahya B. Awang, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Maziah Bt. Mahmoud, PhD

Professor

Faculty of Biotechnology and Biomolecular science

Universiti Putra Malaysia

(Member)

Zakaria B. Wahab, PhD

Professor

Faculty of Agriculture

Universiti Malaysia Perlis

(Member)

Ahmad B. Selamat, PhD

Lecturer

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.

AMIN TAYEBI-MEIGOONI

Date: 11 June 2012



TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xix
LIST OF FIGURES	xx
LIST OF APPENDICES	xxiv
LIST OF ABBREVIATIONS	xxv
 CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Objectives	5
2 REVIEW OF LITERATURE	6
2.1 Effects of salinity on plant growth	6
2.2 Physiological effects of salt stress	9
2.2.1 Effects of salinity on water relations	9
2.2.2 Effects of salinity on photosynthesis	11
2.2.3 Effects of salinity on photosynthesis pigments	14
2.2.4 Effects of salinity on chlorophyll fluorescence	15
2.2.5 Effects of salinity on ion levels and nutrient contents	17
2.3 Biochemical effects of salt stress	19
2.3.1 Effects of salinity on protein	19
2.3.2 Effects of salinity on amino acids	20
2.3.3 Salinity effects on carbohydrates	22
2.4 Salt tolerance in plants	23
2.4.1 Mechanism of salt tolerance in plant	24
2.4.2 Regulation and compartmentalization of ions	25
2.4.3 Induced biosynthesis of compatible solutes	27
2.4.4 Ion uptake and transport control by roots	28
2.4.5 Changes in photosynthetic pathway	29
2.4.6 Induction of plant hormones	30
2.5 Induction of oxidative stress and role of reactive oxygen species scavengers	30
2.6 Chinese kale, origin and nutrition facts	39
2.7 Expectation	39
3 DIFFERENTIAL RESPONSES OF FOUR CHINESE KALE CULTIVARS TO SALINITY	42
3.1 Introduction	42
3.2 Materials and methods	44
3.2.1 Plant material and growth condition	44

3.2.2 NaCl treatment	44
3.2.3 Growth measurements	45
3.2.4 Relative water content	45
3.2.5 Chlorophyll fluorescence	46
3.2.6 Photosynthetic pigments	46
3.2.7 Proline assay	47
3.2.8 Hydrogen peroxide concentration	47
3.2.9 Lipid peroxidation	48
3.2.10 Experimental design and statistical analysis	48
3.3 Results	49
3.3.1 Plant Growth	49
3.3.2 Relative water content (RWC)	52
3.3.3 Maximum quantum yield	53
3.3.4 Photosynthetic pigments	54
3.3.5 Proline, hydrogen peroxide and lipid peroxidation (MDA)	56
3.4 Discussion	60
3.5 Conclusion	64
4 PHYSIOLOGICAL AND BIOCHEMICAL RESPONSES OF CHINESE KALE TO SALINITY	65
4.1 Introduction	65
4.2 Materials and methods	68
4.2.1 Planting materials and growth conditions	68
4.2.2 Salt stress treatment	68
4.2.3 Growth parameters	69
4.2.4 Relative water content	69
4.2.5 Chlorophyll fluorescence	69
4.2.6 Photosynthetic pigments	70
4.2.7 Proline assay	70
4.2.8 Hydrogen peroxide concentration	70
4.2.9 Lipid peroxidation (MDA)	70
4.2.10 Protein and enzymes assay	70
4.2.11 Experimental design statistical analysis	72
4.3 Results	73
4.3.1 Plant growth	73
4.3.2 Relative water content	77
4.3.3 Chlorophyll fluorescence	77
4.3.4 Photosynthetic pigments	78
4.3.5 Proline	80
4.3.6 Hydrogen peroxide	80
4.3.7 Lipid peroxidation	80
4.3.8 Protein	82
4.3.9 Antioxidant enzymes	82
4.4 Discussion	84
4.5 Conclusion	91

5	GROWTH INDICES AND ION COMPOSITION OF CHINESE KALE IN RESPONSE TO INCREASING SALINITY	92
5.1	Introduction	92
5.2	Materials and methods	94
5.2.1	Plant material and growth condition	94
5.2.2	Salt stress treatment	95
5.2.3	Determination of growth parameters	95
5.2.4	Growth analysis	96
5.2.5	Nutrients composition	96
5.2.6	Experimental design and statistical analysis	97
5.3	Results	98
5.3.1	Growth parameters	98
5.3.2	Growth indices	101
5.3.3	Ion composition	105
5.4	Discussion	110
5.5	Conclusion	115
6	ALLEVIATION OF SALINITY DAMAGE OF CHINESE KALE USING EXOGENOUS ASCORBIC ACID	117
6.1	Introduction	117
6.2	Materials and methods	121
6.2.1	Plant material and growth condition	121
6.2.4	Salt stress and ascorbic acid treatments	122
6.2.5	Sampling	123
6.2.6	Extraction for protein and enzymes assay	123
6.2.7	Protein assay	123
6.2.8	Assay of antioxidant enzymes	123
6.2.9	Relative water content	124
6.2.10	Photosynthetic pigments	124
6.2.11	Proline assay	124
6.2.12	Lipid peroxidation (MDA)	125
6.2.13	Experimental design and data analysis	125
6.3	Results	125
6.3.1	Antioxidant enzymes	126
6.3.2	Protein content	128
6.3.3	Relative water content (RWC)	129
6.3.4	Lipid peroxidation (MDA)	130
6.3.5	Proline accumulation	130
6.3.6	Chlorophyll content	131
6.4	Discussion	133
6.5	Conclusion	136
7	ALLEVIATION OF SALINITY DAMAGE OF CHINESE KALE USING HYDROGEN PEROXIDE PRIMING	138
7.1	Introduction	138
7.2	Materials and methods	141

7.2.1	Plant material and growth condition	141
7.2.2	Salt stress and exogenous hydrogen peroxide treatments	141
7.2.3	Sampling	142
7.2.4	Lipid peroxidation (MDA)	142
7.2.5	Hydrogen peroxide concentration	143
7.2.6	Assay of antioxidant enzymes	143
7.2.7	Experimental design and data analysis	144
7.3	results	144
7.3.1	Lipid peroxidation (MDA)	144
7.3.2	Hydrogen peroxide	145
7.3.3	Antioxidant enzymes	146
7.4	Discussion	148
7.5	Conclusion	151
8	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	153
	Recommendations for future research	157
REFERENCES		158
APPENDICES		187
APPENDIX A		187
A-1	Germination and seedling growth	187
A-2	Nutrient solution	187
A-3	Hydroponic system	189
A-3.1	Static aerated technique (Applied in Exp. Chapters 3, 4 and 5)	189
A-3.2	Growth and dynamic hydroponic system (Applied in Exp. Chapter 6 and 7)	191
APPENDIX B		195
BIODATA OF STUDENT		201