

Growth, osmoregulatory responses and changes to the lipid and fatty acid composition of organs from the mud crab, *Scylla serrata*, over a broad salinity range

ABSTRACT

Aquatic animals can often undergo substantial physiological responses to salinity; however, associated lipid/fatty acid alterations to their various tissues have received little attention. To investigate this, we measured the growth of mud crab, *Scylla serrata*, juveniles over two moults (duration of 23–60 days) at salinities of 4, 12, 20, 28, 36 and 44‰ (30 replicates/treatment). After the second moult, 6-day post moult crabs were sampled for hepatosomatic index (HSI), haemolymph osmolality, Na⁺, K⁺ and Ca²⁺ levels, anterior/posterior gill Na⁺/K⁺-ATPase activity as well as the lipid/fatty acid composition of the anterior and posterior gills, hepatopancreas and muscle. High salinities of 36 and 44‰ significantly lowered crab growth and HSI ($p < 0.01$). *S. serrata* strongly hyper-regulated haemolymph osmolality and ions likely due to significantly enhanced posterior gill Na⁺/K⁺-ATPase activity. At decreasing salinities, eicosapentaenoic acid, arachidonic acid, n-3 and long chain polyunsaturated fatty acid (PUFA) significantly increased ($p < 0.05$), likely to maintain Na⁺/K⁺-ATPase activity via increased membrane fluidity. Muscle and hepatopancreas n-3/n-6 PUFA ratios significantly decreased ($p < 0.05$) at increasing salinities indicating possible fatty acid metabolic modifications. Results indicate that *S. serrata* juveniles are well adapted to low salinities, with higher salinities likely reducing their metabolism, reflected by lower growth, HSI and higher posterior gill and hepatopancreatic triacylglycerol.

Keyword: *Scylla serrata*; Osmoregulation; Salinity; Fatty acid; Na⁺/K⁺-ATPase activity