



UNIVERSITI PUTRA MALAYSIA

***ESTABLISHING QUALITY CONTROL OF OIL PALM SUSPENSION
CULTURE VIA FLOW CYTOMETRY, IMAGE CYTOMETRY AND
FLUORESCENCE *in situ* HYBRIDIZATION***

MOHD RAZIK BIN MIDIN

FS 2013 3

**ESTABLISHING QUALITY CONTROL OF OIL PALM SUSPENSION
CULTURE VIA FLOW CYTOMETRY, IMAGE CYTOMETRY AND
FLUORESCENCE *in situ* HYBRIDIZATION**

By

MOHD RAZIK BIN MIDIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

January 2013

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfilment of the requirement for the degree of Master of Science

**ESTABLISHING QUALITY CONTROL OF OIL PALM SUSPENSION
CULTURE VIA FLOW CYTOMETRY, IMAGE CYTOMETRY AND
FLUORESCENCE *in situ* HYBRIDIZATION**

By

MOHD RAZIK BIN MIDIN

January 2013

Chair: Rosimah Nulit, PhD

Faculty: Science

Oil palm industry targets to provide high quantities of clonal palms with agronomic trait *via* clonal propagation of elite palms. Clonal propagation of oil palm through tissue culture process is achieved *via* callus or suspension cultures. The occurrence of *in vitro* culture stress may generate genome variability such as genome size and ploidy level alteration. Hence, several cytological tools were utilized in this study to observe any changes that might occur in adult clonal palms, their respective suspension cultures and regenerant plantlets. The tools used were flow cytometry (FCM), 18S-25S ribosomal DNA-fluorescence *in situ* hybridization (rDNA-FISH), and image cytometry (ICM). The findings of FCM on the nuclear genome sizes of the four adult clonal palms (using leaf samples from Frond-1) varied from 2.59 ± 0.19 pg to 2.91 ± 0.14 pg while for 8-months-old regenerant plantlets (five replicates for each samples) varied from 2.14 ± 0.21 pg to 3.05 ± 0.11 pg. Adult clonal samples and their respective regenerants showed the same ploidy level which is diploid, indicated

by FCM analysis. Oil palm suspension culture materials were limited, hence produced less than 1,000 nuclei and were unable to be analysed by FCM. This led to the utilization of rDNA-FISH and development of ICM technique which used the application available in the PAX-it image analysis software (Midwest Information Systems, USA). Based on the rDNA-FISH analysis on the suspension culture and regenerant plantlet materials, two hybridization signals were observed on the interphase nuclei indicating diploid ploidy level. ICM analysis of the suspension cultures revealed that the nuclear genome size ranged from 4.71 pg to 5.49 pg while the integrated optical density (IOD) values for all suspension cultures ranged from 0.06-0.18 arbitrary unit (a.u.). The pattern showed that the cell cycle of suspension cultures was lagging during the G1-S-G2 phase, hence explaining the slow proliferation rate. In summary, it was observed that diploid ploidy state was maintained throughout the adult clonal palms, their suspension cultures and regenerant plantlets while the genome size of suspension culture materials were higher than their adult clonal palms and regenerants. This study concluded that the quality control of oil palm suspension cultures could be established by the above-mentioned tools based on the estimated nuclear genome size and ploidy level. ICM tool developed in this study can also be applied to analyze the friable suspension calli cultures which are the starting material for suspension cultures. This would ensure the oil palm's clonal fidelity and improve the efficiency and robustness of suspension cultures.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

KAWALAN KUALITI KULTUR AMPAIAN SAWIT MELALUI KAEDAH SITOMETRI ALIRAN, SITOMETRI IMEJ DAN PENGHIBRIDAN *in situ* BERPENDARFLUOR

Oleh

MOHD RAZIK BIN MIDIN

Januari 2013

Pengerusi: Rosimah Nulit, PhD

Fakulti: Sains

Sasaran industri sawit adalah untuk membekalkan kuantiti sawit berciri agronomi pada kadar kuantiti yang tinggi melalui propagasi sawit elit. Propagasi sawit melalui teknik kultur tisu dicapai melalui kalus atau kultur ampaiian sawit. Tekanan *in vitro* yang berlaku berkemungkinan menghasilkan variasi pada genom, seperti saiz genom dan aras ploidi. Justeru itu, beberapa kaedah sitologi telah digunakan dalam kajian ini bagi memerhati sebarang perubahan yang mungkin berlaku pada klon dewasa sawit, kultur ampaiian dan juga planlet regeneran. Kaedah-kaedah tersebut adalah sitometri aliran (FCM), 18S-25S ribosomal DNA-penghibridan *in situ* berpendarfluor (rDNA-FISH) dan sitometri imej (ICM). Hasil analisis FCM terhadap saiz genom klon sawit dewasa (menggunakan pelepah -1) adalah di antara 2.59 ± 0.19 pg hingga 2.91 ± 0.14 pg manakala saiz genom bagi planlet regenerannya yang berusia 8 bulan (5 replikasi bagi setiap sampel) berjulat diantara 2.14 ± 0.21 pg sehingga 3.05 ± 0.11 pg. Klon sawit dewasa dan planlet regenerannya menunjukkan aras ploidi yang sama

iaitu diploid, ditentukan melalui analisis FCM. Material kultur ampaiian sawit adalah terhad, justeru itu menghasilkan populasi nukleus kurang dari 1,000 menyebabkan ia tidak boleh dianalisis melalui FCM. Ini telah membawa kepada penggunaan rDNA-FISH dan pembangunan teknik ICM yang menggunakan aplikasi dalam perisian analisis imej PAX-it (Midwest Information Systems, USA). Berdasarkan kaedah rDNA-FISH pada kultur ampaiian sawit dan planlet regeneran, dua isyarat hibridisasi di atas nukleus interfasa dapat diperhati menandakan aras ploidi diploid. Analisis ICM terhadap kultur ampaiian sawit menunjukkan saiz genomnya berjulat di antara 4.71 sehingga 5.49 pg manakala puncak ketumpatan optik bersepadu (IOD) berjulat di antara 0.06-0.18 arbitrari unit (a.u.). Corak ini menunjukkan perkembangan kitar sel kultur ampaiian adalah perlahan pada fasa G1-S-G2, seterusnya menerangkan kadar proliferasi yang perlahan. Keseluruhannya, dapat diperhati bahawa aras diploid dikekalkan sepanjang klon sawit dewasa, kultur ampaiian dan planlet regeneran manakala saiz genom kultur ampaiian adalah lebih tinggi berbanding klon sawit dewasa dan planlet regenerannya. Kesimpulannya, kualiti kultur ampaiian boleh ditentukan melalui kaedah sitologi yang digunakan dengan merujuk pada saiz genom dan aras ploidi. Kaedah ICM yang dibangunkan dalam kajian ini juga boleh diaplikasi untuk menganalisis kalus 'friable' yang digunakan sebagai bahan pemula untuk kultur ampaiian. Ini akan memastikan ketulenan klon sawit dan meningkatkan keberkesanan kultur ampaiian.

ACKNOWLEDGEMENTS

Praised be to ALLAH S.W.T.

I gratefully acknowledge the Director General of MPOB for offering me a graduate student assistantship during my study. I am deeply grateful to my supervisors, Dr. Rosimah Nulit (Universiti Putra Malaysia) and Dr. Maria Madon (Malaysian Palm Oil Board); supervisory committee members, Associate Professor Dr. Faridah Qamaruz Zaman, for their support, encouragement, patience, comments and suggestions to assist me to make this work more pertinent.

Thanks to my cytogenetic lab members and friends; Madam Norhariza, Madam Zainab, Mr Razali, Madam Nordiana, Masfira, Aizat, Wan, Azuan, Wan Zulqhairi and Nazri for their support throughout this study.

Last but not least, I would like to express my appreciation and indebtedness to my parents (Hj. Midin bin Yaakop and Hjh. Rabiah binti Mokhtar) and family (Mohamed Rawi, Mohamad Razizi, Safiah and Zuriati) who have been always supported me and guided me to achieve my ambitions through my life.

I certify that a Thesis Examination Committee has met on 8th January 2013 to conduct the final examination of Mohd Razik bin Midin on his thesis entitled “Establishing quality control of oil palm suspension culture *via* flow cytometry, image cytometry and fluorescence *in situ* hybridization” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science, Plant and Animal Genetics.

Members of the Thesis Examination Committee were as follows:

Nor Azwady bin Abd Aziz, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Umi Kalsom binti Yusof, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Janna Ong binti Abdullah, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Mahani Mansor Clyde, PhD

Professor
School of Environmental and Natural Resource Sciences
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
Malaysia
(External examiner)



SEOW HENG FONG, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 March 2013

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Rosimah binti Nulit, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Faridah binti Qamaruz Zaman, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Maria binti Madon, PhD

Principle Research Officer
Advanced Biotechnology and Breeding Centre
Malaysian Palm Oil Board
(Member)


BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



MOHD RAZIK BIN MIDIN

Date: 8th January 2013

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xv
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem statement and objectives	2
2 LITERATURE REVIEW	4
2.1 Oil Palm	4
2.2 Oil Palm Fruit Types	4
2.3 Tissue Culture	6
2.4 Oil Palm Planting Materials	9
2.4.1 Production of Oil Palm Clones	10
2.5 Somaclonal Variation in Plant	11
2.6 Causes of Somaclonal Variation	12
2.7 Advantages and Disadvantages of Somaclonal Variation	15
2.8 Tool for Detection of Somaclonal Variation	16
2.9 Flow cytometry (FCM)	19
2.9.1 Isolation Buffer	22
2.9.2 Fluorochrome Selection for DNA staining	24
2.9.3 Nuclear Genome Size Measurement	25
2.10 Ribosomal DNA-fluorescent <i>in situ</i> hybridization (rDNA-FISH)	27
2.10.1 Probes for FISH	30
2.10.2 Ribosomal DNA probe	31
2.11 Image Cytometry (ICM)	32
2.11.1 Integrated Optical Density (IOD)	34
3 MATERIALS AND METHODS	36
3.1 Plant materials	36
3.2 FCM Tool	37
3.2.1 Nuclei Preparation	37
3.2.2 Analysis of Histogram Peak	38
3.2.3 Estimation of Nuclear Genome size	38
3.3 rDNA-FISH Tool	41
3.3.1 Preparation and Fixation of Plant Material	41
3.3.2 Chromosomal Preparation on Glass Slide	41

	3.3.3	Preparation of rDNA Probes	42
	3.3.4	Agarose Gel Electrophoresis and Ethidium Bromide Staining	44
	3.3.5	DNA Fragment Extraction from Agarose Gel	45
	3.3.6	DNA Probe Labelling	45
	3.3.7	FISH Protocol	46
3.4		ICM Tool	51
	3.4.1	Preparation and Fixation of Plant Materials	51
	3.4.2	Preparation of Feulgen Reagent	51
	3.4.3	Feulgen Staining of Nuclear DNA	51
	3.4.4	Image Analysis	52
	3.4.5	Nuclear Genome Size Estimation of Suspension Culture	54
4		RESULTS	55
	4.1	FCM Analysis	55
	4.1.1	Optimizing Nuclei Preparation	55
	4.1.2	Histogram Peak Analysis	55
	4.1.3	Ploidy Level Analysis	58
	4.1.4	Nuclear Genome Size Analysis in Adult Clonal Palm Material	60
	4.1.5	Nuclear Genome Size Analysis in Regenerant Plantlet Material	60
	4.1.6	Variation of Nuclear Genome Size Value in between Adult Clonal Palm with their Respective Regenerant	61
	4.2	rDNA-FISH Analysis	63
	4.2.1	Probe Isolation	63
	4.2.2	Hybridization Signal on Chromosome	63
	4.2.3	Ploidy State Analysis on Suspension Calli and Regenerant Material	64
	4.3	ICM Analysis	67
	4.3.1	Optimizing Nuclei Preparation	67
	4.3.2	Image Analysis	70
	4.3.3	IOD Peak Analysis	72
	4.3.4	Observation on Suspension Culture Growth Rate	75
	4.3.5	Nuclear Genome Size Estimation of Suspension Calli	76
	4.4	Genome Size Trend through each set of P164 Clone	78
	4.5	Consistency of Ploidy Level in each set of P164 Clones	80
5		DISCUSSION	82
	5.1	FCM Tool	82
	5.1.1	The Uses of <i>Glycine max</i> cv polanka as DNA Reference Standard	82
	5.1.2	The selection of Propidium Iodide as DNA fluorochrome	83

5.1.3	Optimizing Nuclei Preparation	84
5.1.4	G0/G1 Histogram Peak	85
5.1.5	Precision in Nuclear Genome Size Estimation	86
5.1.6	The Effect of Stoichiometric Error on Determination of Genome Size	87
5.1.7	Genome Size among Adult Clonal palms and its Regenerants	88
5.1.8	Consistency of Ploidy Level <i>via</i> FCM analysis	88
5.2	rDNA-FISH Tool	89
5.2.1	Metaphase Spread Preparation	89
5.2.2	DNA Probe Isolation and Labelling	91
5.2.3	Signal Detection <i>via</i> Cy3-avidin	92
5.2.4	Ploidy Level Determination of Oil Palm Suspension Culture and Regenerants <i>via</i> FISH analysis	93
5.3	ICM Tool	93
5.3.1	Optimizing Staining and Slide Preparation	93
5.3.2	Image Processing	95
5.3.3	IOD Peak Analysis	96
5.3.4	Nuclear Genome Size of Suspension Calli	98
5.4	Nuclear Genome Size Trend Analysis and its Variation Across Tissue Culture	99
5.5	Ploidy Level Consistency in Oil Palm Tissue Culture	100
6	CONCLUSION	103
	REFERENCES	106
	APPENDICES	122
	BIODATA OF STUDENT	139
	LIST OF PUBLICATIONS	140