



UNIVERSITI PUTRA MALAYSIA

***ANTI-ACUTE INFLAMMATORY EFFECT OF HIBISCUS ROSA-SINENSIS L AND
HIBISCUS ROSA-SINENSIS VAR ALBA FLOWER AND LEAF ETHANOL
EXTRACTS AND ITS MECHANISM OF ACTION IN RATS***

SITI ZALEHA RADUAN

FPSK(m) 2013 28



**ANTI-ACUTE INFLAMMATORY EFFECT OF
Hibiscus rosa-sinensis L. AND *Hibiscus rosa-sinensis*
var. *alba* FLOWER AND LEAF ETHANOL
EXTRACTS AND ITS MECHANISM OF ACTION
IN RATS**

SITI ZALEHA BINTI RADUAN

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2013



**ANTI-ACUTE INFLAMMATORY EFFECT OF *Hibiscus rosa-sinensis* L. AND
Hibiscus rosa-sinensis var. *alba* FLOWER AND LEAF ETHANOL
EXTRACTS AND ITS MECHANISM OF ACTION IN RATS**



Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science

June 2013

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the Degree of Master of Science

**ANTI-ACUTE INFLAMMATORY EFFECT OF *Hibiscus rosa-sinensis* L. AND
Hibiscus rosa-sinensis var. *alba* FLOWER AND LEAF ETHANOL
EXTRACTS AND ITS MECHANISM OF ACTION IN RATS**

By

SITI ZALEHA BINTI RADUAN

June 2013

Chair: Prof Nazrul Hakim Abdullah, PhD

Faculty: Medicine and Health Sciences

Hibiscus rosa-sinensis L. is a plant having medicinal properties especially its flower and leaf but not much on its variant; *alba*. The study was carried out to determine and compare the anti-acute inflammatory activities and its mechanism of action of ethanol extract of flower and leaf of *Hibiscus rosa-sinensis* var *alba* (white) and *Hibiscus rosa-sinensis* L. (red) in rats.

Phytochemical screening was performed with 95% ethanolic crude extracts. Flavanoids, saponins and steroids presence in all extracts. Acute dose response was determined using Fixed Dose Procedure with fixed level of dose. Any signs of toxicity were observed within 14 days. Supplementation of 500 mg/kg of all extracts caused toxicity. The blood samples were collected and liver and kidney were isolated at day 15. No significant changes ($p>0.05$) in liver enzyme levels and histologically no presence of lesions at the organs as response up to 500 mg/kg.

For anti-inflammatory properties, edema and polymorphonuclear leukocyte (PNL) infiltration induced by carrageenan and licking time induced by formalin were studied. 0.1ml of carrageenan was injected subplantarly 30 min before administration of each extracts (5, 50 or 100 mg/kg). Supplementation of 50 and 100 mg/kg of flower and leaf of *Hibiscus rosa-sinensis* L. caused significant inhibition ($p<0.05$) of edema. Flower and leaf of *Hibiscus rosa-sinensis* var *alba* significantly inhibited ($p<0.05$) edema in all range of testing dose. It varies significantly ($p<0.05$) with variant, plant parts and doses. The animals were killed after 6 hrs and PNLs in paw tissues were counted. Supplementation of all extracts at various concentration caused significant reduction ($p<0.05$) on PNL infiltration. It varies significantly ($p<0.05$) with dose but no significant ($p>0.05$) with variant and plant parts. 50 μ l of formalin was injected subplantarly 30 min before administration of 100 mg/kg of each extracts. Supplementation of all extracts showed significant reduction ($p<0.05$) on the duration of licking response. It varies significantly ($p<0.05$) with variant but no significant ($p>0.05$) with plant parts and phase.

For anti-acute inflammatory mechanism, edema and PNL infiltration induced by bradykinin (BK) and histamine were studied. 0.1ml of BK was injected subplantarly 15 min before administration of 100 mg/kg of each extracts. Supplementation of all extracts showed significant inhibition ($p<0.05$) of edema. It varies significantly ($p<0.05$) with variant but no significant ($p>0.05$) with plant parts. The animals were killed after 6 hrs and PNLs in paw tissues were counted. Supplementation of all extracts showed significant reduction ($p<0.05$) on PNL infiltration. It varies significantly ($p<0.05$) with variant and plant parts. 0.1ml of histamine was injected subplantarly immediately before administration of 100 mg/kg of each extracts.

Supplementation of all extracts showed significant inhibition ($p<0.05$) of edema. It varies significantly ($p<0.05$) with variant but no significant ($p>0.05$) with plant parts. The animals were killed after 2 hrs and PNLs in paw tissue were counted. Supplementation of all extracts showed significant reduction ($p<0.05$) on PNL infiltration. It varies significantly ($p<0.05$) with variant but no significant ($p<0.05$) with plant parts.

In conclusion, the study showed flower and leaf of *Hibiscus rosa-sinensis* var *alba* and *Hibiscus rosa sinensis* L. produced anti-acute inflammatory activity. It may involve the inhibition of cyclooxygenase, bradykinin, histamine, and reduce polymorphonuclear infiltration.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

KESAN ANTI-INFLAMASI AKUT EKSTRAK ETANOL BUNGA DAN DAUN *Hibiscus rosa-sinensis* L. DAN *Hibiscus rosa-sinensis* var. *alba* DAN MEKANISME TINDAKANNYA PADA TIKUS.

Oleh

SITI ZALEHA BINTI RADUAN

Jun 2013

Pengerusi: Prof Nazrul Hakim Abdullah, PhD

Fakulti: Perubatan dan Sains Kesihatan

Hibiscus rosa-sinensis L. adalah tumbuhan yang mempunyai ciri-ciri perubatan terutama dalam bunga dan daun tetapi tidak kepada varian; *alba*. Kajian ini dijalankan untuk menentu dan membandingkan aktiviti anti-inflamasi akut dan mekanisme tindakan oleh ekstrak etanol bunga dan daun *Hibiscus rosa-sinensis* var *alba* (putih) dan *Hibiscus rosa-sinensis* L. (merah) pada tikus.

Pemeriksaan fitokimia telah dijalankan dengan menggunakan 95% ekstrak mentah etanol. Terdapat kehadiran flavanoids, saponins dan steroids dalam kesemua ekstrak. Tindak balas akut terhadap dos telah ditentukan dengan menggunakan ‘Fixed Dose Procedure’ dengan tahap dos yang tetap. Tanda toksik diperhatikan dalam tempoh 14 hari. Rawatan pada 500 mg/kg oleh kesemua ekstrak menyebabkan toksik. Sampel darah telah dikumpul manakala hati serta buah pinggang telah diasingkan pada hari ke-15. Tiada perubahan yang signifikan ($p>0.05$) pada takat enzim hati dan tiada tanda kerosakan secara histologi pada organ sebagai tindak balas sehingga 500 mg/kg.

Untuk ciri anti-inflamasi, edema dan penyusupan polimorphonukleus leukosit (PNL) disebabkan oleh carrageenan dan masa penjilatan disebabkan oleh formalin telah dikaji. 0.1ml carrageenan telah disuntik pada subplantar 30 minit sebelum penggunaan setiap ekstrak (5, 50 atau 100 mg/kg). Rawatan oleh 50 dan 100 mg/kg bunga dan daun *Hibiscus rosa-sinensis* L. menyebabkan perencatan edema yang signifikan ($p<0.05$). Bunga dan daun *Hibiscus rosa-sinensis* var *alba* merentangkan edema secara signifikan ($p<0.05$) dalam pelbagai ujian dos. Ia berbeza secara signifikan ($p<0.05$) dengan varian, bahagian tumbuhan dan dos. Haiwan dibunuh selepas 6 jam dan PNL dalam tisu tapak kaki telah dikira. Rawatan oleh kesemua ekstrak pada pelbagai kepekatan menyebabkan pengurangan penyusupan PNL yang signifikan ($p<0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan dos tetapi tidak signifikan ($p>0.05$) dengan varian dan bahagian tumbuhan. 50 μ l formalin telah disuntik pada subplantar 30 minit sebelum penggunaan 100 mg/kg setiap ekstrak. Rawatan oleh kesemua ekstrak menunjukkan pengurangan tempoh penjilatan yang signifikan ($p<0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan varian tetapi tidak signifikan ($p>0.05$) dengan bahagian tumbuhan dan fasa.

Mekanisme anti-inflamasi akut, edema dan penyusupan PNL disebabkan oleh bradikinin (BK) dan histamin telah dikaji. 0.1ml BK telah disuntik pada subplantar 15 minit sebelum penggunaan 100 mg/kg setiap ekstrak. Rawatan oleh kesemua ekstrak menunjukkan perencatan edema yang signifikan ($p <0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan varian tetapi tidak signifikan ($p>0.05$) dengan bahagian tumbuhan. Haiwan dibunuh selepas 6 jam dan PNL dalam tisu tapak kaki telah dikira. Rawatan oleh kesemua ekstrak menunjukkan pengurangan pada penyusupan

PNL yang signifikan ($p<0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan varian dan bahagian tumbuhan. 0.1ml histamin telah disuntik pada subplantar sebelum administrasi 100 mg/kg setiap ekstrak. Rawatan oleh kesemua ekstrak menunjukkan perencatan edema yang signifikan ($p<0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan varian tetapi tidak signifikan ($p>0.05$) dengan bahagian tumbuhan. Haiwan dibunuh selepas 2 jam dan PNL dalam tisu tapak kaki telah dikira. Rawatan oleh semua ekstrak menunjukkan pengurangan pada penyusupan PNL yang signifikan ($p <0.05$). Ia berbeza secara signifikan ($p<0.05$) dengan varian tetapi tidak signifikan ($p<0.05$) dengan bahagian tumbuhan.

Kesimpulannya, kajian menunjukkan bunga dan daun *Hibiscus rosa-sinensis* var *alba* dan *Hibiscus rosa sinensis* L. menghasilkan aktiviti anti-inflamasi akut. Ia mungkin melibatkan perencatan cyclooxygenase, bradykinin, histamine dan mengurangkan penyusupan polimorphonukleus leukosit.

ACKNOWLEDGEMENTS

It was really a laborious task to accomplish a master research. Many people gave me support and help in the process of completing this research. I would like to express my deep sense of gratitude to my research supervisor, Professor Dr. Muhammad Nazrul Hakim bin Abdullah, for his insightful guidance and earnest help all through the information searching, paper-writing stages and invaluable comments. He was being the integral part of this thesis throughout all its stages, both intellectual and the practical level. I really appreciate his generosity in sharing his knowledge and time, and for being attentive to my every question.

Also, I would like to acknowledge my sincere thanks to my co-supervisor, Dr. Roslida Abd Hamid @ Abdul Razak and Associate Prof. Dr Zainul Amirudin Zakaria, for their generous assistance in my research. I am very thankful to Dr Mohamad Aris Mohd Moklas in his guidance throughout histopathology analysis and Dr Mohd Khairi Hussain for his helping mostly in methodology. Million of thanks to Associate Prof Dr Aziz bin Dollah and Associate Prof Dr Saidi bin Moin for helping in the statistical analysis for my research.

My special thanks also goes to my friend, Mr. Muhammad Wahizul Haswan bin Abdul Aziz for lending a hand helping my research. Furthermore, my deep appreciation is sincerely goes to all the staff of Faculty of Medicine and Health Sciences, especially Mr. Ramli (Animal House), Miss Nurhayati Zainal Abidin and Mr. Zain Zailan (Pharmacology and Toxicology Lab), Madam Juita Chupri (Histopathology Lab) and Madam Noridah Mat Top (Research Laboratory of

Anatomy) for providing me with the experimental animals, equipments and reagents needed. Special thanks for their warmhearted helps and cooperation in assisting my research project gone smoothly. I apologized to any unintended inconvenience caused.

Last but not least, I would like to acknowledge my heartfelt appreciation to all my friends. Thanks for sharing their time to facilitate me in various task of experiment. Thanks for being sharing their information and ideas with me in completing this thesis. Finally, I would like to dedicate my thesis to my beloved family. Whenever I was on the verge of giving up, they always stood by me and never hesitated to give me a hand.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the Master of Science. The members of the Supervisory Committee were as follows:

Dr. Muhammad Nazrul Hakim, PhD

Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Chairman)

Dr. Roslida Abd Hamid @ Abdul Razak, PhD

Senior Lecturer,

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Member)

Dr. Zainul Amirudin Zakaria, PhD

Associate Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean

School of Graduate Studies,

Universiti Putra Malaysia

Date: 15 August 2013

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



SITI ZALEHA BINTI RADUAN

Date: 20 June 2013

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxii
 CHAPTER	
1 INTRODUCTION	1
1.1 Introduction	1
1.2 Objectives	3
1.2.1 General objectives	3
1.2.2 Specific objectives	3
1.3 Hypothesis	4
2 LITERATURE REVIEW	5
2.1 Inflammation	5
2.1.1 Introduction	5
2.1.2 Acute Inflammation	7
2.1.3 Chronic Inflammation	12
2.2 Chemical Mediator of Bradykinin	14
2.2.1 Bradykinin and Kinin Receptors Classification	14
2.2.2 Effects of Bradykinin on Polymorphonuclear Leukocytes Infiltration	15
2.3 Chemical Mediator of Histamine	16
2.3.1. Histamine and Receptors Classification	16
2.3.2 Effects of Histamine on Polymorphonuclear Leukocytes Infiltration	21
2.4 Non Steroidal Anti-Inflammatory Drugs	24
2.4.1 Diclofenac and Its Sodium	29
2.4.2 Effects of Non Steroidal Anti-Inflammatory Drugs on Polymorphonuclear Leukocytes Infiltration	31
2.5 Kinin Antagonist Drug	32
2.5.1 HOE-140	32
2.5.2 Effects of HOE-140 on Polymorphonuclear Leukocytes Infiltration	34
2.6 Antihistamine Drug	34
2.6.1 H ₁ -antihistamine; Loratadine	34
2.6.2 Effects of Loratadine on Polymorphonuclear Leukocytes Infiltration	39

2.7	<i>Hibiscus rosa-sinensis</i>	40
2.7.1	Introduction	40
2.7.2	Chemical Compounds in <i>Hibiscus rosa-sinensis</i> L.	42
2.7.3	Previous Study in Biomedical Research	42
2.8	Phytochemicals	43
2.8.1	Alkaloids	43
2.8.2	Flavonoids	45
2.8.3	Saponins	49
2.8.4	Tannins	52
2.8.5	Triterpenes	53
2.8.6	Steroids	55
2.9	Fixed Dose Procedure	57
2.10	Anti-acute Inflammatory Test	60
2.10.1	Paw Edema Test and PNL Infiltration in Carrageenan-induced Paw Edema in Rats	60
2.10.2	Formalin-induced Paw Licking Test	63
2.11	Mechanism of Action Test	63
2.11.1	Paw Edema Test and PNL Infiltration in BK-induced Paw Edema in Rats	63
2.11.2	Paw Edema Test and PNL Infiltration in Histamine-induced Paw Edema in Rats	65
3	METHODOLOGY	67
3.1	Plant Material	67
3.2	Preparation of Ethanol extract of Flower and Leaf of <i>H.rosa-sinensis</i> L. and <i>H.rosa-sinensis</i> var. <i>alba</i>	67
3.3	Phytochemical Screening	68
3.3.1	Test for Alkaloids	68
3.3.2	Test for Flavonoids	68
3.3.3	Test for Saponins	69
3.3.4	Test for Tannins	69
3.3.5	Test for Triterpenes	69
3.3.6	Test for Steroids	70
3.4	STUDY I: Determination of Acute Dose Response	70
3.5	Study II: Anti-Acute Inflammatory Test	72
3.5.1	Paw Edema Test and PNL Infiltration in Carrageenan-induced Paw Edema in Rats	72
3.5.2	Formalin-induced Paw Licking Test in Rats	74
3.6	Study III: Mechanism of Action Test	76
3.6.1	Paw Edema Test and PNL Infiltration in BK-induced Paw Edema in Rats	76
3.6.2	Paw Edema Test and PNL Infiltration in Histamine-induced Paw Edema in Rats	78
4	RESULTS	80
4.1	Phytochemical Screening	80
4.2	Determination of Acute Dose Response	82
4.2.1	Liver Function Test	82

4.2.2	Histopathological Examinations of Liver and Kidney of Rat	85
4.3	Anti-Acute Inflammatory Test	88
4.3.1	Paw Edema Test and PNL Infiltration in Carrageenan-induced Paw Edema in Rats	88
4.3.2	Effects of 100mg/kg of Ethanol Extracts of Flower and Leaf of <i>H.rosa sinensis</i> L. (red) and <i>H.rosa-sinensis</i> var <i>alba</i> (white) on Formalin-induced Paw Licking	112
4.4.	Mechanism of Action Test	115
4.4.1	Paw Edema Test and PNL Infiltration in BK-induced Paw Edema in Rats	115
4.4.2	Paw Edema Test and PNL Infiltration in Histamine-induced Paw Edema in Rats	125
5	DISCUSSION	135
6	CONCLUSION	150
	REFERENCES	152
	APPENDICES	174
	BIODATA OF STUDENT	209