

STUDY OF LACTIC ACID BACTERIA AND *ENTEROBACTERIACEAE* COUNTS AND FATTY ACIDS COMPOSITION IN A FERMENTED PRODUCT

F.L. Law¹, T.C. Loh^{1*}, H. L. Foo^{2,4}, Y.M. Goh³ and I. Zulkifli¹

¹Department of Animal Science, ²Department of Bioprocess Technology, ³Department of Veterinary Preclinical Sciences, ⁴Institute of Bioscience, Universiti Putra Malaysia, 43300 UPM Serdang, Selangor, Malaysia

SUMMARY

This study was conducted to study the characteristics of a fermented product (FP). An evaluation of measurements of consistency and repeatability on microbiological analyses, including lactic acid bacteria (LAB) and *Enterobacteriaceae* counts and fatty acids profiles was performed. Physico-chemical properties (pH, dry matter and ash, crude protein, crude fat and crude fibre) of FP were ascertained. The fermentation process was repeated three times under the same running conditions and using the same amounts of raw materials. The results indicated that the product obtained from the fermentation process increased the LAB population and decreased the *Enterobacteriaceae* populations after the fermentation process. The presence of essential fatty acids (linoleic acid 24%, linolenic acid 1.1%, Eicosapentaenoic acid 3.6%, Docosapentaenoic acid 0.7%, and Docosahexaenoic acid 2.0%) and lime-flavored aroma represents added value attributes to the product and it might have potential as an animal feed additive. Chemical determination showed a net decrease in pH to around 4.5. The dry matter, crude fibre, crude protein, crude fat and ash contents of the three batches of FP were similar. These results indicated that the quality of FP was consistent with repeatable preparation process.

Keywords: Fermented product, fermentation, lactic acid bacteria, *Enterobacteriaceae*, fatty acid profile

INTRODUCTION

Fermentation is one of the oldest technologies used for food and feed preservation. Over the centuries, the fermentation process has been evolved and refined extensively to produce a diversified range of products (Prajapati and Nair, 2003). It is possible to obtain a large variety of fermented products by selecting different types and compositions of raw materials, starter cultures and fermentation conditions (Hansen, 2002). Generally, fermented products contain high numbers of lactic acid bacteria, a low pH and a high concentration of lactic acid. The physical, chemical and biological characteristics have been modified by the activity of microorganisms. Microorganisms, by virtue of their metabolic activities, contribute to the development of characteristic properties such as taste, aroma, texture, visual appearance, shelf life and safety of the product (Holzapfel, 2002). The actions of microorganisms during the preparation of cultured foods play an important role in improving the safety of feeds by removing their natural toxic components, or by preventing the growth of disease-causing microbes (e.g. *Enterobacteriaceae*). Fermentation with LAB cultures has been shown to be able to improve the quality, availability and digestibility of dietary nutrients. LAB improves protein digestibility and micronutrient

bioavailability of fermented feed through the biosynthesis of vitamins and essential amino acids. Thus, they play an essential role in food and feed fermentation and are well known for their acidification of the product (Hammes *et al.*, 1990).

During the early ages of civilisation, fermented products were mainly used for human consumption. The beneficial LAB present in the products is well documented for its antimicrobial compounds such as bacteriocin, as a biopreservative for food products (Schnürer and Magnusson, 2005). Products derived from LAB fermentation have the potential to prevent various human diseases (Minamiyama *et al.*, 2003). Recently, fermented liquid feed has been used as an animal diet. Fermented liquid feed has also shown promising results in enhancing the performance of animals (Mikkelsen and Jensen, 1998; Demecková *et al.*, 2002). However, fermented liquid feed is tedious to prepare and apply. The solid form FP investigated in this study was designed to overcome these limitations. Most of the studies on fermentation had been done on animal and agricultural by-products (Kamra and Srivastava, 1994; Joshi and Sandhu, 1996; El Jalil *et al.*, 2001). Up to date, none of the fermentation studies done on agricultural by-products involved the addition of marine fish in order to enhance the nutritive value of the final product. In the present study, fish

*Corresponding authors: Assoc. Prof. Dr Loh Teck Chwen; E-mail- tcloh@agri.upm.edu.my

was used to increase the levels of polyunsaturated fatty acids (PUFA) as PUFA have been used successfully to manipulate and increase the n-3 levels in animal products for livestock production (Milinsk *et al.*, 2003). Thus, the objectives of this study were to evaluate the chemical composition and nutritive value of the product after the process of fermentation.

MATERIALS AND METHODS

Preparation of fermented product (FP)

The fermentation process was performed according to the method as described by Loh *et al.* (2003a). The raw materials consisted of 9% (wt/wt) lime, 1% (wt/vol) molasses, 53.5% (wt/wt) rice bran, 35% (wt/wt) fish (*Rastrelliger kanagurta*), 1% (wt/vol) vinegar, and 0.5% (wt/wt) starter cultures. The starter culture was a combination of *Lactobacillus sp.* with a population of 3.56×10^4 . The starter culture was obtained from Jia Yi Nutritional Sdn. Bhd (Klang, Malaysia). Lime was blended with molasses and the mixture was then fermented at ambient temperature ($28 \pm 2^\circ\text{C}$) for five days under aerobic conditions. The mixture was then poured into a closed 50L solid fermenter together with rice bran, raw fish, vinegar and starter culture. The fermenting mixture was mixed hourly in order to achieve a homogenous mixing. The fermentation process was carried out for seven days at $35\text{--}38^\circ\text{C}$. Fermentation was considered complete after seven days. The completed FP was allowed to cool to room temperature prior to sampling. The final product was a dry-brownish mash with a distinctive lime-flavoured aroma. The FP was stored in a vacuum-sealed black plastic bag until use.

Chemical and microbiological analyses

The pH of the FP was measured with a Mettler-Toledo pH meter (Mettler-Toledo LTD., England) according to the method as described by Loh *et al.* (2003b). Ten percent (wt/vol) of the sample was mixed homogeneously with deionised distilled water. Dry matter was determined by oven drying at 103°C to a constant weight. Ash was determined by ignition in a muffle furnace at 600°C . The crude protein (micro Kjeldahl method), crude fibre and crude fat (Soxhlet method with petroleum ether as solvent) were analysed as per AOAC (1990) procedure. The microbiological analyses were done according to the method as described by Loh *et al.* (2003a,b). Enumerations of total lactobacilli were performed on MRS-agar (*Lactobacillus*-Agar DE Man, ROGOSA and SHAPE) (Merck). The plates were incubated in an anaerobic jar at 30°C for 48 hours. *Enterobacteriaceae* were counted on EMB-agar (Eosin-methylene-blue Lactose Sucrose Agar) (Merck) and incubated aerobically for 24 hours at 37°C . Numbers of

colony-forming unit (cfu) were expressed as \log_{10} cfu/g. All samples were performed in triplicate.

Determination of fatty acids profiles

The total lipids were extracted from the product using a modified Folch *et al.* (1957) method as described by Rajion *et al.* (2001) using chloroform-methanol 2:1 (vol/vol) solvent system. Transmethylation was carried out using 14% methanolic boron trifluoride (Sigma Chemical Co., St. Louis, Missouri, USA). The derived fatty acid methyl esters (FAME) were separated by gas chromatography using a 5890 Hewlett-Packard Gas-Liquid Chromatograph (Hewlett-Packard, Avondale, PA) fitted with a flame ionisation detector (FID) and a Supelco SP-2330 fused silica capillary Column (30m, 0.25mm ID, 0.20 μm film thickness) (Supelco, Inc., Bellefonte, PA). The column temperature was programmed at $7.2^\circ\text{C}/\text{min}$ from 100 to 190°C . The injector and detector were programmed at 220°C and 220°C , respectively. High purity nitrogen was the carrier gas at a flow rate of 40ml/min. An internal standardisation method was used to quantify the various fatty acids in the FP, where a known concentration of heneicosanoic acid was added to each sample prior to transmethylation. Peak areas were determined by the HP-3993A Integrator (Hewlett-Packard, Avondale, PA). The identification of the peak was made by comparison of equivalent chain lengths (ELC) with those of authentic fatty acid methyl esters (Sigma Chemical Co., St. Louis, Missouri, USA).

Statistical analysis

Results were expressed as mean \pm standard error of the mean. Total lactobacilli and *Enterobacteriaceae* counts were expressed as \log_{10} cfu/g. The entire colony counts, pH value, fatty acid profile, crude protein, crude fibre, crude fat, dry matter were recorded and analysed by one-way ANOVA using SAS (SAS[®], 1991). Duncan's Multiple Range Test was used to compare the differences of mean and the level of significance was $P < 0.05$.

RESULTS AND DISCUSSION

Microbiological characteristics of FP

The fermentation process performed in this study demonstrated reduction of harmful bacteria such as *Enterobacteriaceae* in the end product compared to the raw materials (Table 1). In contrast, the population of LAB increased in the end product as compared to the raw materials. The results obtained in this study were consistent with the findings of other reports (Mikkelsen and Jensen, 2000; van Winsen *et al.*, 2002; Demecková *et al.*, 2002; Loh *et al.*, 2003a; b; Heres *et al.*, 2004). The use of potentially beneficial bacteria such as LAB has been demonstrated to inhibit the *in-vitro* growth of many

Table 1: Total lactobacilli and *Enterobacteriaceae* counts in raw materials and finished FP

Microorganisms Log ₁₀ (cfu/g)	Fermentation time	Batch 1 (n=3)	Batch 2 (n=3)	Batch 3 (n=3)
Total Lactobacilli	Day 0	4.28 ^{a,x} ± 0.01	4.40 ^{b,x} ± 0.01	4.37 ^{b,x} ± 0.02
	Day 12	6.26 ^{a,y} ± 0.03	6.25 ^{a,y} ± 0.11	6.41 ^{a,y} ± 0.02
<i>Enterobacteriaceae</i>	Day 0	3.87 ^{a,x} ± 0.01	3.87 ^{a,x} ± 0.02	3.83 ^{a,x} ± 0.03
	Day 12	0.83 ^{a,y} ± 0.16	0.87 ^{a,y} ± 0.20	1.26 ^{a,y} ± 0.41

The results are presented as mean values ± SEM.

^{a,b}Values with different superscripts within a row differ significantly at P<0.05 due to batch effects;

^{x,y}Values with different superscripts within a column differ significantly at P<0.05 due to day effects

Table 2: Physico-chemical characteristics of the FP

Parameters	Batch 1 (n=3)	Batch 2 (n=3)	Batch 3 (n=3)
pH	4.75 ^a ± 0.02	4.21 ^b ± 0.01	4.55 ^c ± 0.01
GE, Kcal/kg ^{ns}	4490.97 ± 32.20	4509.98 ± 8.70	4415.69 ± 156.00
(%, DM basis)			
DM ^{ns}	90.69 ± 0.83	92.65 ± 0.07	92.18 ± 0.23
Crude fiber ^{ns}	0.07 ± 0.01	0.06 ± 0.01	0.07 ± 0.01
Crude proteins ^{ns}	19.97 ± 0.10	19.00 ± 0.33	19.21 ± 0.22
Crude fat ^{ns}	0.11 ± 0.01	0.13 ± 0.01	0.13 ± 0.01
Ash ^{ns}	1.02 ± 0.01	0.99 ± 0.01	1.00 ± 0.02

The results are presented as mean values ± SEM.

Values with different superscripts within a row differ significantly at P<0.05;

^{ns} No significant difference

enteric pathogens including *Salmonella typhimurium*, *Staphylococcus aureus*, *Escherichia coli*, *Clostridium perfringens* and *Clostridium difficile* (Lindgren *et al.*, 1990; de Vuyst and Vandamme, 1994; Forestier *et al.*, 2001). Furthermore, *in-vivo* study of feeding fermented feed improved the microbial balance in the GIT of animals (Urlings *et al.*, 1993; Mikkelsen and Jensen, 2000; van Winsen *et al.*, 2002; Demecková *et al.*, 2002; Loh *et al.*, 2003a,b; Heres *et al.*, 2004). It may be that the beneficial LAB present in the product produced a range of antimicrobial metabolites such as bacteriocins, ethanol, H₂O₂, and organic acids during fermentation (Ross *et al.*, 2002). Organic acids helped to reduce the pH value to a range of 4-5, which was detrimental for deleterious microbes (Østergaard *et al.*, 1998; Christine *et al.*, 1999; Beal *et al.*, 2002). This in turn, provided a suitable environment to promote the growth and sustenance of the lactic acid-producing microflora (Owen and Mendoza, 1985; Urlings *et al.*, 1993; van Winsen *et al.*, 2002; Loh *et al.*, 2003 a; b).

The undissociated, more hydrophobic form of the organic acids diffuses over the cell membrane. It dissociates inside the cell thus releasing H⁺-ions that acidify the cytoplasm and reducing the intracellular pH. This in turn inhibits the active transport and a variety of metabolic

functions (Doores, 1993; Ross *et al.*, 2002). In addition to the pH effect, the undissociated acid collapses the electro-chemical proton gradient, causing bacteriostasis and finally death of susceptible bacteria (Axelsson *et al.*, 1989 as cited by Schnürer and Magnusson, 2005) such as *Enterobacteriaceae* in the FP. The reduction of *Enterobacteriaceae* in the FP is important to product safety as it indicates a safe product and a successful fermentation process where hazardous microorganisms are inhibited. Thus, this product may be used safely in the animal feed industry for optimal growth performance and as a prophylaxis against infectious agents such as *Salmonella* spp.

Physico-chemical characteristics of FP

The dry matter (DM), crude fibre, ash, crude protein (CP), crude fat and gross energy (GE) contents and the fatty acids compositions were not significantly different (P>0.05) across batches except pH (P<0.05) which ranged from 4.21 to 4.75 (Table 2). These results indicate that the FP could be produced at a consistent quality under a reproducible fermentation process. Presence of beneficial long-chain fatty acids (PUFA) such as

Table 3: Fatty acid compositions of FP in different batches

Fatty Acids	Batch 1		%	Batch 2		%	Batch 3		%
	Mean ± SEM	(mg/100g)		Mean ± SEM	(mg/100g)		Mean ± SEM	(mg/100g)	
C12:0	6.2 ± 1.1		0.1	5.9 ± 0.8		0.1	6.1 ± 1.4		0.1
C14:0	200.6 ± 13.1		2.9	193.7 ± 2.36		2.9	212.3 ± 22.6		2.9
C15:0	28.5 ± 9.0		0.4	26.9 ± 4.3		0.4	30.7 ± 11.6		0.4
C16:0	1444.8 ± 121.0		20.8	1393.5 ± 170.0		20.9	1525.0 ± 31.0		20.8
C16:1 n-9	270.6 ± 21.6		3.9	259.5 ± 34.8		3.9	287.8 ± 34.7		3.9
C17:0	30.2 ± 7.2		0.4	28.7 ± 5.8		0.4	32.3 ± 9.8		0.4
C17:1	40.7 ± 7.1		0.6	38.4 ± 5.6		0.6	43.1 ± 10.1		0.6
C18:0	294.3 ± 20.2		4.2	282.1 ± 37.5		4.2	311.7 ± 29.0		4.2
C18:1 n-9	2216.8 ± 387.0		31.9	2138.9 ± 237.0		32.0	2342.7 ± 231.0		31.9
C18:2 n-6	1671.4 ± 349.0		24.1	1613.2 ± 167.0		24.1	1764.2 ± 246.0		24.0
C18:3 n-3	73.3 ± 12.5		1.1	71.3 ± 5.6		1.1	76.7 ± 10.1		1.0
C20:0	49.7 ± 7.3		0.7	48.8 ± 6.1		0.7	52.2 ± 3.9		0.7
C20:1 n-9	35.2 ± 4.1		0.5	34.6 ± 4.2		0.5	37.0 ± 1.2		0.5
C20:4 n-6	86.3 ± 19.9		1.2	81.3 ± 15.8		1.2	91.9 ± 28.2		1.3
C22:0	24.1 ± 2.7		0.3	22.6 ± 3.4		0.3	25.4 ± 0.9		0.3
C24:0	36.5 ± 4.2		0.5	35.4 ± 4.9		0.5	41.2 ± 1.6		0.6
C20:5 n-3, EPA	246.9 ± 19.8		3.6	230.6 ± 22.4		3.5	261.8 ± 31.4		3.6
C22:5 n-3, DPA	46.4 ± 6.8		0.7	44.4 ± 8.2		0.7	50.1 ± 10.8		0.7
C22:6 n-3, DHA	141.9 ± 18.1		2.0	132.3 ± 13.7		2.0	152.1 ± 25.2		2.1
Total SFA	2114.8 ± 149.0		30.5	2037.5 ± 256.0		30.5	2237.0 ± 102.0		30.5
Total UFA	4829.4 ± 690.0		69.5	4644.5 ± 509.0		69.5	5107.4 ± 350.0		69.5
Total MUFA	2563.3 ± 382.0		36.9	2471.5 ± 282.0		37.0	2710.6 ± 189.0		36.9
Total PUFA n-3	508.4 ± 31.4		7.3	478.6 ± 48.5		7.2	540.7 ± 57.4		7.4
Total PUFA n-6	1757.7 ± 334.0		25.3	1694.5 ± 182.0		25.4	1856.1 ± 218.0		25.3
Overall total	6944.2 ± 802.0			6682.0 ± 765.0			7344.4 ± 255.0		
n-6 : n-3 ratio	3.5		3.5	3.4					
UFA: SFA ratio	2.3		2.3	2.3					
PUFA: SFA ratio	1.1		1.1	1.1					

The results are presented as mean values ± SEM.

All values within a row are not significantly different from each other ($P > 0.05$).

n-3 (linolenic, EPA, DPA and DHA) and n-6 (linoleic, arachidonic) classes fatty acids in this product (Table 3) were a value-added to the product as these are now regarded as nutritionally important fatty acids (Hansen, 1986; Lauritzen *et al.*, 2001; Elmore *et al.*, 2005). The presence of n-3 long chain PUFA in the product was a direct result of incorporation of a sizeable portion of marine fish (*Rastrelliger kanagurta*) as one of the raw materials. Marine fish are rich in n-3 fatty acid because their diet is made up of zooplankton and phytoplankton, a rich source of PUFA (Kyle, 2002). This could offer an added advantage, which helps to improve the content of these fatty acids in animal and animal products when the FP is being fed to livestock. The long chain n-3 fatty acids are prized for the beneficial effects to human health and feedstuffs with added n-3 are definitely more advantageous over conventional feeds. However, the long-chain PUFA particularly the members of n-3 fatty acids are susceptible to lipid oxidation. Surprisingly, these PUFA especially EPA, DPA and DHA were detected in a substantial amount in the FP. EPA was found to be the highest

among all the n-3 long chain fatty acids in the FP. Besides, the high content of n-6 PUFA detected in the FP was probably a direct result of the incorporation of more than 50% rice bran as one of the major components of raw materials. Linoleic acid (the predominant n-6 fatty acid in the product) is the major fatty acid present in the lipid fraction of the rice bran (Ito and Simpson, 1996; Rouanet *et al.*, 1993). Furthermore, rice bran contains high levels of several phytochemicals are claimed to be a good source of natural antioxidants (Krings *et al.*, 2000; Kanaya *et al.*, 2004; Iqbal *et al.*, 2005). Among these natural antioxidants are vitamin E, tocopherols, tocotrienols and the γ -oryzanol fraction, which are thought to exert anti-oxidative and protective effects on fatty acids in animal tissues (Qureshi and Peterson, 2001; Kanaya *et al.*, 2004). In addition, rice bran has been shown to improve the storage stability of foods (Iqbal *et al.*, 2005). Thus, it may have acted as a natural antioxidant during fermentation thereby improving the storage stability of FP and helping to prevent the oxidation of long-chain PUFA in FP.

CONCLUSION

The physical, chemical and microbiological properties of FP obtained in this study showed that the FP was highly consistent and reproducible. FP was rich in LAB and had a low pH. Furthermore, the presence of essential PUFA such as linoleic, linolenic, arachidonic, EPA, DPA and DHA in the FP was another important characteristic of this product. The FP was a dry-brownish mash with distinctive lime-flavoured aroma. This could have an added advantage as it would overcome the problem of unpleasant odour in feed such as fishmeal, as off-flavoured products could be a significant market disadvantage. Based on the results obtained in the present study, it indicates that FP has great potential as an animal feed. The results obtained from this study are essential for large-scale FP production for animal feed applications in the future.

REFERENCES

- AOAC. (1990). Official Methods of Analysis. 15th edn. Association of Official Analytical Chemist, Arlington, Virginia.
- Axelsson, L.T, Chung, T.C., Dobrogosz, W.J. and Lindgren, S.E. (1989). Production of a broad spectrum antimicrobial substance by *Lactobacillus reuteri*. *Microbiol. Ecol. of Health and Disease* 2: 131-136.
- Beal, J.D., Niven, S.J., Campbell, A. and Brooks, P.H. (2002). The effect of temperature on the growth and persistence of Salmonella in fermented liquid pig feed. *Int. J. of Food Microbiol.* 79: 99-104.
- Christine, P.M., Huss H.H. and Gram, L. (1999). Characterization of lactic acid bacteria isolated from a Thai low-salt fermented fish product and the role of garlic as substrate for fermentation. *Int. J. of Food Sci.* 46: 219-229.
- Demecková, V., Kelly, D., Coutts, A.G.P., Brooks, P.H. and Campbell, A. (2002). The effect of fermented liquid feeding on the faecal microbiology and colostrum quality of farrowing sows. *Int. J. of Food Microbiol.* 79: 85-97.
- de Vuyst, L and Vandamme, E.J. (1994). Antimicrobial potential of lactic acid bacteria. In: Bacteriocin of lactic acid bacteria. de Vuyst, L and E.J. Vandamme (Ed.). Glasgow, United Kingdom, Blackie Academic and Professional. pp. 91-142.
- Doores, S. (1993). Organic acids. In: Organic acids. Davidson, P.M. and A.L. Branen (Ed.). Marcel Dekker, New York. pp. 95-136.
- El Jalil, M.H., Faid, M. and M. Elyachioui. (2001). A biotechnological process for treatment and recycling poultry waste manure as a feed ingredient. *Biomass and Bioenergy* 21: 301-309.
- Elmore, J.S., Cooper, S.L., Enser, M., Mottram, D.S., Sinclair, L.A., Wilkinson, R.G and Wood, J.D. (2005). Dietary manipulation of fatty acid composition in lamb meat and its effect on the volatile aroma compounds of grilled lamb. *Meat Sci.* 69:233-242.
- Folch, J. M. and Lees, G.H., and Stanley, S. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chemistry.* 226:497-509.
- Forestier, C., de Champs, C., Vatoux, C. and Joly, B. (2001). Probiotic activities of *Lactobacillus casei rhamnosus*: in-vitro adherence to intestinal cells and antimicrobial properties. *Research in Microbiol.* 152:167-173.
- Hammes, W.P, A. Bantleon and S. Min. (1990). Lactic acid bacteria in meat fermentation. *FEMS Microbiol. Rev.* 87: 165-174.
- Hansen, E.B. (2002). Commercial bacterial starter cultures for fermented foods of the future. *Int. J. Food Microbiol.* 78: 119-131.
- Hansen, H.S. (1986). The essential nature of linoleic acid in mammals. *Trends in Biochem. Sci.* 11:263-265.
- Heres, L., Engel, B., Urlings, H.A.P, Wagenaar J.A. and van Knapen, F. (2004). Effect of acidified feed on susceptibility of broiler chickens to intestinal infection by *Campylobacter* and *Salmonella*. *Vet. Microbiol.* 99: 259-267.
- Holzappel, W.H. (2002). Appropriate starter culture technologies for small-scale fermentation in developing countries. *Int. J. Food Microbiol.* 75: 197-212.
- Iqbal, S., Bhangar, M.I., and Anwar, F. (2005). Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chem.* 93:265-272.
- Ito, M.K and Simpson, K.L. (1996). The biosynthesis of n-3 fatty acids from 18:2n-6 in *Artemia* spp. *Comp. Biochem. and Molecular Biol.* 115: 69-76.
- Joshi, V.K and Sandhu, D.K. (1996). Preparation and evaluation of an animal feed byproduct produced by solid-state fermentation of apple pomace. *Bioresource Technol.* 56: 251-255.

- Kamra, D.D and Srivastava, S.K. (1994). Effect of sugarcane molasses on fermentation of pig faeces and wheat straw inoculated with lactic-acid producing bacteria. *Bioresource Technol.* **47**: 87-88.
- Kanaya, Y., Doi, T., Sasaki, H., Fujita, A., Matsuno, S., Okamoto, K., Nakano, Y., Tsujiwaki, S., Furuta, H., Nishi, M., Tsuno, T., Taniguchi, H. and Nanjo, K. (2004). Rice bran extract prevents the elevation of plasma peroxy lipid in KKAY diabetic mice. *Diabetes Research and Clinical Practice* **66S**: S157-S160.
- Krings, U, El-Saharty, Y.S., El-Zeany, B.A., Pabel, B. and Berger, R.G. (2000). Antioxidant activity of extracts from roasted wheat germ. *Food Chem.* **71**:91-95.
- Kyle, D.J. (2002). Essential fatty acids as food additives. In: Food additives. Brenen, A.L, P.M. Davidson, S. Salminen and J.H. Thorngate (Ed.). Marcel Dekker. New York. pp.277-309
- Lauritzen, L., Hansen, V., Jørgensen, M.H., and Michaelsen, K.F.(2001). The essentiality of long chain n-3 fatty acids in relation to development and function of the brain and retina. *Progress in Lipid Research* **40**: 1-94.
- Lindgren, S.E and Dobrogosz, W.J. (1990). Antagonistic activities of lactic acid bacteria in food and feed fermentations. *FEMS Microbiol. Review* **87**: 149-164.
- Loh, T.C., Foo, H.L., Lee, K.L., Lim, Y.Z. and Kufli, C.N. (2003) a. Effects of fermented fruits on the growth performance, Shedding of *Enterobacteriaceae* and *Lactobacilli* in post-weaning pigs. *Asian-Aust. J. Anim. Sci.* **16**: 1-5.
- Loh, T.C., Foo, H.L., Tan, S.H., Goh, Y.M., Shukriyah, M.H. and Kufli, C.N. (2003) b. Effects of fermented products on performance, faecal pH, enterobacteriaceae and lactic acid bacteria counts and inter-relationships and plasma cholesterol concentration in rats. *J Anim. and Feed Sci.* **12**: 633-644.
- Mikkelsen, L.L and Jensen B.B. (1998). Performance and microbial activity in the gastrointestinal tract of piglets fed fermented liquid feed at feeding. *J. Anim. Sci.* **7**: 211-215.
- Mikkelsen, L.L and Jensen, B.B. (2000). Effects of fermented liquid feed on the activity and composition of the microbiota in the gut of pigs. *Pigs News and Information* **21**: 59N-66N.
- Milinsk, M.C, Murakami, A.E., Gomes, S.T.M, Matsushita, M. and de Souza, N.E. (2003). Fatty acid profile of egg yolk lipids from hens fed diets rich in n-3 fatty acids. *Food Chem.* **83**: 287-292.
- Minamiyama, Y, Takemura, S., Yoshikawa, T. and Okada, S. (2003). Fermented grains products, production, properties and benefits to health. *Pathophysiol.* **9**: 221-227.
- Østergaard, A., Ben Embarek, P.K., Yamprayoon, J., Wedel-Neer-gaard, C., Huss, H.H. and Gram, L. (1998). Fermentation and spoilage of som-fak: a Thai low-salt fish product. *Trop. Sci.* **38**: 105-112.
- Owen, J. D and Mendoza, L.S. (1985). Enzymically hydrolysed and bacterially fermented fishery products. *J. Food Technol.* **20**:273-293.
- Prajapati, J.B and Nair, B.M. (2003). The history of fermented foods. In: Handbook of Fermented Functional Foods. Farnworth, E.R. (Ed.). CRC Press. New York. pp. 1-26.
- Qureshi, A.A and Peterson, D.M. (2001). The combined effects of novel tocotrienols and lovastatin on lipid metabolism in chickens. *Atherosclerosis* **156**: 39-47.
- Rajion, M.A., Goh, Y.M., Dahlan, I. and Salam-Abdullah, A. (2001). Dietary manipulation and increase in plasma unsaturated fatty acids in sheep. *Asian-Australian J. of Anim. Sci.* **14**: 1073-1077.
- Ross, R.P., Morgan, S. and Hill, C. (2002). Preservation and fermentation: past, present and future. *Int. J. Food Microbiol.* **79**: 3-16.
- Rouanet, J.M, Laurent, C. and Besancon, P. (1993). Rice bran and wheat bran: selective effect on plasma and liver cholesterol in high-cholesterol fed rats. *Food Chem.* **47**: 67-71.
- SAS Institute Inc. (1991). SAS Institute Inc., Cary, North Carolina.
- Schnürer, J and Magnusson, J. (2005). Antifungal lactic acid bacteria as biopreservatives. *Trends in Food Sci. and Technol.* **16**:70-78
- Urlings, H.A.P., Mul, A.J., vat't Klooster, A.T., Bijker, P.G., van Logtestijn, J.G. and van Gils, L.G. (1993). Microbial and nutritional aspects of feeding fermented feed (poultry by-products) to pigs. *Vet. Quart.* **15**: 146-151.
- Van Winsen, R.L, Urlings, B.A.P. and Keuzenkamp, D. (2002). Effect of fermented feed on shedding of *Enterobacteriaceae* by fattening pigs. *Vet. Microbiol.* **87**: 267-276.