

## MORPHOLOGICAL AND LECTIN HISTOCHEMICAL STUDIES OF THE SPERMATOZOA OF THE LESSER MOUSE DEER (*Tragulus javanicus*)

W. E. Prasetyaningtyas<sup>1</sup>, S. Agungpriyono<sup>1,2\*</sup>, Y. Yamamoto<sup>3</sup>, A.H. Wahid<sup>4</sup>,  
M. A. Setiadi<sup>5</sup> and M. Fahrudin<sup>1</sup>

<sup>1</sup>Department of Anatomy, Physiology and Pharmacology, Faculty of Veterinary Medicine  
Bogor Agricultural University, Bogor 16680, Indonesia

<sup>2</sup>Department of Preclinical Veterinary Sciences, Faculty of Veterinary Medicine, Universiti Putra Malaysia  
43400 UPM Serdang, Selangor, Malaysia

<sup>3</sup>Department of Anatomy, Faculty of Agriculture, Iwate University  
3-18-8 Ueda, Morioka, Iwate 020-8550, Japan

<sup>4</sup>Department of Clinical Studies, Faculty of Veterinary Medicine, Universiti Putra Malaysia  
43400 UPM Serdang, Selangor, Malaysia

<sup>5</sup>Department of Clinic, Reproduction and Pathology, Faculty of Veterinary Medicine  
Bogor Agricultural University, Bogor 16680, Indonesia

### SUMMARY

The morphology and lectin histochemical property of the spermatozoa of the lesser mouse deer (*Tragulus javanicus*) were studied using light and scanning electron microscopy. Spermatozoa were small in size and measured  $36.52 \pm 5.6 \mu\text{m}$  in length. The head of the spermatozoa was relatively round in shape ( $5.55 \pm 0.8 \mu\text{m}$  in length and  $4.77 \pm 0.5 \mu\text{m}$  in width). Lectin peanut (*Arachis hypogea*) agglutinin (PNA) and lectin wheat germ (*Triticum vulgare*) agglutinin (WGA) were positive in the membrane of the acrosomal region, and in the membrane at the anterior part of the acrosomal region and in the tail, respectively. It is suggested that carbohydrates with galactose  $\beta$ 1-3, D-N acetylgalactosamine, D-N acetylglucosamine and sialic acid sugar residues may be involved and may have significant roles in the spermatozoa in the lesser mouse deer.

Keywords: Sperm morphology, lectin histochemistry, lesser mouse deer, scanning electron microscopy

### INTRODUCTION

There is considerable variation in the size and shape of the head of the spermatozoa among species. For example, it is columnar and elongated in fowl, like a hook in the mouse and rat, or flattened and ovoid in man (Mann, 1964). The entire spermatozoon is covered by plasma membrane, which is rich in carbohydrate molecules. The majority of sugar residues are attached to the sperm membrane proteins. These glycoproteins may be closely related with sperm maturation, protection within the female genital tract, binding of spermatozoa to the zona pellucida and with the fertilisation process (Gilbert, 1988; Nolan and Hammerstedt, 1997; Zara and Naz, 1998; Tulsiani, 2000).

The lesser mouse deer (*Tragulus javanicus*) is regarded as the smallest ruminant in the world (Lekagul and McNeely, 1977) and is an ideal model in ruminant and biomedical research (Sastradipradja, 1978). However, there is a paucity of information on its reproductive biology. In previous studies, the characteristics and quality (Haron *et al.*, 1999) and chemical composition (Prasetyaningtyas *et al.*, 2006) of the sperm of the lesser mouse deer obtained by electroejaculation were reported.

However, the sperm morphology and histochemical properties regarding the distribution of specific carbohydrates have not been studied in the lesser mouse deer.

In the present study, the morphology and the distribution of lectin peanut (*Arachis hypogea*) agglutinin (PNA) and wheat germ (*Triticum vulgare*) agglutinin (WGA) bindings in the spermatozoa of the lesser mouse deer (*Tragulus javanicus*) were investigated in order to provide information on the reproductive biology of this species.

### MATERIALS AND METHODS

#### Semen collection

Four young adult male lesser mouse deer weighing 1.8–2.4 kg were used in this study. The age of the animal was estimated to be around 12–18 months based on the appearance of canine teeth. The animals were housed in the animal facility at the Faculty of Veterinary Medicine, Bogor Agricultural University, Bogor, Indonesia. Each animal was placed in an individual cage measuring 1 x 2

\*Corresponding author: Dr Srihadi Agungpriyono; Email: srihadi@vet.upm.edu.my

m<sup>2</sup>. They were fed twice a day with vegetables which included *ipomoea* leaves, long beans, lettuce, carrots, and potatoes, and fruits such as bananas and apples according to Jumaliah (1999) and water was given *ad libitum*.

The semen samples were collected twice a month for a duration of one year, using an electro-ejaculator (Electric Stimulator, Fujihiro Industry Co., Ltd, Tokyo, Japan) after the animals were anaesthetised using a combination of xylazine (Xylazyl 100<sup>®</sup>, Troy Lab Pty., Ltd., NSW, Australia; 0.1mg/kg BW) and ketamine (Ketavet 100<sup>®</sup>, Delvet Pty., Ltd., NSW, Australia; 11mg/kg BW). The electrical stimulations at low frequency were applied intermittently (Axner and Linde-Forsbeg, 2002) at 5-15 volts for a duration of 5 seconds, followed by 2-3 seconds rest. A total of 16 semen samples were observed in this study.

### Morphological studies

One drop of the semen sample was placed onto the object glass to which a cover slip was mounted to make native specimens. Other semen samples were processed for scanning electron microscopy (SEM). Briefly, a semen sample was dropped onto a cover glass and air dried. The samples were fixed in 2.5% glutaraldehyde for 2 h and were then immersed in 2% tannic acid for 1 h (Murakami, 1974). After washing with 0.01M PBS pH 7.4, the samples were then post fixed using 1% osmium tetroxide (OsO<sub>4</sub>) for 1 h, dehydrated in graded series of alcohol and *t*-butanol (Inoue and Osatake, 1988) and freeze dried using a freeze dryer (VFD-21s, Tokyo, Japan). Samples were then coated with platinum-palladium using an ion coater (Eiko IB-3, Tokyo, Japan) and observed using a SEM (JEOL, LV-5800, Tokyo, Japan) at magnification of 1000-2000x and an accelerating voltage of 10 kV. The measurement of the spermatozoa was taken

on native specimens using a light microscope equipped with measuring gauge and also from the SEM images. In the light microscopy, the measurement was taken at 400x magnification from around 50 spermatozoa per specimen. The length and width of the sperm head and also the total length of the spermatozoa were recorded in the spermatozoa with normal morphology.

### Lectin histochemistry

Semen samples were centrifuged at 640xg for 5 min and the sediment was used to make smear specimens. The specimens were fixed in 2.5% glutaraldehyde for 30 min at 60°C and stained using peroxidase labeled PNA and WGA lectins (Honen Co., Tokyo, Japan). The details of the lectins used are shown in Table 1.

Briefly, after washing with PBS the specimens were treated with 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 min to block the activity of endogenous peroxidase. Specimens were then washed with PBS and stained with peroxidase labeled lectins (10µg/ml of PNA and 15µg/ml of WGA) for 2-3 h at 37°C. Positive reactions were visualised with 0.02% 3, 3'-diaminobenzidine hydrochloride (DAB, Dojindo, Tokyo, Japan) and 0.003% H<sub>2</sub>O<sub>2</sub> in Tris buffer.

## RESULTS

The spermatozoa of the lesser mouse deer were small in size and measured 36.52 ± 5.6 µm in total length. The head of spermatozoa was relatively round, with 5.55 ± 0.8 µm in length and 4.77 ± 0.5 µm in width (Figure 1, Table 2). In the specimens stained by lectin histochemistry, lectin PNA and lectin WGA was positive in the acrosomal membrane (Figure 2a), and in the membrane of the anterior acrosomal region and tail (Figure 2b), respectively.

**Table 1: Dilution and specificity of the lectin used**

Taxonomic name	Acronym	Concentration	Specificity
<i>Arachis hypogea</i>	PNA	10 µg/ml	Galactose β1-3, D-N-acetylgalactosamine
<i>Triticum vulgare</i>	WGA	15 µg/ml	D-N-acetylglucosamine, sialic acid

**Table 2: Measurement of the spermatozoa of the lesser mouse deer**

	Size (µm)
Total length	36.52 ± 5.6
Head	
Length	5.55 ± 0.8
Width	4.77 ± 0.5




Figure 1: Photomicrograph of the spermatozoon of the lesser mouse deer. a. Head, b. Neck, c. Body, d. Principal tail, e. End tail. SEM. 10kV.




Figure 2: Distribution of lectin-bindings in the membrane of the spermatozoa of the lesser mouse deer. PNA (A) is positive in the acrosomal membrane (arrow), while WGA (B) is positive in the membrane of anterior acrosomal region (long arrow) and tail (short arrows). Peroxidase conjugated lectin. Scale bars: 5 $\mu$ m

## DISCUSSION

The shape of the head of the spermatozoon in the lesser mouse deer is relatively more round when compared to the other ruminants (Mann, 1964; Bustos-Obregos and Fléchon, 1975; Salisbury and Van Demark 1985; Evans and Maxwell, 1987). Further, the spermatozoon of the lesser mouse deer was considerably smaller than the values reported in ram (Bustos-Obregon and Fléchon, 1975; Evans and Maxwell, 1987; Rizal, 2005), bull (Bustos-Obregos and Fléchon, 1975; Salisbury and Van Demark 1985) and boar (Bustos-Obregos and Fléchon, 1975).

Lectins are commonly used as a marker to assess carbohydrate composition in the sperm membrane (Bearer and Friend, 1990), sperm acrosomal status and acrosome reaction (Cheng *et al.*, 1996). Sperm with the intact acrosome will undergo acrosome reaction which is essential for sperm penetration and fusion and damages of the acrosome can decrease the fertility rate of the

spermatozoa (Nolan and Hammerstedt, 1997; Hammadah *et al.*, 2001; Ramalho-Santos *et al.*, 2002; Yoshinaga and Toshimori, 2003). This study demonstrated the presence of PNA bindings in the acrosomal membrane and WGA bindings on the membrane of anterior acrosomal region and tail of the spermatozoa of the lesser mouse deer. Lectin PNA indicated carbohydrates with galactose  $\beta$ 1-3 and D-N-acetylgalactosamine sugar residues. These glycoproteins are closely related to various processes in the cell to cell recognition and adhesion of gametes cells, sperm and zona pellucida binding or binding of sperm and epithelia of the female reproductive organ (Gilbert, 1988; Spicer and Schulte, 1992; Thall *et al.* 1995; Clark *et al.*, 1996; Zara and Naz, 1998).

Lectin WGA indicated carbohydrates with D-N acetylglucosamine and sialic acid sugar residues. D-N acetylglucosamine residues function in the binding and fusion of sperm and oocytes (Gougoulidis *et al.*, 1999). Furthermore the N-acetylglucosaminidase (NAG), a

glycosidase-recognising N-acetylglucosamine terminal residue, is also involved in the primary binding to zona pellucida and penetration and polyspermy block (Zitta *et al.*, 2006). Sialic acid residues are secreted by the epithelium of the epididymis as terminal sugar of sialoglycoproteins. These glycoproteins bind to the sperm surface during epididymal transit (Lassalle and Testart, 1994) and cover the spermatozoa surface and protect them from phagocytic activity by leukocyte and vaginal epithelial cells (Harayama *et al.*, 1996; Lassalle and Testart, 1994). Sialic acid may also participate in the binding of spermatozoa and zona pellucida in human (Ozgur *et al.*, 1998).

The PNA bindings in the spermatozoon of the lesser mouse deer are similar to those reported in ram (Flesch *et al.*, 1998), *Bufo arenarum* (Martinez and Cabada 1996), bull (Cross and Watson, 1994), human (Mortimer *et al.*, 1987) and stallion (Cheng *et al.*, 1996) while the WGA bindings are generally similar to those reported in human sperm (Lassalle and Testart, 1994; Gabriel *et al.*, 1995), and boar (Jiménez *et al.*, 2003). This suggests that carbohydrates with galactose  $\beta$ 1-3, D-N-acetyl-galactosamine, D-N-acetylglucosamine and sialic acid sugar residues are involved and may have significant roles in the spermatozoa of the lesser mouse deer.

#### ACKNOWLEDGEMENTS

This work was supported by *Hibah Bersaing* Research Grant (No. 010/P4T/DPPM/PHBXII/III/2004) to S.A from the Directorate General of Higher Education, Ministry of National Education, Indonesia.

#### REFERENCES

- Axner, E. and Linde-Forsbeg, C. (2002). Semen collection and assessment, and artificial insemination in the cat. Recent Advances in Small Animal Reproduction. [www.ivis.org](http://www.ivis.org) (assessed January 2006)
- Bearer, E.L. and Friend, D.S. (1990). Morphology of mammalian sperm membranes during differentiation, maturation, and capacitation. *Journal of Electron Microscopy Technique* **16**: 281-297
- Bustos-Obregon, E. and Fléchon, J.-E. (1975). Comparative scanning electron microscope study of boar, bull and ram spermatozoa. *Cell and Tissue Research* **161**: 329-341
- Cheng, F.P., Fazeli, A., Voorhout, W.F., Marks, A., Bevers, M.M. and Colenbrander, B. (1996). Use of peanut agglutinin to assess the acrosomal status and the zona pellucida-induced acrosome reaction in stallion spermatozoa. *Journal of Andrology* **17**:674-678.
- Clark, G.F., Oehninger, S. and Seppala, M. (1996). Role of glycoconjugates in cellular communication in the human reproductive system. *Molecular Human Reproduction* **2**: 513-517
- Cross, N.L. and Watson, S.K. (1994). Assessing acrosomal status of bovine sperm using fluoresceinated lectin. *Theriogenology* **42**: 88-98.
- Evans, G. and Maxwell, W.M.C. (1987). *Salamon's Artificial Insemination of Sheep and Goats*. Sydney: Butterworth Scientific. pp. 22-30
- Flesch, F.M., Voorhout, W.F., Colenbrander, B., van Golde, L.M.G. and Gadella, B.M. (1998). Use of lectin to characterize plasma membrane preparations from boar spermatozoa: a novel technique for monitoring membrane purity and quantity. *Biology of Reproduction* **59**: 1530-1539.
- Gabriel, L.K., Franken, D.R., Van Der Horst, G. and Kruger, T.F. (1995) Fluorescent isothiocyanate conjugated-wheat germ agglutinin staining of human spermatozoa and fertilization *in vitro*. *Fertility and Sterility* **63**: 894-901
- Gilbert, S.F. (1988). *Developmental Biology*. 2<sup>nd</sup> Ed. Massachusetts: Sinauer Associates, Inc. pp. 37-47
- Gougoulidis, T., Trounson, A. and Dowsing, A. (1999). Inhibition of bovine sperm-oocyte fusion by the carbohydrate GalNAc. *Molecular Reproduction and Development* **54**: 179-85.
- Hammadeh, M.E., Georg, T., Rosenbaum, P. and Schmidt, W. (2001). Association between freezing agent and acrosome damage of human spermatozoa from subnormal and normal semen. *Andrologia* **33**:331-336.
- Harayama, H., Kato, S. and Hammerstedt, R.H. (1996). Electrophoretic characterization of boar epididymal antiagglutinin. *Biology of Reproduction* **55**: 325-332.
- Haron, A.W., Yong, M. and Zainudin, Z.Z. (1999). Evaluation of semen collected by electro ejaculation from captive lesser mouse deer Malay chevrotain (*Tragulus javanicus*). *Journal of Zoo and Wildlife Medicine* **31**:164-167.
- Inoue, T. and Osatake, H. (1988). A new drying method of biological specimens for scanning electron microscopy: the *t*-butyl alcohol freeze-drying method. *Archives of Histology and Cytology* **51**:53-59.

- Jiménez, I., Gonzalez-Marquez, H., Ortiz, R., Herrera, J.A., Garcia, A., Betancourt, M. and Fierro, R. (2003). Changes in the distribution of lectin receptors during capacitation and acrosome reaction in boar spermatozoa. *Theriogenology* **59**:1171-1180.
- Jumaliah, N. (1999). *Pola Perilaku, Estimasi Kuantitatif Konsumsi dan Daya Cerna Kancil (Tragulus javanicus) Terhadap Pakan di Kebun Binatang Ragunan Jakarta [Thesis]: Postgraduate Program, Bogor Agricultural University, Bogor, Indonesia.*
- Lassalle, B. and Testart, J. (1994). Human zona pellucida recognition associated with removal of sialic acid from human sperm surface. *Journal of Reproduction and Fertility* **101**:703-711.
- Lekagul, B. and McNeely, J.A. (1977). Mammals of Thailand. The Association for the Conservation of Wild Life. Bangkok: Kurusapha, Ladprow Press. pp. 665-671.
- Mann, T. 1964. The Biochemistry of Semen. London: Methuen & Co. Ltd. pp.4-15.
- Martinez, M.L. and Cabada, M.O. (1996). Assessment of the acrosome reaction in *Bufo arenarum* spermatozoa by immunostaining: comparison with other methods. *Zygote* **4**: 181-190.
- Mortimer, D., Curtis, E.F. and Miller, R.G. (1987). Specific labeling by peanut agglutinin of the outer acrosomal membrane of the human spermatozoon. *Journal of Reproduction and Fertility* **81**:127-135.
- Murakami, T. (1974). A revised tannin-osmium method for non-coated scanning electron microscope specimens. *Archivum Histologicum Japonicum* **36**: 189-193
- Nolan, J.P. and Hammerstedt, R.H. (1997). Regulation of membrane stability and the acrosome reaction in mammalian sperm. *FASEB Journal* **11**:670-682
- Ozgun, K., Patankar, M.S., Oehninger, S. and Clark, G.F. (1998). Direct evidence for the involvement of carbohydrate sequences in human sperm-zona pellucida binding. *Molecular Human Reproduction* **4**: 318-324.
- Prasetyaningtyas, W.E., Setiadi, M.A., Fahrudin, M., Haron, A.W. and Agungpriyono, S. (2006). Karakteristik dan komposisi semen kancil (*Tragulus javanicus*) yang dikoleksi dengan elektroejakulator. *Indonesia Journal of Anatomy* **1**:30-37
- Ramalho-Santos, J., Schatten, G. and Moreno, R.D. (2002). Control of membrane fusion during spermiogenesis and the acrosome reaction. *Biology of Reproduction* **67**: 1043-1051.
- Rizal, M. (2005). *Fertilitas Spermatozoa Ejakulat dan Epididimis Domba Garut Hasil Kriopreservasi Menggunakan Modifikasi Pengencer Tris dengan Berbagai Krioprotektan dan Antioksidan. PhD thesis. Postgraduate Program, Bogor Agricultural University, Bogor, Indonesia.*
- Salisbury, G.W. and Van Demark, N.L. (1985). Physiology of Reproduction and Artificial Insemination of Cattle. Yogyakarta: Gadjah Mada University Press. pp. 314-343.
- Sastradipradja, D. (1978). The lesser mousedeer (*Tragulus javanicus*) as a model animal for ruminant studies. In: *fondation marcel merieux (eds). Congres international sur l'animal de laboratoire au service de l'homme, ecole nationale veterinaire de lyon.* September 1978. pp 565-573
- Spicer, S.S. and Schulte, B.A. (1992). Diversity of cell glycoconjugates shown histochemically : a perspective. *Journal of Histochemistry and Cytochemistry* **40**: 1-38.
- Thall, A.D., Murphy, H.S. and Lowe, J.B. (1995). Oocytes Gal $\alpha$ 1-3Gal epitopes implicated in sperm adhesion to the zona pellucida glycoprotein ZP3 are not required for fertilization in the mouse. *Journal of Biological Chemistry* **270**: 21437-21440
- Tulsiani, D.R. (2000). Carbohydrates mediate sperm-ovum adhesion and triggering of the acrosome reaction. *Asian Journal of Andrology* **2**: 87-97
- Yoshinaga, K. and Toshimori, K. (2003). Organization and modifications of sperm acrosomal molecules during spermatogenesis and epididymal maturation. *Microscopy Research and Technique* **61**: 39-45
- Zara, J. and Naz, R.K. (1998). The role of carbohydrates in mammalian sperm-egg interactions : How important are carbohydrate epitopes? *Frontiers in Bioscience* **3**: 1028-1038.
- Zitta, K., Wertheimer, E. V. and Miranda, P.V. (2006). Sperm N-acetylglucosaminidase is involved in primary binding to the zona pellucida. *Molecular Human Reproduction* **12**: 557-563