



**UNIVERSITI PUTRA MALAYSIA**

**DRYING AND SOLID-LIQUID EXTRACTION OF  
HYDROXYCHAVICOL AND EUGENOL FROM BETEL LEAVES  
(*PIPER BETLE L.*)**

**PIN KAR YONG**

**FK 2009 34**



**DRYING AND SOLID-LIQUID EXTRACTION OF HYDROXYCHAVICOL  
AND EUGENOL FROM BETEL LEAVES (*PIPER BETLE* L.)**

**By**

**PIN KAR YONG**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of Requirements for the Degree of Doctor of Philosophy**

**JUNE 2009**



*This thesis is specially dedicated to  
my parents  
and  
my late aunty who passed away on 9<sup>th</sup> May 2008*



Abstract of thesis presented to the Senate of Universiti of Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**DRYING AND SOLID-LIQUID EXTRACTION OF HYDROXYCHAVICOL AND EUGENOL FROM BETEL LEAVES (*PIPER BETLE L.*)**

By

**PIN KAR YONG**

**JUNE 2009**

**Chairman: Associate Professor Luqman Chuah Abdullah, PhD**

**Faculty: Engineering**

Betel (*Piper betle L.*) is one of the invaluable medicinal plants originated from Malaysia. Its leaves have been used traditionally for various medication purposes. Scientific research on the leaf of this plant reveals that it possesses many beneficial bioactivities and its extract from betel leaves has a great potential to be used in developing commercial products. However, there is a lack of research on the processing aspects to produce its bioactive extract.

This research studied three key processes including drying, solid-liquid extraction, and freeze drying which are involved in processing of bioactive extract from betel leaves. Different experiments were designed and carried out to look into the effects of various operating parameters on the qualitative and quantitative aspects of betel leaves extract. Hydroxychavicol (HC) and eugenol (EU) were selected as the quality indicators of the product because these two compounds were reported to play an important role in the bioactivities of betel leaves including antioxidant, anti-inflammatory, and anticarcinogenic and antibacterial.



The effect of drying temperature on the quality of betel leaves and drying kinetics were studied in order to determine the optimum drying temperature. Changes in the concentration of HC and EU reveal that the optimum temperature for drying of betel leaves was 70°C because degradation of HC and EU was observed above this temperature. Logarithmic model was found to be the most suitable model among the selected thin layer models in predicting the process.

Water was the most suitable solvent for extracting betel leaves compared to ethanol, ethyl acetate, and hexane. This was because it gave highest yield and the extract from water indicated high antioxidant and anti-inflammatory activities in which the activities were related to HC and EU. The optimum extraction temperature was determined as 60°C to avoid degradation of EU. The ratio of water to solid of 30:1 (ml:g) was found to be optimum based on analysis of Response Surface Methodology (RSM). Extraction kinetics of betel leaves reveals that the optimum extraction time is one hour. A new model named equilibrium driven solid-liquid extraction (EDSLE) model was developed and successfully applied in describing the process.

The study of freeze drying process of betel leaves extract was conducted in two sections namely freezing and drying. The freezing kinetics data shows that the freezing point of betel leaves extract with 20%SC was about -4°C. Prediction of freezing kinetics and freezing time was carried out successfully with numerical model. The results of drying kinetics of betel leaves extract show that the increase of drying temperature increased the drying rate. Midilli *et la.* model was found to be the most effective one among the selected models for modeling of the process.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doctor of Philosophy

**PENGERINGAN DAN PENGEKSTRAKAN PEPEJAL-CECAIR BAGI  
HYDROXYCHAVICOL DAN EUGENOL DARIPADA DAUN SIRIH (*PIPER  
BETLE L.*)**

Oleh

**PIN KAR YONG**

**JUN 2009**

**Pengerusi: Profesor Madya Luqman Chuah Abdullah, PhD**

**Fakulti: Kejuruteraan**

Sirih (*Piper betle L.*) merupakan salah satu tumbuhan ubatan bernilai yang berasal dari Malaysia. Daun sirih telah digunakan secara tradisional untuk pelbagai tujuan perubatan. Penyelidikan ke atas daun pokok ini menunjukkan ianya mempunyai banyak bioaktiviti yang bermanfaat dan ekstrak daripada daun sirih mempunyai potensi yang tinggi untuk digunakan dalam pembangunan produk. Tetapi, penyelidikan dalam aspek pemprosesan untuk menghasilkan ekstrak yang bioaktif ini adalah tidak mencukupi.

Penyelidikan ini merangkumi tiga proses utama termasuklah pengeringan, pengekstrakan pepejal-cecair, dan pengeringan beku yang terlibat dalam pemprosesan bioaktif ekstrak daripada daun sirih. Eksperimen berbeza telah direka dan dijalankan untuk mengkaji kesan pelbagai parameter terhadap aspek kualiti dan kuantiti bagi ekstrak daun sirih. *Hydroxychavicol* (HC) and *eugenol* (EU) dipilih sebagai petunjuk kualiti bagi produk kerana kedua-dua sebatian ini dilaporkan memainkan peranan yang penting dalam bioaktiviti bagi daun sirih termasuklah anti-oksidan, anti-radang, anti-kanser dan anti-bakteria.

Kesan suhu pengeringan terhadap kualiti daun sirih dan kinetik pengeringan dikaji untuk menentukan suhu pengeringan yang optima. Perubahan kepekatan HC dan EU menunjukkan suhu pengeringan optimum adalah 70°C kerana degradasi HC dan EU diperhatikan apabila melebihi suhu ini. Model *Logarithmic* adalah model yang paling sesuai antara model-model yang digunakan untuk peramalan proses itu.

Air merupakan pelarut yang paling sesuai untuk mengekstrak daun sirih berbanding dengan etanol, etil acetate, dan hexana. Ini kerana ia memberikan hasil yang paling tinggi dan ekstrak daripada air menunjukkan aktiviti anti-oksidan dan anti-radang yang tinggi dimana aktiviti itu berkaitan dengan HC dan EU. Suhu pengekstrakan yang optimum ditentukan sebagai 60°C untuk mengelakkan degradasi EU. Nisbah air kepada pepejal yang bernilai 30: 1 (ml:g) didapati optimum berdasarkan analisa *Response Surface Methodology* (RSM). Kajian kinetik pengekstrakan daun sirih menunjukkan tempoh pengekstrakan yang optimum adalah satu jam. Satu model baru dinamakan model *equilibrium driven solid-liquid extraction* (EDSLE) telah dibangunkan dan digunakan dengan berjaya untuk menerangkan proses itu.

Pengkajian pengeringan beku dijalankan dalam dua bahagian termasuklah pembekuan dan pengeringan. Kinetik pembekuan menunjukkan takat beku bagi ekstrak daun sirih dengan kandungan pepejal 20% adalah -4°C. Peramalan bagi kinetik pembekuan dan masa pembekuan dilakukan dengan berjaya dengan model *numerical*. Keputusan kinetik pengeringan menunjukkan penambahan suhu pengeringan dapat meningkatkan kadar pengeringan. Model *Midilli et al.* adalah model yang paling berkesan antara model-model yang digunakan untuk premodelan proses itu.

## ACKNOWLEDGEMENT

First of all, I would like to express my gratitude to my advisor and chairman of the supervisory committee, Associate Professor Dr. Luqman Chuah Abdullah, for his advice and guidance in making this research successful. I would also like to thank my supervisory committee members, Dr. Rasadah Mat Ali, Associate Professor Dr. Thomas Choong Shean Yaw, and Associate Professor Dr. Law Chung Lim for giving useful ideas and comments throughout the project.

I would also like to extend my thanks to my colleagues from Herbal Technology Center and Medicinal Plants Division of Forest Research Institute Malaysia (FRIM) especially Tn. Haji Abdull Rashih Ahmad for their technical assistance and kind knowledge-sharing. I would also like to thank FRIM for sponsoring my study.

Last but not least, I express my deepest gratitude to my parents and brother for their love, encouragement and support. Special thanks to my beloved girlfriend, Miss Teoh Hoe Peng for standing by me through thin and thick.





I certify that an Examination Committee has met on 11<sup>th</sup> June 2009 to conduct the final examination of Pin Kar Yong on his thesis entitled “Drying and Solid-liquid Extraction of Hydroxychavicol and Eugenol from Betel Leaves (*Piper betle* L.)” in accordance with Universities and Universities Colleges Act 1971 and the Constitution of Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Examination Committee were as follows:

**Azni bin Idris, PhD**

Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Chairman)

**Mohd. Ali Hassan, PhD**

Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Internal Examiner)

**Norhafizah bt. Hj. Abdullah, PhD**

Associate Professor  
Faculty of Engineering  
Universiti Putra Malaysia  
(Internal Examiner)

**Abdul Latif Ahmad, PhD**

Professor  
School of Chemical Engineering  
Universiti Sains Malaysia  
(External Examiner)

---

**BUJANG KIM HUAT, PhD**

Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 27 August 2009



This thesis submitted to the Senate of University Putra Malaysia and has been accepted as fulfillment of the requirement of the degree of Doctor of Philosophy. The members of the Supervisor Committee were as follows:

**Luqman Chuah Abdullah, PhD**

Associate Professor  
Faculty of Engineering  
University Putra Malaysia  
(Chairman)

**Thomas Choong Shean Yaw, PhD**

Associate Professor  
Faculty of Engineering  
University Putra Malaysia  
(Member)

**Rasadah Mat Ali, PhD**

Senior Director  
Medicinal Plants Division  
Forest Research Institute Malaysia  
(Member)

**Law Chung Lim, PhD**

Associate Professor  
Faculty of Engineering  
University Nottingham (Malaysia Campus)  
(Member)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 11 September 2009



## **DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

---

**PIN KAR YONG**

Date: 5 August 2009

## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	v
<b>ACKNOWLEDGEMENT</b>	vii
<b>APPROVAL</b>	viii
<b>DECLARATION</b>	x
<b>LIST OF TABLES</b>	xv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xviii

### CHAPTER

<b>1</b>	<b>INTRODUCTION</b>	
1.1	Herbal Industry in Malaysia	1
1.2	Herbal Processing	3
1.3	Betel ( <i>Piper betle</i> L.)	5
1.4	Problem Statement	6
1.5	Objectives	8
1.6	Scope of Research	8
<b>2</b>	<b>LITERATURE REVIEW</b>	
2.1	Betel ( <i>Piper betle</i> Linn)	10
2.1.1	Traditional uses of betel leaves	10
2.1.2	Beneficial bioactivities of betel leaves	11
2.1.3	Hydroxychavicol & Eugenol	14
2.2	Solid – Liquid Extraction (SLE)	16
2.2.1	Extraction temperature	18
2.2.2	Type of solvent	19
2.2.3	Amount of solvent	20
2.2.4	Process duration	21
2.2.5	Particle size of solid	22
2.2.6	Mathematical models for SLE	23
2.2.7	Response surface methodology (RSM)	27
2.3	Drying	31
2.3.1	Key parameter affecting drying process	34
2.3.2	Mathematical models for drying	36
2.4	Freeze Drying	40
2.4.1	Freezing	40
2.4.2	Sublimation in primary drying	44
2.4.3	Desorption in secondary drying	45
2.4.4	Modeling of freeze drying process	46
2.5	Chapter summary	48



<b>3</b>	<b>METHODOLOGY</b>	
3.1	Introduction	50
3.2	Raw material – betel leaves	51
3.3	Drying of betel leaves	
3.3.1	Experiments for effect of drying temperature on product quality	51
3.3.2	High performance liquid chromatography (HPLC)	52
3.3.3	Experiment for effects of drying temperature on drying kinetics	53
3.3.4	Mathematical modeling for drying kinetics	54
3.4	Quantitative and qualitative effects of solvent on solid-liquid extraction (SLE) of betel leaves	
3.4.1	Experiments for effect of solvent on extraction yield	55
3.4.2	Experiments for effect of solvent on quality of extract	56
3.5	Optimization of solid-liquid extraction (SLE) of betel leaves	
3.5.1	Experiments for effect of extraction temperature on extract quality	63
3.5.2	Experiments for effects of extraction temperature and ratio of solvent to solid on the yield of extraction with response surface methodology (RSM)	64
3.6	Mathematical modeling of SLE of betel leaves	
3.6.1	Physical properties of dried betel leaves	67
3.6.2	Extraction kinetics	67
3.6.3	Mathematical modeling for SLE	68
3.7	Freeze drying of betel leaves extract	
3.7.1	Materials	73
3.7.2	Freezing experiments	74
3.7.3	Experiments for effects of drying temperature and extract concentration on freeze drying kinetics	79
3.8	Statistical analysis	81
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	
4.1	Drying of betel leaves	
4.1.1	Effect of drying temperature on product quality	82
4.1.2	Drying kinetics and mathematical modeling	85
4.2	Quantitative and qualitative effects of solvent on solid-liquid extraction (SLE) of betel leaves	
4.2.1	Effect of solvent on extraction yield and phytochemicals content of extract	89
4.2.2	Antioxidant activity of solvent extracts	95
4.2.3	Anti-inflammatory activity of solvent extracts	96

4.3	Optimization of SLE of betel leaves	
4.3.1	Effect of extraction temperature on quality of extract	98
4.3.2	Effects of extraction temperature and ratio of solvent to solid on the yield of extraction	100
4.4	Mathematical modeling of SLE of betel leaves	
4.4.1	Extraction kinetics and mathematical modeling	104
4.5	Freeze drying of betel leaves extract	
4.5.1	Freezing kinetics	108
4.5.2	Effects of drying temperature and extract concentration on freeze drying kinetics	111
<b>5</b>	<b>CONCLUSION</b>	
5.1	Conclusion	117
5.2	Future studies	118
	<b>REFERENCES</b>	119
	<b>APPENDICES</b>	137
	<b>BIODATA OF STUDENT</b>	170
	<b>LIST OF PUBLICATIONS</b>	171

## LIST OF TABLES

Table		Page
1.1	Popular medicinal plants in Malaysia	2
2.1	The time for achieving equilibrium for various extraction systems	22
2.2	Thin layer models commonly used mathematical modeling of drying	38
2.3	Thin layer models selected for different drying process	39
2.4	Freezing points of different juice	41
3.1	The mobile phase changes in HPLC analysis	53
3.2	Thin layer models used in mathematical modeling	55
3.3	Experiments setup for studying the effect of extraction temperature	64
3.4	Central-composite experimental design of the independent variables and experimental data for the response variable	66
3.5	Data recording interval for different freezing temperatures.	74
4.1	The values of correlation coefficient, RMSE, and drying constants for the models.	88
4.2	Results of the antioxidant activity of solvent extracts	95
4.3	Results of the anti-inflammatory activity of solvent extracts	96
4.4	Analysis of variance for the independent variables	101
4.5	The results from EDSLE model and DDSLE model for different extraction temperatures.	106
4.6	Actual and numerical freezing times for the studied freezing temperatures	111
4.7	The values of correlation coefficient, RMSE, and drying constants for drying temperature of $-10^{\circ}\text{C}$	114
4.8	The values of correlation coefficient, RMSE, and drying constants for drying temperature of $-20^{\circ}\text{C}$	115

## LIST OF FIGURES

Figure		Page
1.1.	The processes involved in production of bioactive extract	4
1.2	Well-grown betel plant with heart-shaped leaves	5
2.1	Chemical structure of (a) Hydroxychavicol and (b) Eugenol.	15
2.2	Primary and secondary metabolites produced by plant.	17
2.3	Typical drying curves using moisture content vs. time and drying rate vs. moisture content	33
2.4	Phase diagram for water	45
3.1	Overview of the methodology	50
3.2	Schematic diagram of extraction process and solid	69
3.3	Heat transfer out of the extract during freezing process	75
3.4	Representation of grid system used in solving the problem	77
4.1	HPLC Chromatograms of (a) Standard; (b) Fresh leaves; (c) Drying temperature = 40°C; (d) Drying temperature = 50°C; (e) Drying temperature = 60°C; (f) Drying temperature = 70°C; (g) Drying temperature = 80°C.	83
4.2	Recovery of HC and EU in the fresh leaves and dried leaves from different drying temperatures	84
4.3	The drying rate for different drying temperatures	86
4.4	Comparison between the experimental data and the selected mathematical model for different temperatures	87
4.5	Yield of extraction from different solvent	91
4.6	HPLC Chromatograms of (a) H <sub>2</sub> O extract; (b) EtOH extract; (c) EA extract; (d) Hex extract	93
4.7	Recovery of HC and EU in the extracts from different solvent	94
4.8	HPLC Chromatograms of (a) Extraction temperature = 25°C; (b) Extraction temperature = 40°C; (c) Extraction temperature = 50°C; (d) Extraction temperature = 60°C; (e) Extraction temperature = 70°C; (f) Extraction temperature =	98



80°C.

4.9	Recovery of hydroxychavicol and eugenol in betel leaves extract from different extraction temperatures.	99
4.10	Relationship between the experimental and predicted data	100
4.11	Response surface graph showing the effect of temperature and ratio of solvent to solid on the yield	102
4.12	Comparison between experimental data and EDSLE model for different temperatures	105
4.13	Comparison of rate of extraction for experiments conducted at temperatures of 25 and 60°C	105
4.14	Comparison between experimental data and DDSLE model for different temperatures	107
4.15	Comparison of experimental results and numerical results for freezing temperature of -20°C.	110
4.16	Comparison of experimental results and numerical results for freezing temperature of -25°C.	110
4.17	Comparison of experimental results and numerical results for freezing temperature of -30°C.	111
4.18	Drying curve for betel leaves extract with different concentration for drying temperature of -10 and -20°C	113

## LIST OF ABBREVIATIONS

$\sigma$	Coefficient in $D(C_s)$ relation
$\rho$	Density of the extract ( $\text{kgm}^{-3}$ )
$\omega$	Ice fraction (kg ice/ kg water)
$\varepsilon$	Internal porosity ( $\text{m}^3\text{m}^{-3}$ )
$\alpha$	Shape factor
$\sigma$	Specific surface area of the solid ( $\text{m}^2\text{kg}^{-1}$ )
$\phi$	Water content (kg water/ kg material)
$\rho_d$	Density of dry solid ( $\text{kgm}^{-3}$ )
$\rho_f$	Density of frozen product ( $\text{kgm}^{-3}$ )
$\rho_i$	Density of ice ( $\text{kgm}^{-3}$ )
$\lambda_n$	Eigenvalues
$\rho_p$	Density of the solids ( $\text{kgm}^{-3}$ )
$\Delta t$	Time step (s)
$\rho_w$	Density of water ( $\text{kgm}^{-3}$ )
$\Delta x$	Grid size (m)
$a$	Thickness of slab (m)
$A$	Total surface area of the solids ( $\text{m}^2$ )
$A'$	Fitting parameter
$b_o$	Regression coefficient of intercept term
$b_1$	Linear regression coefficient
$b_2$	Linear regression coefficient
$b_{11}$	Squared regression coefficient
$b_{22}$	Squared regression coefficient

$b_{12}$	Interaction regression coefficient.
$B'$	Fitting parameter
$Bi$	Biot number
$C'$	Fitting parameter
$C_e$	Equilibrium solute concentration in liquid phase ( $\text{gcm}^{-3}$ )
$\text{CH}_3\text{CN}$	Acetonitrile
$C_L$	Solute concentration in liquid phase ( $\text{gcm}^{-3}$ )
$C_{L,exp}$	Solute concentration in liquid phase from experiment result ( $\text{gcm}^{-3}$ )
$C_{L,pre}$	Predicted Solute concentration in liquid phase ( $\text{gcm}^{-3}$ )
$C_p$	Specific heat capacity of the extract ( $\text{Jkg}^{-1}\text{C}^{-1}$ )
$C_{p,fr}$	Specific heat capacity of frozen extract ( $\text{Jkg}^{-1}\text{C}^{-1}$ )
$C_{p,un}$	Specific heat capacity of unfrozen extract ( $\text{Jkg}^{-1}\text{C}^{-1}$ )
$C_S$	Solute concentration in solid phase ( $\text{gcm}^{-3}$ )
$d$	Thickness of product (m)
$D$	Intraparticles diffusion coefficient ( $\text{m}^2\text{s}^{-1}$ )
DDSLE	Diffusion driven solid-liquid extraction
$D_e$	Effective diffusivity coefficient ( $\text{m}^2\text{s}^{-1}$ )
$D_o$	Preexponential factor of Arrhenius equation ( $\text{m}^2\text{s}^{-1}$ )
$E_0$	Parameter of the Elovich equation
$E_1$	Parameter of the Elovich equation
$E_a$	Energy of activation ( $\text{Jmol}^{-1}$ )
EDSLE	Equilibrium driven solid-liquid extraction
EU	Eugenol
FDM	Finite difference method
FEM	Finite element method

FRIM	Forest Research Institute Malaysia
$h$	Heat transfer coefficient ( $\text{Wm}^{-2}\text{K}^{-1}$ )
$\text{H}_3\text{PO}_4$	Orthophosphoric Acid
HC	Hydroxychavicol
HPLC	High Performance Liquid Chromatography
$i,j$	Coordinate in the grid system
$K$	Mass transfer coefficient ( $\text{cms}^{-1}$ )
$k$	Thermal conductivity of the extract ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$K'$	Second-order extraction rate constant
$K_1$	Peleg rate constant
$K_2$	Peleg capacity constant
$k_d$	Thermal conductivity of dry solid ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$k_f$	Thermal conductivity of frozen product ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$k_{fr}$	Thermal conductivity of frozen extract ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$k_i$	Thermal conductivity of ice ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$k_{un}$	Thermal conductivity of unfrozen extract ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$k_w$	Thermal conductivity of water ( $\text{Wm}^{-1}\text{C}^{-1}$ )
$L$	Latent heat of freezing (kJ/kg)
$M$	Weight of total solid particles (g)
$M_{db}$	Moisture content (% dry basis)
$M_e$	Equilibrium moisture content (% wet basis)
$M_o$	Initial moisture content (% wet basis)
$M_p$	Weight of the solid particle (g)
$MR$	Dimensionless moisture ratio
$MR_{exp}$	Dimensionless moisture ratio from experiment result

$MR_{pre}$	Predicted dimensionless moisture ratio
$M_{wb}$	Moisture content (% wet basis)
$P'$	Plank's model constant
PAF	Platelet activating factor
$Pk$	Plank number
$r$	Distance form center axis (m)
$R$	Universal gas constant
$R'$	Plank's model constant
$R^2$	Correlation coefficient
RMSE	Root mean square error
RO	Reverse osmosis
$R_p$	Radius of solid particle (m)
$R_{ss}$	Ratio of solvent to solid ( $\text{mlg}^{-1}$ )
$S$	Weight of solid (kg)
SC	Solid content (%)
SLE	Solid – liquid extraction
$Ste$	Stefan number
$T$	Temperature (K)
$t$	time (s)
$T_f$	Freezing temperature (K)
$T_{a1}$	Temperature of drying chamber (K)
$T_{a2}$	Temperature of thermal shelf (K)
$T_{cr}$	Freezing point (K)
$T_{exp}$	Temperature from experiment result (K)
$T_{fin}$	Final product temperature (K)

$T_{fm}$	Mean freezing temperature (K)
$T_{in}$	Initial product temperature (K)
TPA	Tetradecanoylphorbol acetate
$T_{pre}$	Predicted temperature (K)
$V$	Volume of product (cm <sup>3</sup> )
$V_e$	Volume of aqueous extract used for freeze-drying (cm <sup>3</sup> )
$V_L$	Volume of solvent (cm <sup>3</sup> )
$V_p$	Volume of the solid (cm <sup>3</sup> )
$V_t$	Volume of aqueous extract obtained after filtration (cm <sup>3</sup> )
$W$	Weight of moisture within product (kg)
$W_d$	Weight of dried extract (g)
$W_s$	Weight of solid (g)
$x$	Distance form center axis (m)
X	Coded independent variable
$x'$	Independent variable
Y	Response of RSM

## CHAPTER 1

### INTRODUCTION

#### 1.1 Herbal Industry in Malaysia

The development of the herbal industry in Malaysia is gaining its momentum due to its increasing economic demands. The current value of the local herbal market is RM 5 billion with an annual growth of 15 to 20 percent (Bernama, 2008). It is expected that the local market will achieve a staggering RM 8 billion by 2010. The global market is currently estimated to be US\$ 200 billion with a growth rate between 10 and 20 percent per year and expected to hit US \$5 trillion (RM 19 trillion) by 2050 (Bernama, 2007). This trend is attributed to the hiking awareness on the safety and environmental aspects of consumable products. Consumers are becoming more interested in the products from natural sources which are believed to cause lesser side effects.

Most of herbal products are produced by adding a certain amount of bioactive extract from medicinal plants. Bioactive extract is referred as extract that possesses therapeutic activities (WHO, 2000). There is a wide range of naturally-occurred phytochemicals found in bioactive extracts. These phytochemicals provide balanced and synergic effects which contribute to the beneficial properties of the extracts. This is the crucial part that is lacking in the single-compound synthetic drugs.

Nowadays, bioactive extracts are used in different types of products including herbal medicines, nutritional supplement, functional food and beverage, cosmetic products, health care products and so on. These products offer a more affordable alternative to modern pharmaceuticals. The number of herbal manufacturers in Malaysia is



growing like mushroom after the rain due to the encouraging demand. There are many successful herbal product companies such as HPA Industries Sdn. Bhd., Felda Herbal Corporation, Sendayu Tinggi Corporation (M) Sdn. Bhd., Power Root (M) Sdn. Bhd., Nova Laboratories Sdn. Bhd. and so on. The Malaysian herbal manufacturers have commercialized several well-known herbal products formulated from various medicinal plants such as *Tongkat Ali (Eurycoma, longifolia)*, *Kacip Fatimah (Labisia pumila)* and *Misai Kucing (Orthosiphon stiminues)* to meet the local demands. Table 1.1 listed some of the popular medicinal plants and their herbal products that are available in the local market. These products are widely accepted by local consumers.

**Table 1.1. Popular medicinal plants in Malaysia**

Local Name	Scientific Name	Product
Tongkat ali	<i>Eurycoma longifolia</i>	<ul style="list-style-type: none"> <li>• Energy drinks</li> <li>• Health supplement</li> </ul>
Pegaga	<i>Centella asiatica</i>	<ul style="list-style-type: none"> <li>• Health supplement</li> <li>• Coffee</li> </ul>
Kacip fatimah	<i>Labisia pumila</i>	<ul style="list-style-type: none"> <li>• Herbal Tea</li> <li>• Coffee</li> </ul>
Dukung anak	<i>Andrographis paniculata</i>	<ul style="list-style-type: none"> <li>• Health supplement</li> </ul>
Misai kucing	<i>Orthosiphon aristatus</i>	<ul style="list-style-type: none"> <li>• Herbal Tea</li> </ul>
Asam keping	<i>Garcinia atroviridis</i>	<ul style="list-style-type: none"> <li>• Weight maintenance supplement</li> <li>• Herbal Tea</li> </ul>
Mas cotek	<i>Ficus deltoidea</i>	<ul style="list-style-type: none"> <li>• Herbal Tea</li> </ul>

\*Information above is gathered from Herbal Technology, FRIM and Chemical Engineering Pilot Plant, UTM.

It is reported that there are about 1200 species of medicinal plants found in our forest (Soepadma, 1992). There is a great possibility to develop and produce more beneficial herbal extracts that could be formulated into commercial products. This





would ensure that we are able to share the economic benefits from the rapidly growing local as well as foreign market. Before that, we need to choose the right plant that has great potential and optimize the production process of its bioactive extract.

## **1.2 Herbal Processing**

Herbal processing is part of a larger industry incorporating nutraceuticals, functional foods, nutritional supplements and herbal medicines (Ramlan *et al.*, 2005). In Malaysia, most of the herbal researches mainly focus on the chemistry and biological activities of the medicinal plants while the study on herbal processing from the engineering aspects is relatively limited. In fact, the burgeoning knowledge of herbal processing is the important piece of the puzzle before the large scale production could be carried out. There are two institutions participate actively in research of herbal processing namely Chemical Engineering Pilot Plant (CEPP) of University Technology Malaysia (UTM) and Herbal Technology Center (HTC) of Forest Research Institute Malaysia (FRIM).

The processes involved in producing bioactive extract are shown in Figure 1.1. The key processes are drying, solid-liquid extraction and drying of extract (Ramlan *et al.*, 2000). Drying is known as dehydration process which involves removal of moisture. Most of the drying techniques apply heat on the raw materials which might cause certain level of degradation or loss of quality. It is the phytochemicals that contribute to the beneficial bioactivity of medicinal plants. Thus, the major consideration in herbs drying is to shorten the drying time while preserve the active phytochemicals.