



UNIVERSITI PUTRA MALAYSIA

***EXPRESSION AND CHARACTERIZATION OF RECOMBINANT
THERMOSTABLE L2 LIPASE IN PICHIA PASTORIS***

SURIANA SABRI

FBSB 2007 1



**EXPRESSION AND CHARACTERIZATION OF
RECOMBINANT THERMOSTABLE L2 LIPASE IN
*PICHA PASTORIS***

SURIANA SABRI

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2007



**EXPRESSION AND CHARACTERIZATION OF RECOMBINANT
THERMOSTABLE L2 LIPASE IN *PICHTIA PASTORIS***

By

SURIANA SABRI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

April 2007



A special dedication to

Mak, Abah, K. Long, K. Ngah, Abg. Hosni, Abg. Halim, K. Ani, Abg. Jamal, K. Uji, Abg. Yunus and Sarah, who believe in me,

To my nephews and nieces; Syafiqah, Hafiz, Zafirah, Syahirah, Hazirah, Samirah, Suraya, Safia, Nabil and Idham, for their presence, that light up my life,

To Leow; for his unfaltering support and always being there for me...



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**EXPRESSION AND CHARACTERIZATION OF RECOMBINANT
THERMOSTABLE L2 LIPASE IN *PICHIA PASTORIS***

By

SURIANA SABRI

April 2007

Chairman: Professor Raja Noor Zaliha Raja Abd Rahman, PhD

Faculty: Biotechnology and Biomolecular Sciences

The gene encoding mature thermostable L2 lipase from *Bacillus* sp. L2 was cloned into *Pichia pastoris* expression vectors and placed under the control of the constitutive glyceraldehyde-3-phosphate dehydrogenase (*GAP*) promoter and methanol inducible alcohol oxidase (*AOX*) promoter. In inducible system, recombinant L2 lipase was efficiently secreted into the culture medium driven by the *Saccharomyces cerevisiae* α -factor signal sequence, compared to the constitutive system. The optimization of the recombinant L2 lipase production (from inducible system) in 100 mL culture was done for the best clones pP α S3 and pP α G2 from *Pichia* strains SMD1168H and GS115, respectively. The effect of media formulation, methanol concentration and induction time on L2 lipase production from inducible system was evaluated. A time course profile of recombinant lipase production in 500-mL flasks with the optimized conditions



was performed and 15.3 mg/mL and 14.25 mg/mL of dry cell weight were produced after 144 h of induction time from recombinant pP α S3 and pP α G2, respectively. The lipase activities detected from both clones were 91 U/mL and 125 U/mL for pP α S3 and pP α G2, respectively.

The recombinant L2 lipase was purified to 1.8-fold with 63.2% yield and with specific activity of 458.1 U/mg using affinity chromatography. The enzyme was in a monomeric form, non-glycosylated with a molecular weight of 44.5 kDa. The optimum pH and temperature were 8.0 and 70°C, respectively. The enzyme was stable in the pH range of 8.0-9.0 and at 65°C for 60 min where it retained more than 70% of its residual activity. The metal ions Ca²⁺, Na⁺, Cu²⁺ and Mn²⁺ activated the lipase at 1 mM, whereas Mg²⁺ and Zn²⁺ inhibited it. Lipase showed a notable preference for medium to long chain triacylglycerols (C10-C16), with the highest activity toward tripalmitin (C16). It hydrolyzed all the natural oil tested, with the highest hydrolysis rate on corn oil and the least was on sunflower oil. L2 lipase was inhibited by EDTA, PMSF, pepstatin A and all the surfactants tested. It showed random positional specificity towards triolein. CD spectral analysis of L2 lipase revealed a T_m of around 67.2°C.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGEKSPRESAN DAN PENCIRIAN LIPASE L2 TERMOSTABIL
REKOMBINAN DALAM *PICHIA PASTORIS***

Oleh

SURIANA SABRI

April 2007

Pengerusi: Profesor Raja Noor Zaliha Raja Abd Rahman, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Gen yang mengekod lipase L2 termostabil dari *Bacillus* sp. L2 telah diklonkan di dalam vektor pengepresan *Pichia pastoris* dan diletakkan di bawah kawalan promoter-promoter gliseraldehida-3-fosfat dehydrogenase (GAP) konstitutif dan alkohol oksidase (AOX) teraruh metanol. Di dalam sistem teraruh, lipase L2 rekombinan telah dirembeskan ke dalam media kultur oleh jujukan isyarat α -faktor *Saccharomyces cerevisiae* dengan lebih berkesan berbanding sistem konstitutif. Pengoptimuman penghasilan lipase L2 (dari sistem teraruh) di dalam kultur 100 mL telah dilakukan untuk klon-klon terbaik iaitu pP α S3 dan pP α G2 daripada strain *Pichia* SMD1168H dan GS115, masing-masing. Kesan formulasi media, kepekatan metanol, dan masa aruhan ke atas penghasilan rekombinan lipase L2 dari sistem teraruh telah dinilai. Kajian profil masa terhadap penghasilan lipase L2 dengan keadaan optimum telah dijalankan dengan menggunakan kelalang 500-mL dan sebanyak 15.3 mg/mL dan 14.25



mg/mL berat sel kering telah dihasilkan selepas 144 j masa aruhan dari klon-klon pP α S3 dan pP α G2 masing-masing. Aktiviti lipase untuk kedua-dua klon adalah 91 U/mL dan 125 U/mL untuk pP α S3 dan pP α G2, masing-masing.

Lipase L2 rekombinan telah ditulenkan kepada 1.8 kali ganda, dengan penghasilan sebanyak 63.2% dan aktiviti spesifik sebanyak 458.1 U/mg dengan menggunakan kromatografi afiniti. Enzim tersebut berada dalam bentuk monomer, tidak diglikosilasikan dan mempunyai berat molekul sebanyak 44.5 kDa. pH dan suhu optimum enzim ini adalah 8.0 dan 70°C, masing-masing. Enzim ini stabil pada pH 8.0-9.0 dan pada 65°C selama 60 min di mana ia mengekalkan lebih daripada 70% aktivitinya. Ion-ion logam seperti Ca²⁺, Na⁺, Cu²⁺ dan Mn²⁺ pada kepekatan 1 mM boleh mengaktifkan lipase L2, manakala Mg²⁺ dan Zn²⁺ menyahaktifkannya. Lipase L2 lebih memilih untuk menghidrolisiskan triasilgliserol berantai sederhana ke panjang (C10–C16), dengan aktiviti yang paling tinggi ke atas tripalmitin (C16). Ia juga menghidrolisiskan kesemua minyak semulajadi yang diuji dengan kadar hidrolisis yang tertinggi pada minyak jagung, dan yang terendah pada minyak bunga matahari. Lipase L2 dinyahaktifkan oleh EDTA, PMSF, pepstatin A dan kesemua surfaktan yang telah diuji. Ia menunjukkan kespesifikan posisi rawak terhadap triolein. Analisis spektra CD terhadap lipase L2 menunjukkan nilai T_m sebanyak 67.2°C.

ACKNOWLEDGEMENTS

In the name of Allah, all praise is to Allah the Almighty. Had it not been due to His will, this thesis will not be completed.

A journey is easier when you travel together. Interdependence is certainly more valuable than independence. This thesis is the result of two and half years of work whereby I have been accompanied and supported by many people. It is a pleasant aspect and I have now the opportunity to express my gratitude for all of them.

The first person I would like to thank is my supervisor, Prof. Dr. Raja Noor Zaliha Raja Abd. Rahman, for all the patience, guidance, advice, encouragement and help not only for the sake of the project, but for everything. I would also like to thank my co-supervisors; Prof. Dr. Abu Bakar Salleh and Prof. Dr. Mahiran Basri, who monitored my work and took effort in reading and providing me with valuable comments of this thesis. Not just that, to these three great people, I would like to express lots of gratitude for having shown me to love research. They could not even realize how much I have learned from them. My deepest and sincere gratitude for inspiring and guiding this humble being.

The Enzyme and Microbial Technology Research group also substantially contributed to the completion of this project. Especially the strict and extensive



comments and the many discussions and the interactions during the weekly meeting really had a direct impact on me. Thank you to all the principal lecturers including Assoc. Prof. Dr. Basyaruddin Abdul Rahman and my friends from the Department of Chemistry.

My labmates, who are like my sisters and brothers, thank you for being part of my life; Leow, Tengku, K. Ain, K. Lia, Ada, K. Ferrol, K. Ina, Ghaniee, Shook, Chee Fah, Kok Whye, Wani, Wahida, Rofandi, K. Sha, K. Aiman, Ely, Randa, Afshin and Peiman. Each of you means a lot to me and thank you for making the lab such a wonderful place to be in.

I wish to extend my appreciation to everyone, although not individually named here, who had contributed directly or indirectly to my project and thesis.

This study has been financially aided by National Science Fellowship Scholarship from the Ministry of Science, Technology, and Innovation of Malaysia.

Last but not least, to my parents, brothers, sisters, nephews and nieces for their endless love, care and encouragement.



I certify that an Examination Committee met on 20th April 2007 to conduct the final examination of Suriana Sabri on her Master of Science thesis entitle “Expression and Characterization of Recombinant Thermostable L2 Lipase in *Pichia pastoris*” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the Master of Science.

Members of the Examination Committee were as follows:

Suraini Abdul Aziz, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Janna Ong Abdullah, PhD

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Abdul Rahman Omar, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Sheila Nathan, PhD

Associate Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)

HASANAH MOHD GHAZALI, PhD.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Raja Noor Zaliha Raja Abdul Rahman, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Abu Bakar Salleh, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

Mahiran Basri, PhD

Professor

Faculty of Science

Universiti Putra Malaysia

(Member)

AINI IDERIS, PhD

Professor/Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 17 JULY 2007



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at other institution.

SURIANA SABRI

Date: 22 MAY 2007



TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	x
LIST OF FIGURES	xvi
LIST OF APPENDICES	xviii
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	5
2.1 Lipases	5
2.1.1 Applications of lipases	7
2.1.2 Thermostable lipases	12
2.1.3 Properties of thermostable lipases	12
2.1.4 Recombinant thermostable lipases	14
2.1.5 Thermostable L2 lipase	15
2.2 Comparison of prokaryotic and eukaryotic cell expression	17
2.3 Heterologous protein production in yeast	19
2.4 <i>Pichia pastoris</i> expression systems	20
2.4.1 <i>Pichia pastoris</i>	20
2.4.2 <i>Pichia</i> strains	23
2.4.3 Expression vectors	24
2.4.4 Promoters	25
2.4.5 Intracellular and secretory protein expression	28
2.4.6 Signal sequences	29
2.4.7 Integration of expression vectors into genome	30
2.5 Purification strategy of recombinant proteins in <i>P. pastoris</i>	31
3 MATERIALS AND METHODS	32
3.1 Materials	32
3.2 Strains and plasmids	32
3.3 Extraction and quantification of genomic DNA	34
3.4 Amplification of L2 lipase gene	35
3.5 Cloning of L2 lipase gene into expression vectors	36



3.5.1	Preparation of <i>Escherichia coli</i> competent cells	36
3.5.2	Cloning of L2 lipase gene into <i>Pichia</i> expression vectors	37
3.5.3	Heat-shock transformation of <i>Escherichia coli</i>	38
3.5.4	Analysis of recombinant plasmids	39
3.5.5	Sequencing of and glycosylation site prediction of recombinant L2 lipase gene	39
3.6	Expression of L2 lipase in <i>Pichia pastoris</i>	40
3.6.1	Transformation of recombinant plasmid into <i>Pichia pastoris</i>	40
3.6.2	Direct screening of multicopy transformants	43
3.6.3	Direct PCR analysis of <i>Pichia</i> transformants	43
3.7	Protein expression in <i>Pichia pastoris</i>	44
3.7.1	Inducible expression of recombinant pPICZ α A/L2	44
3.7.2	Constitutive expression of recombinant pGAPZ α A/L2	45
3.8	Optimization studies of inducible L2 lipase expression in shake flask	46
3.8.1	Effect of media on L2 lipase expression	46
3.8.2	Effect of methanol concentration on L2 lipase expression	47
3.8.3	Effect of induction time on L2 lipase expression	48
3.9	Analysis of recombinant L2 lipase expression	48
3.9.1	Determination of lipase activity	48
3.9.2	Measurement of cell biomass	49
3.9.3	Determination of protein concentration	50
3.9.4	SDS-PAGE analysis	50
3.10	Purification of recombinant L2 lipase	51
3.11	Characterization of purified L2 lipase	52
3.11.1	Molecular weight determination of recombinant L2 lipase	52
3.11.2	Protein deglycosylation	53
3.11.3	Effect of pH on lipase activity and stability	54
3.11.4	Effect of temperature on lipase activity and stability	54
3.11.5	Effect of metal ions on lipase activity	55
3.11.6	Effect of surfactants on lipase activity	55
3.11.7	Effect of inhibitors on lipase activity	55
3.11.8	Substrate specificity towards natural oils	56
3.11.9	Substrate specificity towards triacylglycerols	56
3.11.10	Positional specificity	57
3.11.11	Denatured protein analysis of L2 lipase	58
4	RESULTS AND DISCUSSION	59
4.1	Genomic DNA extraction and construction of recombinant plasmids	59
4.2	Transformation of <i>Escherichia coli</i>	62
4.2.1	Sequencing and glycosylation site prediction of	64



	recombinant L2 lipase	
4.3	Cloning of L2 lipase in <i>Pichia pastoris</i>	68
	4.3.1 Transformation and selection of recombinant <i>P. pastoris</i>	68
	4.3.2 Direct PCR screening	72
4.4	Expression of thermostable L2 lipase in <i>Pichia pastoris</i>	74
	4.4.1 Expression under alcohol oxidase (AOXI) promoter	74
	4.4.2 Expression under constitutive glyceraldehyde- 3- phosphate dehydrogenase (GAP) promoter	76
4.5	Optimization of enzyme production	81
	4.5.1 Effect of media on lipase production	81
	4.5.2 Effect of methanol concentration on lipase production	83
	4.5.3 Effect of induction time on lipase production	86
4.6	Purification of recombinant L2 lipase	90
4.7	Characterization of purified recombinant L2 lipase	94
	4.7.1 Molecular weight determination and deglycosylation of recombinant L2 lipase	94
	4.7.2 Effect of pH on lipase activity and stability	97
	4.7.3 Effect of temperature on activity and thermostability profile of lipase	100
	4.7.4 Effect of metal ions on lipase activity	104
	4.7.5 Effect of surfactants on lipase activity	107
	4.7.6 Effect of inhibitors on lipase activity	110
	4.7.7 Substrate specificity of L2 lipase	112
	4.7.8 Positional specificity	117
	4.7.9 Circular dichroism (CD) spectra analysis of L2 lipase	119
5	CONCLUSION AND RECOMMENDATIONS	122
	5.1 Conclusion	122
	5.2 Recommendations	123
	REFERENCES	125
	APPENDICES	141
	BIODATA OF THE AUTHOR	156



LIST OF TABLES

Table		Page
1	Important areas of industrial application of microbial lipases	11
2	List of microorganisms	33
3	List of plasmids	33
4	Sequencing primer	41
5	Different media composition	47
6	Lipase activity of various <i>P. pastoris</i> clones in inducible system	75
7	Lipase activity of various <i>P. pastoris</i> clones in constitutive system	78
8	Summary of the purification of His-tagged recombinant L2 lipase from pP α G2	93
9	Effect of metal ions on L2 lipase activity	106
10	Effect of surfactant on L2 lipase activity	108
11	Effect of inhibitor on L2 lipase activity	111
12	Specificity of L2 lipase towards natural oils	116



LIST OF FIGURES

Figure		Page
1	Enzymatic reaction of a lipase catalyzing hydrolysis or synthesis of a triacylglycerol substrate	5
2	Gel electrophoresis of genomic DNA from <i>Bacillus</i> sp. strain L2	60
3	Gel electrophoresis of PCR product of gene encoding mature L2 lipase	61
4	Analysis of recombinant plasmids harboring gene encoding mature L2 lipase	63
5	Nucleotide and amino acid sequence encoded the recombinant L2 lipase in pPICZ α A and pGAPZ α A	65
6	Putative N-glycosylation sites in recombinant L2 lipase in <i>Pichia pastoris</i>	67
7	Gel electrophoresis of linearized plasmids	70
8	Gel electrophoresis of PCR products from <i>Pichia</i> transformants	73
9	Growth curve of recombinant GS115 integrated with constitutive vector (pGAPZ α A)	80
10	Effect of various media on yeast growth and L2 lipase production	82
11	Effect of methanol concentration on lipase production	85
12	Time course of recombinant L2 lipase expression of <i>P. pastoris</i> clone pP α S3 and pP α G2.	87
13	Time course of L2 lipase expression in recombinant pP α G2 analysed by SDS-PAGE.	89
14	Immobilized metal affinity chromatography of His-tagged recombinant L2 lipase.	92
15	SDS-PAGE (12%) of His-tagged recombinant L2 lipase purified	93



	through Ni-Sepharose 6 Fast Flow affinity chromatography	
16	Molecular weight determination of native recombinant L2 lipase by Sephadex G100 gel filtration chromatography	95
17	Effect of deglycosylation on the mobility of purified recombinant L2 lipase from <i>Pichia pastoris</i> under denaturing conditions	96
18	pH profile of L2 lipase	99
19	pH stability of L2 lipase	99
20	Temperature profile of L2 lipase	101
21	Thermostability profile of L2 lipase	103
22	Substrate specificity of L2 lipase towards different chain length of triacylglycerols.	114
23	Thin layer chromatography (TLC) analysis of hydrolysis products after incubation of L2 lipase on triolein as substrate at 70°C for 1 h	118
24	Denatured protein analysis of L2 lipase	121



LIST OF APPENDICES

Appendix		Page
A	Materials and equipments used in the project	142
B	Map and Multiple Cloning site of pPICZ α A vector	146
C	Map and Multiple Cloning site of pGAPZ α A vector	148
D	Calibration curve for determination of oleic acid colorimetrically	150
E	Calibration curve for determination of dry cell weight of <i>Pichia pastoris</i>	151
F	Calibration curve for determination of protein content by Bradford assay	152
G	Composition for SDS-PAGE	153
H	Fatty acid composition of natural oils	154
I	Thermodynamic parameter: T_m , ΔH and ΔS of L2 lipase	155



LIST OF ABBREVIATIONS

APS	ammonium persulphate
bp	base pair
BSA	bovine serum albumin
CTAB	cetyltrimethylammonium bromide
dH ₂ O	distilled water
DNA	deoxyribonucleic acid
dNTPs	deoxyribonucleotide triphosphate
DTT	dithiothreitol
EDTA	ethylenediaminetetraacetic acid
Da	dalton
kDa	kilo dalton
g/L	gram per liter
pmol	picomole
N	normal
rpm	rotation per minute
xg	gravity
UV	ultraviolet
PCR	polymerase chain reaction
PMSF	phenylmethylsulfonyl fluoride
SDS-PAGE	sodium dodecyl sulphate polyacrylamide gel electrophoresis



LB	Luria Bertani
TCA	trichloroacetic acid
OD	optical density
OD ₆₀₀	optical density at 600 nm
A ₂₆₀	absorbance at 260 nm
A ₂₈₀	absorbance at 280 nm
ms	milisecond
SLS	sodium lauryl sulphonate
TEMED	N,N,N,N-Tetramethylenediamide
TSB	tripticase soy broth
YNB	yeast nitrogen base
U/mL	unit per milliliter
U/mg	unit per milligram
v/v	volume per volume
w/v	weight per volume
V	Volt
V/cm	volt per centimeter
μF	Microfarad
MD	minimal dextrose
MM	minimal methanol
kb	kilo base
Mut ⁺	methanol utilization phenotype plus



Mut ^s	methanol utilization phenotype slow
BMGY	buffered glycerol-complex medium
BMMY	buffered methanol-complex medium
RT	room temperature
sp	species
U	unit
YPD	yeast extract, peptone and dextrose media
YPDS	yeast extract, peptone, dextrose and sorbitol media
CD	circular dichroism
MW	molecular weight



CHAPTER 1

INTRODUCTION

The global market for industrial enzymes has already achieved the USD 2 billion mark, and it is sure to grow. A report from McKinsey & Co., recently indicated that the future for sustainable development is clearly a bright one, and enzyme technology will play a major role, along with the use of microorganisms, both natural and engineered (Wood and Scott, 2004). To date, approximately 80% of all industrial enzymes are hydrolytic in nature and used for depolymerization of natural substances. Of these enzymes, 60% are proteolytic enzymes used by the detergent, dairy and leather industries. Thirty percent are carbohydrases used in baking, distilling, brewing, starch, and textile industries. This leaves lipases and highly specialized enzymes for use in pharmaceutical, oleochemical, and analytical industries (Kirk *et al.*, 2002). However, this share has the potential to increase dramatically via a wide range of lipases' new applications (Jaeger and Eggert, 2002; Pandey *et al.*, 1999).

Lipases are efficient catalysts for lipolytic reactions initiating the catabolism of fats and oils by hydrolyzing the fatty acyl ester bonds of acylglycerols (Vulfson, 1994). Lipases have tremendous potential for further exploitation in biotechnology. Their ability to catalyze a wide variety of reactions allow numerous applications in industry such as the removal of oils and fats from



fabrics, machinery and waste water, the production of mono- and diglycerides for food emulsifiers and stereospecific synthesis of compounds including precursors for biologically active therapeutics, herbicides or pesticides (van Kuiken and Behnke, 1994; Haas *et al.*, 1992).

Enzymes from thermophiles have been found to be the most practical commercial used biocatalysts to date because of their overall inherent stability which are better suited to the harsh conditions of industrial processes (Kirk *et al.*, 2002). There are many efforts directed at improving enzymes involved in industrial processes in order to decrease cost and increase energy efficiency. One of the most promising methods to obtain better enzymes is via recombination DNA technology to produce the enzymes in large quantities with desired properties which will make them economically viable.

Cloning and characterization of lipases from thermophilic bacteria and the expression of the biologically active proteins in *Escherichia coli* had been reported (Rahman *et al.*, 2005, Sinchaikul *et al.*, 2001). This protein shows high activity at high temperature and this feature offers several interesting advantages in term of biotechnological applications. Although the protein obtained from recombinant *E. coli* was sufficient to perform a variety of experiments, the low production together with the complex purification procedures were proven unsuitable for industrial production of the enzyme.

