



UNIVERSITI PUTRA MALAYSIA

***CO-COMPOSTING OF CHICKEN MANURE WITH RICE HUSK BIOCHAR
AND ITS EFFECTS ON THE PERFORMANCE OF CUCUMIS SATIVUS L.***

THEEBA A/P MANICKAM

FP 2013 66



**CO-COMPOSTING OF CHICKEN MANURE WITH RICE HUSK
BIOCHAR AND ITS EFFECTS ON THE PERFORMANCE
OF *CUCUMIS SATIVUS* L.**

THEEBA A/P MANICKAM

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2013



**CO-COMPOSTING OF CHICKEN MANURE WITH RICE HUSK
BIOCHAR AND ITS EFFECTS ON THE PERFORMANCE OF
*CUCUMIS SATIVUS L.***

By

THEEBA A/P MANICKAM

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the requirements for the Degree of Master of Science**

December 2013

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is dedicated to my beloved husband, children, parents and sisters for your unfailing support, love, care and prayers throughout my journey of completion. Without all of you this would not be possible. May god bless you all.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

CO-COMPOSTING OF CHICKEN MANURE WITH RICE HUSK BIOCHAR AND ITS EFFECTS ON THE PERFORMANCE OF *CUCUMIS SATIVUS* L.

By

THEEBA A/P MANICKAM

December 2013

Chairman : Associate Professor Ahmad Husni bin Mohd. Hanif, PhD

Faculty : Agriculture

Rice husk biochar (RHC) in this study is a major by-product of the rice processing mill, abundantly available (estimated about 32,000 tonne/ year) and very low in cost. It is found underutilized and often paid, approximately RM8,000 annually to be disposed of by mill owners in some area especially Kelantan state. RHC was observed to increase crop growth and yield when used as polybag or potting media by farmers. However, there are inadequate detailed quantification on local mill RHC have been documented in relation to its capability in increasing crop production and soil improvement in Malaysia. This study was therefore carried out to, (i) determine physico-chemical properties of locally milled RHC, (ii) evaluate the effects of selected rates of RHC amendmend in composting process and compost properties of chicken manure and (iii) assess the effects of RHC and developed RHC compost on fruity vegetable crop production and soil properties.

RHC in this study was collected from a local mill which operates cyclone husk furnace to generate heat for rice drying activity. RHC was characterized for its physico-chemical properties such as to its particle size, ash content, nutrient contents, Methylene Blue adsorption capacity, Brunauer Emmett and Teller (BET) surface area, functional groups using FTIR and Scanning electron micrograph image (SEM). Selected RHC application rates of 5% and 10% were evaluated for its effects on chicken manure composting through determination of composting temperature and CO₂ release pattern, moisture, pH, nutrient content and microbial population. The finished compost were analysed for its leaching rates through laboratory column leaching test and the leached composts were viewed under SEM/EDX. RHC compost and fresh RHC were assessed for its efficacy through field trial on *Cucumis sativus* for two cropping cycles. The evaluation was based on crop growth and yield

performances, changes in soil physico-chemical properties, crop nutrient concentrations and crop nutrient use efficiency.

Results from the study showed that RHC is highly alkaline with pH 8.9, with a total carbon content of 16%, low bulk density of 0.17 g/cm^3 , particle size $< 1 \text{ mm}$, possess moderately high surface area of $401 \text{ m}^2\text{g}^{-1}$ with numerous meso to macropores ranging from (2.7 nm- 10 μm), presence of functional groups such as carboxylic acids and Methylene Blue adsorption capacity of 38.5 mg/g. RHC amendments at 5% and 10% was found to accelerate the composting process. The composts with RHC amendments showed early drops from the plateaued thermophilic temperature stage by 10 days supported by higher CO_2 respiration. The final RHC composts contained neutral pH characteristics (7-7.1). It was identified that higher moisture retention of 8% and 4% was obtained in the matured RHC compost 10% and RHC compost 5% respectively as compared to control. RHC was also found to minimize nitrogen loss during composting in which minimum losses of 0.4% for RHC compost 10% and 0.47% for RHC compost 5% was quantified at 10 days of composting as compared to 10.3% for compost without RHC (control). Laboratory leaching tests and SEM micrograph on RHC compost at both rates has proven that RHC does physically trapped and chemically sorb organic and inorganic molecules containing C, N, P and K and retained them against leaching losses.

Field assessment for two seasons on *Cucumis sativus* showed the plots treated with 10% RHC compost gave the highest yield in season 1 (51.2 t/ha) followed by 5% RHC compost (49.5 t/ha) and fresh RHC 3% applied into soil (49.4 t/ha). Meanwhile for season 2, the highest crop yield production was in 5% RHC compost (45.9 t/ha), followed by 10% RHC compost (45 %) and fresh RHC 3% (41 %). The yield achievement is strongly correlated with crop nutrient concentration of N, P, and K as well dry root mass which were significantly higher in plots under 5% RHC compost and 10% RHC compost and followed by fresh RHC at 3% amendment as compared to non-amended RHC compost plot. Meanwhile soil CEC, pH, water holding capacity and porosity were found the highest in plots treated with 3% RHC. Treatment of RHC 1% into soil showed lesser positive effects and plots without RHC treatment was showing significantly lowest values for crop yield, growth and changes in soil physico-chemical characteristics.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan ijazah Master Sains

**PENGGOMPOSAN TINJA AYAM BERSAMA ARANG-BIO SEKAM PADI
DAN KEBERKESANNYA TERHADAP PERTUMBUHAN *CUCUMIS
SATIVUS L.***

Oleh

THEEBA A/P MANICKAM

Disember 2013

Pengerusi : Profesor Madya Ahmad Husni bin Mohd. Hanif, PhD

Fakulti : Pertanian

Arang-bio sekam padi (RHC) dalam kajian ini, merupakan hasil sampingan utama bagi kilang pemprosesan padi, yang mana ia boleh didapati dalam kuantiti yang banyak dan harganya sangat murah. Ia didapati kurang diguna-pakai malah pemilik kilang di sesetengah kawasan khususnya di Kelantan terpaksa mengeluarkan kos untuk ia dilupuskan. Terdapat juga RHC yang diguna-pakai oleh petani khususnya di Kedah yang mana ia digunakan sebagai media semaian benih padi untuk sistem 'transplanting' dan media polibeg untuk sistem fertigasi. Namun, masih kurang terdapatnya kajian mengenai kadar penggunaan RHC dalam meningkatkan hasil tanaman dan pembaikan tanah di Malaysia. Justeru itu, kajian ini telah di rekabentuk dan dijalankan dengan beberapa objektif iaitu (i) mengkaji ciri-ciri kimia-fizik RHC tempatan, (ii) mengkaji kesan kadar campuran RHC terpilih ke atas proses pengkomposan dan ciri-ciri kompos tinja ayam dan (iii) mengkaji kesan RHC dan pembuatan kompos RHC terhadap pengeluaran tanaman sayur buah (mentimun) dan perubahan ke atas ciri-ciri kimia- fizik tanah.

RHC untuk kajian ini, diperolehi dari kilang padi tempatan yang diusahakan oleh BERNAS yang menggunakan mesin pembakaran 'Cyclone Furnace' yang mengeluarkan haba panas untuk aktiviti pengeringan padi. RHC tersebut di analisa untuk ciri-ciri kimia-fizik seperti saiz partikel, kuantiti abu, kandungan nutrien, kapasiti penyerapan 'Methylene Blue', luas permukaan BET, kumpulan berfungsi FTIR dan imej mikrograf SEM/EDX. Kadar terpilih RHC pada nilai 10% dan 5% telah diuji keberkesannya melalui proses pengkomposan dengan tinja ayam dan semasa aktiviti pengkomposan data-data seperti suhu kompos, kadar pelepasan karbon dioksida, kelembapan, pH, kandungan nutrien dan populasi mikrob telah di ambil. Kompos matang telah dikaji untuk kandungan nutrien semasa ujian tahap larut resap di dalam makmal dan kompos tersebut diteliti di bawah SEM/EDX. Bagi kajian lapangan pula, dua rawatan iaitu Kompos RHC (dua kadar berbeza 10% dan 5%) dan RHC sahaja (3% dan 1%) diuji ke atas tanaman mentimun (*Cucumis sativus*) untuk dua musim berturut. Penilaian ujikaji adalah berdasarkan kesan ke

atas pertumbuhan tanaman, pengeluaran hasil, perubahan kimia-fizik tanah, kuantiti nutrien dalam tanaman dan keefisienan penggunaannya terhadap tanaman.

Keputusan analisa menunjukkan, RHC sangat beralkali dengan pH 8.9, kandungan karbonnya ialah 16%, mempunyai ketumpatan yang rendah iaitu 0.17 g/cm^3 , saiz partikel $< 1\text{mm}$, dengan luas permukaan $401 \text{ m}^2\text{g}^{-1}$ yang meliputi pelbagai liang meso dan makro (berukuran anggaran 2.7nm - $10\mu\text{m}$), kedapatan kehadiran kumpulan berfungsi seperti asid karboksilik dan kapasiti penyerapan 'Methylene Blue' sebanyak 38.5 mg/g . Campuran RHC pada kadar 5% dan 10% didapati boleh mempercepatkan proses pengkomposan (pematangan awal). Campuran kompos dengan RHC menunjukkan penurunan suhu awal 10 hari dari keadaan termofilik dengan dibantu peningkatan lebih tinggi kadar respirasi CO_2 . Kompos matang mempunyai nilai pH yang stabil (7-7.1). Nilai kelembapan yang tinggi pada kadar 8% dan 4% di dapati pada kompos RHC yang matang 10% dan kompos RHC 5%. RHC juga diukur kadar kehilangan nitrogen semasa proses pengkomposan dengan nilai 0.4% untuk kompos RHC 10% dan nilai 0.47% untuk kompos RHC 5%, manakala 10.3% untuk kompos tanpa RHC. Sementara itu, ujikaji tahap larut resap dan imej mikrograf SEM/EDX dalam makmal juga menunjukkan kedua-dua kadar RHC menunjukkan RHC secara fizikalnya mampu memerangkap molekul dan secara kimia mampu menyerap molekul organik dan tak organik seperti C, N, P, dan K serta mengekalkannya dari hilang melalui proses larut resap.

Hasil kajian lapangan ke atas tanaman mentimun menunjukkan pada musim pertama Kompos RHC 10% mempunyai hasil pengeluaran yang tertinggi (51.2 t/ha), diikuti oleh Kompos RHC 5% (49.5 t/ha) dan RHC 3% (49.4 t/ha). Manakala pada musim ke dua hasil tertinggi didapati dari plot Kompos RHC 5% (45.9 t/ha), diikuti Kompos RHC 10% (45 t/ha) dan RHC 3% (41 %). Pengeluaran hasil didapati sangat berkaitan (korelasi positif) dengan kuantiti nutrien N, P, dan K dalam tanaman dan berat kering biomas akar yang mana ia menunjukkan nilai lebih tinggi pada plot yang menggunakan rawatan berikut mengikut hierarki: Kompos RHC 5% > Kompos RHC 10% > RHC sahaja 3%. Namun begitu, nilai CEC tanah, nilai pH, kadar pegangan air dalam tanah dan porositi didapati tertinggi dalam plot rawatan RHC sahaja 3%. Penggunaan RHC 1% pula tidak menunjukkan peningkatan begitu signifikan berbanding dengan plot kompos tanpa RHC manakala plot rawatan kompos tanpa RHC pula mempunyai nilai kedua paling rendah untuk hasil tanaman, pertumbuhan dan perubahan dalam sifat kimia fizik tanah untuk dua musim penanaman.

ACKNOWLEDGEMENTS

Praise to God Almighty for His blessings, which provided me the strength and spirit that enabled me to complete this thesis. I would like to express my deepest and sincerest appreciations with a heartfelt gratitude to Associate Professor Dr. Ahmad Husni bin Mohd Hanif, the Chairman of Supervisory Committee for his full support, expert guidance encouragement and understanding throughout my research and study. I extend my sincere gratitude to Dr. Samsuri Abd Wahid and Associate Professor Dr. Robert Thomas Bachmann as members of the Supervisory Committee, for their guidance, suggestions, invaluable advice and comments throughout the duration of this study and preparation of this thesis.

Also with deep sense of honour, I wish to express my sincere gratitude to Malaysian Agriculture Research & Development Institute in providing me the opportunity to obtain my master's degree and for making sure that i have received the financial support in completing my study. My sincere thanks are extended to all supporting staffs of the Department of Land Management, University Putra Malaysia and Malaysian Agriculture Research & Development Institute for their time and assistance in the completion of laboratory and field works. Special thanks and appreciation go to my beloved family members for all the encouragement given and sacrifices made throughout this journey. Last but not least, my sincere appreciation to my colleagues especially Pn. Aini Zakaria, Pn. Illani Ibrahim, Mr. Muhd. Syahren Adzhar and Pn. Norziana ZZ for their help and moral support in completing this thesis.

I certify that a Thesis Examination Committee has met on 24th December 2013 to conduct the final examination of Theeba a/p Manickam on her thesis entitled “Co-Composting of Chicken Manure with Rice Husk Biochar and its Effects on The Performance of *Cucumis sativus* L.” in accordance with Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Hamdan bin Jol, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Aminuddin bin Hussin, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Shamshuddin bin Jusop, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Adzemi Mat Arshad, PhD

Senior Lecturer
Universiti Malaysia Terengganu
Malaysia
(External Examiner)

NORITAH OMAR, PhD

Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 January 2013

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Ahmad Husni Mohd. Hanif, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Samsuri Abd Wahid, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Robert Thomas Bachmann, PhD

Associate Professor
University Kuala Lumpur
Malaysia
(Member)


BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 13 February 2014

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.



THEEBA A/P MANICKAM


Date: 24 Dec 2013

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENT	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS/NOTATIONA/GLOSSARY OF TERMS	xvii
CHAPTER	
1 INTRODUCTION	1
2 LITREATURE REVIEW	4
2.1 Biochar	
2.1.1 Biochar in General	4
2.1.2 Biochar Effects on Soil Physico-Chemical and Biological Properties	5
2.1.3 Biochar Composting	6
2.2 Biochar from Rice Husk	
2.2.1 Utilization of Rice Husk and Biochar Production	7
2.2.2 Cylone Gasifier for RHC Production	8
2.2.3 RHC Properties	10
2.3 Advantages of RHC in Agriculture	
2.3.1 RHC for Soil Improvement and Crop Production	12
2.3.2 Local Mill RHC Utilizations in Malaysia	13
2.4 Composting of Animal Manure	
2.4.1 Animal Manure Composting for Crop Production	16
2.4.2 Drawbacks of Animal Manure Composting	17
3 MATERIALS AND METHODS	18
3.1 Characterization of RHC	
3.1.1 Collection and Preparation of RHC Samples	18
3.1.2 RHC pH and Particle Size	18
3.1.3 RHC Ash Content	20
3.1.4 RHC ultimate (C, H, N, O and S) Analysis	20
3.1.5 RHC Methylene Blue Adsorption Capacity	20

3.1.6	RHC FTIR (Fourier Transform Infrared) Analysis	22
3.1.7	RHC Cation Exchange Capacity Determination	22
3.1.8	SEM/EDX Image and BET Surface Area of RHC	22
3.1.9	RHC Water Holding Capacity	23
3.2	Co-Composting Chicken Manure with RHC	
3.2.1	RHC Co-composting	23
3.2.2	Determination of Composting Temperature and CO ₂ Flux	24
3.2.3	Determination of compost pH and Moisture Content	25
3.2.4	Total Nutrient Contents in Compost	25
3.2.4.1	Total Carbon and Nitrogen	25
3.2.4.2	Total Phosphorus	25
3.2.4.3	Exchangeable Bases	26
3.2.4.4	Total silica	26
3.2.5	Compost Microbial Population	26
3.2.6	Compost SEM/EDX Spectra	27
3.2.7	Compost Laboratory Column Leaching Test	27
	Field Assessment on <i>Cucumis sativus</i>	
3.3.1	Study Site	28
3.3.2	Experimental Set up and Design	28
3.3.3	Crop Cultivation and Establishment	30
3.3.4	RHC Compost and RHC Applications	31
3.3.5	Soil Samplings and Analysis	31
3.3.5.1	Soil pH (H ₂ O)	31
3.3.5.2	Soil Total Carbon and Nitrogen	32
3.3.5.3	Available Phosphorus	32
3.3.5.4	Cation Exchange Capacity and Exchangeable Bases	32
3.3.5.5	Soil Porosity	33
3.3.5.6	Water Holding Capacity	33
3.3.6	Crop Growth Data, Samplings and Analysis	33
3.3.6.21	Crop Nutrient Use Efficiency	34
3.4	Statistical Data Analysis	34
4	RESULTS AND DISCUSSIONS	35
4.1	Characterization of RHC	
4.1.1	Physical Properties of RHC	35
4.1.2	Chemical Properties of RHC	40
4.2	RHC Composting and Properties	
4.2.1	Composting Temperature and CO ₂ Flux	44
4.2.2	Compost pH and Moisture Content	47
4.2.3	Compost Nutrient Contents	51
4.2.4	Compost Microbial Population	56
4.2.5	Nutrient-RHC Interaction in Compost	60

4.3	Field Assessment on <i>Cucumis sativus</i>	
4.3.1	Crop Growth Performances	66
4.3.2	Crop Yield Performances	68
4.3.3	Crop Nutrient Concentrations	72
4.3.4	Soil Properties Changes	73
4.3.5	Correlation between Crop and Soil Variables	75
5	SUMMARY, CONCLUSION AND FUTURE RESEARCH RECOMMENDATIONS	79
	REFERENCES	82
	APPENDICES	94
	BIODATA OF STUDENT	101
	LIST OF PUBLICATIONS	102



LIST OF TABLES

Table		Page
2.1	Physical and economical flows through rice mill (Padi Beras Nasional Berhad, 2011)	9
2.2	Physical and economical flows through the mill (BERNAS, 2007)	10
2.3	BERNAS Cyclone Husk Furnace specifications (BERNAS, 2007)	10
2.4	Characteristics of gasified and rapid pyrolysed rice husk biochar	12
3.1	Main characteristics of the adsorbate (Zhang et al., 2006)	21
4.1	Physico-chemical characteristics of RHC in this study	36
4.2	Methylene blue adsorption capacity of various biomass carbon (Kannan and Sundram, 2001)	43
4.3	Other nutrient contents in final compost at day 50	55
4.4	Correlation matrix of variables during composting period using Pearson Correlation Coefficient (n=45)	59
4.5	Total percentage of nutrient elements leached from compost	62
4.6	Mean values of root biomass dry weight for two cropping cycles	68
4.7	Mean values of crop nutrient concentrations for two cropping cycles	73
4.8	Soil properties at experimental plot (before cropping cycle)	74
4.9	Mean values of soil chemical properties after cropping cycles	75
4.10	Mean values of soil pH, CEC and porosity after cropping cycle	75
4.11	Relationships between different variables under field assessment of <i>Cucumis sativus</i> by Pearson Correlation Coefficient (n=24)	78

LIST OF FIGURES

Figure		Page
2.1	A basic model of a complex biochar particle from soil showing pores of various sizes and chrystalline–graphene-like sheets (Nguyen et al, 2004; Yu et al, 2006)	6
2.2	The compartments of Cyclonic Husk Furnace (CHF) (BERNAS, 2007)	8
2.3	RHC used as media for rice transplanting seedling bed	14
2.4	RHC used as 100% polybag media for crops planted under fertigation system	14
2.5	Crop rooting development under 100% RHC media compared to cocopeat and soil	15
2.6	RHC used as soil amendment for rice cropping	15
3.1	Cyclone Husk Furnace (CHF) in Seong Heng Rice Mill, Kuala Kedah	19
3.2	Produced RHC at furnace outlet	19
3.3	CO ₂ flux sampling and temperature recording in compost pile	26
3.4	The layout of experimental plot (complete randomized design)	29
3.5	<i>Cucumis sativus</i> planted in rows on raised soil beds	30
4.1	Particle size distribution for RHC	36
4.2	BET Isotherm obtained and showing pore volume at $p/p^{\circ} 0.95$ for RHC	37
4.3	Water retention curve of RHC in this study	39
4.4	Atomic ratio of O/C and H/C of RHC in this study	40
4.5	FTIR spectrum of RHC samples in this study	42
4.6	Changes in composting temperature	42
4.7	Changes in composting temperature and CO ₂ flux during thermophilic stage	46
4.8	Changes in pH between treatments during composting process	47
4.9	Changes in moisture content between treatments during composting process	49
4.10	Total carbon preserved in composts throughout composting process	50
4.11	Total nitrogen preserved in composts throughout composting process	52
4.12	Changes in C/N ratio of composts throughout composting process	54
4.13	Changes in C/N ratio of composts throughout composting process	55
4.14	Changes in total fungi count between treatments during composting process	57
4.15	Changes in total bacteria count between treatments during composting process	58

4.16	Cumulative leaching value (%) of NO ₃ for estimated average rainfall volume Peninsular Malaysia	61
4.17	Cumulative leaching value (%) of P for estimated average rainfall volume Peninsular Malaysia	61
4.18	Cumulative leaching value (%) of K for estimated average rainfall volume Peninsular Malaysia	62
4.19	SEM-EDX spectrum for RHC compost T2	64
4.20	SEM-EDX spectrum for RHC compost T3	64
4.21	Mean values of SPAD chlorophyll index and plant height for two cropping cycles	67
4.22	Mean values of total fresh weight of <i>Cucumis sativus</i> and nutrient use efficiency (partial factor) for two cropping seasons	70
4.23	Mean values of fresh weight of <i>Cucumis sativus</i> at different harvesting time for two cropping cycles	71

LIST OF ABBREVIATIONS/NOTATIONS/GLOSSARY OF TERMS

ANOVA	Analysis of Variance
BERNAS	Padi Beras Nasional
BET	Brunauer–Emmett–Teller
CEC	Cation Exchange Capacity
CHF	Cyclone Husk Furnace
CRD	Completely Randomized Design
DMRT	Duncan Multiple Range Test
EDX	Energy Dispersive X-ray
FTIR	Fourier Transform Infrared
HCl	Hydrochloric Acid
HSD	Honestly Significant Different
ICP-OES	Inductively Couple Plasma-Optical Emission Spectrometer
ISO	International Organization for Standardization
IUPAC	International Union for Pure and Applied Chemistry
K ₂ SO ₄	Potassium Sulphate
MB	Methylene Blue
mL	millilitre
NA	Nutrient Agar
NH ₄	Ammonium
NH ₄ OAC	Ammonium Acetate
PDA	Potato Dextrose Agar
SAS	Statistical Analysis System
SEM	Scanning Electron Microscopy

CHAPTER 1

INTRODUCTION

In the year 2011 Malaysia's paddy production was 2,575,988 tonnes obtained from 687,516 hectares of the total rice cultivated area. The total production of milled rice was 1,648,632 tonne which comprised about 64% of the total paddy production in nation (Department of Agriculture, 2011). The remaining 36% consist of biomass waste that was removed during the rice processing operations in the mill. Rice husk constitute 23% of the removed biomass waste (Padi Beras Nasional Berhad, 2011). Thus, it is estimated that, in 2011 alone, a total amount of 592,477 tonnes of rice husk were generated from rice milling operation.

In major rice processing mills such as BERNAS owned rice mills, the rice husks generated are utilized as fuel. They are gasified using cyclone furnace to generate heat for the rice drying activity. Rice drying is a batch process and carried out over a period of three months per annum. The cyclone furnace system produces a byproduct namely rice husk biochar (RHC) or rather widely known as rice husk black ash at a RHC recovery rate of 30%. Shackley et al., (2012) had stated that rice husk biomass is widely used as a renewable energy worldwide in developing and industrialized countries. The fuel generation is done through various processes from downdraft gasification, rapid pyrolysis to fluidized bed reactors in the rice processing mills. Beagle (1978) had done a comprehensive survey on the utilization of rice husk for energy purposes and identified gasification as one of the viable routes for effective use of rice husk for thermal and mechanical/electrical applications.

During the gasification process approximately 30 kg of rice husk biochar (RHC) are produced/ hour per rice mill from 100 kg rice husks per hour per mill at temperatures ranging from 350-600°C. A total of about 650 tonnes of RHC per mill yearly are produced. Based on the surveys done in Kedah, Kelantan, and Selangor, there are more than 50 RHC producing mills identified, thus an estimated amount of 32,000 tonne of RHC are generated in these states every year. RHC in Kelantan rice mills can be obtained free of charge up to 300kg. On the other hand, rice mill operators in Kedah and Sekinchan have started to charge approximately RM 10 for 300 kg, since RHC has potential as an efficient cropping medium. Local mill RHC shares some of the characteristics of biochar, which has been defined as the porous carbonaceous solid produced by thermochemical conversion suitable for the safe and long-term storage of carbon (Shackley and Sohi, 2010).

In Kedah, RHC are widely utilized as polybag medium and in transplanting beds for rice seedlings as it encourages high rooting capacity. On the other hand in Kelantan, RHC is paid to be disposed of by the rice mills costing approximately RM 8000 /mill/year (BERNAS, 2011). The disposal of RHC is mainly due to ineffective utilization of RHC for any beneficial purposes because Kelantan farmers normally practise direct seeding which does not make use of RHC in any way.

Through observation on farmer's crops which used RHC as media, the ability of RHC in crop improvement especially with regards to its high rooting development, water and nutrient retention was identified. Thus there is a need to further investigate the potential of RHC in retaining nutrients of fertilizers such as compost. There are some studies done on other biomass biocharcoal such as bamboo biocharcoal and wood biocharcoal that showed some properties of biocharcoal that enhanced the composting properties (Yoshizawa et al., 2007). Glaser (2007) and Glaser and Birk (2011) explained that the world's most fertile soil located at *Terra Preta* was most likely formed by the mixing of charred residues (biochar) with biogenic wastes from human settlements (excrements and food wastes including bones and ashes) which were converted by microbes to a biochar-compost-like substrate. This explains why co-composting of biochar and fresh organic material is likely to have more benefits as compared to the mixing of biochar or compost separately into soil. The benefits include, enhanced nutrient use efficiency, higher and long-term C sequestration potential, biological activation of biochar and better material flow management compared to individual compost and biochar applications (negative priming effect).

In Malaysia, composting of biomass and animal waste is being widely practised mainly as fertilizers for crop production. Currently, waste generated from chicken farms keeps increasing as a result from rapid growth of the chicken industry. Based on report by Department of Agriculture (2012), price per kg for composted chicken manure is RM 1.40 and untreated is RM 0.36. Chicken manure is still preferred and widely used by farmers as organic fertilizers as it is cheaper than other locally produced commercial organic fertilizers such as Complehumus[®] (RM 2.50 /kg), Wonderful Organic[®] (RM2.00/kg) and AmmoGold[®] (RM 1.40/kg). The utilization of the chicken farm waste mainly chicken manure as sources of nutrients for food crop production is becoming very important as it appears to be the most cost-effective source. However, direct applications of chicken manure onto the soil as fertilizers may cause environmental problems such as release of foul odors and may trigger pest infestations which can be minimised if chicken manure is composted. The disadvantages of composting include the inefficiency of the composting process due to nutrient losses through volatilization and leaching (Eghball et al., 1997); possible odor (Walker, 1993) equipment, labour requirement and process time. The biological transformation of chicken manure via

composting takes several weeks and may have nutrient losses through leaching during the process. Thus, it is expected that the additions of RHC will enhance the composting process and quality of compost as achieved by other studies using various types of biochars.

In summary, RHC produced by local rice mill is either a waste or low in cost and found in abundance. Most importantly, it was observed to increase crop growth and yield when used as polybag or potting medium. However, there is a lack of quantitative data on the capability of RHC to increase crop production and soil improvement in Malaysia. Apart from that there is an urgent need to increase the utilization of RHC to solve the waste disposal problem in some states by the use of RHC in composting of animal manure and as soil amendment for other crops. Therefore, a study was embarked with the following hypothesis and objectives.

Hypothesis:

Local mill rice husk biochar gives positive effects in enhancing composting process of chicken manure, improves crop production and soil physico-chemical properties

Objectives:

1. To determine physico-chemical properties of locally milled RHC.
2. To evaluate the effects of selected rates of RHC amendment in composting process and compost properties of chicken manure.
3. To assess the effects of RHC and developed RHC compost on fruity vegetable crop production and soil properties

REFERENCES

- Abe, F. (1988). The thermochemical study of forest biomass. *Forest Produce Chemistry*. 45: 1-95.
- Ahiduzzaman, M. (2007). Rice husk energy technologies in Bangladesh. *Agricultural Engineering International: the CIGR E journal Invited Overview* No. 1. Vol. IX.
- Aini, Z., Sivapragasam, P., Vimala, P. and Mohamed Roff, M.N. (2005). Compost and composting, in book chapter: Organic Vegetable Cultivation in Malaysia. Malaysian Agriculture Research and Development Institute (MARDI). pp. 73-92
- Amonette, J.E. and Joseph, S. (2009). Characteristics of biochar: Microchemical properties. In *Biochar for Environmental Management: Science and Technology*, ed. J. Lehmann and S. Joseph, (pp. 33-52). Earthscan, London, United Kingdom.
- Ankerman, D. and Large, R. (1992). *Plant Analysis*, In *Agronomy Handbook* (pp. 99-130), Midwest Laboratories.
- Anuradha, G., Prem, D. G. and Ramachandaran Iyer, P. V. (1992). Combustion and gasification characteristics of rice, *Journal of FUEL*. 71: 889-894.
- Ashoka, G., Ben, B., Janendra, C. and Eranjani, M. (2012). Impact of rice husk charcoal application on soil properties and growth and yield of rice. vol 17, pp 202, International Forestry and Environment Symposium. Department of Forestry and Environment Science, University of Sri Jayewardenepura.
- Bain, R.L., Overend, R.P. and Craig, K.R. (1998). Biomass – fired Power Generation. *Fuel Process Technology*. 54:1 - 16
- Beagle, E.C. (1978). Rice husk conversion to energy. Rome, Italy: FAO Consultant.
- BERNAS. (2011). Annual paddy production report, BERNAS Daerah Peringat. pp. 50.
- BERNAS. (2007). Operasi “Cylonic Husk Furnace”, Pekeliling BERNAS Production Sdn Bhd (BEPRO-1). pp.17.
- Bishop, P.L. and Godfrey, C. (1983). Nitrogen transformations during sludge composting. *Biocycle*. 24:34–39.
- Blasi, C., Signorelli, G., Russo, C. and Rea, G. (1999). Product distribution from pyrolysis of wood and agricultural residues. *Industrial Engineering Chemistry Resources*. 38(6):2216–2224.

- Brady, N. C. and Weil, R.R. (2004). *Elements of the Nature and Properties of Soils* 2nd Ed. Pearson Prentice Hall. Upper Saddle River NJ. pp.111-112.
- Brady, N.C. and Weil, R.R. (2002). *The Nature and Properties of Soils, 13th edition*. Prentice Hall, Upper Saddle River, N.J
- Bray, R.H. and Kurtz, L.T. (1945). Determination of total, organic, and available forms of phosphorus in soils. *Soil Science*. 59: 39-45.
- Bridgewater, A.V., Meier, D. and Radlein, D. (1999). An overview of fast pyrolysis of biomass. *Journal of Organic Geochemistry*. 30:8918-892.
- Brennan, J.K., Bandosz, T.J., Thomson, K.T. and Gubbins, K.E. (2001). Water in porous carbons. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. 187-188: 539-568.
- Brunauer, S., Emmett, P. H. and Teller, E. (1938). Adsorption of gases in multimolecular layers. *Journal of the American Chemical Society* . 60:309-319.
- Bruno, O. Dias., Carlos, A. Silva., Fabio, S., Higashikawa, R. and Miguel A.S. (2010). Use of biochar as bulking agent for the composting of poultry manure. Effect on organic matter degradation and humification. *Bioresource Technology*. 101:1239-1246.
- Buijsman, E., Maas, H.F.M. and Asman, W.A.H. (1987). Anthropogenic NH₃ emissions in Europe. *Atmospheric Environment*. 21:1009–1022. doi: 10.1016/0004-6981(87)90230-7.
- Chapman, H. D. (1965). Cation Exchange Capacity. In C. A. Black et al. (ed.) *Methods of soil analysis*. American Society of Agronomy, Inc., Madison, Wisconsin. Agronomy. 9:891-901
- Chan, K. Y., van Zwiten, B.L., Meszaros, I., Downie, D. and Joseph, S. (2007). Agronomic values of greenwaste biochars as soil amendments. *Australian Journal of Soil Research*. 46:437-444.
- Christopher, T.B.S. and Jamal, T. (2006). *Bulk density, Water Content and Porosity*, In Chapter 3, *Soil Physics Analysis*, Volume 1, University Putra Malaysia, pp 11-13.
- Cornelissen, G., Shitumbanuma, V. and Martinsen, V. (2011). Biochar in conservation farming in Zambia: Improving crop yield and storing carbon. End report phase 1, report no: 20100920-006-R, Conservation Farming Unit, pp.19. http://www.ngi.no/upload/Projektweb/Biochar/NGI%20report%20no%2020100920-00-6-R%20End%20report%20Phase1_15%20Okt1_web.pdf. (accessed on 14th March 2013).

- David, H. S. (1999). Ash behavior during combustion and gasification, Volume 24 of Report (IEA Coal Research). pp. 38.
- Demirbas, A. (2001). Carbonization ranking of selected biomass charcoal, liquid and gaseous products. *Energy Conversion and Management*. 42:1229-1238.
- Department of Agriculture. (2011). Principal Statistics of Paddy and Rice by All Seasons, 2007-2011.
http://www.doa.gov.my/c/documentlibrary/get_file?uuid=35c14631-ca10-4a26-a191-fdd64708dce4&groupId=38371 (accessed on 15th Nov 2012)
- Department of Agriculture. (2012). Purata Harga Runcit Beberapa Jenis Baja di Malaysia Bagi Tahun 2012.
- Dias, B. O., Silva, C., Higashikawa, F. S., Roig, A. and Sánchez-Monedero, M. (2010). Use of biochar as bulking agent for the composting of poultry manure: effect on organic matter degradation and humification. *Bioresource Technology*, 101, 1239–1246.
- Downie, A., Crosky, A. and Munroe, P. (2009). *Physical properties of biochar*. In *Biochar for Environmental Management: Science and Technology*, ed. J. Lehmann and S. Joseph, (pp. 13-32). Earthscan, London, United Kingdom.
- Downie, A., Klatt, P., and Munroe, P. (2007). Slow pyrolysis: Australian demonstration plant successful on multi-feedstocks. Jyvaskyla, Finland, 3-6 September. The Bioenergy Association of Finland.
- Dunisch, O., Lima, V.C., Seehann, G., Donath, J., Montoia, V.R. and Schwarz, T. (2007). Retention properties of wood residues and their potential for soil ameliorant. *Wood Science and Technology*. 41:169-189.
- Eiland, F., Klamer, M., Lind, A., Leth, M. and Baath, E. (2001). Influence of initial C/N ratio on chemical and microbial composition during long term composting of straw. *Microbial Ecology*.41:272–280.
- Eghball, B., Power, J.F., Gilley, J.E. and Doran, M. (1997). Nutrient, carbon and mass loss during composting of beef cattle feedlot manure. *Journal Environment Quality*.126: 189-193.
- Eklind, Y. and Kirchmann, H. (2000). Composting and storage of organic household waste with different litter amendments. II: Nitrogen turnover and losses. *Bioresource Technology*. 74:125–133. doi: 10.1016/S0960-8524(00)00005-5.
- Errikson, K.E.L., Blanchette, R.A. and Ander, P. (1990). *Microbial and Enzymatic Degradation of Wood and Wood Components*. Springer, Berlin, Germany. pp.112-132.

- Fadelah, A. A. (1992). Penyediaan tanah dan pengurusan ladang, In Chapter 4, *Panduan Pengeluaran Sayur-Sayuran*, Institut Penyelidikan dan Kemajuan Pertanian Malaysia. pp. 23-28.
- FAO. (2011). Food and agricultural organization of United Nation, FAOSTAT, <http://faostat.fao.org/site/339/default.aspx> (accessed on 17th Jan 2013).
- FAO. (2002). Statistical database. Available at <http://apps.fao.org>
- Fernandes, L. and Sartaj, M. (1997). Comparative study of static pile composting using natural, forced and passive aeration method. *Compost Science and Utilization*. 5(4):65-67.
- Fredriksson, J. (1999). Exploratory experimental and theoretical studies of cyclone gasification of wood powder. Doctoral thesis, ISSN 1402-1544, Lulea University of Technology, Sweden.
- Gajalakshmi, S., and Abbasi. S.A. (2008). Solid waste management by composting: state of the art. *Critical Reviews in environmental Science and Technology* 38.5:311-400.
- Giltrap, D.L., Mckiblin, R. and Barnes G.R.G. (2003). A steady state model of gas char reactions in a down draft biomass gasifier. *Solar Energy*. 74: 85-91.
- Glaser, B. and Birk, J. J. (2011). State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio). *Geochimica Et Cosmochimica Acta* doi:10.1016/j.gca.2010.11.029.
- Glaser, B., Balashov, E., Haumaier, L., Guggenberger, G. and Zech, W. (2000). Black carbon in density fractions of anthropogenic soils of the Brazilian Amazon region. *Organic Geochemistry*. 31:669–678.
- Glaser, B. (2007). Prehistorically modified soils of central Amazonia: a model for sustainable agriculture in the twenty-first century. *Philosophical Transactions of the Royal Society B-Biological Sciences*. 362: 187-196.
- Glaser, B., Lehman, J. and Zech, W. (2002). Ameliorating physical and Chemical properties of highly weathered soil in tropics with charcoal: a review. *Biology Fertile Soil*. 35 : 219-230
- Guerrero, M., Ruiz, M.P., Millera, A., Alzueta, M. and Bilbao, R. (2008). Characterization of biomass chars formed under different devolatilization conditions: Differences between rice husk and eucalyptus. *Energy and Fuels*. 22: 1275-1284.
- Helyar, K. R. (1976). Nitrogen cycling and soil acidification. *J. Aust. Inst. Agric. Sci.* 42: 217-21.

- Hockaday, W. C., Grannas, A.M., Kim, S. and Hatcher, P.G. (2007). The transformation and mobility of charcoal in a fire impacted water shed. *Geochimica et Cosmochimica Acta*. 71: 3432-3445.
- Horwitz, W and Latimer, G.W. (2007). Ash of plants method 930.05. Official Methods of Analysis of AOAC International, AOAC-Int. Gaithersburg, MD, USA.
- Hua, L., Wu, W., Liu, Y., McBride, M.B. and Chen, Y. (2009). Reduction of nitrogen loss and Cu and Zn mobility during sludge composting with bamboo charcoal amendment. *Environmental Science Pollution Resources*. 16:1-9.
- Hugh, M.L., Paul, S. A., Frank E. S. and Thomas B. R. (2009). All Biochars are Not Created Equal, and How to Tell Them Apart Version 2, which supercedes the digital reprint issued at the North American Biochar Conference, Boulder, CO – August 2009.
- Igarashi, T. (1996) Soil Improvement effects of FMP & CRH in Indonesia. JICA Pamphlet. pp.30
- International Biochar Initiative. (2012). Standardized Product Definition and Product Testing Guidelines for Biochar that is Used in Soil, International Biochar Initiative 2012. Available online: http://www.biocharinternational.org/sites/default/files/guidelines_for_biochar_that_is_used_in_soil_final.pdf (accessed on 14 May 2013)
- ISO 11277. (1998). Soil quality -- Determination of particle size distribution in mineral soil material -- Method by sieving and sedimentation. International Organization for Standardization (ISO).
- Iwahashi, M., Tachibana, Y. and Ohta, Y. (1982). Accumulation of calcium, magnesium, potassium and sodium with growth of individual leaves, petioles, and stems of cucumber plants. *Soil Science Plant Nutrition*. 28:441-449.
- Jackson, R.B., Mooney, H.A. and Schulze, E.D. (1997). A global budget for fine root biomass surface area and mineral nutrients. *Proc. Natl. Acad. Sci. USA*., 94: 7360-7366.
- Jain, A. (2006) Design parameters for a rice husk throatless gasifier. *Agricultural Engineering International: the CIGR Ejournal*, Manuscript EE 05 012. Vol VIII.
- Jindo, K., Suto, K., Matsumoto, K., Garcia, C., Sonoki, T. and Sanchez-Monedero, M.A. (2012). Chemical and biochemical characterisation of biochar-blended composts prepared from poultry manure. *Bioresource Technology* 110:396-404
- Jones, J. B., Wolf Jr, B. and Mills, H. (1991). Plant analysis handbook, Micro-Macro Publishing, Inc., Athens, GA, pp. 23-26

- Jonker, M.T. and Koelmans, A.A. (2002). Sorption of polycyclic aromatic hydrocarbons and polychlorinated biphenyls to soot and soot like materials in the aqueous environment: mechanistic consideration. *Environmental Science Technology* 36:3725-3734.
- Kaiser, K. and Guggenberger, G. (2000). The role of DOM sorption to mineral surfaces in the preservation of organic matter in soils. *Organic Geochemistry*. 31:711-725.
- Kannan, N. and Sundram, M.M. (2001). Kinetics and Mechanism of removal of Methylene Blue by adsorption on various carbons - a comparative study. *Dyes and pigments*. 51:25-40.
- Karkamar, S., Mitra, B.N. and Gosh, B.C. (2009). Influence of industrial solid waste on soil-plant interaction in rice under acid Lateritic soils. Paper presented at World of Coal Ash (WOCA) Conference, 4 – 7 May 2009. Lexington, USA.
- Karlsson, S. and Jeppson, K.H. (1995). Deep litter in livestock buildings and field storage, Legends and Summaries in English, JTI Rapport 204, Swedish Institute of Agricultural Engineering, Ultuna Uppsala, Sweden.
- Kleegberg, K.K., Schlegelmich, M., Stress, J., Steinhart, H. and Stegmann, R. (2005). Odour abatement strategy for a sustainable odour management, In Proceeding of Tenth International Waste Management and Landfill Symposium, Sardinia, 3-7 October 2005 by CISA, Environmental Sanitary Engineering Centre, Italy
- Kwon, S. and Pignatello, J.J. (2005). Effect of natural organic substances on the surface and adsorptive properties of environmental black carbon (char): Pseudo pore blockage by model lipid components and its implication for N₂-probed surface properties of natural sorbents. *Environmental Science and Technology*. 39:7932-7939.
- Laird, D. A. (2008). The Charcoal Vision: A Win-Win-Win Scenario for Simultaneously Producing Bioenergy, Permanently Sequestered Carbon, while Improving Soil and Water Quality. *Agronomy Journal* 100:178-181.
- Lal, R. (2006). Encyclopedia of Soil Science, CRC Press, Boca Raton, FL
- Lambers, H. (1987). Growth Respiration Exudation and Symbiotic Association the Fate of Carbon Translocated to Roots. In: *Root Development and Function*, Greosy, P.J., J.V. Lake and D.A. Rose (Eds.). Cambridge University Press, London, UK.
- Lehmann, J. and Joseph, S. (2009). *Biochar for Environmental Management: Science and Technology*. Earthscan, London.

- Lehman, J. (2007). 'Bioenergy in the black' frontiers in ecology and the environment. 5:381-387. Department of Crop and Soil Sciences, College of Agriculture and Life Sciences, Cornell University, Ithaca, NY 14853 (CL273@ cornell.edu).
- Lebo, J. A., Huckins, J. N., Petty, J.D., Cranor, W.L. and Ho, K.T. (2003). Comparisons of coarse and fine versions of two carbons for reducing the bioavailability of sediment- bound hydrophobic organic contaminants. *Chemosphere*. 50:1309-1317.
- Leinweber, P. and Schulten, H.R. (1999). Advances in analytical pyrolysis of soil organic matter. *Journal of Analytical and Applied Pyrolysis*. 49:359-383.
- Liang, B., Lehmann, J., Solomon, D., Sohi, S., Thies, J.E., Skjemstad, J.O., Luizao, F.J., Engelhard, M.H., Neves E.G. and Wirick, S. (2008). Stability of biomass-derived black carbon in soils. *Geochimica et Cosmochimica Acta*. 72:6069-6078.
- Liang, B., Lehmann, J., Peterson, J. and Neves, E.G. (2006). Black carbon increases cation exchange capacity in soils. *Soil Science Society of American Journal*. 45:71-672.
- Lua, A.C., Yang, T. and Guo, J. (2004). Effects of pyrolysis conditions on the properties of activated carbons prepared from pistachio-nut shells. *Journal of Analytical and Applied Pyrolysis*. 72:279-287.
- Maher, M. J. (1998). The effect of type and rate of controlled release fertiliser on the performance of hardy nursery stock in containers. Combined Proceedings of the International Plant Propagators Society, 46:178-183.
- Maiti, S., Dey, S., Purakayastha, S. and Ghosh, B. (2005). Physical and thermochemical characterization of rice husk char as potential biomass energy source. *Bioresource Technology*. 97:2065-2070.
- Masulili, A. (2010). Rice Husk Biochar for Rice Based Cropping System in Acid Soil 1 . The Characteristics of Rice Husk Biochar and Its Influence on the Properties of Acid Sulfate Soils and Rice Growth in West Kalimantan, Indonesia, *Journal of Agriculture Science*. 2:139-47.
- Melgar, R.J., Smyth, T.J., Sanchez, P. A. and Cravo, M.S. (1992). Fertilizer nitrogen movement in a central Amazon Oxisol and Entisol cropped to maize. *Fertilizer Research*. 31:241-252.
- Mosier, A.R., J.K. Syers and J.R. Freney. (2004). *Agriculture and the Nitrogen Cycle. Assessing the Impacts of Fertilizer Use on Food Production and the Environment*. Scope-65. Island Press, London.

MS677: Pt I to VIII: 1980, Part II: Ashing and Preparation of Ash solution, Recommended Methods for Plant Chemical Analysis, SIRIM

MS677: Pt I to VIII: 1980, Part IV: Determination of Phosphorus, Recommended Methods for Plant Chemical Analysis, SIRIM

MS677: Pt I to VIII: 1980, Part V: Determination of Potassium, Recommended Methods for Plant Chemical Analysis, SIRIM

Martin, H. (2005). Manure composting as a pathogen reduction strategy. Factsheet, Ministry of Agriculture and Food, Ontario, Order No: 05-021. <http://www.omafra.gov.on.ca/english/engineer/facts/05-021.pdf> (accessed on 31st Dec 2013)

Martin, D.L. and Gershuny eds. G. (1992). The Rodale Book of Composting. Rodale Press. Emmaus, PA.

Mukunda, H. S., Shrinivasa, U., Paul, P. J., Dasappa, S. and Rajan, N. K. S. (2002). Indian Institute of Science, Bangalore, 2002. <http://fmb.no/arnt/elspa/stand-alone.html> (accessed on 12th Jan 2013).

Munshower, F.F. (1994). Practical Handbook of Disturbed Land Revegetation. CRC Press, Inc., Boca Raton, Florida, USA.

Natarajan, E., Nordin, A. and Rao, A. N. (1998). Overview of combustion and gasification of rice husk in fluidized bed reactors. *Biomass Bioenergy*. 14: 533-546.

Netto, A. T., Eliemar, C., Jurandi, G. O. and Ricardo E.B. (2005). Photosynthetic pigments, nitrogen, chlorophyll a fluorescence and SPAD 502 readings in coffee leaves. *Scientia Horticulturae*. 104:199–209.

Neumann, G. and Martinoia, E. (2002). Survival cluster roots an underground adaptation for survival in extreme environment. *Trends Plant Science*. 7: 162-167.

Nguyen, T.H., Brown, R. A. and Ball, W.P. (2004). An evaluation of thermal resistance as a measure of black carbon content in diesel soot, wood char and sediment. *Organic Geochemistry*. 35:217-234.

Odette V.M. and Wu, J. H. (2010). Effects of rice husk biochar under slow pyrolysis on growth of water spinach, presentation slides, International symposium on environmental behaviour and effects of biomass derived carbon. National Pingtung University of Science and Technology Pingtung, Taiwan. http://www.biochar-international.org/sites/default/files/Odette_Varela.pdf (accessed on 11th Oct 2012).

- Ogawa M. (1991). Effective utilization of charcoal as a material for soil amendment. *AICAF Expert Bulletin* 12(3):1-13.
- Ogawa, M. (1999). Utilization of symbiotic microorganisms and charcoal in tropical agriculture and forestry and CO₂ fixation. *Soil Microorganisms*. 53:73-79.
- Oka, H. (1993). Improvement of sandy soil in the northeast by using carbonized rice husks. *JICA Technical Report* 13: 42-40.
- Ott, S. H. and Vogtmann, H. (1983). Nitrates in relation to composting and use of farmyard manures. p. 145-154. In *Environmentally Sound Agriculture*. 4th Conference of International Federation of Organic Agriculture Movements, Proeger, New York, 1983.
- Padi Beras Nasional Berhad. (2011). Rice Pedia, Standard sequence of rice processing. http://www.bernas.com.my/index.php?option=com_content&view=article&id=90&Itemid=103 (accessed on 2nd Feb 2013).
- Paramanathan, S. (2000). Soils of Malaysia: Their characteristic and identification. 1st ed., Academy of Science Malaysia and Param, Agriculture Soil Science Surveys, p. 616.
- Parr, J. F., Epstein, E. and Willson, G.B. (1978). Composting sewage sludge for land application, *Agriculture Environmental*. 4:123-137.
- Pastor- Villeges, J., Pastor -Valle, J.F., Meneses Rodriguez, J. M. and Garcia, M. (2006). Study of commercial wood charcoals for the preparation of carbon adsorbents. *Journal of Analytical and Applied Pyrolysis*. 76:103-108.
- Pietikainen. T., Kiikkila, O. and Fritze, H. (2000). Charcoal as habitat for microbes and its effects on the microbial community of the underlying humus. *Oikos*. 89: 231-242.
- Ray, O. G. and Rush., R. (2007). *The Terra Preta phenomenon*, Managing the carbon cycle. Katanning Workshop 21-22 March 2007. www.amazingcarbon.com.
- Rajvir, S., Maheshwari, R.C. and Ojha, T.P. (1980). Development of Husk fired furnace. *Journal of Agriculture Engineering Resources*. 25:109-120.
- Richards, L.A. (1947). Pressure- membrane apparatus-construction and use. *Agric. Eng.* 28: 451-454.
- Rynk, R.M., van de Kamp, G.B., Willson, M.E., Singley, T. L., Richard, J.J., Kolega, F.R., Gouin, L., Laliberty Jr, D., Kay, D.W., Murphy, H.A.J., Hoitink and Brinton, W.F. (1992). *On Farm Composting*. Northeast Regional Agric. Eng. Service, Ithaca, NY.

- Rouquerol, F., Rouquerol, I. and Sing, K. (1999). Adsorption by Powders and Porous solids. Academic Press, London UK.
- Rozainee, M., Ngo, S. P., Salema, A.A. and Tan, K.G. (2010). Effect of feeding method on rice husk ash quality in a fluidised bed combustor. *Emirates Journal for Engineering Research*. 15:1-12.
- Rozainee, M. (2007). Production of rice husk amorphous silica form fluidized bed system. Research Vot No. 74526, Universiti Teknologi Malaysia.
- Rufatullah, M., Sulaiman, O., Hashim, R and Ahmad, A. (2010). Adsorption of methylenen blue on low cost adsorbents: A review, *Journal of Hazardous Materials*. 177(1-3): 70-80.
- Saowakan, H. and George C. (2011). *Charcoal amendments to improve soil fertility and rice production in North East Thailand*, FSC Brief No.3, Food security Center, Germany, pp 1-4.
- Shackley, S., Carter, S., Knowles, T., Middleink, K., Haefele, S., Sohi, S., Cross, A. and Haszeldine, S. (2012). Sustainable gasification–biochar systems. A case-study of rice-husk gasification in Cambodia, Part I: Context, chemical properties, environmental and health and safety issues. *Energy Policy* (42): 49-58.
- Shackley, S. and Sohi, S.P. (2010) An Assessment of the Benefits and Issues Associated with the Application of Biochar to Soil, Report to the Department for Environment, Food and Rural Affairs and the Department of Energy and Climate Change: London, UK, 2010, pp. 14–132.
- Sing, K.S.W., Everett D.H., Haul R.A.W., Moscou L., Pierotti R.A. and Rouqueroul, T.S.(1985). Reporting physisorption data gas/solid systems with special reference to the determination of surface area and porosity. *Pure and Applied Chemistry*. 57:603-619.
- Song, H., Xiang, J., Sun, L., Minhau X, Jianny Q and Peng F. (2008). Characterization of char from rapid pyrolysis of rice husk. *Fuel Processing Technology*. 89:1096-1105.
- Steiner, C., Das K.C., Melear, N. and Lakly, D.(2010). Reducing nitrogen loss during poultry litter composting using biochar. *Journal of Environmental Quality*. 39:1236-1242.
- Steiner, C., Teixeira, W.G., Lehmann, J., Nehls, T., De Macêdo, J. L. V., Blum, W.E.H. and Zech, W. (2007). Long term effects of manure, charcoal and mineral fertilization on crop production and fertility on a highly weathered Central Amazonian upland soil. *Plant and Soil*. 291(1): 275-290.

- Steiner, C., Teixeira W.G., Lehman J and Zech W. (2004). *Microbial Response to Charcoal Amendment of Highly Weathered Soils and Amazonian Dark Earth: Explorations in Time and Space*. Springer, Berlin, Germany, pp.195-212.
- Sogbedji, J. M., Van es, H.M., Klausner, S.D., Bouldin, D.R. and Cox, W.J. (2001). Spatial and temporal proceses affecting nitrogen availability at the landscape scale. *Soil and Tillage Research*. 58:233-244.
- Thomas, G.W. 1982. Exchangeable cations. In A.L. Page (ed.). *Methods of soil analysis, Part 2, Second Edition, Agronomy Monograph 9, American Society of Agronomy, Madison, WI*.
- Thies, E. and Rillig, M. C. (2009). Characteristics of Biochar : Biological properties, In Chapter 6 in *Biochar Environmental Management, Science and Technology*, Earthscan pp 86-105.
- Tiquia, S.M., Richard, T.L. and Honeyman, M.S. (2000). Carbon, nutrient, and mass loss during composting. *Journal of Nutrient Cycling in Agroecosystem* 62: 15-24.
- Trautmann, N.M. and Krasny. M. E (1997). Compost Properties. In Chapter 5: *Composting in Classroom* (pp 70-92). Cornell University.
- Tsai, W.T., Lee, M.K. and Chang, Y. (2007). Fast pyrolysis of rice husk: product yield and composition. *Bioresources Technology*. 98: 2-28.
- Tseng, R. L. and Tseng, S. K. (2006). Characterization and use of high surface area activated carbons prepared form cane pith for liquid-phase adsorptions. *Journal of Hazardous Materials*. B136: 671-680.
- Tuaprakone, T., Wongphaet, N., and Wasanapiarnpong, T. (2011). Fabrication of Activated Rice Husk Charcoal by Slip Casting as a Hybrid Material for Water Filter Aid. *IOP Conference Series: Materials Science and Engineering*, 18(18), 182009. <http://doi:10.1088/1757-899X/18/18/182009> (accessed on 16th Feb 2013).
- Tryon, E. H. (1948). Effect of charcoal on certain physical, chemical, and biological properties of forest soils. *Ecological Monographs* 18:81–115. <http://dx.doi.org/10.2307/1948629> (accessed on 11th Jan 2013).
- Van Krevelen, D.W. (1950). Graphical-statistical method for the study of structure and reaction processes of coal. *Fuel*. 29:269-84.
- Vimala, P. and Chan, K. (1992). Tanah dan Pembajaan, In Chapter 5, *Panduan Pengeluaran Sayur- Sayuran* (pp 29-40). Institut Penyelidikan dan Kemajuan Pertanian Malaysia.
- Walker, J.M. (1993). Control of composting odors. In H.A.J. Hoitink and H.M. Keener (eds). In *Science and Engineering of Composting: Design*,

Environmental, Microbiological and Utilization Aspects, (pp 185-218). Worthington, Ohio. USA: Renaissance Publications.

Wander, M.M., Traina, S.J., Stinner, B.R. and Peters, S.E. (1994). Organic and conventional management effects on biologically active soil organic matter pools. *Soil Science Society of America Journal*. 58:1130-113.

Warnock, D.D., Lehmann, J., Kuyper, T.W and Rillig, M.C. (2007). Mycorrhizal response to biochar in soil- concept and mechanisms. *Plant and Soil*. 300: 9-20.

Wiederholt, R., Franzen, D. and Johnson, B.(2005). Livestock manure utilization in non-till cropping system, Report by North Dakota State extension Service. Pp.4.

Wildman, J. and Derbyshire, F. (1991). Origins and functions of macroporosity in activated carbons form coal and wood precursors. *Fuel*. 70:655-661.

Wollum II, A.G. (1985). Cultural methods for soil microorganisms. In *Methods of soils analysis Part 2. Chemical and Microbiological Properties Second Edition*, A.L Page et al (ed.) *American Society of Agronomy ., Inc ., Madison, Wisconsin*. 37:781-802.

Wong C.L., Venneker R., Uhlenbrook S., Jamil A.B.M and Zhou Y. (2009). Variability of rainfall in Peninsular Malaysia. *Hydrology Earth System Science Discussion*. 6:5471–5503

Yoshizawa, S., Tanakan, S. and Ohata, M. (2007). Proliferation effects of aerobic microorganisms during composting of rice bran by addition of biomass charcoal, In *Proceeding of the International Agrichar Conference, Terrigal , Australia, May 2007, 26. www.iaiconference.org*.

Yu, X. Y., Ying, G.G. and Kookana, R. (2006). Sorption and desorption behaviors of diuron in soils amended with charcoal. *Journal of Agriculture and Food Chemistry*. 54:8545-8550.

Zhang, X., Rojas, J.C. and Gonzalez-Lima, F. (2006). Methylene blue prevents neurodegeneration caused by rotenone in the retina. *Neurotox Res* 9: 47–57. doi: 10.1007/BF03033307.