UNIVERSITI PUTRA MALAYSIA

BIOLOGY AND DEVELOPMENT OF PROTOCOLS FOR CULTURING SOIL-DWELLING ENDOGEIC WORM Pontoscolex corethrurus MÜLLER FOR POTENTIAL USE IN SOIL REHABILITATION

KOK HO YIN

FS 2014 26

BIOLOGY AND DEVELOPMENT OF PROTOCOLS FOR CULTURING SOIL-DWELLING ENDOGEIC WORM*Pontoscolex corethrurus*MÜLLER FOR POTENTIAL USE IN SOIL REHABILITATION

By

KOK HO YIN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfillment of the Requirements for the Degree of Doctor of Philosophy

May 2014

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

BIOLOGY AND DEVELOPMENT OF PROTOCOLS FOR CULTURING SOIL-DWELLING ENDOGEIC WORM *Pontoscolex corethrurus* MÜLLER FOR POTENTIAL USE IN SOIL REHABILITATION

By
KOK HO YIN
May 2014

Chairman : Nor Azwady Abd. Aziz, PhD
Faculty : Science

Pontoscolex corethrurus (Müller, 1857) is the dominant tropical soil-dwelling endogeic worm and has the ability to live in a variety of soils differing in pH and organic matter content. This suggestedthat this species may serve as a potential agent for soil rehabilitation particularly in the tropics. Unfortunately, the relevant information about *P. corethrurus* is very limited in Malaysia. A detailed understanding of life history of *P. corethrurus* is required because the biological characteristics of earthworms (e.g., growth and reproduction rates) could affect their potential in soil rehabilitation programs. The rates of growth and reproduction of *P. corethrurus*on different stocking densities were also evaluated to determine the optimum stocking density for culturing this species. This study discovered that the optimum stocking for culturing *P. corethrurus*was around 200 earthworms per m^2. This study also reported that *P. corethrurus* has a high survival rate and has the capability to reproduce *via* parthenogenesis. These biological characteristics enable *P. corethrurus* to colonize new environments hence, increasing its potential for the use in worm inoculation activities. In order to obtain a large and consistent supply of earthworms for worm inoculation activities in soil rehabilitation, indoor culturing of *P. corethrurus* is crucial. However, there are difficulties in using natural soil as the medium for culturing endogeic worms due to the heterogeneous nature of soils in different localities that will surely affect the culture performance. Thus, a protocol was developed using artificial soil to replace nature soil for culturing *P. corethrurus*. This study found that clay content can affect the growth of *P. corethrurus* and the manipulation of the clay content in artificial medium (consists of kaolinite clay, industrial quartz sand and decomposed cow dung) could result it as an alternative medium for culturing *P. corethrurus*. Furthermore, the properties of earthworm surface cast, worm-worked soil and bulk soil (soil without earthworm inoculation) were analyzed to study the influence of *P. corethrurus* activities on soil properties. Earthworm surface cast and worm-worked soil recorded significantly higher amounts of available nutrients than bulk soil. Soil enzymes (protease and acid phosphatase) activities, microbial (bacteria) population and humic acid content were also significantly

higher in earthworm cast than in bulk soil. A subsequent polybag-plant experiment was conducted to determinehow the presence of *P. corethrurus* in soil could benefit plant productivity. With the application of organic amendment, the inoculation of *P. corethrurus* into the planting medium (+E+O treatment) significantly increased plant height (20%), root (27%) and shoot (37%) dry weights as well as crop yield (15%) compared to treatment without earthworm inoculation (–E+O treatment). The present study suggested that the tropical soil-dwelling endogeic worm *P.corethrurus* can be considered as a good soil rehabilitation agent for rehabilitating soil and increasing plant productivity.

Abstract tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluaan untuk ijazah Doktor Falsafah

BIOLOGI DAN PEMBANGUNAN PROTOKOL PENGKULTURAN CACING TANAH ENDOGEIK, *Pontoscolex corethrurus* MÜLLER YANGBERPOTENSI DALAM PEMULIHAN TANAH

Oleh
KOK HO YIN
Mei 2014

Pengerusi : Nor Azwady Abd. Aziz, PhD
Fakulti : Sains

Pontoscolex corethrurus (Müller, 1857) adalah cacing tanah endogeik tropika dominan dan mempunyai keupayaan untuk hidup dalam pelbagai jenis tanah yang berbeza dalam pH and kandungan bahan organik. Ini mencadangkan bahawa spesies ini boleh bertindak sebagai ejen yang berpotensi dalam pemulihan tanah terutamanya di kawasan tropika.Malangnya, maklumat berkaitan dengan *P. corethrurus* adalah sangat terhad di Malaysia.Satu pemahaman yang terperinci tentang kitaran hidup *P. corethrurus* adalah diperlukan kerana ciri biologi cacing tanah (contohnya, kadar pertumbuhan dan pembiakan) boleh mempengaruhi potensi cacing dalam program pemulihan tanah.Kadar pertumbuhan dan pembiakan *P. corethrurus* pada kepadatan yang berbeza juga dinilai untuk menentukan kepadatan yang optimum untuk pengkulturan spesies ini.Kajian ini menunjukkan bahawa kepadatan optimum untuk pengkulturan *P. corethrurus* adalah kira-kira 200 cacing tanah per m^2. Kajian ini juga menunjukkan bahawa *P. corethrurus* mempunyai kadar kemandirian yang tinggi dan bersifat parthenogenetik. Ciri biologi ini membolehkan *P. corethrurus* menguasai persekitaran baru dan meningkatkan potensinya dalam aktiviti inokulasi cacing. Pengkulturan *P. corethrurus*adalah pentinguntuk menghasilkan bekalan cacing tanah yang banyak dan konsisten untuk aktiviti inokulasi cacing. Penggunaan tanah semulajadi sebagai media kultur cacing kurang sesuai disebabkan sifat tanah yang berbeza di sesuatu kawasan akan mempengaruhi keputusan pengkulturan. Maka, protokol telah dibangunkan dengan menggunakan media buatan bagi menggantikan tanah semulajadi dalam pengkulturan *P. corethrurus*. Kajian ini mendapati bahawa kandungan tanah liat boleh menjejaskan pertumbuhan *P. corethrurus* dan memanipulasi kandungan tanah liat dalam media buatan (terdiri daripada tanah liat kaolinit, pasir dan tahi lembu reput) boleh dijadikan media alternatif dalam pengkulturan *P. corethrurus*.Sifat tinja cacing, tanah dengan kehadiran cacing dan tanah tanpa cacing juga telah dianalisa untuk mengkaji pengaruh aktiviti *P. corethrurus* ke atas tanah. Kajian ini mendapati tinja cacing dan tanah dengan kehadiran cacing mengandungi kandungan nutrien yang lebih tinggi berbanding dengan tanah tanpa cacing. Aktiviti enzim (protease dan asid fosfatase), populasi mikrob

(bakteria) dan kandungan asid humik dalam tinja cacing juga lebih tinggi daripada tanah tanpa cacing. Seterusnya,eksperimen polibeg-tumbuhan dijalankan untuk menentukan sama ada kehadiran *P. corethrurus* dalam tanah boleh memberi manfaat kepada produktiviti tanaman.Dengan pembajaan,inokulasi *P. corethrurus* ke dalam tanah (+E+O)meningkatkan tinggi tumbuhan (20%), berat kering akar (27%) dan pucuk (37%) serta peningkatan hasil tanaman (15%) berbanding dengan rawatan tanpa kehadiran cacing(–E+O). Kajian ini mencadangkan cacing tanah endogeik tropika *P. corethrurus* boleh berfungsi sebagai ejen pemulihan tanah yang baik untuk memulihkan tanah dan meningkatkan productiviti tanaman.

ACKNOWLEDGEMENTS

Firstly, I would like to express my deepest appreciation to my Supervisor, Dr. Nor Azwady Abd. Aziz for his supervision, advice on research methodology, the art of scientific writing, and his valued support and friendship throughout this study. Without his guidance and persistent help, this thesis would not have been possible. Thanks should also be extended to my Co-Supervisors, Associate Professor Dr. Muskhazli Mustafa and Dr. Syaizwan Zahmir Zulkifli, for their guidance and valuable suggestions.

Special thanks to Encik Baharuddin and Puan Zaleha from Organic and Aroma Section, Taman Pertanian Universiti (TPU), for their assistance and friendship throughout the field experiment of my study. I would also like to thank all the Laboratory Assistants in the Department of Biology for their help and encouragement.

In addition, I would like to take this opportunity to record my thanks to my fellow postgraduate students in the Department of Biology, especially Khye Er, Suk Kuan and Yi Wei from Vermitechnology Laboratory for their help and unselfish sharing of knowledge and skills in overcoming problems that arose in my study.

Lastly, I would like to express my heartfelt thanks to my beloved family,especially my dad for his patience and moral support during my days at the university.

I certify that a Thesis Examination Committee has met on 8 May 2014 to conduct the final examination of Kok Ho Yin on her thesis entitled "Biology and Development of Protocols for Culturing Soil-Dwelling Endogeic Worm *Pontoscolex corethrurus* Müller for Potential Use in Soil Rehabilitation" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Shamarina binti Shohaimi, PhD
Senior Lecturer
Faculty of Science
Univeristi Putra Malaysia
(Chairman)

Hishamuddin bin Omar, PhD
Senior Lecturer
Faculty of Science
Univeristi Putra Malaysia
(Internal Examiner)

Mohamed Hanafi bin Musa, PhD
Professor
Faculty of Agriculture
Univeristi Putra Malaysia
(Internal Examiner)

José Adolfo Amador, PhD
Professor
University of Rhode Island
United States
(External Examiner)

NORITAH OMAR, PhD
Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 July 2014

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Nor Azwady Abd Aziz, PhD
Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Muskhazli Mustafa, PhD
Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Syaizwan Zahmir Zulkifli, PhD
Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

_____
BUJANG KIM HUAT, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: 7July 2014

Name and Matric No.: Kok Ho Yin (GS24393)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman of Supervisory Committee: Dr. Nor Azwady Abd Aziz
Senior Lecturer
Faculty of Science
Universiti Putra Malaysia

Signature: _____
Name of Member of Supervisory Committee: Assoc. Prof. Muskhazli Mustafa
Associate Professor
Faculty of Science
Universiti Putra Malaysia

Signature: _____
Name of Member of Supervisory Committee: Dr. Syaizwan Zahmir Zulkifli
Lecturer
Faculty of Science
Universiti Putra Malaysia

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

LIST OF PLATES

LIST OF ABBREVIATIONS

AA	Actinomycetes Isolation Agar
Al	Aluminium
ANOVA	Analysis of Variance
AS	Artificial Soil with 70% Industrial Quartz Sand, 20% Kaolinite Clay and 10% Decomposed Cow Dung
$AS_{30\%clay}$	Artificial Soil with 60% Industrial Quartz Sand, 30% Kaolinite Clay and 10% Decomposed Cow Dung
$AS_{40\%clay}$	Artificial Soil with 50% Industrial Quartz Sand, 40% Kaolinite Clay and 10% Decomposed Cow Dung
$AS_{50\%clay}$	Artificial Soil with 40% Industrial Quartz Sand, 50% Kaolinite Clay and 10% Decomposed Cow Dung
C	Carbon
Ca	Calcium
CEC	Cation Exchange Capacity
cm	Centimeter
Fe	Iron
g	Gram
IBS	Institute of Bioscience
ICP-MS	Inductively Coupled Plasma-Mass Spectrometry
h	Hour
ha	Hectare
K	Potassium
K_2O	Potash
kg	Kilogram
L	Liter

m	Meter
m^2	Square Meter (area)
MARDI	Malaysian Agricultural Research and Development Institute
Mg	Magnesium
mg	Miligram
mL	Mililiter
mm	Milimeter
n	Number of Replicates
N	Nitrogen
NA	Nutrient Agar
NH_4-N	Ammonium
NO_3-N	Nitrate
OECD	Organization for Economic Co-operation and Development
P	Phosphorous
P_2O_5	Phosphorous Pentoxide
ppm	Part per Million
RBA	Rose Bengal Chloromphenicol Agar
SPSS	Statistical Package for Social Science
rpm	Revolutions per Minute
µg	Microgram
µmole	Micromole
TPU	Tanam Pertainan Universiti
UPM	Univeristi Putra Malaysia
V	Volume
Wt	Weight

WUE	Water Use Efficiency
%	Percentage

CHAPTER 1

INTRODUCTION

Malaysia with the tropical climate has a fairly high, but uniform temperature and large amounts of rain delivering in intense bursts each year. Because of the weather, local agricultural soils contain low organic matter and highly compact soil texture resulting in lower crop productivity (Sabtu, 2006; APEC, 2012). A greater amount of chemical fertilizers is used each year to maintain soil fertility and boost crop productivity. However, the use of chemical fertilizers only restores soil nutrient content and does not improve the soil's physical structure and microbiological activities. Long-term use of chemical fertilizers also results in many negative effects on the soil such as acidification, decline in organic matter content and reduction of microbial communities (Sinha *et al*., 2010). Therefore, the rehabilitation of local agricultural soils is necessary and important to reduce the loss of soil nutrients as well as to promote sustainable agriculture.

One of the approaches to create a healthy agro-ecosystem is introducing soil-dwelling endogeic worms into the soil (Butt *et al*., 1997). Endogeic worms are known as "ecological engineers" as their burrowing and casting activities result in beneficial chemical, microbiological and physical changes in the soil that increase the fertility of the soil (Jones *et al*., 1994; Scheu, 2003). Efforts have been made to transplant endogeic worms into degraded pastures in New Zealand which have significantly improved pasture productivity (Stockdill, 1982). In grasslands on reclaimed industrial cutaway peat in Ireland, herbage production was significantly improved by earthworm colonization through improvement in the physical structure of the soil (Curry and Boyle, 1987).

The use of earthworms to improve soil conditions is a common practice in temperate regions like New Zealand (Stockdill, 1982), The Netherlands (Hoogerkamp *et al*., 1983), Ireland (Curry and Boyle, 1987), and the United Kingdom (Butt *et al*., 1997). However, this approach is rarely practiced in the tropics including Malaysia. Available information on tropical earthworms is limited to litter-dwelling epigeic species while knowledge about the biology and ecology of tropical endogeic worms is very scarce. The differences between temperate and tropical climates are likely to influence the activities of earthworms in soils (Kale and Karmegam, 2010). The higher temperature and humidity in tropical ecosystems may increase the overall activities of earthworms in soils. The higher burrowing and casting activities may accelerate soil rehabilitation processes by improving the physicochemical and biological properties of soil and in turn create a healthy environment for plant growth. On the other hand, intense bursts of rain in tropical ecosystems may destroy earthworm surface casts. Nutrients within earthworm casts may therefore be lost through surface runoff (Jouquet *et al*., 2013). The overall effects of these processes may cause the effects of earthworms on soil properties

in tropical ecosystems to differ from those in temperate regions. More experimental research is necessary to determine the roles of tropical earthworms in their ecosystems.

Obtaining a sufficient number of earthworms for soil rehabilitation is a fundamental aspect to be achieved prior to the practice. Most earthworm inoculation activities conducted in temperate regions are accomplished *via* the mass collection of earthworms from densely populated areas and transplanted into lower density areas (Butt *et al*., 1995). This technique is expensive, labor-intensive and time-consuming (Butt *et al*., 1992). Earthworm collection sites must be amended with a lot of organic matter over a long period before further collection in order to maximize the number of earthworms present in the area (Butt, 2011). In addition, under natural conditions, earthworm availability is strongly affected by seasonal changes, organic resource distribution and farm management practices (Ghafoor *et al*., 2008; Pauli *et al*., 2010). Thus, it may be difficult to acquire a sufficient number of earthworms for inoculation purposes. An alternative method for obtaining a large number of earthworms is indoor cultivation (Butt *et al*., 1992). For example, successful production of epigeic worms such as *Eisenia fetida*, *Lumbricus rubellus*, *Perionyx excavatus*, and *Eudrilus eugeniae* under indoor conditions resulted in the widespread use of these species in vermicomposting, vermiculture for protein source and ecotoxicological studies.

Unlike epigeic worms, soil-dwelling endogeic worms are fairly difficult to cultivate indoors. Endogeic worms are geophagous deriving their nutrition through the consumption of large amounts of soil. They require culturing in a mineral soil-based medium (Butt and Lowe, 2011). Butt and his co-workers used soil collected from earthworm-sampling sites as the medium to culture endogeic worms (Butt *et al*., 1992; Butt, 1993; Lowe and Butt, 2005). However, there are difficulties in using natural soil as the medium for culturing endogeic worms. This is because field-collected soils are highly heterogeneous. Its texture, water-holding capacity, organic matter content, nutrient availability, and microbial activities differ on the basis of sampling locations, land management and plant communities (Huggins and Uberuaga, 2010). Differences in the physiochemical and biological properties of field-collected soils could affect the survival, growth, sexual maturity, and cocoon production rates of earthworms (Klok *et al*., 2007). Therefore, it is difficult to reach a considerable level of standardization with field-collected soil as the earthworm culture medium. Field-collected soils may also contain harmful compounds such as heavy metals, polycyclic aromatic hydrocarbons and pesticides that could adversely affect earthworm cultures. An alternative culture medium should be established in order to achieve standardization for indoor cultivation of endogeic worms and also to produce a large and consistent supply of earthworms for soil rehabilitation programs.

The soil-dwelling endogeic worm *Pontoscolex corethrurus* (Müller, 1857) is one of the few persistent species capable of surviving under tropical agricultural tillage conditions (Lavelle *et al*., 1987; Sabrina *et al*., 2009; Nath and Chaudhuri, 2010). In Malaysia, *P. corethrurus* is the only endogeic species found in local rubber and oil palm plantations

(Loh, 2009; Sabrina *et al*., 2009). It is widely distributed in a variety of soil types including Serdang series, Bungor series and Jawa series (Vijiandran and Gurmit, 2009). Previous studies have shown that adaptation to tropical climates, capability to tolerate a wide range of environmental factors (e.g., soil pH and organic matter content) and the ability to enter diapause under drought conditions are the adaptive strategies that enable *P. corethrurus* to be widely distributed in tropical ecosystems (Lavelle *et al.*, 1987; James and Hendrix, 2004; Chaudhuri *et al*., 2008; Nath and Chaudhuri, 2010). The fact that it is widely distributed in tropical ecosystems and shows good environmental plasticity suggests that *P. corethrurus* may serve as a potential candidate for soil rehabilitation in tropical regions.

Unfortunately, in Malaysia, other than ecological data (distribution and abundance), relevant information about *P. corethrurus* is very limited. A detailed study on the effects of *P. corethrurus* on soil chemical and biological properties as well as crop productivity is required to determine its role in local soil ecosystems. Prior to determining its effects on soil properties, fundamental aspects of earthworm biology (e.g., life history) need to be studied since the biological characteristics of earthworms will affect their potential in soil rehabilitation programs. Nevertheless, the life history of earthworms is strongly affected by earthworm stocking density (Hait and Tare, 2011; Suthar, 2012). To date, there is still no study on how stocking density will influence the life history of *P. corethrurus*. In addition, a suitable culture medium for culturing *P. corethrurus* should be established to provide a large and consistent supply of *P. corethrurus* for worm inoculation activities.

Therefore, the objectives of the present research were:

i. To determine the effect of stocking density on life history of the endogeic worm *P. corethrurus*,

ii. To establish a suitable medium for culturing the endogeic worm *P. corethrurus*,

iii. To investigate the effect of the endogeic worm *P. corethrurus* on nutrient availability, humic acid content, microbial population, and enzyme activity in soil,

iv. To evaluate the effect of inoculating the endogeic worm *P. corethrurus* on plant production under field condition.

CHAPTER 1

INTRODUCTION

Malaysia with the tropical climate has a fairly high, but uniform temperature and large amounts of rain delivering in intense bursts each year. Because of the weather, local agricultural soils contain low organic matter and highly compact soil texture resulting in lower crop productivity (Sabtu, 2006; APEC, 2012). A greater amount of chemical fertilizers is used each year to maintain soil fertility and boost crop productivity. However, the use of chemical fertilizers only restores soil nutrient content and does not improve the soil's physical structure and microbiological activities. Long-term use of chemical fertilizers also results in many negative effects on the soil such as acidification, decline in organic matter content and reduction of microbial communities (Sinha *et al*., 2010). Therefore, the rehabilitation of local agricultural soils is necessary and important to reduce the loss of soil nutrients as well as to promote sustainable agriculture.

One of the approaches to create a healthy agro-ecosystem is introducing soil-dwelling endogeic worms into the soil (Butt *et al*., 1997). Endogeic worms are known as "ecological engineers" as their burrowing and casting activities result in beneficial chemical, microbiological and physical changes in the soil that increase the fertility of the soil (Jones *et al*., 1994; Scheu, 2003). Efforts have been made to transplant endogeic worms into degraded pastures in New Zealand which have significantly improved pasture productivity (Stockdill, 1982). In grasslands on reclaimed industrial cutaway peat in Ireland, herbage production was significantly improved by earthworm colonization through improvement in the physical structure of the soil (Curry and Boyle, 1987).

The use of earthworms to improve soil conditions is a common practice in temperate regions like New Zealand (Stockdill, 1982), The Netherlands (Hoogerkamp *et al*., 1983), Ireland (Curry and Boyle, 1987), and the United Kingdom (Butt *et al*., 1997). However, this approach is rarely practiced in the tropics including Malaysia. Available information on tropical earthworms is limited to litter-dwelling epigeic species while knowledge about the biology and ecology of tropical endogeic worms is very scarce. The differences between temperate and tropical climates are likely to influence the activities of earthworms in soils (Kale and Karmegam, 2010). The higher temperature and humidity in tropical ecosystems may increase the overall activities of earthworms in soils. The higher burrowing and casting activities may accelerate soil rehabilitation processes by improving the physicochemical and biological properties of soil and in turn create a healthy environment for plant growth. On the other hand, intense bursts of rain in tropical ecosystems may destroy earthworm surface casts. Nutrients within earthworm casts may therefore be lost through surface runoff (Jouquet *et al*., 2013). The overall effects of these processes may cause the effects of earthworms on soil properties

in tropical ecosystems to differ from those in temperate regions. More experimental research is necessary to determine the roles of tropical earthworms in their ecosystems.

Obtaining a sufficient number of earthworms for soil rehabilitation is a fundamental aspect to be achieved prior to the practice. Most earthworm inoculation activities conducted in temperate regions are accomplished *via* the mass collection of earthworms from densely populated areas and transplanted into lower density areas (Butt *et al*., 1995). This technique is expensive, labor-intensive and time-consuming (Butt *et al*., 1992). Earthworm collection sites must be amended with a lot of organic matter over a long period before further collection in order to maximize the number of earthworms present in the area (Butt, 2011). In addition, under natural conditions, earthworm availability is strongly affected by seasonal changes, organic resource distribution and farm management practices (Ghafoor *et al*., 2008; Pauli *et al*., 2010). Thus, it may be difficult to acquire a sufficient number of earthworms for inoculation purposes. An alternative method for obtaining a large number of earthworms is indoor cultivation (Butt *et al*., 1992). For example, successful production of epigeic worms such as *Eisenia fetida*, *Lumbricus rubellus*, *Perionyx excavatus*, and *Eudrilus eugeniae* under indoor conditions resulted in the widespread use of these species in vermicomposting, vermiculture for protein source and ecotoxicological studies.

Unlike epigeic worms, soil-dwelling endogeic worms are fairly difficult to cultivate indoors. Endogeic worms are geophagous deriving their nutrition through the consumption of large amounts of soil. They require culturing in a mineral soil-based medium (Butt and Lowe, 2011). Butt and his co-workers used soil collected from earthworm-sampling sites as the medium to culture endogeic worms (Butt *et al*., 1992; Butt, 1993; Lowe and Butt, 2005). However, there are difficulties in using natural soil as the medium for culturing endogeic worms. This is because field-collected soils are highly heterogeneous. Its texture, water-holding capacity, organic matter content, nutrient availability, and microbial activities differ on the basis of sampling locations, land management and plant communities (Huggins and Uberuaga, 2010). Differences in the physiochemical and biological properties of field-collected soils could affect the survival, growth, sexual maturity, and cocoon production rates of earthworms (Klok *et al*., 2007). Therefore, it is difficult to reach a considerable level of standardization with field-collected soil as the earthworm culture medium. Field-collected soils may also contain harmful compounds such as heavy metals, polycyclic aromatic hydrocarbons and pesticides that could adversely affect earthworm cultures. An alternative culture medium should be established in order to achieve standardization for indoor cultivation of endogeic worms and also to produce a large and consistent supply of earthworms for soil rehabilitation programs.

The soil-dwelling endogeic worm *Pontoscolex corethrurus* (Müller, 1857) is one of the few persistent species capable of surviving under tropical agricultural tillage conditions (Lavelle *et al*., 1987; Sabrina *et al*., 2009; Nath and Chaudhuri, 2010). In Malaysia, *P. corethrurus* is the only endogeic species found in local rubber and oil palm plantations

(Loh, 2009; Sabrina *et al*., 2009). It is widely distributed in a variety of soil types including Serdang series, Bungor series and Jawa series (Vijiandran and Gurmit, 2009). Previous studies have shown that adaptation to tropical climates, capability to tolerate a wide range of environmental factors (e.g., soil pH and organic matter content) and the ability to enter diapause under drought conditions are the adaptive strategies that enable *P. corethrurus* to be widely distributed in tropical ecosystems (Lavelle *et al.*, 1987; James and Hendrix, 2004; Chaudhuri *et al*., 2008; Nath and Chaudhuri, 2010). The fact that it is widely distributed in tropical ecosystems and shows good environmental plasticity suggests that *P. corethrurus* may serve as a potential candidate for soil rehabilitation in tropical regions.

Unfortunately, in Malaysia, other than ecological data (distribution and abundance), relevant information about *P. corethrurus* is very limited. A detailed study on the effects of *P. corethrurus* on soil chemical and biological properties as well as crop productivity is required to determine its role in local soil ecosystems. Prior to determining its effects on soil properties, fundamental aspects of earthworm biology (e.g., life history) need to be studied since the biological characteristics of earthworms will affect their potential in soil rehabilitation programs. Nevertheless, the life history of earthworms is strongly affected by earthworm stocking density (Hait and Tare, 2011; Suthar, 2012). To date, there is still no study on how stocking density will influence the life history of *P. corethrurus*. In addition, a suitable culture medium for culturing *P. corethrurus* should be established to provide a large and consistent supply of *P. corethrurus* for worm inoculation activities.

Therefore, the objectives of the present research were:

i. To determine the effect of stocking density on life history of the endogeic worm *P. corethrurus*,

ii. To establish a suitable medium for culturing the endogeic worm *P. corethrurus*,

iii. To investigate the effect of the endogeic worm *P. corethrurus* on nutrient availability, humic acid content, microbial population, and enzyme activity in soil,

iv. To evaluate the effect of inoculating the endogeic worm *P. corethrurus* on plant production under field condition.

REFERENCES

Adianto, Safitri, D.U. and Yuli, N. 2004. Pengaruh inokulasi cacing tanah (*Pontoscolex corethrurus* Fr Mull) terhadap sifat fisika kimia tanah dan pertumbuhan tanaman kacang hijau (*Vigna radiata* L. Wilczek) varietas Walet. *Jurnal Matematika dan Sains* 9: 175-182.

Aira, M., Monroy, F. and Domínguez, J. 2003. Effects of two species of earthworms (*Allolobophora* spp.) on soil systems: a microfaunal and biochemical analysis. *Pedobiologia* 47: 877-881.

Amador, J.A., Glucksman, A.M., Lyons, J.B. and Gores, J.H. 1997. Spatial distribution of soil phosphatase activity within a riparian forest. *Soil Science* 162: 808-825.

Amador, J.A., Görres, J.H. and Savin, M.C. 2003. Carbon and nitrogen dynamics in *Lumbricus terrestris* (L.) burrow soil: relationship to plant residues and macropores. *Soil Science Society of America Journal* 67: 1755-1762.

Amador, J.A., Görres, J.H. and Savin, M.C. 2006. Effects of *Lumbricus terrestris* L. on nitrogen dynamics beyond the burrow. *Applied Soil Ecology* 33: 61-66.

Anđelković, T., Nikolić, R., Bojić, A., Anđelković, D. and Nikolić, G. 2010. Binding of cadmium to soil humic acid as a function of carboxyl group content. *Macedonian Journal of Chemistry and Chemical Engineering* 29: 215-224.

Asia-Pacific Economic Cooperation (APEC). 2012. *Summary report of APEC International Conference on Natural Resources and Infrastructure Management for Agriculture*, (6-8 August 2012), Bangkok, Thailand. pp. 1-7.

Arancon, N.Q., Edwards, C.A., Bierman, P., Metzger, J.D., Lee, S. and Welch, C. 2003. Effects of vermicomposts on growth and marketable fruits of field-grown tomatoes, peppers and strawberries. *Pedobiologia* 47: 731-735.

Arancon, N.Q., Edwards, C.A., Bierman, P., Welch, C. and Metzger, J.D. 2004. Influences of vermicomposts on field strawberries: 1. Effects on growth and yields. *Bioresource Technology* 93: 145-153.

Arancon, N.Q., Edwards, C.A., Lee, S. and Byrme, R. 2006. Effects of humic acids from vermicomposts on plant growth. *European Journal of Soil Biology* 42: 65-69.

Araujo, Y., Luizão, F.J. and Barros, E. 2004. Effect of earthworm addition on soil nitrogen availability, microbial biomass and litter decomposition in mesocosms. *Biology and Fertility of Soils* 39: 146-152.

Atlavinyte, O. and Pociene, C. 1973. The effect of earthworms and their activity on the amount of algae in soil. *Pedobiologia* 13: 445-455.

Baker, G. 2007. Differences in nitrogen release from surface and incorporated plant residues by two endogeic species of earthworms (Lumbricidae) in a red-brown earth soil in southern Australia. *European Journal of Soil Biology* 43: 165-170.

Baker, G.H., Brown, G., Butt, K., Curry, J.P. and Scullion, J. 2006. Introduced earthworms in agricultural and reclaimed land: their ecology and influences on soil properties, plant production and other soil biota. *Biological Invasions* 8: 1301-1316.

Baker, G.H., Cartera, P.J., Curry, J.P., Cultreric, O. and Beck, A. 1998. Clay content of soil and its influence on the abundance of *Aporrectodea trapezoides* Dugès (Lumbricidae). *Applied Soil Ecology* 9: 333-337.

Baker, G.H. and Whitby, W.A. 2003. Soil pH preferences and the influences of soil type and temperature on the survival and growth of *Aporrectodea longa* (Lumbricidae). *Pedobiologia* 47: 745-753.

Barois, I. and Lavelle, P. 1986. Changes in respiration rate and some physicochemical properties of a tropical soil during transit through *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta). *Soil Biology and Biochemistry* 18: 539-541.

Barois, I., Lavelle, P., Brossard, M., Tondoh, J., Martinez, M., Rossi, J.P., Senapati, B.K., Angeles, A., Fragoso, C., Jiménez, J.J., Decaëns, T., Lattaud, C., Kanyonyo, J., Blanchart, E., Chapuis, L., Brown, G.G. and Moreno, A. 1999. Ecology of earthworm species with large environmental tolerance and/or extended distributions. In *Earthworm Management in Tropical Agroecosystems*, eds. P. Lavelle, L. Brussaard and P. Hendrix, pp. 57-86. CABI International Press, Wallingford, UK.

Barois, I., Villemin, G., Lavelle, P. and Toutain, F. 1993. Transformation of the soil structure through *Pontoscolex corethrurus* (Oligochaeta) intestinal tract. *Geoderma* 56: 57-66.

Basker, A., Macgregor, A.N. and Kirkman, J.H. 1993. Exchangeable potassium and other cations in non-ingested soil and casts of two species of pasture earthworms. *Soil Biology and Biochemistry* 25: 1673-1677.

Bhattacharjee, G. and Chaudhuri, P.S. 2002. Cocoon production, morphology, hatching pattern and fecundity in seven tropical earthworm species – a laboratory-based investigation. *Journal of Bioscience* 27: 283-294.

Bhole, R.J. 1992. Vermiculture biotechnology basis, scope for application and developments. In *Proceedings of the National Seminar on Organic Farming*, 18-19 April, 1992, Mahatma Phule Krishi Vidyapeetha, College of Agriculture, Pune. pp: 50-51.

Bisht, R., Pandey, H., Bharti, D., Bisht, S.P.S. and Kaushal, B.R. 2007. Reproductive potential of the earthworm *Metaphire posthuma* (Oligochaeta) in different food substrates. *Tropical Ecology* 48: 107-114.

Blanchart, E., Albrecht, A., Alegre, J., Duboisset, A., Pashanasi, B., Lavelle, P. and Brussaard, L. 1999. Effects of earthworms on soil structure and physical properties. In *Earthworm Management in Tropical Agroecosystems*, eds. P. Lavelle, L. Brussaard and P. Hendrix, pp. 139-162. CAB International Press, Wallingford, UK.

Bossuyt, H., Six, J. and Hendrix, P.F. 2005. Protection of soil carbon by microaggregates within earthworm casts. *Soil Biology and Biochemistry* 37: 251–258.

Bostrom, U. and Lofs-Holmin, A. 1986. Growth of earthworms *(Allolobophora caliginosa)* fed shoots and roots of barley, meadow fescue and lucerne. Studies in relation to particle size, protein, crude fibre content and toxicity. *Pedobiologia* 29: 1-12.

Bouché, M.B. 1977. Strategies lombriciennes. In *Soil Organisms as Components of Ecosystems*, eds. U. Lohm and T. Persson, pp. 122-132 . Ecological Bulletins. (Stockholm).

Brown, G.G. 1995. How do earthworms affect microfloral and faunal community diversity? *Plant and Soil* 170: 209-231.

Brown, G.G., Barois, I. and Lavelle, P. 2000. Regulation of soil organic matter dynamics and microbial activity in the drilosphere and the role of interactions with other edaphic functional domains. *European Journal of Soil Biology* 36: 177-198.

Brown, G.G. and Doube, B.M. 2004. Functional interactions between earthworms, microorganisms, organic matter, and plant. In *Earthworm Ecology* (2nd edition), ed. CA. Edwards, pp. 213-262. CRC Press, Boca Raton, London, New York, Washington.

Brown, G.G., Edwards, C.A. and Brussaard, L. 2004. How earthworms affect plant growth: burrowing into the mechanisms. In *Earthworm Ecology* (2nd edition), ed. C.A. Edwards, pp. 13-48. CRC Press, Boca Raton, London, New York, Washington.

Brown, G.G., Pashanasi, B., Villenave, C., Patrón, J.C., Senapati, B.K., Giri, S., Barois, I., Lavelle, P., Blanchart, E., Blakemore, R.J., Spain, A.V. and Boyer, J. 1999. Effects of earthworms on plant production in the tropics. In *Earthworm Management in Tropical Agroecosystems*, eds. P. Lavelle, L. Brussaard and P. Henrix, pp. 87-147. CAB International Press, Wallingford, UK.

Brussaard, L., Behan-Pelletier, V.M., Bignell, D.E., Brown, V.K., Didden, W.A.M., Folgarait, P.J., Fragoso, C., Freckman, D.W., Gupta, V.V.S.R., Hattori, T., Hawksworth, D.L., Klopatek, C., Lavelle, P., Walloch, D., Rusek, J., Söderström, B., Tiedje, J.M. and Virginia, R.A. 1997. Biodiversity and ecosystem functioning in soil. *Ambio* 26: 563-570.

Buch, A.C., Brown, G.G., Niva, C.C., Sautter, K.D. and Lourencato, L.F. 2011. Life cycle of *Pontoscolex corethrurus* (Müller, 1857) in tropical artificial soil. *Pedobiologia* 54: 19-25.

Butt, K.R. 1993. Reproduction and growth of three deep-burrowing earthworms (Lumbricidae) in laboratory culture in order to assess production for soil restoration. *Biology and Fertility of Soils* 16: 135-138.

Butt, K.R. 1997. Reproduction and growth of the earthworm *Allolobophora chlorotica* (Savigny, 1826) in controlled environments. *Pedobiologia* 41: 369-374.

Butt, K.R. 2011. The use of vermiculture for land improvement. In *Vermiculture Technology: Earthworms, Organic Wastes and Environmental Management*, eds. C.A. Edwards, N.Q. Arancon and R. Sherman, pp. 335–348. CRC Press, Boca Rotan, Florida.

Butt, K.R., Frederickson, J. and Morris, R.M. 1992. The intensive production of *Lumbricus terrestris* L. for soil amelioration. *Soil Biology and Biochemistry* 24: 1321-1325.

Butt K.R., Frederickson, J. and Morris, R.M. 1993. Investigations of an earthworm inoculation experiment, London Borough of Hillingdon. *Waste Planning* 7: 9-12.

Butt, K.R., Frederickson, J. and Morris, R.M. 1994. Effects of earthworm density on the growth and reproduction of *Lumbricus terrestris* L. (Oligochaeta: Lumbricidae) in culture. *Pedobiologia* 38: 254-261.

Butt, K.R., Frederickson, J. and Morris, R.M. 1995. An earthworm cultivation and soil inoculation technique for land restoration. *Ecological Engineering* 4: 1-9.

Butt, K.R., Frederickson, J. and Morris, R.M. 1997. The earthworm inoculation unit technique: an integrated system for cultivation and soil inoculation of earthworms. *Soil Biology and Biochemistry* 29: 251-257.

Butt, K.R. and Lowe, C.N. 2011. Controlled cultivation of endogeic and anecic earthworms. In *Biology of Earthworms*, ed. A. Karaca, pp. 107-122. Springer-Verlag Berlin Heidelberg.

Cahill, J.F. and Casper, B.B. 1999. Growth consequences of soil nutrient heterogeneity for two old-field herbs, *Ambrosia artemisiifolia* and *Phytolacca Americana*, grown individually and in combination. *Annals of Botany* 83: 471-478.

Cappuccino, J.G. and Sherman, N. 2005. *Microbiology: A Laboratory Manual* (7th edition). Pearson Education Press, USA.

Chamberlain, E.J. and Butt, K.R. 2008. Distribution of earthworms and influence of soil properties across a successional sand dune ecosystem in NW England. *European Journal of Soil Biology* 44: 554-558.

Chan, K.Y. 2003. Using earthworms to incorporate lime into subsoil to ameliorate acidity. *Communication in Soil Science and Plant Analysis* 34: 985-997.

Chan, K.Y. and Barchia, I. 2007. Soil compaction controls the abundance, biomass and distribution of earthworms in a single dairy farm in south-eastern Australia. *Soil and Tillage Research* 94: 75-82.

Chang, C.H., Yang, K.W., Wu, J.H., Chuang, S.C. and Chen, J.H. 2001. Species composition of earthworms on the main campus of National Taiwan University. *Acta Zoologica Taiwanica* 12: 75-81.

Chapuis-Lardy, L., Brossard, M., Lavelle, P. and Schouller, E. 1998. Phosphorus transformations in a ferralsol through ingestion by *Pontoscolex corethrurus*, a geophagous earthworm. *European Journal of Soil Biology* 34: 61-67.

Chaudhuri, P.S. and Bhattacharjee, S. 2011. Reproductive biology of eight tropical earthworm species of rubber plantations in Tripura, India. *Tropical Ecology* 52: 49-60.

Chaudhuri, P.S., Bhattacharjee, S., Dey, A., Chattopadhyay, S. and Bhattacharya, D. 2013. Impact of age of rubber (*Hevea brasiliensis*) plantation on earthworm communities of West Tripura (India). *Journal of Environmental Biology* 34: 59-65.

Chaudhuri, P.S. and Nath, S. 2011. Community structure of earthworms under rubber plantations and mixed forests in Tripura, India. *Journal of Environmental Biology* 32: 537-541.

Chaudhuri, P.S., Nath, S., Pal, T.K. and Dey, S.K. 2009. Earthworm casting activities under rubber (*Hevea brasiliensis*) plantations in Tripura (India). *World Journal of Agriculture Sciences* 5: 515-521.

Chaudhuri, P.S., Nath, S. and Paliwal, R. 2008. Earthworm population of rubber plantations (*Hevea brasiliensis*) in Tripura, India. *Tropical Ecology* 49: 225-234.

Chaudhuri, P.S., Pal, T.K., Bhattacharjee, G. and Dey, S.K. 2000. Chemical changes during vermicomposting (*Perionyx excavatus*) of kitchen wastes. *Tropical Ecology* 41: 107-110.

Clapperton, M.J., Lee, N.O., Binet, F. and Conner, R.L. 2001. Earthworms indirectly reduce the effects of take-all (*Gaeumannomyces graminis* var. Tritici) on soft white spring wheat (*Triticum aestivum* cv. Fielder). *Soil Biology and Biochemistry* 33: 1531-1538.

Cluzeau, D., Fayolle. L. and Hubert, M. 1992. The adaptation value of reproductive strategy and mode in three epigeous earthworm species. *Soil Biology and Biochemistry* 24: 1309-1315.

Cook, S.M.F. and Linden, D.R. 1996. Effect of food type and placement on earthworm (*Aporrectodea tuberculata*) burrowing and soil turnover. *Biology and Fertility of Soils* 21: 201–206.

Crommentuijn, T., Doodeman, C.J.A.M., Pol, J.J.C., Doornekamp, A., Rademaker, M.C.J. and Van Gestel, C.A.M. 1995. Sublethal sensitivity index as an ecotoxicity parameters measuring energy allocation under toxicant stress: application to cadmium in soil anthropods. *Ecotoxicology and Environmental Safety* 31: 92-200.

Curry, J.P. 2004. Factors affecting the abundance of earthworms in soils. In *Earthworm Ecology* (2nd edition), ed. C.A. Edwards, pp: 91-113. CRC Press, Boca Raton, London, New York, Washington.

Curry, J.P. and Boyle, K.E. 1987. Growth rates, establishment, and effects on herbage yield of introduced earthworms in grassland on reclaimed cutover peat. *Biology and Fertility of Soils* 3: 95-98.

Daily, G.C., Alexander, S. and Ehrlich, P.R. 1997. Ecosystem services: benefits supplied to human societies by natural ecosystems. *Issues in Ecology* 2: 18-24.

Daniel, O. and Anderson, J.M. 1992. Microbial biomass and activity in contrasting soil materials after passage through the gut of the earthworm *Lumbricus rubellus* Hoffmeister. *Soil Biology and Biochemistry* 24: 465-470.

Daniel, O., Kohli, L. and Bieri, M. 1996. Weight gain and weight loss of the earthworm *Lumbricus terrestris* at different temperatures and body weights. *Soil Biology and Biochemistry* 28: 1235-1240.

De Silva, P.M.C.S. and van Gestel, C.A.M. 2009. Comparative sensitively of *Eisenia andrei* and *Perionyx excavatus* in earthworm avoidance tests using two soil types in the tropics. *Chemosphere* 77: 1609-1613.

Dedeke, G.A., Omemu, O., Aladesida, A.A. and Museliu, F. 2010. Comparative microbial analysis of earthworm casts collected from Ikenne, Ogun State, Nigeria. *Ethiopian Journal of Environmental Studies and Management* 3: 57-63.

Dempsey, M.A., Fisk, M.C. and Fahey, T.J. 2011. Earthworms increase the ratio of bacteria to fungi in northern hardwood forest soils, primarily by eliminating the organic horizon. *Soil Biology and Biochemistry* 43: 2135-2141.

Demspey, M.A., Fisk, M.C., Yavitt, J.B., Fahey, T.J. and Balser, T.C. 2013. Exotic earthworms alter soil microbial community composition and function. *Soil Biology and Biochemistry* 67: 263-270.

Devi, S.H., Vijayalakshmi, K., Jyotsna, K.P., Shaheen, S.K., Jyothi, K. and Rani, M.S. 2009. Comparative assessment in enzyme activities and microbial populations during normal and vermicomposting. *Journal of Environmental Biology* 30: 1013-1017.

Devliegher, W. and Verstraete, W. 1997. The effect of *Lumbricus terrestris* on soil in relation to plant growth: effect of nutrient-enrichment processes (NEP) and gut-associated processes (GAP). *Soil Biology and Biochemistry* 29: 341-346.

Dix, N.J. and Webster, J. 1995. *Fungal Ecology*. Chapman & Hall, London.

Dom ńguez, J., Briones, M.J.I. and Mato, S. 1997. Effect of the diet on growth and reproduction of *Eisenia andrei* (Oligochaeta, Lumbricidae). *Pedobiologia* 41: 566-576.

Dom ńguez, J. and Edwards, C.A. 1997. Effects of stocking rate and moisture content on the growth and maturation of *Eisenia andrei* (oligochaeta) in pig manure. *Soil Biology and Biochemistry* 29: 743-746.

Dom ńguez, J. and Edwards, C.A. 2004. Vermicomposting organic wastes: a review. In *Soil Zoology for Sustainable Development in the 21st century*, eds. S.H. Shakir and W.Z.A. Mikha ĭ, pp. 369-396. EI Cairo.

Dom ńguez, J. and Edwards, C.A. 2010. Biology and ecology of earthworms species used for vermicomposting. In *Vermiculture Technology: Earthworms, Organic Waste and Environmental Management*, eds. C.A. Edwards, N.Q. Arancon and R.L. Sherman, pp. 25-37. CRC Press, Boca Raton, Florida.

Doube, B.M. and Brown, G.G. 1998. Life in a complex community: functional interactions between earthworms, organic matter, microorganisms and plants. In *Earthworm Ecology*, ed. C.A. Edwards, pp. 179-212. CRC Press, Boca Raton, Florida.

Doube, B.M., Williams, P.M.L. and Willmott, P.J. 1997. The influence of two species of earthworm (*Aporrectodea trapezoides* and *Aporrectodea rosea*) on the growth of wheat, barley and faba beans in three soil types in the greenhouse. *Soil Biology and Biochemistry* 29: 503-509.

Edwards, C.A. 2004. The importance of earthworms as key representatives of the soil fauna. In *Earthworm Ecology* (2nd edition), ed. C.A. Edwards, pp. 3-11. CRC Press, Boca Raton, London, New York, Washington.

Edwards, C.A. and Bohlen, P.J. 1996. *Biology and Ecology of Earthworms* (3rd edition). Chapman & Hall, London, UK.

Edwards, C.A. and Lofty, J.R. 1977. *Earthworm Ecology and Biology*. Chapman & Hall, London, UK.

Edwards, C.A. and Lofty, J.R. 1980. Effects of earthworm inoculation upon the root growth of direct drilled cereals. *Journal of Applied Ecology* 17: 533-543.

Ellis, S.R., Hodson, M.E. and Wege, P. 2007. The influence of different artificial soil types on the acute toxicity of carbendazim to the earthworm *Eisenia fetida* in laboratory toxicity tests. *European Journal of Soil Biology* 43: 239-245.

Fahramand, M., Moradi, H., Noori, M., Sobhkhizi, A., Adibian, M., Abdollahi, S. and Rigi, K. 2014. Influence of humic acid on increase yield of plants and soil properties. *International Journal of Farming and Allied Sciences* 3: 339-341.

Fernández, R., Novo, M., Gutiérrez, M., Almodóvar, A. and Díaz Cosín, D.J. 2010. Life cycle and reproductive traits of the earthworm *Aporrectodea trapezoides* (Dugès, 1828) in laboratory cultures. *Pedobiologia* 53: 295-299.

Fragoso, C., Kanyonyo, J., Moreno, A., Senapati, B.K., Blanchart, E. and Rodríguez, C. 1999. A survey of tropical earthworms: taxonomy, biogeography and environmental plasticity. In *Earthworm Management in Tropical Agroecosystems*, eds. P. Lavelle, L. Brussaard and P. Hendrix, pp. 1-26. CAB International Press, Wallingford, UK.

Fragoso, C. and Lozano, N. 1992. Resource allocation strategies imposed by caudal amputation and soil moisture in the tropical earthworm *Pontoscolex corethrurus*. *Soil Biology and Biochemistry* 24: 1237-1240.

Fraser, P.M., Beare, M.H., Butler, R.C., Harrison-Kirk, T. and Piercy, J.E. 2003. Interactions between earthworms (*Aporrectodea caliginosa*), plants and crop residues for restoring properties of a degraded arable soil. *Pedobiologia* 47: 870-876.

Frey, S.D., Six, J. and Elliot, E.T. 2003. Reciprocal transfer of carbon and nitrogen by decomposer fungi at the soil-litter interface. *Soil Biology and Biochemistry* 35: 1001-1004.

García, J.A. and Fragoso, C. 2002. Growth, reproduction and activity of earthworms in degraded and amended tropical open mined soils: laboratory assays. *Applied Soil Ecology* 20: 43-56.

García, J.A. and Fragoso, C. 2003. Influence of different food substrates on growth and reproduction of two tropical earthworm species (*Pontoscolex corethrurus* and *Amynthas corticis*). *Pedobiologia* 47: 754–763.

Garg, V.K., Kaushik, P. and Yadav, Y.K. 2008. Effect of stocking density and food quality on the growth and fecundity of an epigeic earthworm (*Eisenia fetida*) during vermicomposting. *Envionmentalist* 28: 483-488.

Garvín, M.H., Trigo, D., Hernández, P., Ruiz, M.P. and Díaz Cosín, D.J. 2002. Intercations of *Hormogaster elisae* (Oligochaeta, Hormogastridae) with other earthworm species from Reduena (Madrid, Spain). *Applied Soil Ecology* 20: 163-169.

Gates, G.E. 1972. Burmese earthworms: an introduction to the systematic and biology of megadrile oligochaetes with special reference to Southeast Asia. *Transactions of the American Philosophical Society* 62: 1-326.

Ghafoor, A., Hassan, M. and Hussainalvi, Z. 2008. Biodiversity of earthworm species from various habitats of district Narowal, Pakistan. *International Journal of Agriculture and Biology* 10: 681-684.

Giraddi, R.S. 2008. Effect of stocking rate of *Eudrillus eugeniae* (Kinberg) on vermicompost production. *Karnataka Journal of Agriculture and Science* 21: 49-51.

Glaser, B. 2007. Prehistorically modified soils of central Amazonia: a model for sustainable agriculture in the twenty-first century. *Philosophical transactions of the Royal Society* 362: 187-196.

Greacen, E.L., Barley, K.P. and Farrell, D.A. 1968. The mechanics of root growth on soils with particular reference to the implications for root distribution. In *Root growth*, ed. W.J. Whittington, pp. 256-269. Proceedings of the 15th Easter School in Agricultural Science, University of Nottingham.

Guenet, B., Leloup, J., Hartmann, C., Barot, S. and Abbadie, L. 2011. A new protocol for an artificial soil to analyse soil microbiological processes. *Applied Soil Ecology* 48: 243-246.

Guild, W.J.McL. 1948. Effect of soil type on populations. *Annals of Applied Biology* 35: 181-192.

Gupta, P.K. 2007. *Soil, Plant, Water and Fertilizer Analysis* (2nd edition). Agrobios, India.

Hait, S. and Tare, V. 2011. Optimizing vermistabilization of waste activated sludge using vermicompost as bulking material. *Waste Management* 31: 502-511.

Hauser, S., Asawalam, D. and Vanlauwe, B. 1998. Spatial and temporal gradients of earthworm casting activity in alley cropping systems. *Agroforestry Systems* 41: 127-137.

Hawkins, C.L., Rutledge, M., Savin, M.C., Shipitalo, M.J. and Brye, K.R. 2008. A sand layer deters burrowing by *Lumbricus terrestris* L. *Soil Science* 173: 186-194.

Haynes, R.J., Fraser, P.M., Piercy, J.E. and Tregurtha, R.J. 2003. Casts of *Aporrectodea caliginosa* (Savigny) and *Lumbricus rubellus* (Hoffmeister) differ in microbial activity, nutrient availability and aggregate stability. *Pedobiologia* 47: 882-887.

Hendriksen, N.B. 1991. Gut load and food-retention time in the earthworms *Lumbricus festivus* and *L. castaneus*: a field study. *Biology and Fertility of Soils* 11: 170-173.

Hirth, J.R., McKenzie, B.M. and Tisdall, J.M. 1997. Do the roots of perennial ryegrass elongate to biopores filled with the casts of endogeic earthworms? *Soil Biology and Biochemistry* 29: 529-531.

Hoben, H.J. and Somasegaran, P. 1982. Comparison of the pour, spread, and drop plate methods for enumeration of *Rhizobium* spp. in inoculants made from presterilized peat. *Applied and Environmental Microbiology* 44: 1246-1247.

Hodge, A. 2004. The plastic plant: root responses to heterogeneous supplies of nutrients. *New Phytologist* 162: 9-24.

Hofman, J., Hovorková, I. and Machát, J. 2009. Comparison and characterization of OECD artificial soils. In *Ecotoxicological Characterization of Waste*, eds. H. Moser and J Römbke, pp. 223-229. Springer Science+Business Media, LLC.

Holland. E.A. and Coleman, D.C. 1987. Litter placement effects on microbial and organic-matter dynamics in an agroecosystem. *Ecology* 68: 425-433.

Holmstrup, M., Østergaard, I.K., Nielsin, A. and Hansen, B.T. 1991. The relationship between temperature and cocoon incubation time for some lumbricid earthworm species. *Pedobiologia* 35: 179-184.

Hoogerkamp, M., Rogaar, H. and Eijsackers, H.J.P. 1983. Effect of earthworms on grassland on recently reclaimed polder soils in the Netherlands. In *Earthworm Ecology from Darwin to Vermiculture*, ed. J.E. Satchell, pp. 85-105. Chapman and Hall, London, New York.

Huggins, D.R. and Uberuaga, D.P. 2010. *Field Heterogeneity of Soil Organic Carbon and Relationships to Soil Properties and Terrain Attributes*. CSANR Research Report. pp. 1-23.

Hugenholtz, P. 2002. Exploring prokaryotic diversity in the genomic era. *Genome Biology* 3:Reviews0003.

Hyvönen, R., Andersson, S., Clarholm, M. and Persson, T. 1994. Effect of lumbricids and enchytraeids on nematodes in limed and umlimed coniferous mor humuc. *Biology and Fertility of Soils* 17: 201-205.

Ibrahim, B.U., Auta, J. and Adebote, D.A. 2010. Effects of soil types and enhanced nutrient levels on the productivity of earthworm (*Eudrilius eugeniae*, Kinberg). *Bayero Journal of Pure and Applied Science* 3: 59-62.

Idowu, A.B., Edema, M.O. and Adeyi, A.O. 2006. Distribution of bacteria and fungi in the earthworm *Libyodrillus violaceous* (Annelida: Oligochaeta), a native earthworm from Nigeria. *Revista de Biologia Tropical* 54: 49-58.

Jabatan Pertanian Malaysia, 2007. Pakej Teknologi Cili (1st edition). Jabatan Pertanian, Kementerian Pertanian dan Industri Asas Tani, Malaysia.

James, S.W. and Hendrix, P.F. 2004. Invasion of exotic earthworms into North America and other regions. In *Earthworm Ecology* (2nd edition), ed. C.A. Edwards, pp. 75-88. CRC Press, Boca Raton, London, New York, Washington.

Jayasinghe, B.A.T.D and Parkinson, D. 2009. Earthworms as the vectors of actinomycetes antagonistic to litter decomposer fungi. *Applied Soil Ecology* 43: 1-10.

Jégou, D., Schrader, S., Diestel, H. and Cluzeau, D. 2001. Morphological, physical and biochemical characteristics of burrow walls formed by earthworms. *Applied Soil Ecology* 17: 165–174.

Jiménez, J.J., Cepada, A., Decaëns, T., Oberson, A. and Friesen, D.K. 2003. Phosphorus fractions and dynamics in surface earthworm casts under native and improved grasslands in a Colombian savanna Oxisol. *Soil Biology and Biochemistry* 35: 715-727.

Jones, C.G., Lawton, J.H. and Shachak, M. 1994. Organisms as ecosystem engineers. *Oikos* 69: 373–386.

Joshi, N. and Dabral, M. 2008. Life cycle of earthworms *Drawida nepalensis*, *Metaphire houlleti* and *Perionyx excavatus* under laboratory controlled conditions. *Life Science Journal* 5: 83-86.

Jouquet, P., Blanchart, E. and Capowiez, Y. 2014. Utilization of earthworms and termites for the restoration of ecosystem functioning. *Applied Soil Ecology* 73: 34-40.

Jouquet, P., Bottinelli, N., Kerneis, G., Henry-des-Tureaux, T., Thuy, D.T., Planchon, O. and Toan, T.D. 2013. Surface casting of the tropical *Metaphire posthuma* increases soil erosion and nitrate leaching in a laboratory experiment. *Geoderma* 204-205: 10-14.

Jouquet, P., Phuong, N.T., Hanh, N.H., Henry-des-Tureaux, T., Chevallier, T. and Tran Duc, T. 2011. Laboratory investigation of organic matter mineralization and nutrient leaching from earthworm casts produced by *Amynthas khami*. *Applied Soil Ecology* 47: 24-30.

Jouquet, P., Plumere, T., Thuy, D.T., Rumpel, C., Toan, T.D. and Orange, D. 2010. The rehabilitation of tropical soils using compost and vermicompost is affected by the presence of endogeic earthworms. *Applied Soil Ecology* 46: 125-133.

Kale, R.D. and Karmegam, N. 2010. The role of earthworms in tropics with emphasis on Indian Ecosystems. *Applied and Environmental Soil Science* 1-16. (doi: 10.1155/2010/414356)

Karmegan, N. and Daniel, T. 2009. Growth, reproductive biology and life cycle of the vermicomposting earthworm, *Perionyx ceylanensis* Mich. (Oligochaeta: Megascolecidae). *Bioresource Technology* 100: 4790-4796.

Kaushal, B.R., Bisht, S.P.S. and Kalia, S. 1994. Effect of diet on cast production by the megascolecid earthworm *Amynthas alexandri* in laboratory culture. *Biology and Fertility of Soils* 17: 14-17.

Keeney, D.R. and Nelson, D.W. 1982. Nitrogen-inorganic forms. In *Methods of Soil Analysis, Part 2. Chemical and Microbial Properties*, eds. A.L. Page, R.H. Miller and D.R. Keeney, pp. 643-698. ASA, SSSA, Madison.

Khalaf El-Duweini, A. and Ghabbour, S.I. 1965. Population density and biomass of earthworms in different types of Egyptian soils. *Journal of Applied Ecology* 2: 271-287.

Kirby, E.C. and Baker, S.W. 1995. Earthworm populations, casting and control in sports turf areas: a review. *The Journal of the Sports Turf Research Institute* 71: 84–98.

Kirk, J.L., Beaudette, L.A., Hart, M., Moutoglis, P., Klironomos, J.N., Lee, H. and Trevors, J.T. 2004. Methods of studying soil microbial diversity. *Journal of Microbiological Methods* 58: 169-188.

Kizilkaya, R. and Hepşen, Ş. 2007. Microbiological properties in earthworm cast and surrounding soil amended with various organic wastes. *Communications in Soil Science and Plant Analysis* 38: 2861-2876.

Kizilkaya, R., Karaca, A., Turgay, O.C. and Cetin, S.C. 2011. Earthworm interactions with soil enzymes. In *Biology of Earthworms*, ed. A. Karaca. pp. 141-158. Springer-Verlag Berlin Heidelberg.

Klok, C. 2007. Effects of earthworm density on growth, development, and reproduction in *Lumbricus rubellus* (Hoffm.) and possible consequences for the intrinsic rate of population increase. *Soil Biology and Biochemistry* 39: 2401-2407.

Klok, C. and de Roos, A.M. 1996. Population level consequences of toxicological influences on individual growth and reproduction in *Lumbricus rubellus* (Lumbricidae Oligochaeta). *Ecotoxicology and Environmental Safety* 33: 118-127.

Klok, C., Faber, J., Heijmans, G., Bodt, J. and van der Hout, A. 2007. Influence of clay content and acidity of soil on development of the earthworm *Lumbricus rubellus* and its population level consequences. *Biology and Fertility of Soils* 43: 549-556.

Kondapa, D., Radder, B.M., Patil, P.L., Hebsur, N.S. and Alagundagi, S.C. 2009. Effect of integrated nutrient management on growth, yield and economics of chili (cv. Byadgi dabbi) in a vertisol. *Karnataka Journal of Agricultural Science* 22: 430-440.

Kostka, J.E., Wu, J., Nealson, K.H. and Stucki, J.W. 1999. The impact of structural Fe(III) reduction by bacteria on the surface chemistry of smectite clay minerals. *Geochimica et Cosmochimica Acta* 63: 3705-3713.

Kretzschmar, A. 2004. Effects of earthworms on soil organization. In *Earthworm Ecology* (2nd edition), ed. C.A Edwards, pp. 201-210. CRC Press, Boca Raton, London, New York, Washington.

Kristufek, V., Ravasz, K. and Pizl, V. 1992. Changes in density of bacteria and microfungi during gut transit in *Lumbricus rubellus* and *Aporrectodea caliginosa* (Oligochaeta: Lumbricidae). *Soil Biology and Biochemistry* 24: 1499-1500.

Kuczak, C.N., Fernandes, E.C.M., Lehmann, J., Rondon, M.A. and Luizão, F.J. 2006. Inorganic and organic phosphorus pools in earthworm casts (Glossoscolecidae) and a Brazilian rainforest Oxisol. *Soil Biology and Biochemistry* 38: 553-560.

Ladd J.N. and Butler J.A.H. 1972. Short-term assays of soil proteolytic enzyme activities using proteins and dipeptide derivatives as substrates. *Soil Biology and Biochemistry* 4: 19-30.

Laffan, M.D. and Kingston, T.J. 1997. Earthworms in some Tasmanian forest soils in relation to bioturbation and soil texture profile. *Australian Journal of Soil Research* 35: 1231-1243.

Lattaud, C., Zhang, B., Locati, S., Rouland, C., Lavelle, P. 1997. Activities of the digestive enzymes in the gut and in tissue culture of a tropical geophagous earthworm, *Polypheretima elongata* (Megascolecidae). *Soil Biology and Biochemistry* 29: 335-339.

Lavelle, P. 1983. The structure of earthworm communities. In *Earthworm Ecology from Darwin to Vermiculture*, ed. J.E. Satchell, pp. 449-466. Chapman & Hall, London, New York.

Lavelle, P., Barois, I., Cruz, C., Hernandez, A., Pineda, A. and Rangel, P. 1987. Adaptative strategies of *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta), a peregrine geophagous earthworm of the humid tropics. *Biology and Fertility of Soils* 5: 188-194.

Lavelle, P., Blanchart, E., Martin, A., Spain, A.V. and Martin, S. 1992a. Impact of soil fauna on the properties of soils in the humid tropics. In *Myths and Science of Soils of the Tropics*, eds. R. Lal and P.A. Sanchez, pp. 157-185. Soil Science Society of America, Madison, SSSA Special Publication.

Lavelle, P., Brussaard, L. and Hendrix, P. 1999. *Earthworm Management in Tropical Agroecosystems*. CAB International Press, Wallingford, UK.

Lavelle, P., Decaëns, T., Aubert, M., Barot, S., Blouin, M., Bureau, F., Margerie, P., Mora, P. and Rossi, J.P. 2006. Soil invertebrates and ecosystem services. *European Journal of Soil Biology* 42: 3-15.

Lavelle, P. and Martin, A. 1992. Small-scale and large-scale effects of endogeic earthworms on soil organic matter dynamics in soils of the humid tropics. *Soil Biology and Biochemistry* 24: 1491–1498.

Lavelle, P., Melendez, G., Pashanasi, B. and Schaefer, R. 1992b. Nitrogen mineralization and recognization in casts of the geophagous tropical earthworm *Pontoscolex corethrurus* (Glossoscolecidae). *Biology and Fertility of Soils* 14: 49-53.

Lazcano, C. and Domínguez, J. 2010. Effects of vermicompost as a potting amendment of two commercially-grown ornamental plant species. *Spanish Journal of Agricultural Research* 8: 1260-1270.

Lazcano, C. and Domínguez, J. 2011. The use of vermicompost in sustainable agriculture: impact on plant growth and soil fertility. In: *Soil Nutrients*, ed. M. Miransari, pp. 230-254. Nova Science Publishers, New York.

Le Bayon, R.C. and Binet, F. 2006. Earthworms change the distribution and availability of phosphorous in organic substrates. *Soil Biology and Biochemistry* 38: 235-246.

Lee, K.E. *1985. Earthworms: Their Ecology and Relationships with Soils and Land Use*. Academic Press, Sydney.

Li, H.B., Zhang, F.S. and Shen J.B. 2012. Contribution of root proliferation in nutrient-rich soil patches to nutrient uptake and growth of maize. *Pedosphere* 22: 776-784.

Logsdon, G. 1994. Worldwide progress in vermicomposting. *Biocycle* 35: 63-65.

Loh, K.E. 2009. *The ecology of endogeic earthworm in selected agriculture area in Universiti Putra Malaysia*, Degree Thesis, Universiti Putra Malaysia.

Loh, K.E., Nor Azwady, A.A., Intan, S.I. Alvin, H.K.W, Muskhazli, M. and Kok, H.Y. 2013. Characterization of *Bactrocera dorsalis* (fruit fly) emergency inhibitor from neem leaf-empty fruit bunch-based vermicompost using metabolomics approach. *Acta Biologica Malaysiana* 2: 54-63.

Loh, K.E., Nor Azwady, A.A., Kok, H.Y., Muskhazli, M., Intan, S.I. and Nur Ain Izzati, M.Z. 2012. Potential of neem leaf-empthy fruit bunch-based vermicompost as biofertilizer-cum-biopesticide: chemical properties, humic acid content and ezymes (protease and phosphatase) activity in vermicompost (Part 1). *Scientific Research and Essays* 7: 3657-3664.

Loranger-Merciris, G., Cabidoche, Y.M., Deloné, B., Quénéhervé, P. and Ozier-Lafontaine, H. 2012. How earthworm activities affect banana plant response to nematodes parasitism. *Applied Soil Ecology* 52: 1-8.

Lowe, C.N. and Butt, K.R. 1999. Interspecific interactions between earthworms: potential applications in soil amelioration. *Pedobiologia* 43: 808-817.

Lowe, C.N. and Butt, K.R. 2005. Culture techniques for soil dwelling earthworms: a review. *Pedobiologia* 49: 401-413.

Lowe, C.N. and Butt, K.R. 2007. Earthworm culture, maintenance and species selection in chronic ecotoxicological studies: a review. *European Journal of Soil Biology* 43: 281-288.

Lowe, C.N. and Butt, K.R. 2008. Life cycle traits of the parthenogenetic earthworm *Octolasion cyaneum* (Savigny, 1826). *European Journal of Soil Biology* 44: 541-544.

López-Hernández, D., Lavelle, P., Fardeau, J.C. and Niño, M. 1993. Phosphorous transformations in two P-sorption contrasting tropical soils during transit through *Pontoscolex corethrurus* (Glossoscolecidae: Oligochaeta). *Soil Biology and Biochemistry* 25: 789-792.

Marichal, R., Grimaldi, M., Mathieu, J., Brown, G.G., Desjardins, T., da Silva Junior, M.L., Praxedes, C., Martins, M.B., Velasquez, E. and Lavelle, P. 2012. Is invasion of deforested Amazonia by the earthworm *Pontoscolex corethrurus* driven by soil texture and chemical properties? *Pedobiologia* 55: 233-240

Marichal, R., Martinez, A.F., Praxedes, C., Ruiz, D., Carvajal, A.F., Oszwald, J., Hurtoda, M.d.P., Brown, G.G., Grimaldi, M., Desjardins, T., Sarrazin, M., Decaëns, T., Velasquez, E. and Lavelle, P. 2010. Invasion of *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta) in landscapes of the Amazonian deforestation arc. *Applied Soil Ecology* 46: 443-449.

Mehlich, A. 1984. Mehlich-3 soil test extractant: a modification of Mehlich-2 extractant. *Communications in Soil Science and Plant Analysis* 15: 1409-1416.

Monroy, F., Aira, M., Gago, J.A. and Domínguez, J. 2007. Life cycle of the earthworm *Octodrilus complanatus* (Oligochaeta, Lumbricidae). *Comptes Rendus Biologies* 330: 389-391.

Muscolo, A., Bovalo, F., Gionfriddo, F. and Nardi, S. 1999. Earthworm humic matter produces auxin-like effects on *Daucus carota* cell growth and nitrate metabolism. *Soil Biology and Biochemistry* 31: 1303-1311.

Nath, S. and Chaudhuri, P.S. 2010. Human-induced biological invasions in rubber (*Hevea brasiliensis*) plantations of Tripura (India) – *Pontoscolex corethrurus* as a case study. *Asian Journal of Experimental Biological Science* 1: 360-369.

Nath, S. and Chaudhuri, P.S. 2014. Growth and reproduction of *Pontoscolex corethrurus* (Müller) with different experimental diets. *Tropical Ecology* 55: 305-312.

Nielson, R.L. 1965. Presence of plant growth substances in earthworms demonstrated by paper chromatography and the Went pea test. *Nature* 208: 1113-1114.

Neilson, R., Boag, B. and Smith, M. 2000. Earthworm $\delta^{13}C$ and $\delta^{15}N$ analyses suggest that putative functional classifications of earthworms are site specific and may also indicate habitat diversity. *Soil Biology and Biochemistry* 32: 1053–1061.

Noguera, D., Rondón, M., Laossi, K.R., Hoyos, V., Lavelle, P., de Carvalho, M.H.C. and Barot, S. 2010. Contrasted effect of biochar and earthworms on rice growth and resource allocation in different soils. *Soil Biology and Biochemistry* 42: 1017-1027.

Nordström, S. and Rundgren, S. 1974. Environmental factors and lumbricid associations in southern Sweden. *Pedobiologia* 14: 1-27.

Nurshazeela, A.R. 2009. *The culture of indigenous earthworm, Pontoscolex corethrurus by manipulating the ratio of soil and sheep manure under laboratory condition*. Degree Thesis, Universiti Putra Malaysia.

OECD. 1984. *Earthworm, acute toxicity test, Test Guideline No. 207, Guidelines for the testing of chemicals*. OECD, Paris.

OECD. 2004. *Earthworm reproduction test (Eisenia fetida/Eisenia andrei), Test Guideline No. 222, Guidelines for the testing of chemicals.* OECD, Paris.

Olander, L.P. and Vitousek, P.M. 2000. Regulation of soil phosphatase and chitinase activity by N and P availability. *Biogeochemistry* 49: 175-190.

Ortiz-Ceballos, A.I., Fragoso, C. and Brown, G.G. 2007. Synergistic effect of a tropical earthworm *Balanteodrilus pearsei* and velvetbean *Mucuna pruriens* var. *utilis* on maize growth and crop production. *Applied Soil Ecology* 35: 356-362.

Ortiz-Ceballos, A.I., Fragoso, C., Equihua, M. and Brown, G.G. 2005. Influence of food quality, soil moisture and the earthworm *Pontoscolex corethrurus* on growth and reproduction of the tropical earthworm *Balanteodrilus pearsei*. *Pedobiologia* 49: 89-98.

Ortiz-Ceballos, A.I., Hernández-García, M.E.C. and Galindo-González, J. 2009. Nest and feeding chamber construction for cocoon incubation in the tropical earthworm: *Pontoscolex corethrurus*. *Dynamic Soil, Dynamic Plant* 3: 115-118.

Owojori, O.J., Reineicke, A.J. and Rozanov, A.B. 2009. Role of clay content in partitioning, uptake and toxicity of zinc in the earthworm *Eisenia fetida*. *Ecotoxicology and Environmental Safety* 72: 99-107.

Owojori, O.J., Reineicke, A.J. and Rozanov, A.B. 2010. Influence of clay content on bioavailability of copper in the earthworm *Eisenia fetida*. *Ecotoxicology and Environmental Safety* 73: 407-414.

Palm, J., van Schaik, N.L.M.B. and Schröder, B. 2013. Modeling distribution patterns of anecic, epigeic and endogeic earthworms at catchment-scale in agro-ecosystems. *Pedobiologia* 56: 23-31.

Parmelee, R.W. and Crossley, D.A.Jr. 1988. Earthworm production and role in the nitrogen cycle of a no-tillage agroecosystem on the Georgia piedmont. *Pedobiologia* 32: 351-361.

Parthasarathi, K. 2007. Life cycle of *Lampito mauritii* (Kinberg) in comparison with *Eudrilus eugeniae* (Kinberg) cultured on different substrates. *Journal of Environmental Biology* 28: 803-812.

Parthasarathi, K., Ranganathan, L.S., Anandi, V. and Zeyer, J. 2007. Diversity of microflora in the gut and casts of tropical composting earthworms reared on different substrates. *Journal of Environmental Biology* 28: 87-97.

Paul, E.A. and Clark, F.E. 1996. *Soil Microbiology and Biochemistry* (2nd edition). Academic Press, San Diego, CA.

Pauli, N., Oberthür, T., Barrios, E. and Conacher, A.J. 2010. Fine-scale spatial and temporal variation in earthworm surface casting activity in agroforestry fields, western Honduras. *Pedobiologia* 53: 127-139.

Pelosi, C., Toutous, L., Chiron, F., Dubs, F., Hedde, M., Muratet, A., Ponge, J.F., Salmon, S. and Makowski, D. 2013. Reduction of pesticide use can increase earthworm populations in wheat crops in a European temperate region. *Agriculture, Ecosystems and Environment* 181: 223-230.

Piccolo, A., Cozzolino, A., Conte, P. and Spaccini, R. 2000. Polymerization of humic substances by an enzyme-catalyzed oxidative coupling. *Naturwissenschaften* 87: 391-394.

Rastogi, G. and Sani, R.K. 2011. Molecular Techniques to assess microbial community structure, function, and dynamics in the environment. In *Microbes and Microbial Technology*, eds. I. Ahmad, F. Ahmad and J. Pichtel, pp. 29-57. Springer Science+Business Media, LLC.

Robinson, C.H., Ineson, P., Piearce, T.G. and Parrington, J. 1996. Effects of earthworms on cation and phosphate mobilization in limed peat soils under *Picea sitchensis*. *Forest Ecology and Management* 86: 253-258.

Ross, D.J. and Cairns, A. 1982. Effects of earthworms and ryegrass on respiratory and enzyme activities of soil. *Soil Biology and Biochemistry* 14: 583-587.

Ruzicka, D.R., Barrios-Masias, F.H., Hausmann, N.T., Jackson, L.E. and Schachtman, D.P. 2010. Tomato root transcriptome response to a nitrogen-enriched soil patch. *BMC Plant Biology* 10: 75.

Sabrina, D.T., Hanafi, M.M., Gandahi, A.W., Mahmud, T.M.M. and Nor Azwady, A.A. 2013. Effect of mixed organic-inorganic fertilizer on growth and phosphorus uptake of setaria grass (*Setaria splendida*). *Australian Journal of Crop Science* 7: 75-83.

Sabrina, D.T., Hanafi, M.M., Mahmud, T.M.M. and Nor Azwady, A.A. 2011. Evaluation of nutrients released from phosphorous-enriched empty oil palm fruit bunches as growing media using *Setaria splendida*. *Compost Science and Utilization* 19: 61-68.

Sabrina, D.T., Hanafi, M.M., Nor Azwady, A.A. and Mahmud, T.M.M. 2009. Earthworm populations and cast properties in the soils of oil palm plantations. *Malaysian Journal of Soil Science* 13: 29-42.

Sabtu, M. 2006. *National report on the implementation of UNCCD: combating land degradation and promoting sustainable land resource management in Malaysia*. Third National Report for the Fifth Session of the Committee for the Review of the Implementation of the Convention to Combat Desertification (CRIC5). pp. 1-9.

Sadeghi-Shoae, M., Paknejad, F., Darvishi, H.H., Mozafari, H., Moharramzadeh, M. and Tookalloo, M.R. 2013. Effect of intermittent furrow irrigation, humic acid and deficit irrigation on water use efficiency of sugar beet. *Annals of Biological Research* 4: 187-193.

Samaranayake, J.W.K. and Wijekoon, S. 2010. Effect of selected earthworms on soil fertility, plant growth and vermicomposting. *Tropical Agricultural research and Extension* 13: 33-40.

Sastry, K.S. 2013. Methods of combating seed-transmitted virus diseases. In: *Seed-borne Plant Virus Diseases*, ed. K.S. Sastry, pp. 185-284. Springer, India.

Scheu, S. 1992. Automated measurement of the respiratory response of soil micro-compartments: Active microbial biomass in earthworm faeces. *Soil Biology and Biochemistry* 24: 1113-1118.

Scheu, S. 2003. Effects of earthworms on plant growth: patterns and perspectives. *Pedobiologia* 47: 846-856.

Schimel, J.P. and Bennet, J. 2004. Nitrogen mineralization: challenges of a changing paradigm. *Ecology* 85: 591-602.

Schmidt, O., Scrimgeour, C.M. and Curry, J.P. 1999. Carbon and nitrogen stable isotope ratios in body tissue and mucus of feeding and fasting earthworms (*Lumbricus festivus*). *Oecologia* 118: 9-15.

Shairul R.O., Osumanu, H.A., Shaharudin, S. and Nik Muhamad, A.M. 2008. Gamma Radiosensitivity Study on Chili (*Capsicum annuum*). *American Journal of Applied Science*s 5: 67-70.

Shen, H.P. and Yeo, D.C.J. 2005. Terrestrial earthworms (Oligochaeta) from Singapore. *The Raffles Bulletin of Zoology* 53: 13-33.

Shipitalo, M.J., Dick, W.A. and Edwards, W.M. 2000. Conservation tillage and macropore factors that affect water movement and the fate of chemicals. *Soil Tillage Research* 53: 167-183.

Shivakumar, K.M., Nagaraja, M.S., Champa, B.V. and Kuligod, V.B. 2010. Response of chili to applied nutrients and its influence on important soil properties in the upper Krishna project command. *Karnataka Journal of Agricultural Science* 23: 437-441.

Simek, M., Pizl, V. and Chalupsky, J. 1991. The effect of some terrestrial Oligochaeta on nitrogenase activity in soil. *Plant and Soil* 137: 161-165.

Sinha, R.K. and Heret, S. 2012. Organic farming: producing chemical-free, nutritive and protective food for the society while also protecting the farm soil by earthworms and vermicompost – reviving the dreams of sir Charles Darwin. *Agricultural Science Research Journal* 2: 217-239.

Sinha, R.K., Valani, D., Chauhan, K. and Agarwal, S. 2010. Embarking on a second green revolution for sustainable agriculture by vermiculture biotechnology using earthworms: Reviving the dreams of Sir Charles Darwin. *Journal of Agricultural Biotechnology and Sustainable Development* 2: 113-128.

Siti Khadijah, C.A., Misri, K. and Hishammuddin Omar, R.N. 2011. The effect of goat dung on growth of maize. In *Proceeding of the Fundamental Science Congress*, 5–6th July 2011, Universiti Putra Malaysia, Selangor, Malaysia. pp. 215-216.

Stockdill, S.M.J. 1982. Effects of introduced earthworms on the productivity of New Zealand pastures. *Pedobiologia* 24: 29-35.

Suthar, S. 2009. Impact of vermicompost and composted farmyeard manure on growth and yield of garlic (*Allium stivum* L.) field crop. *International Journal of Plant Productions* 3: 27-38.

Suthar, S. 2012. Earthworm production in cattle dung vermicomposting system under different stocking density loads. *Environmental Science and Pollution Research* 19: 748-755.

Suthar, S. and Singh, S. 2008. Vermicomposting of domestic waste by using two epigeic earthworms (*Perionyx excavatus* and *Perionyx sansibaricus*). *International Journal of Environmental Science and Technology* 5: 99-106.

Tabatabaei, S.J., Fatemi, L.S. and Fallahi, E. 2006. Effect of ammonium:nitrate ratio on yield, calcium concentration, and photosynthesis rate in strawberry, *Journal of Plant Nutrition* 29: 1273-1285.

Tabatabai, M.A. 1982. Soil enzymes. In *Methods of Soil Analysis, Part 2. Chemical and Microbial Properties*, eds. A.L. Page, R.H. Miller and D.R. Keeney, pp. 501-538. ASA, SSSA, Madison.

Tao, J., Griffiths, B., Zhang, S.J., Chen, X.Y., Liu, M.Q., Hu, F. and Li, H.X. 2009. Effects of earthworms on soil enzyme activity in an organic residue amended rice-wheat rotation agro-ecosystem. *Applied Soil Ecology* 42: 221-226.

Tapia-coral, S.C., Luizão, F., Barros, E., Pashanasi, B. and Castillo, D.D. 2006. Effect of *Pontoscolex corethrurus* Muller, 1857 (Oligochaeta: Glossoscolecidae) inoculation on litter weight loss and soil nitrogen in mesocosms in the Preruvian Amazon. *Carribean Journal of Science* 3: 410-418.

Tayel, M.Y., Abd El-Hady, M. and Eldardiry, E.I. 2010. Soil structure affected by some soil characteristics. *American-Eurasian Journal of Agricultural and Environmental Science* 7: 705-712.

Teng, S.K., Nor Azwady, A.A., Muskhazli, M., Suraini, A.A. and Yan, Y.W. 2012. Evaluation on physical, chemical and biological properties of casts of geophagous earthworm, *Metaphire tschiliensis tschiliensis*. *Scientific Research and Essays* 7: 1169-1174.

Teng, S.K., Nor Azwady, A.A., Noraziha, A., Muskhazli, M., Ahmad, I. and Yan, Y.W. 2013. Earthworm diversity and population density in the Kaki Bukit agroecosystem, Perlis, Peninsular Malaysia. *Tropical Ecology* 54: 289-296.

Tisdall, J.M. 1978. Ecology of earthworms in irrigated orchards. In *Modification of Soil Structure*, eds. W.W. Emerson, R.R. Bond and A.R. Dexter. pp. 297-303. Wiley, Chichester, UK.

Tiunov, A.V. and Scheu, S. 1999. Microbial respiration, biomass, biovolume and nutrient status in burrow walls of *Lumbricus terrestris* L. (Lumbricidae). *Soil Biology and Biochemistry* 31: 2039-2048.

Topoliantz, S. and Ponge, J.F. 2003. Burrowing activity of the geophagous earthworm *Pontoscolex corethrurus* (Oligochaeta: Glossoscolecidae) in the presence of charcoal. *Applied Soil Ecology* 23: 267-271.

Topoliantz, S., Ponge, J.F. and Ballof, S. 2005. Manioc peel and charcoal: a potential organic amendment for sustainable soil fertility in the tropics. *Biology and Fertility of Soils* 41: 15-21.

Toriyama, J., Ohnuki, Y., Ohta, S., Kosugi, K., Kabeya, N., Nobuhiro, T., Shimizu, A., Tamai, K., Araki, M., Keth, S. and Chann, S. 2013. Soil physicchemical properties and moisture dynamics of a large soil profile in a tropical monsoon forest. *Geoderma* 198: 205-211.

Troeh, F.R and Thompson, L.M. 2005. *Soils and Soil Fertility*. Blackwell Publishing, Iowa, US.

Tsai, C.F., Shen, H.P. and Tsai, S.C. 2000. Occurrence of the exotic earthworm *Pontoscolex corethrurus* (Müller) (Glossoscolecidae: Oligochaeta) in Taiwan. *Endemic Species Research* 2: 68-73.

Uchida, R. 2000. Essential nutrients for plant growth: nutrient functions and deficiency symptoms. In *Plant Nutrient Management in Hawaii's Soils, Approaches for Tropical and Subtropical Agriculture*, eds. J.A. Silva and R. Uchida, pp. 31-55. College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa.

Valdrighi, M.M., Pera, A., Agnolucci, M., Frassinetti, S., Lunardi, D. and Vallini, G. 1996. Effects of compost-derived humic acids on vegetable biomass production and microbial growth within a plant (*Cichorium intybus*) – soil system: a comparative study. *Agriculture, Ecosystems and Environment* 58: 133-144.

van der Werff, P.A., Noordhuis, R. and Bekkers, Th.B.M. 1998. Introduction of earthworms into an organic arable farming system. *Applied Soil Ecology* 9: 311-317.

van Gestel, C.A.M. and Koolhaas, J.E. 2004. Water-extractability, free ion activity and pH explain cadmium sorption and toxicity to *Folsomia candida* (Collembola) in seven soil–pH combinations. *Environmental Toxicology and Chemistry* 23: 1822–1833.

van Gestel, C.A.M., van Dis, W.A., Dirven-Van Breemen, E.M., Sparenburg, P.M. and Baerselman, R. 1991. Influence of cadmium, copper and pentachlorophenol on growth and sexual development of *Eisenia andrei* (Oligochaeta: Annelida). *Biology and Fertility of Soils* 12: 117-121.

Van Ranst, E., Verloo, M., Demeyer, A. and Pauwels, J.M. 1999. *Manual for Soil Chemistry and Fertility Laboratory: Analytical Methods for Soils and Plants Equipment and Management of Consumables*. University of Ghent, Belgium. pp. 166-168.

Vijiandran, J.R. and Gurmit, S. 2009. Earthworm population dynamics in the oil palm ecosystem on coastal and inland soils of United Plantation Berhad. In *Soil Health: Preserving Resources for Sustainable Agriculture*, eds. Z. Malik, C.F. Ishak, K.J. Goh, R. Abdullah, J. Vijayanathan, U.K. Md. Shah and Z. Hashim, pp. 49-53. Malaysian Society of Soil Science (MSSS) and Malaysia Agriculture Research & Development Institute (MARDI).

Vinotha, S.P., Parthasarathi, K. and Ranganathan, L.S. 2000. Enhanced phosphatase activity in earthworm casts in more of microbial origin. *Current Science* 79: 1158-1159.

Welke, S.E. and Parkinson, D. 2003. Effect of *Aporrectodea trapezoides* activity on seedling growth of *Pseudotsuga menziesii*, nutrient dynamics and microbial activity in different forest soils. *Forest Ecology and Management* 173: 169-186.

Whalen, J.K. and Parmelee, R.W. 1999. Quantification of nitrogen assimilation efficiencies and their use to estimate organic matter consumption by the earthworms *Aporrectodea tuberculata* (Eisen) and *Lumbricus terrestris* L. *Applied Soil Ecology* 13: 199-208.

Widarto, T.H., Holmstrup, M. and Forbes, V.E. 2004. The influence of nonylphenyl on life-history of the earthworm *Dendrobaena octaedra* Savigny: linking effects from the individual- to the population-level. *Ecotoxicology and Environmental Safety* 58: 147-159.

Wilkinson, G.E. 1975. Effect of grass fallow rotations on the infiltration of water into a savanna zone of northern Nigeria. *Tropical Agriculture* 52: 97-103.

Wolters, V. 2000. Invertebrate control of soil organic matter stability. *Biology and Fertility of Soils* 31: 1-19.

Wurst, S. and Jones, T.H. 2003. Indirect effects of earthworms (*Aporrectodea caliginosa*) on an above-ground tritrophic interaction. *Pedobiologia* 47: 91–97.

Xu, H.L., Wang, X.J. and Fujita, M. 2001. Effects of organic farming practices on photosynthesis, transpiration and water relations, and their contributions to fruit yield and the incidence of leaf-scorch in pear trees. *Journal of Crop Production* 3: 127-138.

Yan, Y.W., Nor Azwady, A.A., Shamsuddin, S.Z.H., Muskhazli, M., Suraini, A.A. and Teng, S.K. 2012. Vermicomposting potential and plant nutrient contents in rice straw vermicast of *Perionyx excavatus* and *Eudrilus eugeniae*. *Scientific Research and Essays* 7: 3639-3645.

Zhang, B.G., Li, G.T., Shen, T.S., Wang, J.K. and Sun, Z. 2000. Changes in microbial biomass C, N and P and enzyme activities in soil incubated with the earthworm *Metaphire guillelmi* or *Eisenia fetida*. *Soil Biology and Biochemistry* 32: 2055-2062.

Zhang, H., Yang, X.D., Du, J. and Wu Y.X. 2008. Influence of soil temperature and moisture on cocoon production and hatching of the exotic earthworm *Pontoscolex corethrurus*. *Zoological Research* 29: 305-312.

Zhang, M., Zou, X.M. and Schaefer, D.A. 2010. Alteration of soil labile organic carbon by invasive earthworms (*Pontoscolex corethrurus*) in tropical rubber plantations. *European Journal of Soil Biology* 46: 74-79.

Zhang, Yi., Wang, G.C., Wu, Y.P., Zhao, H., Zhang, Y.F. and Sun, Z.J. 2013. PCR-DGGE analysis of earthworm gut bacteria diversity in stress of *Escherichia coli* O157:H7*. *Advances in Bioscience and Biotechnology* 4: 437-441.

Zund, P.R., Pillai-Mc Garry, U., Mc Garry, D. and Bray, S.G., 1997. Repair of a compacted Oxisol by the earthworm *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta). *Biology and Fertility of Soils* 25: 202-208.