



**UNIVERSITI PUTRA MALAYSIA**

***BIOLOGY AND CHEMICAL COMPOSITIONS OF *Plotosus canius*  
HAMILTON, 1822 IN THE COASTAL WATERS OF PORT DICKSON,  
MALAYSIA***

**BINTA ISYAKU USMAN**

**FP 2014 32**



**BIOLOGY AND CHEMICAL COMPOSITIONS OF *Plotosus canius*  
HAMILTON, 1822 IN THE COASTAL WATERS OF PORT DICKSON,  
MALAYSIA**

**By**

**BINTA ISYAKU USMAN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the Requirement for the Degree of Doctor of Philosophy**

**November 2014**

## **COPYRIGHT**

All material contained within the thesis including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of all material contained within the thesis for non-commercial purposes from the copyright holder. Commercial Use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia

Copyright © Universiti Putra Malaysia



## DEDICATIONS

*This thesis is dedicated to my Husband, Dr. Sabo Wada Dutse and my children;  
Umar, Aisha, Maryam and Usman*



© “Allah is indeed The Most Generous for it is He Who teaches by the pen and teaches man that which he knew not”

(QUR’AN96:3-5)

Abstract of thesis was presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Doctor of Philosophy

**BIOLOGY AND CHEMICAL COMPOSITIONS OF *Plotosus canius*  
HAMILTON, 1822 IN THE COASTAL WATERS OF PORT DICKSON,  
MALAYSIA**

By

**BINTA ISYAKU USMAN**

**November 2014**

**Chairperson: S. M. Nurul Amin, PhD**

**Faculty: Agriculture**

*Plotosus canius*, the Gray-eel catfish and locally known as “Sembilang or Semilan” in Malaysia, is a popular and well-known marine catfish with high commercial value and usually sold fresh in the markets. Literature on the fish is very scant, and the fish has now been reported to be declining in the wild. The present study was undertaken to investigate the population stock status, fatty acid and amino acid composition, feeding habits, reproductive cycle, sex ratio and fecundity of *P. canius* in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia between January and December 2012. The samples were collected from the local fishermen of the area during the “full moon” and were immediately stored in ice chest and transported to the laboratory for the various analyses. The data were analyzed using SPSS version 20, Minitab version 16 and FiSAT II software.

The length frequency distribution of *P. canius* implied that the total length ranged from 22.80 to 62.90 cm (mean  $\pm$  SD,  $38.34 \pm 8.19$  cm) for males and from 24.60 to 60.00 cm ( $39.06 \pm 7.72$  cm) for females. There was no significant difference observed between males and females in the size frequency distribution of *P. canius* (t-test,  $p > 0.05$ ; Kolmogorov-Smirnov test:  $d_{\max} = 1.409$ ,  $p > 0.01$ ). The negative allometric nature of growth was observed in both males ( $b = 2.707$ ) and females ( $b = 2.879$ ). Analysis of the relationships between TL and various morphometric characteristics of the fish showed that there was strong and significant ( $R^2 > 0.805$ ,  $p < 0.001$ ) relationship. The length frequency data of *P. canius* analyzed revealed that asymptotic length ( $L_{\infty}$ ) and growth coefficient (K) were estimated at 67.20 cm and  $0.95 \text{ yr}^{-1}$  respectively. The growth performance index ( $\phi'$ ) was calculated as 3.63. Total mortality (Z), natural mortality (M) and fishing mortality (F) were estimated at  $2.73 \text{ yr}^{-1}$ ,  $1.43 \text{ yr}^{-1}$  and  $1.31 \text{ yr}^{-1}$  respectively. The exploitation level (E) of *P. canius* was calculated as 0.48.

Examination of the stomach fullness revealed that 61.54% of the stomachs contained food at various degrees of fullness, while the remaining (34.46%) were empty. According to index of preponderance ( $I_p$ ), the stomach contents of *P. canius* were composed of six major groups viz; fish and fish parts (38.00%), crustaceans (26.69%), molluscs (25.58%), sand and mud (6.68%), debris and detritus (2.99%) and unidentified items (0.06%). In the fish and fish parts group, fish eggs (37.49%)

was the dominant food item and this was followed by scales and other partly digested fish parts (0.36%) and small fish (0.15 %) respectively. Among the crustaceans group, *Portunus* spp was the dominant (1.59%), followed by *Neopisserma* spp (0.50%), *Charabdis* spp (0.42%), *Acetes* spp (0.39%) and *Sesarma* spp (0.07%). *Pholas* spp (1.87%) and *Anadara* spp (1.64%) were the dominant molluscs genera observed in the stomach of *P. canius*.

Examination of the proximate composition of the eggs, juvenile and adult of *P. canius* revealed that protein ranged between 16.39 and 25.04% (mean =  $19.68 \pm 4.68\%$ ) and fat was in the range of 3.65 and 5.10% ( $4.68 \pm 0.93\%$ ). Nine essential amino acids namely; histidine, threonine, valine, methionine, isoleucine, leucine, phenylalanine arginine and lysine were observed in both the eggs, juvenile and adult of *P. canius*. Saturated fatty acids were the highest (61.62 – 77.25%,  $66.95 \pm 8.92\%$ ) followed by monounsaturated fatty acids (17.57 – 33.82%,  $27.92 \pm 8.99\%$ ), while polyunsaturated fatty acids were the least (4.56 – 5.64%,  $5.13 \pm 0.53\%$ ) in all the eggs, juvenile and adult.

The overall sex ratio (males: females) of *P. canius* was observed to be 1:0.98 and did not differ significantly from the hypothetical ratio 1:1. Analysis of the annual variation of gonadosomatic index (GSI) showed that the major spawning activity in *P. canius* lasts between May and June in both sexes. 50% of males and females of *P. canius* were found to mature at sizes between 44 - 48 cm and 40 - 44 cm of total lengths respectively. Fecundity of the fish ranged from 642.26 to 1140.34 ( $943.05 \pm 38.40$ ) eggs per fish. The average fecundity per 1 kg of body weight was found to be 1225.89 eggs, and was established to show positive and significant relationship with body length ( $R^2 = 0.8662$ ,  $p < 0.05$ ), body weight ( $R^2 = 0.8556$ ,  $p < 0.05$ ) and ovary weight ( $R^2 = 0.9527$ ,  $p < 0.05$ ). The mean monthly GSI of females *P. canius* indicated positive and significant correlation with salinity ( $r = 0.613$ ;  $p < 0.05$ ). No significant correlation was observed between the mean monthly GSI and the remaining four variables (temperature, pH, dissolved oxygen and total suspended solids).

Overall it could be concluded that *P. canius* is carnivorous bottom feeder. , The ratio of essential to non-essential amino acids was found to be healthy. The major spawning period was between May and June and the status of exploitation ( $E = 0.48$ ) was slightly below the optimum level in the coastal waters of Port Dickson, Malaysia.

Abstrak tesis dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**BIOLOGI DAN KOMPOSISI KIMIA *Plotosus canius* HAMILTON, 1822 DI PERAIRAN PANTAI PORT DICKSON, MALAYSIA**

Oleh

**BINTA ISYAKU USMAN**

**November 2014**

**Pengerusi: S. M. Nurul Amin, PhD**

**Fakulti: Pertanian**

*Plotosus canius*, dikenali dengan nama tempatan “Sembilang or Semilan” di Malaysia, adalah ikan keli laut yang disukai dan terkenal dengan nilai pasaran yang tinggi serta bisanya dijual segar di pasar. Rujukan berkenaan dengan ikan ini adalah agak kurang, dan ikan ini dilaporkan semakin berkurangan di habitat semulajadi mereka. Kajian ini telah dijalankan bagi menyelidik status stok populasi, komposisi asid lemak dan asid amino, tabiat pemakanan, kitaran pembiakan, nisbah jantina dan fekunditi *P. canius* di perairan pantai Kg. Telok, Port Dickson, Semenanjung Malaysia antara Januari dan Diember 2012. Sampel diperoleh dari nelayan tempatan ketika “bulan penuh” and segera disimpan di dalam kotak air batu dan dipindahkan ke makmal bagi pelbagai analisis. Data tersebut dianalisis dengan menggunakan SPSS versi 20, Minitab versi 16 dan perisian FiSAT II.

Taburan kekerapan panjang *P. canius* menunjukkan bahawa jumlah panjang adalah dari 22.80 hingga 62.90 cm ( $\text{min} \pm \text{SD}$ ,  $38.34 \pm 8.19$  cm) untuk jantan dan dari 24.60 hingga 60.00 cm ( $39.06 \pm 7.72$  cm) untuk betina. Tidak ada perbezaan ketara didapati antara jantan dan betina dalam taburan kekerapan *P. canius* (ujian-t,  $p > 0.05$ ; ujian Kolmogorov-Smirnov:  $d_{\text{max}} = 1.409$ ,  $p > 0.01$ ). Keadaan alometrik negatif pada tumbesaran didapati pada kedua-dua jantan ( $b = 2.707$ ) dan betina ( $b = 2.879$ ). Analisis hubungan antara TL dan pelbagai ciri morfometrik ikan ini menunjukkan hubungan yang kuat dan ketara ( $R^2 > 0.805$ ,  $p < 0.001$ ). Data kekerapan panjang *P. canius* yang telah dianalisis mendedahkan bahawa panjang asimtotik ( $L_{\infty}$ ) dan koefisien pertumbuhan ( $K$ ) dianggarkan sebanyak 67.20 cm dan  $0.95 \text{ yr}^{-1}$ . Indeks prestasi pertumbuhan ( $\phi'$ ) dianggarkan sebanyak 3.63. Jumlah kematian ( $Z$ ), kematian semulajadi ( $M$ ) dan kematian tangkapan ( $F$ ) telah dianggarkan kira-kira  $2.73 \text{ yr}^{-1}$ ,  $1.43 \text{ yr}^{-1}$  dan  $1.31 \text{ yr}^{-1}$ . Tahap eksploitasi *P. canius* dikira sebanyak 0.48.

Pemeriksaan kepenuhan isi perut mendedahkan bahawa 61.54% dari kandungan perut terdiri daripada pelbagai tahap kepenuhan, manakala baki (34.46%) adalah kosong. Mengikut indeks praponderan ( $I_p$ ), kandungan perut *P. canius* adalah terdiri daripada enam kumpulan utama; ikan dan bahagian badan ikan (38.00%), krustasia (26.69%), moluska (25.58%), pasir dan lumpur (6.68%), debris dan detritus (2.99%) dan benda tidak dikenali (0.06%). Dalam kumpulan ikan dan bahagian badan faktor ikan, telur ikan (37.49%) adalah dominan dan ini diikuti dengan sisik dan bahagian ikan yang separa hadam (0.36%) dan ikan kecil (0.15 %). Di antara kumpulan krustasia, *Portunus* spp merupakan dominan (1.59%), diikuti oleh *Neopisserma* spp



(0.50%), *Charabdis* spp (0.42%), *Acetes* spp (0.39%) dan *Sesarma* spp (0.07%). *Pholas* spp (1.87%) dan *Anadara* spp (1.64%) pula merupakan genera moluska dominan yang dikesan di dalam perut *P. canius*.

Pemeriksaan komposisi proksimat telur, juvenil dan induk *P. canius* menunjukkan protein berjulat di antara 16.39 dan 25.04% (min =  $19.68 \pm 4.68\%$ ) dan lemak berjulat di antara 3.65 dan 5.10% ( $4.68 \pm 0.93\%$ ). Sembilan asid amino; histidin, threonin, valin, methionin, isoleusin, leusin, phenylalanin, arginin dan lysin telah dikesan pada kedua-dua telur, juvenil dan induk *P. canius*. Asid lemak tepu merupakan yang tertinggi (61.62 – 77.25%,  $66.95 \pm 8.92\%$ ) diikuti asid lemak mono tak tepu (17.57 – 33.82%,  $27.92 \pm 8.99\%$ ), manakala asid lemak poli tak tepu merupakan yang paling sedikit (4.56 – 5.64%,  $5.13 \pm 0.53\%$ ) pada telur, juvenil dan induk.

Nisbah keseluruhan jantina (jantan: betina) *P. canius* diperhatikan sebanyak 1:0.98 and dan tidak ketara perbezaannya dari nisbah hipotetikal 1:1. Analisis tahunan kepelbagaian indeks gonadosomatik (GSI) menunjukkan bahawa aktiviti pembiakan utama bagi *P. canius* berlangsung dari Mei dan Jun bagi kedua-dua jantina. 50% jantan dan betina *P. canius* didapati matang pada saiz antara jumlah panjang 44 - 48 cm dan 40 - 44 cm. Fekunditi ikan ini adalah sekitar 642.26 hingga 1140.34 ( $943.05 \pm 38.40$ ) telur per ikan. Purata fekunditi per 1 kg berat badan adalah didapati sebanyak 1225.89 telur, dan menunjukkan hubungan positif dan ketara dengan panjang badan ( $R^2 = 0.8662$ ,  $p < 0.05$ ), berat badan ( $R^2 = 0.8556$ ,  $p < 0.05$ ) dan berat ovari ( $R^2 = 0.9527$ ,  $p < 0.05$ ). Min bulanan GSI bagi betina *P. canius* menunjukkan korelasi positif dan ketara dengan kemasinan ( $r = 0.613$ ;  $p < 0.05$ ). Tiada korelasi ketara didapati di antara min bulanan GSI dan empat baki parameter (suhu, pH, oksigen terlarut dan jumlah pepejal termendap).

Keseluruhannya dapat disimpulkan bahawa *P. canius* ialah pemakan dasar karnivor. Nisbah antara asid amino penting dengan tidak penting didapati berada dalam keadaan baik. Tempoh pembiakan utama adalah antara Mei dengan Jun status eksploitasi ( $E = 0.48$ ) adalah berada bawah sedikit dari tahap optimum di kawasan perairan pantai Port Dickson, Malaysia.



## ACKNOWLEDGEMENTS

In the name of Allah, the most beneficent, the most eternally merciful. Thank you Dear Lord for enabling me to submit this thesis.

Firstly, my deep gratitude and whole hearted appreciations to the chairman supervisory committee and advisor, Dr. S. M. Nurul Amin for his tireless support, patience, guidance, advice and encouragement throughout the period of my studies at UPM. My profound thanks also goes to my co-supervisors; Prof. Dr. Aziz Arshad and Prof. Dr. Mohd Salleh Kamarudin for their constructive comments and suggestions on many aspects of this work which enabled me to pursue this research work to completion.

I will like to offer my thanks and appreciation to Universiti Putra Malaysia (UPM) firstly for providing partial financial support through the International Graduate Research Fellowship (IGRF) and secondly for the research grant (Grant No. 05-02-12-1703RU, Vote No. 9327300) under the Research University Grant Scheme (RUGS). I would also like to offer my sincere gratitude to my employer, Federal College of Education Kano-Nigeria, for giving me a chance to further my education. In addition thanks to my laboratory mates Mohd Hazmadi Zakariya and Mohd Azim Bin Mohd Khatib who are always readily available to introduce me to the various fishermen and fishing sites around Peninsular Malaysia. Thank you again may Allah reward you abundantly. My thanks also go to Dr. Mahmoud Danaee, who always help and guide me with the statistical analyses. The fishermen, Zul and Amad can also not be forgotten for their valuable contributions during the field sampling. Also, my thanks to Mr Kamarul Arifin Hadithon of the Chromatography unit, Faculty of Food Science and Technology, UPM for his assistance during the chemical composition analyses.

My special appreciation goes to my family especially my mother for the love and taking care of my kids while I was away from home (Nigeria) for the entire period of my studies at Malaysia. I am also thankful to all my well-wishers.

Last, but not the least my heartfelt thanks and love goes to my dear and beloved husband, Dr. Sabo Wada Dutse for his encouragement, patience, support and unremitting love throughout these years. I certainly would not be able to get through without my pillars of strength. Thank you and may Allah takes you to greater heights.

I certify that a Thesis Examination Committee has met on 24 November 2014 to conduct the final examination of Binta Isyaku Usman on her thesis entitled "Biology and Chemical Compositions of *Plotosus canius* Hamilton, 1822 in the Coastal Waters of Port Dickson, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Rosimah bt Nulit, **PhD**  
Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

Annie Christianus, **PhD**  
Senior Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Internal Examiner)

Hishamuddin b Omar, **PhD**  
Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

A. Jesu Arockia Raj, **PhD**  
Professor  
Faculty Science and Humanities  
Srm University, Kattankulathur  
(External Examiner)



---

**ZULKARNAIN ZAINAL, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 12 March 2015

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**S. M. NURUL AMIN, PhD**

Senior Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**AZIZ BN ARSHAD, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**MOHD SALLEH KAMARUDIN, PhD**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

### **Declaration by graduate student**

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No: Binta Isyaku Usman (GS30858)



## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: \_\_\_\_\_

Name of

Chairman of

Supervisory

Committee: \_\_\_\_\_

  
Dr. S. M. Nurul Amin  
Lecturer

Department of Aquaculture

Faculty of Agriculture

Universiti Putra Malaysia

43400 UPM Serdang, Selangor

AMIN

Signature: \_\_\_\_\_

Name of

Member of

Supervisory

Committee: \_\_\_\_\_



Prof. Dr. Aziz Arshad

Jabatan Akuakultur

Fakulti Pertanian

Universiti Putra Malaysia

43400 UPM Serdang, Selangor

Signature: \_\_\_\_\_

Name of

Member of

Supervisory

Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	<b>ABSTRACT</b>	<b>Page</b>
	<b>ABSTRAK</b>	i
	<b>ACKNOWLEDGEMENTS</b>	iii
	<b>APPROVAL</b>	v
	<b>DECLARATION</b>	vi
	<b>LIST OF TABLES</b>	viii
	<b>LIST OF FIGURES</b>	xiii
	<b>LIST OF ABBREVIATIONS</b>	xv
	<b>LIST OF ABBREVIATIONS</b>	xvii
	<b>CHAPTER</b>	
1	<b>GENERAL INTRODUCTION</b>	1
	1.1 Background of the study	1
	1.2 Statement of the problem	2
	1.3 Significance of the study	2
	1.4 Objectives of the study	3
	1.5 Outline of the thesis	3
2	<b>LITERATURE REVIEW</b>	5
	2.1 Family plotosidae	5
	2.2 Distribution of <i>Plotosus canius</i>	5
	2.3 Taxonomy and identification key	7
	2.4 External morphology	8
	2.5 Morphometric study	10
	2.6 Morphology of the digestive tract	11
	2.7 Reproductive biology	12
	2.8 Food and feeding habits	15
	2.9 Length-weight relationship (LWR)	16
	2.10 Chemical compositions	16
	2.11 The fisheries of <i>P. canius</i>	18
3	<b>GENERAL METHODOLOGY</b>	21
	3.1 Introduction	21
	3.2 Materials and methods	21
	3.2.1 Study location	21
	3.2.2 Sampling procedure	21
	3.2.3 Sample identification and measurements	24
	3.2.4 Population dynamics	24
	3.2.5 Food habits	25
	3.2.6 Chemical compositions	25
	3.2.7 Reproductive biology	25
	3.2.8 Statistical analyses	25
	3.2.9 Physico-chemical parameters	26
	3.3 Results of physico-chemical parameters	26

3.3.1	Temperature	26
3.3.2	pH	26
3.3.4	Dissolved oxygen	27
3.3.4	Salinity	27
3.3.5	Total suspended solids (TSS)	27
4	<b>POPULATION DYNAMICS AND STOCK STATUS OF <i>Plotosus canius</i> IN THE COASTAL WATERS OF KG. TELOK, PORT DICKSON, PENINSULAR MALAYSIA</b>	28
4.1	Introduction	28
4.2	Materials and methods	28
4.2.1	Collection of samples	28
4.2.2	Laboratory measurements	28
4.2.3	Data analysis	29
4.3	Results	33
4.3.1	Population structure	33
4.3.2	Length-weight relationship	34
4.3.3	Relationship between total length and the various morphometric characteristics	37
4.3.4	Condition factor (K)	44
4.3.5	Population parameters of <i>P. canius</i>	44
4.4	Discussion	51
4.6	Conclusions	53
5	<b>FOOD HABITS OF <i>Plotosus canius</i> IN THE COASTAL WATERS OF KG. TELOK, PORT DICKSON, PENINSULAR MALAYSIA</b>	55
5.1	Introduction	55
5.2	Materials and Methods	56
5.2.1	Sample collection	56
5.2.2	Laboratory process	56
5.2.3	Stomach examination	56
5.2.4	Stomach contents analysis	57
5.3	Results	57
5.3.1	Feeding intensity	57
5.3.2	Diet composition	57
5.3.3	Monthly variation in diet composition	61
5.4	Discussion	66
5.5	Conclusions	68
6	<b>PROXIMATE AND CHEMICAL COMPOSITIONS OF EGGS AND MUSCLE OF <i>Plotosus canius</i> IN THE COASTAL WATERS OF KG. TELOK PORT DICKSON, PENINSULAR MALAYSIA</b>	69
6.1	Introduction	69
6.2	Materials and Methods	70
6.2.1	Sample collection and processing	70
6.2.2	Proximate composition	70
6.2.3	Amino acid composition	71



6.2.4	Fatty acid composition	73
6.3	Results	74
6.3.1	Proximate composition	74
6.3.2	Amino acid composition	74
6.3.3	Fatty acid composition	75
6.4	Discussion	76
6.5	Conclusions	82
7	<b>SOME ASPECTS OF REPRODUCTIVE BIOLOGY OF <i>Plotosus canius</i> IN THE COASTAL WATERS OF KG. TELOK, PORT DICKSON, PENINSULAR MALAYSIA</b>	83
7.1	Introduction	83
7.2	Materials and methods	83
7.2.1	Sample collection	83
7.2.2	Laboratory preparation	83
7.2.3	Sex ratio	83
7.2.4	Gonad examination	84
7.2.5	Gonadosomatic index (GSI)	84
7.2.6	Size at first sexual maturity	84
7.2.7	Fecundity	85
7.2.8	Statistical analysis	85
7.3	Results	85
7.3.1	Sex ratio	85
7.3.2	Gonad maturity stage	87
7.3.3	Reproductive cycle	93
7.3.4	Size at first sexual maturity	96
7.3.5	Fecundity and type of spawning	98
7.4	Discussion	102
7.5	Conclusions	105
8	<b>CONCLUSIONS AND RECOMMENDATIONS</b>	106
8.1	Conclusions	106
8.2	Recommendations	107
	<b>REFERENCES</b>	109
	<b>APPENDICES</b>	132
	<b>BIODATA OF STUDENT</b>	139
	<b>LIST OF PUBLICATIONS</b>	140

## LIST OF TABLES

Table		Page
2.1	Distribution and occurrence of <i>P. canius</i> by country (Froese and Pauly, 2011)	6
2.2	Review of reproductive biology of <i>P. canius</i>	14
2.3	Landings (tonnes) of <i>P. canius</i> by state (Negeri Sembilan), East and West coast of Peninsular Malaysia from 1990 – 2013 based on Malaysian Fisheries Statistics (1990 – 2013)	20
3.1	Monthly variation of the different physico-chemical parameters in the coastal waters of Kg. Telok, Port Dickson from January to December, 2012	27
4.1	The morphometric characteristics of <i>P. canius</i> used in the study and their descriptions	31
4.2	Bimonthly length frequency data of <i>P. canius</i> samples collected between January and December from the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	32
4.3	Length-weight relationship parameters of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	35
4.4	Summary of the descriptive statistics of the various morphometric characters of males and females <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	38
4.5	Summary of regression analysis of the various morphometric characters of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	39
4.6	Estimated population parameters of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	46
5.1	Empirical scale of <i>P. canius</i> stomach fullness separated into five classes as outlined for this study	56
5.2	Percentage of stomach fullness of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	59
5.3	Overall diet composition of <i>P. canius</i> ranked by index of preponderance (Ip) in the coastal waters of Kg. Telok, Port Dickson	60
5.4	Index of preponderance (Ip) of prey items in 234 stomachs of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	63
6.1	The proximate composition of the eggs, juvenile and adult of <i>P. canius</i>	74
6.2	Amino acid compositions (mg/g wet weight) of eggs, juvenile and adult of <i>P. canius</i>	75
6.3	Fatty acid composition (% wet weight) of eggs, juvenile and adult of <i>P. canius</i>	76
6.4	A comparison of chemical compositions in <i>P. canius</i> with other fishes and red meats	78
6.5	Comparison of the EAA/ NEAA ratio of amino acids in <i>P. canius</i> with other fishes	80
6.6	Comparison of the ratios of PUFA/ SFA and n-3/ n-6 of fatty acids in <i>P. canius</i> with other fishes	81
7.1	Monthly variation in sex ratio of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	86
7.2	Variation in sex ratio of different size groups of <i>P. canius</i>	86

7.3	Mean oocyte size and GSI of different maturity stages in <i>P. canius</i>	87
7.4	Fecundity of <i>P. canius</i> at various length groups	99



## LIST OF FIGURES

Figure		Page
2.1	Yellow dot; showing the geographical distribution of <i>P. canius</i> around the world	6
2.2	Diagram showing the maxillary barbel of <i>P. canius</i> extending well behind the eye	7
2.3	Diagram showing external features of <i>P. canius</i>	8
2.4	Ventral view of anterior portion of <i>P. canius</i>	9
2.5	Bucco- pharyngeal region of <i>P. canius</i>	10
2.6	Nature and disposition of alimentary tract of <i>P. canius</i>	11
3.1	Geographical location of the sampling station in Peninsular Malaysia	22
3.2	Photograph of <i>Plotusus canius</i> (Gray-eel catfish)	23
3.3	Photograph of local fish trap 'Lukah Ikan' which is used to catch <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	23
3.4	Photograph of set net which is used to catch <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	24
4.1	Morphometric characteristics on (a) body and (b) head used for <i>P. canius</i> in the study	30
4.2	Annual size frequency distribution of male and female <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	34
4.3	Length-weight relationship of male <i>P. canius</i> (logarithmic and arithmetic scale) in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	36
4.4	Length-weight relationship of female <i>P. canius</i> (logarithmic and arithmetic scale) in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	36
4.5	Length-weight relationship of male and female <i>P. canius</i> (logarithmic and arithmetic scale) in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	37
4.6	Relationship between total length and various morphometric characteristics in males (red dot) and females (blue dot) <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	40- 41
4.7	Monthly Fulton (K) condition factor of male and female <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia	44
4.8	Predicted maximum lengths of <i>P. canius</i> based on extreme value theory (Formacion <i>et al.</i> , 1991). The predicted maximum length value and the 95% confidence interval is obtained from the intersection of overall maximum length with the line y and x, z respectively	45
4.9	K-scan routine for determination of best growth curvature giving best value of asymptotic length ( $L_{\infty}$ ) with growth performance index of <i>P. canius</i>	45
4.10	von Bertalanffy growth curve of <i>P. canius</i> superimposed on the restructured length-frequency histograms. The black and white bars are positive and negative deviation from the "weighed" moving average of three length classes and they represent psedo-cohorts	47
4.11	Plot of age and growth based on computed mean growth parameters of <i>P. canius</i> ( $L_{\infty}$ = 67.20 cm and K = 0.95)	48

4.12	Length converted catch curve of <i>P. canius</i> , the darkened full dots represent the points used in calculating through least square linear regression and the open dots represent the points either not fully recruited or nearing to $L_{\infty}$	49
4.13	Logistic selection curve for probability of capture showing 25%, 50% and 75% selection length of <i>P. canius</i>	49
4.14	Recruitment pattern of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	50
4.15	Length-structured virtual population analysis of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	50
4.16	Yield-per-recruit and biomass-per-recruit model, showing level of yield index in <i>P. canius</i> from the coastal waters of Kg. Telok, Port Dickson	51
5.1	Digestive tract of <i>P. canius</i> showing the position of the stomach	58
5.2	The index of preponderance (Ip) of the prey items of <i>P. canius</i> from the coastal waters of Port Dickson	59
5.3	Some prey items observed in the stomachs of <i>P. canius</i>	61
5.4	Index of preponderance (%Ip) of three most dominant groups of food items in the stomach of <i>P. canius</i> in the coastal waters of Kg. Telok, Port Dickson	65
7.1	Part of immature testis of male <i>P. canius</i> , stage I	88
7.2	Part of maturing testis of male <i>P. canius</i> , stage II	88
7.3	Part of mature testis of male <i>P. canius</i> , stage III	89
7.4	Part of ripe testis of male <i>P. canius</i> , stage IV	89
7.5	Part of spent testis of male <i>P. canius</i> , stage V	90
7.6	Part of immature ovary of female <i>P. canius</i> , stage I	91
7.7	Part of maturing ovary of female <i>P. canius</i> , stage II	91
7.8	Part of mature ovary of female <i>P. canius</i> , stage III	92
7.9	Part of ripe ovary of female <i>P. canius</i> , stage IV	92
7.10	Part of spent ovary of female <i>P. canius</i> , stage V	93
7.11	Monthly changes in the percentage of occurrence of gonad stage of males <i>P. canius</i>	94
7.12	Monthly changes in the percentage of occurrence of gonad stage of females <i>P. canius</i>	94
7.13	Monthly variation of GSI of males <i>P. canius</i> during January to December, 2012	95
7.14	Monthly variation of GSI of females <i>P. canius</i> during January to December, 2012	95
7.15	A plot of GSI and egg size against the various gonadal development stages of females <i>P. canius</i>	96
7.16	Length at 50% maturity of males <i>P. canius</i>	97
7.17	Length at 50% maturity of females <i>P. canius</i>	97
7.18	Relationship between fecundity and total length of <i>P. canius</i>	98
7.19	Relationship between fecundity and body weight of <i>P. canius</i>	100
7.20	Relationship between fecundity and ovary weight of <i>P. canius</i>	101
7.21	Frequency of occurrences of eggs of different class intervals of mature females <i>P. canius</i>	102

## LIST OF ABBREVIATIONS

a	Intercept/ constant
AA	Amino acid
ANOVA	Analysis of variance
AOAC	Association of Official Analytical Chemists
b	Coefficient of length-weight relationship/ slope
B	Both sexes
BDA	Body depth at anus
B/R	Relative biomass-per-recruit
CA	Cluster analysis
CC	Chemical composition
CI	Confident of interval
cm	Centimeter
Cn	Composition by number
CRD	Completely randomized design
Cw	Composition by weight
DFA	Discriminant function analysis
DO	Dissolved oxygen
E	Exploitation rate
$E_{\max}$	Maximum allowable limit of exploitation
E-50	Exploitation level at which 50% of the relative biomass-per-recruit
EAA	Essential amino acid
ED	Eye diameter
EL	Eye length
ELEFAN	Electronic Length Frequency Analysis
ES	Egg size
F	Fishing mortality
FAA	Flavour amino acids
FAMEs	Fatty acid methyl esters
FAO	Food and agriculture organization
FiSAT	FAO ICLARM Stock Assessment Tools
FO	Frequency of occurrence
GC	Gas chromatography
GSI	Gonadosomatic index
GW	Total weight of ovary
HL	Head length
HPLC	High pressure liquid chromatography
HW	Head width
IOL	Inter orbital length
Ip	Index of preponderance
IUCN	International Union for Conservation Of Nature
K	Growth coefficient of VBGF
Kg.	'Kampung'
$L_c$	Length at first capture
$L_{\infty}$	Asymptotic length
$L_{\max}$	Predicted extreme length
LLR	Length-length relationship
LWR	Length-weight relationship

M	Natural mortality
mm	Millimeter
MUFA	Monounsaturated fatty acid
MG	Mouth gape
MSY	Maximum Sustainable Yield
N	Sample size
n-3	Omega-3
n-6	Omega-6
NEAA	Non-essential amino acid
PAD	Pre anal distance
PCA	Principal component analysis
PDD	Pre dorsal distance
POL	Post orbital length
PPD	Pre pectoral distance
PUFA	Polyunsaturated fatty acid
PVD	Pre ventral distance
R <sup>2</sup>	Coefficient of determination
Rn	Response surface
SD	Standard deviation
SE	Standard error
SFA	Saturated fatty acid
SL	Standard length
SNL	Snout length
Sp.	Species
SPSS	Statistical package for social science
TL	Total length
TSS	Total suspended solids
TW	Total weight
T <sub>max</sub>	Maximum life span
UPM	Universiti Putra Malaysia
USFA	Unsaturated fatty acid
Y	Estimated annual catch
Y/F	Average standing stock
Y/R	Relative yield per recruit
Y/U	Total annual catch
Z	Total mortality
φ'	Growth performance index
%	Percentage
<	Less than
>	More than
°C	Degree Celsius



## CHAPTER 1

### GENERAL INTRODUCTION

#### 1.1 Background of the study

*Plotosus canius* Hamilton, 1822 (Gray-eel catfish or catfish eel), is a member of the family plotosidae locally called 'Sembilang or Semilan' in Malaysia. The fish is native to South-east Asia and Australia (Mohsin and Ambak, 1996; Ferraris, 2007) and is easily recognized by its long, eel-like and sub-cylindrical body that is tapered and flattened near the tail region. It possesses four long pairs of barbels; with the nasal barbel extending well behind the eye and almost reaching the nape and has no stripes on its body (Mohsin and Ambak, 1996; Gupta and Gupta, 2006). The fish primarily found in marine habitat but sometimes can be caught in brackish or fresh water habitats (Riede, 2004). According to Kottelat (2001), *P. canius* occur in coastal seas and fresh or brackish waters. Juveniles are commonly found to form compact aggregates, thus resulting in very tight shoals with about 50 juvenile fish (Mohsin and Ambak, 1996; Ambak *et al.*, 2010). It has been found to live on or near the bottom of the sea and migrate between sea and fresh water (Riede, 2004).

There has been high rate of consumption of marine fishes among the rural and urban adults in Malaysia. The daily prevalence consumption rate of marine fish among rural adults has been estimated to be at 51% and 34% for urban adults (Norimah Jr *et al.*, 2008). This is an indication that consumption of marine fish in Malaysia is very high. *P. canius* is important both commercially, being sold fresh at the market (Gomon, 1984; Mohsin and Ambak, 1996; Ambak *et al.*, 2010) and as a source of food (Nurnadia *et al.*, 2011). In Malaysia, it has been reported to be mostly harvested by small scale fishermen and being usually sold fresh in villages and coastal towns, hence the real harvest data may not be revealed in the official statistics as such the official catch statistics could be regarded as gross underestimation (Leh *et al.*, 2012). This fish was mentioned to be among the ten most commonly marine fishes preferred to be eaten by Malaysians (Osman *et al.*, 2001). Also, Nurnadia *et al.* (2011) revealed *P. canius* to be among the five fish that are commonly consumed from Malacca Straits of Malaysia. The fish is highly priced and good for eating (Gupta and Gupta, 2006; Ambak *et al.*, 2010).

The fishery sector has been recognized as a source of income and livelihood for lots of people around the globe (Chowdhury *et al.*, 2011). It has provided full time employment to about 1.2 million people globally, out of which the marine fishery sub-sector provided employment to about 0.5 million people (DOF, 2010). There is increased dependence on marine and coastal resources by humans (Berkas *et al.*, 2001), this has thus results to the dwindling of the captured marine fisheries resources (DOF Bangladesh, 2010). There is also need of increasing concern on basic aspects of biology in marine fish (Mace, 1994; Murawski *et al.*, 2001). Knowledge of reproduction, feeding and other basic aspects of biology of any fish according to McAllister *et al.* (2000) are very useful parameters in the management of any fish stock .

## 1.2 Statement of the problem

The population of the earth is increasing at all times, as such food demand is increasing. To sustain this demand, the exploitation of wild stocks of marine organisms has intensified substantially. The fish production has attained a level that no further increase from the coastal waters is feasible. Also, natural stocks have been exploited to their utmost limits and this resulted to over fishing (Godfray *et al.*, 2010; Burgess *et al.*, 2013; Islam *et al.*, 2013, Molfese and Hall-Spencer, 2014). Marine fisheries are very vital worldwide source of food and livelihoods. However, in recent times issues related to habitat destruction and fish stock depletion have threatened many fishery resources and the fishery industry at large. This is particularly true in the case of the *P. canius* and the related fishery stocks. In such circumstances, the conservation and fishery management turn out to be the need of the time.

It has been reported that *P. canius* stocks throughout their range have declined due to overfishing, illegal fishing, indiscriminate fishing of brood stocks and juveniles, fast degradation of marine habitat, land development, agricultural activities and introduction of the alien species (Khan *et al.*, 2002; Ahmed and Haque, 2007). Although, the status of this fish has not been assessed in the IUCN red list (IUCN, 2013), it has been declared endangered in Bangladesh and India (Mukhopadhyay, 1994; IUCN Bangladesh, 2000; Mijkherjee *et al.*, 2002).

Availability of information on the biology of any fish species plays a vital role in its conservation and management. However, very little information in terms of literature on this fish is available both from the national and international perspective. Most studies on basic biology of this fish have only been carried out in a small number of areas from other countries of its distribution (Sinha, 1981, 1984, 1986a, 1986b; Khan *et al.*, 2002; Ahmed and Haque, 2007). Presently, only the work of Leh *et al.* (2012) on the feeding habit of *P. canius* was found from Malaysia. Effective management and conservation of *P. canius* stocks require knowledge on aspects of its biology. This is clear indication for more research on this fish, so as to understand some aspect of its basic biology which could be very useful in the management and conservation of this fish.

## 1.3 Significance of the study

Availability of information on the biology of any fish species plays a vital role in its conservation and management. Effective management and conservation of *P. canius* stocks require knowledge of aspects of its biology. Understanding the length-weight and length-length relationships, condition factor, age and growth and population dynamics of *P. canius* are essential in describing its general life history and are more important from a management view point. Data on age and growth of this fish will be useful for understanding the age composition of the fish and the position of different class-years in its fisheries. It will also be vital in determining the mortality and survival rate of the different year-classes.

Information on diet and feeding of *P. canius* is vital for understanding of its ecological role and life history such as growth, breeding and migration. It is also important fisheries management and conservation. Without such knowledge, it may not be possible to apprehend the predicted changes that could rise from any natural

or anthropogenic intervention. Information on food habits of the fish could also be useful in the aquaculture of the fish in order to obtain the best growth at least time. Furthermore, understanding the chemical composition of the fish will reveal its quality and nutritional status. It will also be useful in the formulation of the fish diets in aquaculture.

Investigation of reproductive biology is vital and required for the conservation and propagation of the fish. Several aspects of the reproductive procedure are usually employed either to catch the fish or to protect them if they are excessively vulnerable. Therefore, investigating aspects of reproductive biology of *P. canius* is indispensable and necessary prior to the measures adopted for its conservation and propagation. Understanding of its spawning period is one of the most essential requisites in its management and rational exploitation. Moreover, knowing the fecundity of *P. canius* can be used to evaluate its reproductive potential and will be very vital for successful commercialization. The information will also help greatly in offering advice for possible management possibilities to the fishery industry, fisheries managers, fishermen as well as the planners and policy makers.

In view of the commercial importance of *P. canius* to the Malaysians, study on age, growth, population dynamics, food habits, chemical compositions and reproductive biology of the fish from coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia is very essential in order to offer the much needed information for exploitation of its full potential for both capture and culture fisheries. Lack of attempt to manage the *P. canius* fishery resources in Malaysia can lead to its extinction, and this can affect the livelihood of the fishermen and the Malaysian coastal population at large.

#### **1.4 Objectives of the study**

1. To examine the population parameters and determine the stock status of *P. canius* from the coastal waters of Kg. Telok Port Dickson, Peninsular Malaysia.
2. To investigate the stomach contents and determine the food habits of *P. canius*.
3. To determine the proximate and chemical compositions of *P. canius*.
4. To determine sex ratio, sexual maturity, spawning season and fecundity of *P. canius*.

#### **1.5 Outline of the thesis**

The thesis is structured in the form of eight chapters.

- i. Chapter 1: This chapter discussed the general introduction and background of the subject, the statement of the problem, significance of the study and finally ended with the thesis outline.
- ii. Chapter 2: The chapter critically reviewed related and relevant literature on the basic aspects of biology, as well as chemical compositions of *P. canius*.

- iii. Chapter 3: The general thesis methodology was presented in this chapter. The chapter explained the various methods used in this research and the statistical techniques used in the data analyses.
- iv. Chapter 4: This chapter was based on the first objective of the research. It investigated the population parameters and stock status of *P. canius* from the coastal waters of Kg. Telok, Port Dickson, Peninsular Malaysia.
- v. Chapter 5: Objective two of this research work was the basis of this chapter. It reported aspects of food habits of *P. canius*. An investigation of the stomach contents and seasonal variation in the diet of the fish was made and reported in this chapter.
- vi. Chapter 6: This chapter answered the third objective of the study by explaining the chemical compositions of the fish. It reported the proximate, amino and fatty acid compositions of the egg, juvenile as well as the adult of *P. canius*.
- vii. Chapter 7: This chapter was based on the fourth objective of the research. It examined aspects of reproductive biology, such as sex ratio, size at sexual maturity and spawning of the fish.
- viii. Chapter 8: This chapter draws the conclusions of various chapters of the work and gives overall conclusions on this work. It also contained prospective suggestions and recommendations for future studies.

## REFERENCES

- Abascal, F. J. and Medina, A., 2005. Ultrastructure of oogenesis in the bluefin tuna, *Thunnus thynnus*. *Journal of Morphology*, 264 (2): 149-160.
- Adebiyi, F., Siraj, S., Harmin, S. and Christianus, A., 2013. Plasma sex steroid hormonal profile and gonad histology during the annual reproductive cycle of river catfish *Hemibagrus nemurus* (Valenciennes, 1840) in captivity. *Fish Physiology and Biochemistry*, 39 (3): 547-557.
- Adebiyi, F. A., Siraj, S. S., Harmin, S. A. and Christianus, A., 2011. Ovarian development of a river catfish *Hemibagrus nemurus* (Valenciennes, 1840) in captivity. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315 (9): 536-543.
- Ahmed, S. U. and Haque, A., 2007. Studies on the fishery biology and domestication of gang magur (*Plotosus canius*). Bfri Annual progress report. Bangladesh: Bangladesh Fisheries research institute, pp. 76-79.
- Alhassan, E. H. and Ansu-Darko, M., 2011. Food and feeding habits of a potential aquaculture candidate, the black Nile catfish, *Bagrus bajad* in the Golinga reservoir. *Australian Journal of Basic and Applied Sciences*, 5 (5): 354-359.
- Alimon, A. R., Roustaian, P., Saad, C. R. and Kamarudin, M. S., 2003. Lipid content and fatty acid composition during early and late embryonic development of redclaw crayfish, *Cherax quadricarinatus* (Crustacea, decapoda). *Journal of Applied Ichthyology*, 19 (6): 397-398.
- Allison, M. E. and Sikoki, F. D., 2013. Food and feeding habits of *Parailia pellucida* (Boulenger, 1901) (Schilbeidae) in the freshwater reaches of the nun river of the Niger Delta, Nigeria. *International Journal of Advanced Fisheries and Aquatic Science*, 1 (1): 1-14.
- Ama-Abasi, D. and Ogar, A., 2013. Proximate Analysis of snakehead fish, *Parachanna obscura*, (Gunther 1861) of the Cross River, Nigeria. *Journal of Fisheries and Aquatic Science*, 8 (1): 295-298.
- Amani, A. A., Amin, S. M. N. and Arshad, A., 2011. Stomach contents of sergestid shrimp *Acetes japonicus* from the estuary of Tanjung Dawai Peninsular Malaysia. *Journal of Fisheries and Aquatic Sciences*, 6 (7): 771-779.
- Ambak, M. A., Isa, M., Zakaria, Z. and Ghaffar, M. A., 2010. *Fishes of Malaysia* (First ed.). Malaysia: Pernebit Universiti Malaysia Terengganu.
- Amin, S. M. N., 2001. Studies on age and growth, VPA analysis and relative condition factor of *Harpodon nehereus* (Ham-Buch) from the neritic water of Bangladesh. *Online Journal of Biological Sciences*, 1: 192-194.
- Amin, S. M. N., Arshad, A., Bujang, J. S., Siraj, S. S. and Goddard, S., 2009. Reproductive biology of the sergestid shrimp *Acetes indicus* (Decapoda:



Sergestidae) in coastal waters of Malacca, Peninsular Malaysia. Zoological Studies, 48: 753-760.

Amin, S. M. N., Arshad, A., Haldar, G. C., Shohaimi, S. and Ara, R., 2005. Estimation of size frequency distribution, sex ratio and length-weight relationship of Hilsa (*Tenualosa ilisha*) in the Bangladesh water. Research Journal of Agriculture and Biological Sciences, 1 (1): 61-66.

Anthony, J. A., Roby, D. D. and Turco, K. R., 2000. Lipid content and energy density of forage fishes from the northern Gulf of Alaska. Journal of Experimental Marine Biology and Ecology, 248 (1): 53-78.

AOAC, 1990. Official Methods of Analysis. Vol. I. 15th ed. AOAC, Arlington, VA.

Arafa, S., Chouaibi, M., Sadok, S. and El Abed, A., 2012. The influence of season on the gonad index and biochemical composition of the sea urchin *Paracentrotus lividus* from the Gulf of Tunis. The Scientific World Journal, 2012. 8 pp.

Arai, S., 1981. A purified test diet for coho salmon, *Oncorhynchus kisutch*, fry. Bulletin of the Japanese Society of Scientific Fisheries, 47: 547-550.

Araújo, A. S. d., do Nascimento, W. S., Yamamoto, M. E. and Chellappa, S., 2012. Temporal dynamics of reproduction of the Neotropical fish, *Crenicichla menezesi* (Perciformes: Cichlidae). The Scientific World Journal, vol. 2012, Article ID 579051, 10 pages.

Arendt, M. D., Olney, J. E. and Lucy, J. A., 2001. Stomach content analysis of cobia, *Rachycentron canadum*, from lower Chesapeake Bay. Fishery bulletin-national oceanic and atmospheric administration, 99 (4): 665-670.

Armstrong, M. J. and Witthames, P. R., 2012. Developments in understanding of fecundity of fish stocks in relation to egg production methods for estimating spawning stock biomass. Fisheries Research, 117-118 (2012): 35-47.

Arafa, S., Chouaibi, M., Sadok, S. and El Abed, A., 2012. The influence of season on the gonad index and biochemical composition of the sea urchin *Paracentrotus lividus* from the Gulf of Tunis. The Scientific World Journal, 2012, 8 pp.

Arshad, A., Amin, S. M., Nuradiella, Y. L. Z., Cob, Z. C., Ara, R. and Aziz, D., 2012. Population characteristics of *A. japonicus* from the Kedah coastal waters of Peninsular Malaysia. Journal of Fisheries and Aquatic Science, 7 (2): 162-172.

Aytekin, Y. and Yüce, R., 2008. Ovary maturation stages and histological investigation of ovary of the Zebrafish (*Danio rerio*). Brazilian Archives of Biology and Technology, 51 (3): 513-522.

- Azadi, M. A., Nasiruddin, M. and Rahman, A., 2013. Food and feeding habits of the clupeid, *Gonialosa manmina* (Ham.) from the Kaptai Lake, Bangladesh. Chittagong University Journal of Biological Sciences, 4 (1): 53-61.
- Azadi, M. A. and Rahman, A. S. M. S., 2013a. Morphometric and meristic study of *Gudusia chapra* (Ham. 1822) and *Gonialosa manmina* (Ham. 1822) (Clupeidae) from the Kaptai lake, Bangladesh. Chittagong University Journal of Biological Sciences, 3 (1): 21-31.
- Azadi, M. A. and Rahman, A. S. M. S., 2013b. Reproduction biology of the clupeid. *Gonialosa manmina* (Hamilton, 1822) from the Kaptai lake Bangladesh. Chittagong University Journal of Biological Sciences, 3 (1): 139-148.
- Azadi, M. A. and Ullah, M., 2009. Food and feeding habits of the ribbon fish, *Lepturacanthus savala* (Cuvier, 1829) from the bay of Bangladesh. Journal of the Asiatic Society of Bangladesh, 35 (1): 57-64.
- Azadi, M. A. and Ullah, M., 2013. Length-weight relationship and relative condition factor of the ribbon fish, *Lepturacanthus savala* (Cuvier, 1829) from the bay of Bengal, Bangladesh. Chittagong University Journal of Biological Sciences, 3 (1): 119-126.
- Bae, J. H., Yoon, S. H. and Lim, S. Y., 2011. Heavy metal contents and chemical compositions of atlantic (*Scomber scombrus*), blue (*Scomber australasicus*), and chub (*Scomber japonicus*) mackerel muscles. Food Science and Biotechnology, 20 (3): 709-714.
- Balan, V., 1984. The Indian oil sardine fishery: A review. Marine Fisheries Information Service, Technical and Extension Series, 60: 1-10.
- Barbin, D. F., El Masry, G., Sun, D.W. and Allen, P., 2013. Non-destructive determination of chemical composition in intact and minced pork using near-infrared hyperspectral imaging. Food Chemistry, 138 (2-3): 1162-1171.
- Berkes, F., Mahon, R. and McConney, P., 2001. *Managing small-scale fisheries: alternative directions and methods*: International Development Research Center. Ottawa, ON Canada K1G 3H9, pp. 308.
- Beverton, R. J. H. and Holt, S. J., 1966. *Manual of methods for fish stock assessment: Part 2-tables of yield functions*: Food and Agriculture Organization of the United Nations.
- Beverton, R. J. H. and Holt, S. J., 1993. *On the dynamics of exploited fish populations*. Vol. VI. In: Fish and Fisheries Series. Springer Netherlands. 538p.
- Bhavan, P. S., Radhakrishnan, S., Seenivasan, C., Shanthi, R., Poongodi, R. and Kannan, S., 2010. Proximate composition and profiles of amino acids and fatty acids in the muscle of adult males and females of commercially viable



prawn species *Macrobrachium rosenbergii* collected from natural culture environments. International Journal of Biology, 2 (2): 107-119.

- Bindu, L., Padmakumar, K. G., Sreerekha, P. S. and Joseph, N., 2012. Reproductive biology of the golden catfish, *Horabagrus brachysoma* (Günther, 1864), an endemic species of the Western Ghats, India. Journal of Applied Ichthyology, 28 (5): 772-777.
- Bligh, E. and Dyer, W. J., 1959. A rapid method of total lipid extraction and purification. Canadian Journal of Biochemistry and Physiology, 37 (8): 911-917.
- Body, D. R., 1985. The composition of orange roughy (*Hoplostethus atlanticus*) roe lipids. Journal of the Science of Food and Agriculture, 36 ( 8): 679-684.
- Bombata-Fashina, H. A., Megbowon, I., Olumide, O., Ozor, P. A., Ibrahim, A. O., Adejonwo, O. A. and Kolade, O. Y., 2013. Comparative study of the proximate composition of some wild tilapiine fishes in Epe lagoon, Lagos, Nigeria. Journal of Fisheries and Aquatic Science, 8 (1): 265-267.
- Boulenger, G. A., 1895. *Catalogue of the perciform fishes in the British Museum*. Vol. 1: Printed by order of the Trustees.
- Bowman, W. C. and Rand, M. J., 1980. *Textbook of Pharmacology*: Blackwell Scientific Publications.
- Burgess, M. G., Polasky, S. and Tilman, D., 2013. Predicting overfishing and extinction threats in multispecies fisheries. Proceedings of the National Academy of Sciences, 110 (40): 15943-15948.
- Cadrin, S. X., 2000. Advances in morphometric identification of fishery stocks. Reviews in Fish Biology and Fisheries, 10 (1): 91-112.
- Cao, M., Duan, J., Cheng, N., Zhong, X., Wang, Z., Hu, W. and Zhao, H., 2012. Sexually dimorphic and ontogenetic expression of *dmrt1*, *cyp19a1a* and *cyp19a1b* in *Gobiocypris rarus*. Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology, 162 (4): 303-309.
- Çek, S., Bromage, N., Randall, C. and Rana, K., 2001. Oogenesis, hepatosomatic and gonadosomatic indexes, and sex ratio in rosy barb (*Puntius conchonius*). Turkish Journal of Fisheries and Aquatic Sciences, 1: 33-41.
- Chelemal, M., Jamili, S. and Sharifpour, I., 2009. Reproductive biology and histological studies in Abu Mullet, *Liza abu* in the water of the Khozestan province. Journal of Fisheries and Aquatic Science, 4 (1): 1-11.
- Chowdhury, M. S. N., Hossain, M. S., Das, N. G. and Barua, P., 2011. Environmental variables and fisheries diversity of the Naaf River Estuary, Bangladesh. Journal of Coastal Conservation, 15 (1): 163-180.

- Chyun, J.H. and Griminger, P., 1984. Improvement of nitrogen retention by arginine and glycine supplementation and its relation to collagen synthesis in traumatized mature and aged rats. *The Journal of Nutrition*, 114 (9): 1697.
- Clarke, A., Skadsheim, A. and Holmes, L. J., 1985. Lipid biochemistry and reproductive biology in two species of Gammaridae (Crustacea: Amphipoda). *Marine Biology*, 88 (3): 247-263.
- Conceição, L. E. C., Grasdalen, H., and Rønnestad, I., 2003. Amino acid requirements of fish larvae and post-larvae: new tools and recent findings. *Aquaculture*, 227 (1): 221-232.
- Conner, W. E., Lin, D. S. and Colvis, C., 1996. Differential mobilization of fatty acids from adipose tissue. *Journal of Lipid Research*, 37 (2): 290-298.
- Cortez-Vega, W. R., Pizato, S. and Prentice, C., 2012. Quality of raw chicken breast stored at 5C and packaged under different modified atmospheres. *Journal of Food Safety*, 32 (3): 360-368.
- Corriero, A., Karakulak, S., Santamaria, N., Deflorio, M., Spedicato, D., Addis, P. and Vassallo-Agius, R., 2005. Size and age at sexual maturity of female bluefin tuna (*Thunnus thynnus* L. 1758) from the Mediterranean sea. *Journal of Applied Ichthyology*, 21 (6): 483-486.
- Cuvier, G. and Valenciennes, A., 1840. *Histoire Naturelle des Poissons*. Tome 14. Pitois-Levrault, Paris. 464 pp.
- Cowey, C. B. and Luquet, P., 1983. Physiological basis of protein requirements of fishes. Critical analysis of allowances. In: Arnal, M., Pion, R. and Bonin, D. (eds). *Protein Metabolism and Nutrition*, 1: 365-84.
- Das, M. K. and Bordoloi, S., 2013. Length-weight relationship and condition factor of *Lepidocephalichthys goalparensis* Pillai and Yazdani, 1976 in Assam, India. *Journal of Applied Ichthyology*, 30: 246-247.
- Discoverlife.org., 2011. World map showing the distribution of *P. canius*. Retrieved 12 October, 2011 from Discover Life website <http://www.discoverlife.org/>.
- De Bruin, G. H. P., Russell, B. C. and Bogusch, A., 1995. *FAO species identification field guide for fishery purposes*. The Marine Fishery Resources of Sri Lanka: 411 pp.
- De Lestang, S., Hall, N. G. and Potter, I. C., 2003. Reproductive biology of the blue swimmer crab (*Portunus pelagicus*, Decapoda: Portunidae) in five bodies of water on the west coast of Australia. *Fishery Bulletin*, 101(4): 745-757.
- DeMartini, E. E., Uchiyama, J. H. and Williams, H. A., 2000. Sexual maturity, sex ratio, and size composition of swordfish, *Xiphias gladius*, caught by the Hawaii-based pelagic longline fishery. *Fishery Bulletin*, 98(3): 489-506.

- DOF, Bangladesh, 2010. Fishery Statistical Yearbook of Bangladesh: 2008-2009: Department of Fisheries, Ministry of Fisheries and Livestock Dhaka.
- DOF, Malaysia, 2013. Annual Fisheries Statistics, Ministry of Agriculture, Malaysia.
- Dineshbabu, A. P. and Manissery, J. K., 2008. Reproductive biology of ridgeback shrimp *Solenocera choprai* (Decapoda, Penaeoidea, Solenoceridae) off Mangalore coast, south India. *Fisheries Science*, 74 (4): 796-803.
- Dobriyal, A. K. and Singh, H. R., 1993. Reproductive biology of a hillstream catfish, *Glyptothorax madraspatanum* (Day), from the Garhwal, Central Himalaya, India. *Aquaculture Research*, 24 (6): 699-706.
- Dupuis, A. W. and Sutton, T. M., 2011. Reproductive biology of female humpback whitefish *Coregonus pidschian* in the Chatanika River, Alaska. *Journal of Applied Ichthyology*, 27 (6): 1365-1370.
- Ecoutin, J.M., Albaret, J.J. and Trape, S., 2005. Length-weight relationships for fish populations of a relatively undisturbed tropical estuary: The Gambia. *Fisheries Research*, 72 (2): 347-351.
- Elamin, S. M., Ambak, M. A., Samoilys, M. A. and Hamza, M. E., 2011. Some morphometric relationships of coral trouts *Plectropomus pessuliferus* and *Plectropomus areolatus* inhabiting Sudanese red sea. *Advances in Environmental Biology*, 5 (9): 2860-2865.
- Erdem, M. E., Baki, B. and Samsun, S., 2009. Fatty acid and amino acid compositions of cultured and wild sea bass (*Dicentrarchus labrax* L., 1758) from different regions in Turkey. *Journal of Animal and Veterinary Advances*, 8: 1959-1963.
- Ferraris, C. J., 2007. *Checklist of catfishes, recent and fossil (osteichthyes, siluriformes) and catalogue of siluriform primary type*. USA: Magnolia press.
- Ferriz, R. A., Fernández, E. M., Bentos, C. A. and López, G. R., 2007. Reproductive biology of *Pseudocorynopoma doriai* (Pisces: Characidae) in the High Basin of the Samborombón River, province of Buenos Aires, Argentina. *Journal of Applied Ichthyology*, 23 (3): 226-230.
- Formacion, S. P., Rongo, J. M. and Sambilay Jr, V. C., 1991. Extreme value theory applied to the statistical distribution of the largest lengths of fish. *UP Research Digest*, Philippines.
- Froese, R., 1998. Length-weight relationships for 18 less-studied fish species. *Journal of Applied Ichthyology*, 14 (1-2): 117-118.
- Froese, R., 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22 (4): 241-253.

- Froese, R. and Pauly, D., 2011. FishBase 2011. world wide web electronic publication.
- Fry, F. E. J., 1949. Statistics of a lake trout fishery. *Biometrics*, 5 (1): 27-67.
- Gautam, K., Marichamy, G., Nazar, A. R. and Anand Ganesh, E., 2013. Fatty acid profiles of four marine edible fishes Parangipettai coast, India. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 3 (22): 71-74.
- Gayanilo, F. C., Soriano, M. and Pauly, D., 1988. A draft guide to the complete ELEFAN. ICLARM Software 2. International Center for Living Aquatic Resources Management, Manila, Philippines.
- Gayanilo, F. C., Sparre, P. and Pauly, D., 2005. *FAO-ICLARM stock assessment tools II: User's guide*. Vol. 8. Food and Agriculture Organization.
- Gibson, R. A., 1983. Australian fish-An excellent source of both arachidonic acid and  $\omega$ -3 polyunsaturated fatty acids. *Lipids*, 18 (11): 743-752.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F. and Toulmin, C., 2010. Food security: the challenge of feeding 9 billion people. *science*, 327 (5967): 812-818.
- Gomes, I. D., Araujo, F. G., Uehara, W. and Sales, A., 2011. Reproductive biology of the armoured catfish *Loricariichthys castaneus* (Castelnau, 1855) in Lajes reservoir, southeastern Brazil. *Journal of Applied Ichthyology*, 27 (6) 1322-1331.
- Gomes, I. D. and Gerson Araújo, F., 2004. Reproductive biology of two marine catfishes (Siluriformes, Ariidae) in the Sepetiba Bay, Brazil. *Revista de Biología Tropical*, 52 (1): 143-156.
- Gomiero, L. M. and Braga, F. M. S., 2004. Reproduction of species of the genus *Cichla* in a reservoir in southeastern Brazil. *Brazilian Journal of Biology*, 64 (3B): 613-624.
- Gomon, J. R., 1984. *Plotosidae*. Vol. 3, Western Indian Ocean fishing area 51: W. Fischer and G. Bianchi (eds).
- Gomon, J. R. and Taylor, W. R., 1982. *Plotosus Mkunga, a New Species of Catfish from South Africa, with a Redescription of Plotosus Limbatus Valenciennes and Key to the Species of Plotosus (Siluriformes: Plotosidae)*: JLB Smith Institute of Ichthyology. Special Publication, 22: 1-16.
- Gonçalves, J. M. S., Bentes, L., Lino, P. G., Ribeiro, J., Canario, A. V. M. and Erzini, K., 1997. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. *Fisheries Research*, 30 (3): 253-256.

- Grigorakis, K., 2007. Compositional and organoleptic quality of farmed and wild gilthead sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) and factors affecting it: A review. *Aquaculture*, 272 (1): 55-75.
- Guilliams, T. G., 2000. Fatty acids: essential therapeutic. A Concise Update of Important Issues Concerning Natural Health Ingredients, 3 (2): 1-8.
- Gulland, J. A., 1969. Manual of methods for fish stock assessment, part I. Fish population analysis. FAO Manual of Fisheries Science, 4: 154p.
- Gulland, J. A., 2013. The fish resources of the ocean. Fishing News Books, West Byfleet, England. 255pp.
- Gupta, S., and Banerjee, S., 2013. Studies on reproductive biology of *Mystus tengara* (Ham.-Buch., 1822), a freshwater catfish of West Bengal, India. *International Journal of Aquatic Biology*, 1 (4): 175-184.
- Gupta, S. K. and Gupta, P. C., 2006. *General and Applied Ichthyology: (Fish and Fisheries)* (First ed.). New Delhi, India: S. Chand and Company Ltd.
- Gustafsson, I.-B., Öhrvall, M., Ekstrand, B. and Vessby, B., 1996. Moderate amounts of n-3 fatty acid enriched seafood products are effective in lowering serum triglycerides and blood pressure in healthy subjects. *Journal of Human Nutrition and Dietetics*, 9 (2): 135-145.
- Hamilton, F., 1822. *An account of the fishes found in the river Ganges and its branches*. Edinburgh: Archibald Constable, vii: 405 pp.
- Harrison, T. D., 2001. Length-weight relationships of fishes from South African estuaries. *Journal of Applied Ichthyology*, 17 (1): 46-48.
- Hart, P. J. B. and Reynolds, J. D., 2008. *Handbook of fish biology and fisheries* (Vol. 2): Wiley. com.
- Haruna, A. B., 2003. *Aquaculture in the tropics: theory and practice*. Kano, Nigeria: Al-Hassana publishers.
- Hazmadi, M. Z., Amin, S. M. N., Arshad, A., Rahman, M. A. and Al-Barwani, S. M., 2011. Size frequency and length-weight relationships of spined anchovy, *Stolephorus tri* from the coastal waters of Besut, Terengganu, Malaysia. *Journal of Fisheries and Aquatic Science*, 6: 857-861.
- Hearn, T. L., Sgoutas, S. A., Hearn, J. A. and Sgoutas, D. S., 1987. Polyunsaturated fatty acids and fat in fish flesh for selecting species for health benefits. *Journal of Food Science*, 52 (5): 1209-1211.
- Hedgecock, D. and Pudovkin, A. I., 2011. Sweepstakes reproductive success in highly fecund marine fish and shellfish: a review and commentary. *Bulletin of Marine Science*, 87 (4): 971-1002.



- Henderson, J. R. and Tocher, D. R., 1987. The lipid composition and biochemistry of freshwater fish. *Progress in Lipid Research*, 26 (4): 281-347.
- Hennessy, R. J. and Stringer, C. B., 2002. Geometric morphometric study of the regional variation of modern human craniofacial form. *American Journal of Physical Anthropology*, 117 (1): 37-48.
- Heydarnejad, M. S., 2009. Length-weight relationships for six freshwater fish species in Iran. *Chinese Journal of Oceanology and Limnology*, 27 (1): 61-62.
- Hisar, O., Sönmez, A. Y., Hisar, Ş. A., Budak, H. and Gültepe, N., 2013. The sexually dimorphic adipose fin is an androgen target tissue in the brown trout (*Salmo trutta fario*). *Fish Physiology and Biochemistry*, 39: 233-241.
- Hoese, D. F., Bray, D. J., Allen, G. R., Paxton, J. R., Wells, A. and Beesley, P. L., 2006. Leiognathidae. Fishes. *Zoological Catalogue of Australia*, 35: 1-3.
- Hossain, M. Y., Ahmed, Z. F., Islam, A. B., Jasmine, S. and Ohtomi, J., 2010. Gonadosomatic index-based size at first sexual maturity and fecundity indices of the Indian River shad *Gudusia chapra* (Clupeidae) in the Ganges River (NW Bangladesh). *Journal of Applied Ichthyology*, 26 (4): 550-553.
- Hossain, M. Y., Ahmed, Z. F., Leunda, P. M., Jasmine, S., Osoz, J., Miranda, R. and Ohtomi, J., 2006. Condition, length-weight and length-length relationships of the Asian striped catfish *Mystus vittatus* (Bloch, 1794) (Siluriformes: Bagridae) in the Mathabhanga River, southwestern Bangladesh. *Journal of Applied Ichthyology*, 22 (4): 304-307.
- Hyndes, G. A. and Potter, I. C., 1997. Age, growth and reproduction of *Sillago schomburgkii* in south-western Australian, nearshore waters and comparisons of life history styles of a suite of *Sillago* species. *Environmental Biology of Fishes*, 49 (4): 435-447.
- Hynes, H. B. N., 1950. The food of fresh-water sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*), with a review of methods used in studies of the food of fishes. *The Journal of Animal Ecology*, 19 (1): 36-58.
- Hyslop, E. J., 1980. Stomach contents analysis-a review of methods and their application. *Journal of Fish Biology*, 17 (4): 411-429.
- Isa, M. M., Noor, N. S. M., Yahya, K. and Nor, S. A. M., 2012. Reproductive biology of estuarine catfish, *Arius argyropleuron* (Siluriformes: Ariidae) in the northern part of Peninsular Malaysia. *Journal of Biology, Agriculture and Healthcare*, 2 (3): 14-27.
- Islam, G. M. N., Noh, K. M., Yew, T. S. and Noh, A. F. M., 2013. Assessing environmental damage to marine protected area: a case of Perhentian marine park in Malaysia. *Journal of Agricultural Science*, 5 (8): 132.

- IUCN, 2013. IUCN red list of threatened species. Retrieved 15 August, 2013, from IUCN [www.iucnredlist.org](http://www.iucnredlist.org).
- IUCN Bangladesh, 2000. *Red book of threatened fish of Bangladesh*. The world conservation Union Appendix, 1, 61.
- Iwasaki, M. and Harada, R., 1985. Proximate and amino acid composition of the roe and muscle of selected marine species. *Journal of Food Science*, 50 (6): 1585-1587.
- Jacquot, R., 1961. Organic constituents of fish and other aquatic animal foods, in: Borgstrom, G. (Ed.) (1961). *Fish as food: 1. Production, Biochemistry and Microbiology*, pp. 145-209.
- Jensen, I.J., Dort, J. and Eilertsen, K.E., 2014. Proximate composition, antihypertensive and antioxidative properties of the semimembranosus muscle from pork and beef after cooking and *in vitro* digestion. *Meat Science*, 96 (2): 916-921.
- John, D. G., 1971. *Malay Poisons and Charm Cures*. London, Oxford University Press.
- Jones, R., 1984. *Assessing the effects of changes in exploitation pattern using length composition data (with notes on VPA and cohort analysis)*. Vol. 256, FAO Roma.
- Joyce, W. N., Campana, S. E., Natanson, L. J., Kohler, N. E., Pratt, H. L. and Jensen, C. F., 2002. Analysis of stomach contents of the porbeagle shark (*Lamna nasus* Bonnaterre) in the northwest Atlantic. *ICES Journal of Marine Science: Journal du Conseil*, 59 (6): 1263-1269.
- Kadye, W. T. and Booth, A. J., 2012. Integrating stomach content and stable isotope analyses to elucidate the feeding habits of non-native sharptooth catfish *Clarias gariepinus*. *Biological Invasions*, 14 (4): 779-795.
- Kailola, P. J. and Hoese, D. F., 1988. *The fishes of Papua New Guinea: a revised and annotated checklist* (Vol. 2). Myxinidae to Symbranchidae. Research Bulletin, No. 41, Port Moresby: Department of Fisheries and Marine Resources.
- Kalayci, F., Samsun, N., Bilgin, S. and Samsun, O., 2007. Length-weight relationship of 10 fish species caught by bottom trawl and midwater trawl from the middle black sea, Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 7: 33-36.
- Kamaruddin, I. S., Mustafa-Kamal, A. S., Christianus, A., Daud, S. K., Aminand, S. M. N. and Yu-Abit, L., 2011. Length-weight relationship and condition factor of three dominant species from the lake Tasik Kenyir, Terengganu, Malaysia. *Journal of Fisheries and Aquatic Science*, 6 (7): 1-5.



- Kamarudin, M. S., Mohsin, A. K. M., Mastol, M. R., Hutagalung, R. I., Chen, C. P., Wan Mohamed, W. E., Law, A. T. and Sivarajasingam, S., 1987. Chemical compositions of Malaysian freshwater catfish (Suborder-Siluroidea). In: Proceedings Conference of the Malaysian Society of Animal Production. Pp. 201-205. Malaysian Society of Animal Production, Serdang Malaysia.
- Kamrani, E., Sabili, A. N. and Yahyavi, M., 2010. Stock Assessment and reproductive biology of the blue swimming crab, *Portunus pelagicus* in Bandar Abbas coastal waters, Northern Persian Gulf. Journal of the Persian Gulf, 1 (2): 11-22.
- Kanwal, B. P. S. and Pathani, S. S., 2012. Food and feeding habits of a hill stream fish, *Garra lamta* (Hamilton-Buchanan) in some tributaries of Suyal River, Kumaun Himalaya, Uttarakhand (India). International Journal of Food and Nutrition Science, 1 (2): 16-22.
- Khan, M. S., Ambak, M. A., Ang, K. J. and Mohsin, A. K. M., 1990. Reproductive biology of a tropical catfish, *Mystus nemurus* Cuvier and Valenciennes, in Chenderoh reservoir, Malaysia. Aquaculture Research, 21 (2): 173-180.
- Khan, M. S., Ambak, M. A. and Mohsin, A. K. M., 2011. Food and feeding biology of a tropical freshwater catfish, *Mystus nemurus* Cuvier and Valenciennes with reference to its functional morphology. Indian Journal of Fisheries, 35 (2): 78-84.
- Khan, M. S. A., Alam, M. J., Rheman, S., Mondal, S. and Rahman, M. M., 2002. Study on the fecundity and GSI of brackishwater catfish *Plotosus canius* (Hamilton-Buchanan). OnLine Journal of Biological Sciences, 2 (4): 232-234.
- Kim, S. 2005. Population structure, growth, mortality, and size at sexual maturity of *Palaemon gravieri* (Decapoda: Caridea: Palaemonidae). Journal of Crustacean Biology, 25 (2): 226-232.
- Kinsella, J. E., Shimp, J. L., Mai, J. and Weihrauch, J., 1977. Fatty acid content and composition of freshwater finfish. Journal of the American Oil Chemists' Society, 54 (10): 424-429.
- Kottelat, M., 2001. *Fishes of Laos*: WHT Publication (Pty) Ltd, Cotta, Colombo. pp198.
- Kuiter, R. H. and Tono-zuka, T., 2001. *Pictorial guide to indonesian reef fishes*: Zoonetics, Seaford, Vic.
- Kumar, M. S., Xiao, Y., Venema, S. and Hooper, G., 2003. Reproductive cycle of the blue swimmer crab, *Portunus pelagicus*, off southern Australia. Journal of the Marine Biological Association of the UK, 83 (5): 983-994.
- Kumaran, R., Ravi, V., Gunalan, B., Murugan, S. and Sundramanickam, A., 2012. Estimation of proximate, amino acids, fatty acids and mineral composition of

mullet (*Mugil cephalus*) of Parangipettai, Southeast Coast of India. *Advances in Applied Science Research*, 3 (4); 2015-2019.

Lawson, E. O., 2010. Morphometric measurements and meristic counts in mudskipper (*Periophthalmus papilio*) from mangrove swamps of Lagos lagoon, Nigeria. *Journal of Applied Bioscience*, 34: 2166-2172.

Lawson, E. O., Akintola, S. L. and Awe, F. A., 2013. Length-weight relationships and morphometry for eleven (11) fish species from Ogudu creek, Lagos, Nigeria. *Advances in Biological Research*, 7 (4): 122-128.

Lawson, E. O., Thomas, A. E. and Nwabueze, A. A., 2011. Seasonal abundance, morphometric measurements and growth patterns in frill fin goby, *Bathygobius soporator* from Badagry creek, Lagos, Nigeria. *Asian Journal of Biological Sciences*, 4 (4): 325-339.

Le Cren, E. D., 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *The Journal of Animal Ecology*, 20: 201-219.

Leh, M. U. C., Sasekumar, A., and Chew, L. L., 2012. Feeding biology of eel catfish *Plotosus canius* Hamilton in a Malaysian mangrove estuary and mudflat. *The Raffles Bulletin of Zoology*, 60 (2): 551-557.

Liang, S.H., Wu, H.P. and Shieh, B.S., 2005. Size structure, reproductive phenology, and sex ratio of an exotic armored catfish (*Liposarcus multiradiatus*) in the Kaoping River of southern Taiwan. *Zoological Studies*, 44 (2): 252.

Lima-Junior, S. E. and Goitein, R., 2001. A new method for the analysis of fish stomach contents. *Acta Scientiarum*, 23 (2): 421-424.

Lovett, D. L., 1981. *A guide to the shrimps, prawns, lobsters, and crabs of Malaysia and Singapore*: Faculty of Fisheries and Marine Science, Universiti Pertanian Malaysia.

Lucifora, L. O., Valero, J. L. and García, V. B., 1999. Length at maturity of the green eye spurdog shark, *Squalus mitsukurii* (Elasmobranchii: Squalidae), from the SW Atlantic, with comparisons with other regions. *Marine and Freshwater Research*, 50 (7): 629-632.

Lungren, R., Staples, D., Funge-Smith, S., and Clausen, J., 2006. *Status and potential of fisheries and aquaculture in Asia and the Pacific*. RAP Publication. FAO Regional Office for Asia and the Pacific, Bangkok.

Mablouké, C., Kolasinski, J., Potier, M., Cuvillier, A., Potin, G., Bigot, L. and Jaquemet, S., 2013. Feeding habits and food partitioning between three commercial fish associated with artificial reefs in a tropical coastal environment. *African Journal of Marine Science*, 35 (3): 323-334.

- Mace, P. M., 1994. Relationships between common biological reference points used as thresholds and targets of fisheries management strategies. *Canadian Journal of Fisheries and Aquatic Sciences*, 51 (1): 110-122.
- Mambrini, M. and Kaushik, S. J., 1995. Indispensable amino acid requirements of fish: correspondence between quantitative data and amino acid profiles of tissue proteins. *Journal of Applied Ichthyology*, 11 (3-4): 240-247.
- Mamun, A., Tareq, K. M. A. and Azadi, M. A., 2004. Food and feeding habits of *Amblypharyngodon mola* (Hamilton) from Kaptai Reservoir, Bangladesh. *Pakistan Journal of Biological Sciences*, 7 (4): 584-588.
- Manon, M. R. and Hossain, M. D., 2013. Food and feeding habit of *Cyprinus carpio* var. *specularis*. *Journal of Science Foundation*, 9 (1-2): 163-169.
- Marimuthu, K., Arumugam, J., Sandragasan, D. and Jegathambigai, R., 2009. Studies on the fecundity of native fish climbing perch (*Anabas testudineus*, Bloch) in Malaysia. *American-Eurasian Journal of Sustainable Agriculture*, 3 (3): 266-274.
- Marriott, M. S., Booth, A. J. and Skelton, P. H., 1997. Reproductive and feeding biology of the Natal mountain catfish, *Amphilius natalensis* (Siluriformes: Amphiliidae). *Environmental Biology of Fishes*, 49 (4): 461-470.
- Mat Isa, M., Doroh, N.-I.-A., Ambri, F. K. and Wing Keong, N., 2012. Reproductive biology of the sleeper goby, *Butis gymnopomus* (Bleeker, 1853) from the Merbok estuary, Kedah, Malaysia. *Indian Journal of Fisheries*, 59 (4): 147-155.
- McAllister, M. K., Babcock, E. A., Pikitch, E. K. and Prager, M. H., 2000. Application of a non-equilibrium generalized production model to South and North Atlantic swordfish: combining Bayesian and demographic methods for parameter estimation. *Collective Volumes of Scientific Papers ICCAT*, 51 (5): 1523-1550.
- Meador, M. R. and Brown, L. M., 2014. Life history strategies of fish species and biodiversity in eastern USA streams. *Environmental Biology of Fishes*, 1-15.
- Mijkherjee, M., Praharaj, A. and Das, S., 2002. Conservation of endangered fish stocks through artificial propagation and larval rearing technique in West Bengal, India. *Aquaculture Asia*, 7 (2): 8-11.
- Minnis, R., Haq, I., Jackson, P., Yeo, W. and Ramsay, L., 1998. Oily fish and fish oil supplements in the prevention of coronary heart disease. *Journal of Human Nutrition and Dietetics*, 11 (1): 13-19.
- Mohan, M. V. and Sankaran, T. M., 1988. Two new indices for stomach content analysis of fishes. *Journal of Fish Biology*, 33 (2): 289-292.

- Mohsen, E. R., 1985. NIH Launching major research program on fish oils and health. Food Chemical News, 6: 34-39.
- Mohsin, A. K. M., Ambak, M. A. and Salam, M. N. A., 1993. Malay, English and scientific names of the fishes of Malaysia. Serdang: Faculty of Fisheries and Marine Science, Universiti Pertanian Malaysia.
- Mohsin, A. K. M. and Ambak, M. A., 1996. *Marine fishes and fisheries of Malaysia and neighbouring countries*. Vol. 495: Universiti Pertanian Malaysia Press Serdang.
- Molfese, C., Beare, D. and Hall-Spencer, J. M., 2014. Overfishing and the replacement of demersal finfish by shellfish: an example from the english channel. PloS One, 9 (7): e101506.
- Montero, D., Robaina, L., Caballero, M. J., Ginés, R. and Izquierdo, M. S., 2005. Growth, feed utilization and flesh quality of European sea bass (*Dicentrarchus labrax*) fed diets containing vegetable oils: A time-course study on the effect of a re-feeding period with a 100% fish oil diet. Aquaculture, 248 (1): 121-134.
- Morris, C. A., Haynes, K. C., Keeton, J. T. and Gatlin, D. M., 1995. Fish oil dietary effects on fatty acid composition and flavor of channel catfish. Journal of Food Science, 60 (6): 1225-1227.
- Morris Jr, J. A., Sullivan, C. V. and Govoni, J. J., 2011. Oogenesis and spawn formation in the invasive lionfish, *Pterois miles* and *Pterois volitans*. Scientia Marina, 75 (1): 147-154.
- Muchlisin, Z. A., Musman, M. and Azizah, M. N. S., 2010a. Spawning seasons of *Rasbora tawarensis* (Pisces: Cyprinidae) in Lake Laut Tawar, Aceh Province, Indonesia. Reproductive Biology and Endocrinology, 8 (1): 49.
- Muchlisin, Z. A., Musman, M. and Azizah, S., 2010b. Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. Journal of Applied Ichthyology, 26 (6): 949-953.
- Muhamad, N. A. and Mohamad, J., 2012. Fatty acids composition of selected Malaysian fishes. Sains Malaysiana, 41 (1): 81-94.
- Mukhopadhyay, M. K., 1994. Some threatened estuarine fishes of India. Paper presented at the national seminar on endangered fishes of India. Barrackpore: Central Inland Capture Fisheries Research Institute, pp. 229-235.
- Murawski, S. A., Rago, P. J. and Trippel, E. A., 2001. Impacts of demographic variation in spawning characteristics on reference points for fishery management. ICES Journal of Marine Science: Journal du Conseil, 58 (5): 1002-1014.

- Murua, H. and Saborido-Rey, F., 2003. Female reproductive strategies of marine fish species of the North Atlantic. *Journal of Northwest Atlantic Fishery Science*, 33: 23-31.
- Murua, H., Kraus, G., Saborido-Rey, F., Witthames, P. R., Thorsen, A. and Junquera, S., 2003. Procedures to estimate fecundity of marine fish species in relation to their reproductive strategy. *Journal of Northwest Atlantic fishery science*, 33: 33-54.
- Mushahida-Al-Noor, S., Samad, M. A. and Bhuiyan, N. I. M. A. S., 2013. Food and feeding habit of the critically endangered catfish *Rita rita* (Hamilton) from the Padda river in the north-western region of Bangladesh. *International Journal of Advancements in Research and Technology*, Volume 2 (1): 155-166.
- Nautiyal, P. and Lal, M. S., 1984. Preliminary observations on the migratory behaviour of the Garhwal Himalayan Mahseer. *Journal of the Bombay Natural History Society*, 81: 204-208.
- Needham, P. R., 1962. *A guide to the study of fresh-water biology*. Holden-Day. Inc. San Francisco, California.
- Nejed, S., Petrinc, Z., Kuzir, S. and Srebocan, E., 2004. Annual oscillation of ovarian morphology in European pilchard (*Sardina pilchardus* Walbaum) in the Northern Adriatic Sea. *Veterinarski arhiv*, 74 (2): 97-106.
- Newman, S. J., 2002. Growth, age estimation and preliminary estimates of longevity and mortality in the Moses perch, *Lutjanus russelli* (Indian Ocean form), from continental shelf waters off north-western Australia. *Asian Fisheries Science*, 15 (3): 283-294.
- Ng, H. H. and Sparks, J. S., 2002. *Plotosus fisadoha*, a new species of marine catfish (Teleostei: Siluriformes: Plotosidae) from Madagascar. *Proceedings of the Biological Society of Washington*, 115 (3): 564-569.
- Nguyen, H. P., Nguyen, N. T., Nguyen, P. D. and Do, T. N. N., 1994. Checklist of marine fishes in Vietnam. Hanoi: Science and Technics Publishing House, vol. 1: Amphioxii and Chondrichthyes, 64 p.
- Nielsen, S. S. (2010). *Food analysis* (fourth ed.). New York: Springer.
- Nogueira, N., Cordeiro, N. and Aveiro, M. J., 2013. Chemical composition, fatty acids profile and cholesterol content of commercialized marine fishes captured in Northeastern Atlantic. *Journal of Fisheries Sciences.Com*, 7 (3): 271-286.
- Nordøy, A., Marchioli, R., Arnesen, H. and Videbæk, J., 2001. n-3 Polyunsaturated fatty acids and cardiovascular diseases: To whom, how much, preparations. *Lipids*, 36 (1): S127-S129.



- Norimah Jr, A. K., Safiah, M., Jamal, K., Haslinda, S., Zuhaida, H., Rohida, S. and Kandiah, M., 2008. Food consumption patterns: findings from the Malaysian adult nutrition survey (MANS). *Malaysian Journal of Nutrition*, 14 (1): 25.
- Norse, E. A., Brooke, S., Cheung, W. W. L., Clark, M. R., Ekeland, I., Froese, R. and Morato, T., 2012. Sustainability of deep-sea fisheries. *Marine Policy*, 36 (2): 307-320.
- Nurnadia, A. A., Azrina, A. and Amin, I., 2011. Proximate composition and energetic value of selected marine fish and shellfish from the West coast of Peninsular Malaysia. *International Food Research Journal*, 18 (1): 137-148.
- Obeten Offem, B., Akegbejo-Samsons, Y. and Tunde Omoniyi, I., 2008. Diet, size and reproductive biology of the silver catfish, *Chrysichthys nigrodigitatus* (Siluformes: Bagridae) in the Cross River, Nigeria. *Revista de Biología Tropical*, 56 (4): 1785-1799.
- Offem, B. O., Akegbejo-Samsons, Y. and Omoniyi, I. T., 2007. Biological assessment of *Oreochromis niloticus* (Pisces: Cichlidae; Linne, 1958) in a tropical floodplain river. *African Journal of Biotechnology*, 6 (16): 1966-1971.
- Olele, N. F., 2010. Reproductive Biology of *Sarotherodon galilaeus* (Artedi, 1757) in Onah Lake, Delta State, Nigeria. *Journal of Applied Sciences Research*, 6 (12): 1981-1987.
- Oluwaniyi, O. O. and Dosumu, O. O., 2009. Preliminary studies on the effect of processing methods on the quality of three commonly consumed marine fishes in Nigeria. *Biokemistri*, 21 (1): 1-7.
- Orban, E., Nevigato, T., Masci, M., Di Lena, G., Casini, I., Caproni, R. and Rampacci, M., 2007. Nutritional quality and safety of European perch (*Perca fluviatilis*) from three lakes of Central Italy. *Food Chemistry*, 100 (2): 482-490.
- Oribhabor, B. J. and Ogbeibu, A. E., 2012. The food and feeding habits of fish species assemblage in a Niger Delta mangrove creek, Nigeria. *Journal of Fisheries and Aquatic Science*, 7 (2): 134-149.
- Osibona, A. O., 2011. Comparative study of proximate composition, amino and fatty acids of some economically important fish species in Lagos, Nigeria. *African Journal of Food Science*, 5 (10): 581-588.
- Osman, H., Suriah, A. R. and Law, E. C., 2001. Fatty acid composition and cholesterol content of selected marine fish in Malaysian waters. *Food Chemistry*, 73 (1): 55-60.
- Ota, K., Kohda, M. and Sato, T., 2010. Unusual allometry for sexual size dimorphism in a cichlid where males are extremely larger than females. *Journal of Biosciences*, 35 (2): 257-265.



- Özogul, Y., Özogul, F., Çiçek, E., Polat, A. and Kuley, E., 2008. Fat content and fatty acid compositions of 34 marine water fish species from the Mediterranean Sea. *International Journal of Food Sciences and Nutrition*, 60 (6): 464-475.
- Özyurt, G. and Polat, A., 2006. Amino acid and fatty acid composition of wild sea bass (*Dicentrarchus labrax*): a seasonal differentiation. *European Food Research and Technology*, 222 (3-4): 316-320.
- Pandey, K. and Shukla, J. P. A., 2005. *A textbook of Fish and Fisheries*. Rastogi Publications.
- Pathak, B. C., Zahid, M. and Serajuddin, M., 2013. Length-weight, length-length relationship of the spiny eel, *Macrogathus pancalus* (Hamilton 1822) sampled from Ganges and Brahmaputra river basins, India. *Iranian Journal of Fisheries Sciences*, 12 (1): 170-182.
- Pauly, D. and Caddy, J. F., 1985. A modification of Bhattacharya's methods for the analysis of mixtures of normal distributions. *FAO Fisheries Circulation* (781), 16pp.
- Pauly, D. and David, N., 1981. ELEFAN I, a basic program for the objective extraction of growth parameters from length-frequency data. *Meeresforschung*, 28 (4): 205-211.
- Pauly, D., 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal of Conservation and International Exploration of Maritime*, 39 (2): 175-192.
- Pauly, D. and Munro, J. L., 1984. Once more on the comparison of growth in fish and invertebrates. *Fishbyte*, 2 (1): 21.
- Pauly, D. and Soriano, M. L., 1986. Some practical extensions to Beverton and Holt's relative yield-per-recruit model. In: Maclean, J.L., Dizon, L. B., Hosillo, L. V. (Eds). *The first Asian Fisheries Forum*. Asian Fisheries Society, Manila, Philippines. PP. 491-496.
- Pauly, D., 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil*, 39 (2): 175-192.
- Pauly, S., Soriano-Bartz, M., Moreau, J. and Jarre-Teichmann, A., 1992. A new model accounting for seasonal cessation of growth in fishes. *Marine and Freshwater Research*, 43 (5): 1151-1156.
- Payne, S. A., Johnson, B. A. and Otto, R. S., 1999. Proximate composition of some north-eastern Pacific forage fish species. *Fisheries Oceanography*, 8 (3): 159-177.

- Pervaiz, K., Iqbal, Z., Mirza, M. R., Javed, M. N., Naeem, M. and Ishtiaq, A., 2012. Length-weight, length-length relationships and feeding habits of wild Indus Mahseer, *Tor macrolepis*, from Attock, Pakistan. *Journal of Applied Ichthyology*, 28 (4): 673-676.
- Prasad, G. and Ali, P. H. A., 2008. Morphology of the diet in the gut of threatened yellow catfish *Horabagrus brachysoma* (Gunther 1864) at two life stages. *Fish Physiology and Biochemistry*, 34 (4): 385-389.
- Praveena, S. M., Siraj, S. S., Suleiman, A. K. and Aris, A. Z., 2011. A brush up on water quality studies of Port Dickson, Malaysia. *Research Journal of Environmental Sciences*, 5 (12): 841-849.
- Prokofiev, A. M., 2008. A new species of eel catfishes of the genus *Plotosus* *nha* Trang Bay (South China Sea, Central Vietnam) (Siluriformes: Plotosidae). *Journal of Ichthyology*, 48 (8): 671-675.
- Qasim, S. Z., 1973. Some implications of the problem of age and growth in marine fishes from the Indian waters. *Indian Journal of Fisheries*, 20 (2): 351-371.
- Rahman, A. K. A., 1989. *Freshwater fishes of Bangladesh*: Zoological Society of Bangladesh Department of Zoology, University of Dhaka.
- Rahnan, S. A., Huah, T. S., Nassan, O. and Daud, N. M., 1995. Fatty acid composition of some Malaysian freshwater fish. *Food Chemistry*, 54 (1), 45-49.
- Rajan, P. T., Sreeraj, C. R. and Immanuel, T., 2011. *Fish fauna of coral reef, mangrove, freshwater offshore and seagrass beds of Andaman and Nicobar Islands*. Zoological survey of India, Andaman and Nicobar Regional center. Haddo, Port Blair.
- Rao, D. M. and Rao, K. S., 1991. Food and feeding behaviour of *Nemipterus japonicus* (Bloch) populations off Visakhapatnam, South India. *Journal of the Marine Biological Association of India*. Cochin, 33 (1): 335-345.
- Ravaglia, M. A. and Maggese, M. C., 2002. Oogenesis in the swamp eel *Synbranchus marmoratus* Bloch, 1795 (Teleostei; synbranchidae). Ovarian anatomy, stages of oocyte development and micropyle structure. *Biocell*, 26 (3): 325-337.
- Ravi, V., 2013. Food and feeding habits of the mudskipper, *Boleophthalmus boddarti* (Pallas, 1770) from Pichavaram Mangroves, Southeast Coast of India. *International Journal of Marine Science*, 3 (12): 98-104.
- Reale, A., Ziino, M., Ottolenghi, F., Pelusi, P., Romeo, V., Condurso, C. and Sanfilippo, M., 2006. Chemical composition and nutritional value of some marine species from the Egadi Islands. *Chemistry and Ecology*, 22(sup1): S173-S179.

- Riede, K., 2004. Global register of migratory species: from global to regional scales. Final report for Research and Development-project, Federal agency for nature conservation, Bonn Germany, pp. 329.
- Rueda, F. M., Lopez, J. A., Martinez, F. J., Zamora, S., Divanach, P. and Kentouri, M., 1997. Fatty acids in muscle of wild and farmed red porgy, *Pagrus pagrus*. Aquaculture Nutrition, 3 (3): 161-165.
- Saini, A., Dua, A. and Mohindra, V., 2008. Comparative morphometrics of two populations of giant river catfish (*Mystus seenghala*) from the Indus river system. Integrative Zoology, 3 (3): 219-226.
- Sallam, K. I., Ahmed, A. M., Elgazzar, M. M. and Eldaly, E. A., 2007. Chemical quality and sensory attributes of marinated Pacific saury (*Cololabis saira*) during vacuum-packaged storage at 4° C. Food Chemistry, 102 (4): 1061-1070.
- Sarker, P. K., Pal, H. K. and Rahman, M. M., 2002. Observation on the fecundity and gonado-somatic index of *Mystus gulio* in brackishwater of Bangladesh. Online Journal of Biological Sciences, 2 (4): 235-237.
- Satoh, S., Poe, W. E. and Wilson, R. P., 1989. Effect of dietary n-3 fatty acids on weight gain and liver polar lipid fatty acid composition of fingerling channel catfish. The Journal of Nutrition, 119 (1): 23.
- Seeto, J. and Baldwin, W. J., 2010. *A Checklist of the Fishes of Fiji and a Bibliography of Fijian Fish*. Division of Marine Studies Technical Report. The University of the South Pacific.
- Seng, Y. T., 1994. *Molluscs in Malaysia* (2nd ed.). Kuala Lumpur Malaysia: Department of Fisheries, Ministry of Agriculture Malaysia.
- Serajuddin, M. and Ali, R., 2011. Food and feeding habits of striped spiny eel, *Macrognathus pancalus* (Hamilton). Indian Journal of Fisheries, 52 (1): 81-86.
- Serajuddin, M. and Mustafa, S., 1994. Feeding specialization in adult spiny eel, *Mastacembelus armatus*. Asian Fisheries Science, 7: 63-65.
- Shaji, S. A. and Kannan, H. C., 2013. Chemical composition and amino acid profile of *Sardinella longiceps* collected from western coastal areas of Kerala, India. Journal of Biology and Earth Sciences, 3 (1): B129-B134.
- Shekhar, C., 2011. Changes in muscle biochemical composition of *Labeo rohita* (Ham.) in relation to season. Indian Journal of Fisheries, 51 (3): 319-323.
- Shinkafi, B. A. and Daneji, A. I., 2011. Morphology of the gonads of *Synodontis eupterus* (Boulenger) from River Rima, North-western Nigeria. International Journal of Zoological Research, 7: 382 -392.

- Shinkafi, B. A., Ipinjolu, J. K., Argungu, L. A. and Abubakar, U., 2002. Length-weight relationship and fecundity of *Synodontis clarias* (Linnaeus) in River Rima, Nigeria. *Journal of Agriculture and Environment*, 3 (1): 147-154.
- Simon, K. D., Bakar, Y., Mazlan, A. G., Zaidi, C. C., Samat, A., Arshad, A. and Brown-Peterson, N. J., 2012. Aspects of the reproductive biology of two archer fishes *Toxotes chatareus* (Hamilton 1822) and *Toxotes jaculatrix* (Pallas 1767). *Environmental Biology of Fishes*, 93 (4): 491-503.
- Sinclair, H. M., 1990. *History of essential fatty acids. Omega-6 Essential Fatty Acids: Pathophysiology and Roles in Clinical Medicine*. In D. F. Horrobin (Ed.), *Omega-6 Essential Fatty Acids. Pathophysiology and Roles in Clinical Medicine*. New York: Wiley-Liss, pp. 1-20.
- Sinha, M., 1981. Length-weight relationship and relative condition factor of the canine catfish-eel *Plotosus canius* Hamilton. *Journal of the Marine Biological Association of India*, 23 (1): 39-43.
- Sinha, M., 1984. Food preference studies of *Plotosus canius* Hamilton and its cultural suitability. *Proceedings: Animal Sciences*, 93 (5): 437-443.
- Sinha, M., 1986a. Age and growth of *Plotosus canius* Hamilton from Hooghly-Malah estuary, India. *Indian Journal of Animal Sciences*, 56 (1): 116-126.
- Sinha, M., 1986b. Functional morphology, anatomy and histology of the digestive organs of the catfish *Plotosus canius* Hamilton. *Proceedings: Animal Sciences*, 95 (1): 23-44.
- Sivan, G., 2009. Fish venom: pharmacological features and biological significance. *Fish and Fisheries*, 10 (2): 159-172.
- Solak, K., Solak, K. and Akyurt, U., 2001. Certain reproductive characteristics of the catfish (*Clarias gariepinus* Burchell, 1822) living in the River Asi, Turkey. *Turkish Journal of Zoology*, 25: 453-460.
- Solomon, F. N. and Ramnarine, I. W., 2007. Reproductive biology of white mullet, *Mugil curema* (Valenciennes) in the Southern Caribbean. *Fisheries Research*, 88 (1): 133-138.
- Sparre, P. and Venema, S. C., 1998. Introduction to tropical fish stock assessment-Part 1: Manual: FAO Fisheries Technical Paper 306/1, 376 pp.
- Sree, V. J., Parulekar, A. H., Wahidulla, S. and Kamat, S. Y., 1994. Seasonal changes in biochemical composition of *Holothuria leucospilota* (Echinodermata). *Indian Journal of Marine Sciences*, 23: 117-119.
- Sreeraj, N., Raghavan, R. and Prasad, G., 2006. The diet of *Horabagrus brachysoma* (Gunther), an endangered bagrid catfish from Lake Vembanad (South India). *Journal of Fish Biology*, 69 (2): 637-642.

- Srivastava, C. B. L., 1999. *A text book of fishery science and Indian fisheries*: Kitab Mahal Publications, Allahabad, India, pp. 307.
- Srivastava, S. M., Singh, S. P. and Pandey, A. K., 2012. Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *Journal of Experimental Zoology, India*, 15 (2): 395-402.
- Ssali, W. M., 1989. Chemical composition data for Nile perch (*Lates niloticus*) and its application to the utilization of the species. In: *Proceedings of FAO Expert Consultation Paper on Fish Technology in Africa Held in Abidjan Cote d'Ivoire, 25–28 April 1988*. FAO Rome, Italy. pp. 17-23.
- Stansby, M.E., 1962. Proximate composition of fish, in: Heen, E. *et al.* (1962). *Fish in Nutrition*. pp. 55-61.
- Standis, M. E., 1982. Properties of fish oils and their application to handling of fish and to nutritional and industrial use. In R. E. Martin, G. J. Flick, C. E. Hedbard, and D. R. Ward (Eds.), *Chemistry and biochemistry of marine food products*, pp. 75-92. Westport Connecticut: AVI Publishing Co.
- Suvatti, C., 1981. *Fishes of Thailand*: Ratchabandit Sathan Publishers, Thailand.
- Talwar, P. K., 1992. *Inland fishes of India and adjacent countries*. 2 (1992.) Vol. 2, CRC Press.
- Tan, C. G. S. and Ng, P. K. L., 1994. An annotated checklist of mangrove brachyuran crabs from Malaysia and Singapore. *Hydrobiologia*, 285 (1-3): 75-84.
- Tee, E. S., Kuladevan, R., Young, S. I., Khor, S. C. and Zakiah, H. O., 1996. *Laboratory procedures in nutrient analysis of foods*. Division of Human Nutrition, Institute of Medical Research, Kuala Lumpur, Malaysia.
- Templeman, W., 1987. Differences in sexual maturity and related characteristics between populations of thorny skate (*Raja radiata*) in the northwest Atlantic. *Journal of Northwest Atlantic Fisheries Science*, 7: 155-167.
- Teugels, G. G., 1984. The nomenclature of African *Clarias* species used in aquaculture. *Aquaculture*, 38 (4): 373-374.
- Thunberg, C.P., 1787. *Donation. Thunbergianae 1785 Continuat.* 2. *Museum Naturalium Academiae Upsaliensis*, 4: 43-58.
- Tulli, F. and Tibaldi, E., 1997. Changes in amino acids and essential fatty acids during early larval rearing of dentex. *Aquaculture International*, 5 (3): 229-236.
- Turan, C., Yalcin, S., Turan, F., Okur, E. and Akyurt, I., 2005. Morphometric comparisons of African catfish, *Clarias gariepinus*, populations in Turkey. *Folia Zoologica*, 54 (1-2): 165-172.



- Ugoala, C., 2008. The fatty and amino acids profiles of Chichlidae and Claridae finfish species. *Internet Journal of Food Safety*, 10: 18-25.
- Ujjania, N. C., Kohli, M. P. S. and Sharma, L. L., 2012. Length-weight relationship and condition factors of Indian major carps (*C. catla*, *L. rohita* and *C. mrigala*) in Mahi Bajaj Sagar, India. *Research Journal of Biology*, 2 (1): 30-36.
- Ujjania, N. C., Sharma, L. L. and Balai, V. K., 2013. Length-weight relationship and condition factor of Indian major carp (*Labeo rohita* Ham., 1822) from southern Rajasthan, India. *Applied Biological Research*, 15 (2): 1-5.
- Urban, D., 2012. Food habits of Pacific cod and walleye pollock in the northern Gulf of Alaska. *Marine Ecology Progress Series*, 469: 215-222.
- Utoh, T., Horie, N., Okamura, A., Yamada, Y., Tanaka, S., Mikawa, N. and Oka, H. P., 2003. Oogenesis in the common Japanese conger *Conger myriaster*. *Fisheries Science*, 69 (1): 181-188.
- Valverde, J. C., Martínez-Llorens, S., Vidal, A. T., Jover, M., Rodríguez, C., Estefanell, J. and García, B. G., 2013. Amino acids composition and protein quality evaluation of marine species and meals for feed formulations in cephalopods. *Aquaculture International*, 21 (2): 413-433.
- Van Der Molen, S. and Matallanas, J., 2004. Reproductive biology of female Antarctic spiny plunderfish *Harpagifer spinosus* (Notothenioidei: Harpagiferidae), from Îles Crozet. *Antarctic Science*, 16 (2): 99-106.
- Vieira, R. P., Bentes, L., Monteiro, P., Ribeiro, J., Erzini, K. and Goncalves, J. M. S., 2013. Weight-length relationships for five fish species from a temperate coastal lagoon of southern Europe. *Journal of Applied Ichthyology*, 29 (1): 292-293.
- Vlieg, P., 1984. Proximate analysis of commercial New Zealand fish species. 2. *New Zealand Journal of Science*, 27 (4): 427-433.
- Volcan, M. V., Gonçalves, Â. and Guadagnin, D. L., 2013. Length-weight relationship of three annual fishes (Rivulidae) from temporary freshwater wetlands of southern Brazil. *Journal of Applied Ichthyology*, 29: 1188-1190.
- Wang, F., Ma, X., Wang, W. and Liu, J., 2012. Comparison of proximate composition, amino acid and fatty acid profiles in wild, pond- and cage-cultured longsnout catfish (*Leiocassis longirostris*). *International Journal of Food Science and Technology*, 47 (8): 1772-1776.
- Wang, S. P., Sun, C. L. and Yeh, S. Z., 2003. Sex ratios and sexual maturity of swordfish (*Xiphias gladius* L.) in the waters of Taiwan. *Zoological Studies*, 42 (4): 529-539.



- Waters, 2007. *Pico-tag amino acid analysis system-operators manual*. USA: Waters chromatography division.
- Wilson, R. P. and Cowey, C. B., 1985. Amino acid composition of whole body tissue of rainbow trout and Atlantic salmon. *Aquaculture*, 48 (3): 373-376.
- Wilson, R. P. and Poe, W. E., 1985. Relationship of whole body and egg essential amino acid patterns to amino acid requirement patterns in channel catfish, *Ictalurus punctatus*. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, 80 (2): 385-388.
- Weber, M., 1910. Neue Fische aus Niederländisch Süd-Neu-Guinea. *Notes Leyden Museum*, 32: 225-240.
- Wright, J., 2009. Diversity, phylogenetic distribution, and origins of venomous catfishes. *BMC Evolutionary Biology*, 9 (1): 282.
- Yatuha, J., Kang'ombe, J. and Chapman, L., 2012. Diet and feeding habits of the small catfish, *Clarias liocephalus* in wetlands of Western Uganda. *African Journal of Ecology*, 51: 385-392.
- Yoshino, T. and Kishimoto, H., 2008. *Plotosus japonicus*, a new eeltail catfish (Siluriformes: Plotosidae) from Japan. *Bulletin of the Natural Museum of Nature and Science. Serial A Supplementary*, 2: 1-11.
- Yu-Abit, I. L., 2011. Length-weight relationship and condition factor of three dominant species from the lake Tasik Kenyir, Terengganu, Malaysia. *Journal of Fisheries and Aquatic Science*, 6 (7): 852-856.
- Zenebe, T., Ahlgren, G. and Boberg, M., 1998. Fatty acid content of some freshwater fish of commercial importance from tropical lakes in the Ethiopian Rift Valley. *Journal of Fish Biology*, 53 (5): 987-1005.
- Zhao, F., Zhuang, P., Zhang, L. and Shi, Z., 2010. Biochemical composition of juvenile cultured vs. wild silver pomfret, *Pampus argenteus*: determining the diet for cultured fish. *Fish Physiology and Biochemistry*, 36 (4): 1105-1111.
- Zorica, B., Sinovčić, G., Pallaoro, A. and Čikeš Keč, V., 2006. Reproductive biology and length-weight relationship of painted comber, *Serranus scriba* (Linnaeus, 1758), in the Trogir Bay area (middle-eastern Adriatic). *Journal of Applied Ichthyology*, 22 (4): 260-263.
- Zuraini, A., Somchit, M. N., Solihah, M. H., Goh, Y. M., Arifah, A. K., Zakaria, M. S. and Mat Jais, A. M., 2006. Fatty acid and amino acid composition of three local Malaysian *Channa spp.* fish. *Food Chemistry*, 97 (4): 674-678.