



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF NEW METHODS FOR SYNTHESIZING REACTIVE
NANOSTRUCTURED VANADIUM PHOSPHATE CATALYSTS FOR
PARTIAL OXIDATION OF *N*-BUTANE TO MALEIC ANHYDRIDE**

ALI ASGHAR ROWNAGHI

FS 2007 58



**DEVELOPMENT OF NEW METHODS FOR SYNTHESIZING REACTIVE
NANOSTRUCTURED VANADIUM PHOSPHATE CATALYSTS FOR PARTIAL
OXIDATION OF N-BUTANE TO MALEIC ANHYDRIDE**

ALI ASGHAR ROWNAGHI

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2007



Especially Dedicated To

My Mother, Who is the rarity of my life

*The great soul of my father,
Which my knowledge is indebted to him*

My beloved wife

My lovely sisters

and

My Dear brothers



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

DEVELOPMENT OF NEW METHODS FOR SYNTHESIZING REACTIVE NANOSTRUCTURED VANADIUM PHOSPHATE CATALYSTS FOR PARTIAL OXIDATION OF *N*-BUTANE TO MALEIC ANHYDRIDE

By

ALI ASGHAR ROWNAGHI

November 2007

Chairman: Professor Taufiq Yap Yun Hin, PhD

Faculty : Science

Vanadium phosphate catalysts are well known as the active phase and are commercially used for the selective oxidation of *n*-butane to maleic anhydride. They represent the sole example of a commercialized material for the catalytic oxidation of an alkane. In this study, vanadium phosphate catalysts were synthesised via three methods including two new methods *i.e.* new organic and solvothermal method and via dihydrate method. These catalysts were modified by microwave irradiation, addition of dopants and reducing agent.

The effect of reducing agents (*i.e.* isobutyl alcohol, 1-butanol and ethylene glycol), distilled water, microwave irradiation and conventional heating were investigated. Both heating methods gave similar XRD patterns, however microwave irradiated sample showed higher crystallinity which indicated by the high intensity of the peaks. The microwave treated catalysts exhibit a more homogeneous distribution of the rosette-



shape surface species and thin structure which consequently enhanced the specific surface area. However, microwave irradiation and the used of water treatment to enhance the crystalline phase, morphology and also increased the total amount of the oxygen species removed. The introduction of Nb, La and Ce as dopants to $(VO)_2P_2O_7$ catalysts prepared via reduction of $VOPO_4 \cdot 2H_2O$ gives significantly increased the surface area and subsequently improved the conversion efficiency of the catalyst.

A new organic route for preparation of high surface area ($>50 \text{ m}^2\text{g}^{-1}$) of vanadium phosphate catalyst was developed, by reducing the time from 26 to only 8 h. This together with incorporation of dopants such as Nb, La, and Ce can increase the surface area by reducing the particle size to nanoparticle. The high surface area contributed to higher mobility and better activity of the lattice oxygen and enhanced the conversion and selectivity for achieve a significant yield of maleic anhydride. The new organic route also increased the amount of V^{5+} phase in the catalyst. An appropriate amount of V^{5+} phase presence significantly enhanced the activity of VPO catalysts.

A novel solvothermal method was also developed by using a number of primary alcohols (C_3 - C_{10}) at various temperatures ($<423 \text{ K}$, 72 h) via a mild solvothermal synthesis to prepare phase-pure crystalline $(VO)_2P_2O_7$ with specific surface area up to three fold higher than that can be achieved by hydrothermal method. Furthermore, $(VO)_2P_2O_7$ solid solution can be crystallized in alcohols at temperature lower than that required by hydrothermal conversion. The intensity of (020) phase of nanocrystalline VPO catalyst prepared by the solvothermal were remarkably higher than the hydrothermal method. Another advantage of this new method for the synthesis of the



solid solution of vanadium phosphate catalyst is the low consumption of organic media in a close autoclave system. This eliminates the usage of water and yielded higher *n*-butane conversion catalyst as compared to conventional organic VPO method.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMBANGUNAN KAEDAH BARU DALAM SINTESIS MANAKIN
VANADIUM FOSFAT REAKTIF BERSTRUKTUR NANO UNTUK
PENGOKSIDAAN SEPARA *N*-BUTANA KE MALEIK ANHIDRIDA**

Oleh

ALI ASGHAR ROWNAGHI

November 2007

Pengerusi : Profesor Taufiq-Yap Yun Hin, PhD

Fakulti : Sains

Mangkin vanadium fosfat terkenal sebagai fasa aktif dan digunakan secara komersil dalam pengoksidaan terpilih *n*-butana ke maleik anhidrida. Manglain ini merupakan satu-satunya bahan yang telah dikomersilkan sebagai mangkin dalam pengoksidaan alkana. Dalam kajian ini, mangkin vanadium fosfat telah disintesis melalui tiga kaedah termasuk dua kaedah baru iaitu kaedah organik dan solvoterma dan melalui kaedah dihidrat. Mangkin ini juga telah diubahsuai melalui pelbagai pra-rawatan iaitu, (i) pemancaran gelombang mikro (ii) penambahan dopan dan (iii) agen penurunan.

Kesan agen penurunan (iaitu isobutil alkohol, 1-butanol dan etilena glikol), air suling, pemancaran gelombang mikro dan kaedah pemanasan konvensional telah dikaji. Kedua-dua bahan memberikan corak perghabluran yang sama, bagaimanapun sampel yang telah disinarkan dengan gelombang mikro menunjukkan struktur keamatan kehabluran yang tinggi yang ditunjukkan melalui keamatan puncak yang tinggi. Mangkin yang dihasilkan melalui kaedah gelombang mikro menunjukkan taburan permukaan bercorak ros yang



lebih homogen, berstruktur nipis dan seterusnya meningkatkan luas permukaan spesifik. Walaubagaimanapun, mangkin yang disinarkan oleh gelombang mikro dengan menggunakan air sebagai pelarut telah meningkatkan fasa kehabluran, morfologi dan juga meningkatkan jumlah keseluruhan spesies oksigen yang dibebaskan. Penambahan Nb, La dan Ce sebagai dopan ke atas mangkin $(VO)_2P_2O_7$ yang dihasilkan melalui penurunan $VOPO_4 \cdot 2H_2O$ telah meningkatkan luas permukaan yang ketara dan seterusnya meningkatkan penukaran mangkin.

Mangkin vanadium fosfat yang mempunyai luas permukaan yang tinggi ($> 50 \text{ m}^2\text{g}^{-1}$) telah dibangunkan dengan menggunakan medium organik memerlukan masa penyediaan yang singkat (8 jam) berbanding kaedah organik konvensional (26 jam). Pengurangan masa tindakbalas kepada 8 jam dan penggunaan dopan seperti Nb, La dan Ce dapat meningkatkan luas permukaan melalui pengurangan saiz partikel mangkin kepada berskala nano. Luas permukaan yang tinggi menyumbang ke arah mobiliti dan aktiviti oksigen kekisi yang meningkatkan penukaran dan keterpilihan untuk pengoksidaan *n*-butana. Kaedah organik baru ini juga meningkatkan bilangan fasa V^{5+} mangkin. Dengan nisbah V^{5+}/V^{4+} yang bersesuaian sebenarnya dapat meningkatkan aktiviti mangkin ini dengan ketara.

Kaedah baru solvoterma juga telah dibangunkan dengan menggunakan beberapa jenis alkohol primer (C_3 - C_{10}) dalam pelbagai suhu melalui sintesis solvoterma lembut ($< 423 \text{ K}$, 72 jam) telah digunakan untuk menyediakan fasa hablur $(VO)_2P_2O_7$ yang tulen dengan peningkatan luas permukaan spesifik dua kali ganda berbanding penukaran yang diperlukan dalam hidroterma. Keamatan fasa (020) mangkin hablur nano VPO yang disediakan melalui solvoterma menunjukkan peningkatan berbanding kaedah

hidroterma. Kebaikan kaedah baru ini dapat digunakan untuk mensintesis larutan pepejal mangkin vanadium fosfat disamping penggunaan media organik yang kurang di dalam sistem tertutup "autoclave". Kaedah ini juga dapat mengurangkan sisa air dan meningkatkan penukaran berbanding dengan mangkin yang disediakan melalui kaedah organik VPO konvensional.



ACKNOWLEDGEMENTS

I would like to take this opportunity to express my utmost gratitude to the chairman of my supervisory committee Prof. Dr. Taufiq-Yap Yun Hin, for his invaluable guidance throughout the course of this study. Without his kind advice and support, it would not be possible for me to complete my research. His guidance and advice throughout the project have been very indispensable for completion of this thesis. Sincere thanks are due to Prof. Dr. Mohd Zobir Hussein and Assoc. Prof. Dr. Irmawati Ramli for their kind support and guidance.

Heartfelt thanks are extended to all the laboratory assistants in Department of Chemistry and Institute of Bioscience for their favorable help and advice.

I am also very grateful to Tang Wen Jiun and Sharmeele, Rina, Murni and other lab mates, whose help, suggestions, encouragement and companion are of great help in sustaining the morale and enthusiasm.

Last but not least, I would like to thank all those who had contributed to the success of this work in one way or another especially my wife Fateme Rezaei for her kind support and encouragements. Finally, my sincere appreciation also goes to my mother, sisters, brothers and friends for their deep support, understanding and sacrifices.



I certify that an Examination Committee has met on 12th November 2007 to conduct the final examination of Ali Asghar Rownaghi on his Doctor of Philosophy thesis entitled “Development of New Methods for Synthesizing Reactive Nanostructured Vanadium Phosphate Catalysts for Partial Oxidation of *N*-Butane to Maleic Anhydride” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Doctor of Philosophy.

Members of the Examination Committee were as follows:

Anuar Kassim, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Zulkarnain Zinal, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Mohd Zaizi Desa, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Mohd Ambar Yarmo, PhD

Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)

HASANAH MOHD. GHAZALI, PhD

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Taufiq-Yap Yun Hin, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Mohd Zobir Hussein, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Irmawati Ramli, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 22 January 2008



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

ALI ASGHAR ROWNAGHI

Date: 27 November 2007



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvii
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS	xxvii
CHAPTER	
1 INTRODUCTION	1
1.1 Background of Study	1
1.1.1 The Oxidation of <i>n</i> -Butane Catalyzed by VPO	4
1.1.2 The Modification of Precursor's Properties by the Addition of Dopants	7
1.1.3 Industrial Processes of Maleic Anhydride Synthesis from <i>n</i> -Butane	8
1.1.4 Vanadium Phosphate Oxide Catalysts Synthesis and Characteristics	9
1.2 Significance of Study	10
1.2.1 Catalysis and Sustainable (Green) Chemistry	11
1.2.2 Solvothermal Synthesis	12
1.3 Objectives of Study	13
1.4 Structure of Thesis	14
2 LITERATURE REVIEW	17
2.1 Basic Concepts of Heterogeneous Catalysis	17
2.2 Vanadium in Catalysis	17
2.2.1 Vanadium Oxides	18
2.2.2 Vanadium Phosphorus Oxide Catalysts	19
2.3 Maleic Anhydride: Production and Consumption	20
2.3.1 Maleic Anhydride Consumption	21
2.3.2 Maleic Anhydride Industrial Production	24
2.4 Catalytic System	28
2.4.1 Structural Aspect of Vanadium Phosphate Catalyst	28
2.4.2 The Role of V ⁵⁺ and V ⁴⁺ Phases on the VPO Catalyst under <i>n</i> -Butane Oxidation	30
2.4.3 Different Routes in Preparation of Catalyst Precursors	36
2.4.4 Topotactic Transformation of the Precursor	40



2.4.5	Activation Procedure	42
2.4.6	Role of the P/V Ratio	47
2.4.7	Proposed Active Sites and Mechanisms of <i>n</i> -Butane Oxidation	49
2.4.8	The Role of Lattice Oxygen	53
2.4.9	Role of Dopant/Promoter	58
2.5	Conversion, Yield and Selectivity	60
2.6	Microwave-Enhanced Chemistry	63
2.6.1	Microwaves Are Energy	63
2.6.2	Microwaves Can Interact with Matter	65
2.6.3	Two Principal Mechanisms for Interaction with Matter	66
2.6.4	Applications of Microwaves in Catalytic Chemistry	67
2.6.5	Advantages of Microwave Assisted Synthesis	68
2.7	Solvothermal Synthesis Processes	69
2.7.1	Advantages of Solvothermal Synthesis	74
3	CATALYSTS PREPARATION AND CHARACTERISATIONS	76
3.1	Preparations of Bulk Vanadium Phosphate Catalysts via Different Routes	76
3.1.1	Reduction of VOPO ₄ ·2H ₂ O Phase	76
3.1.2	The Conventional Organic (VPO) method	79
3.1.3	New Organic Route (NVPO)	79
3.2	Preparation of Doped Vanadyl Pyrophosphate Catalysts	81
3.2.1	Reduction of VOPO ₄ ·2H ₂ O Phase	81
3.2.2	Organic Route (New VPO)	82
3.3	Solvothermal Route	83
3.4	Catalysts Characterization	85
3.4.1	X-Ray Diffraction	86
3.4.2	BET Surface Area Measurements	88
3.4.3	Redox Titration Method	89
3.4.4	Energy Dispersive X-ray (EDX)	91
3.4.5	Inductively Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES)	91
3.4.6	Scanning Electron Microscopy (SEM)	92
3.4.7	Temperature Programmed Analyze	93
3.5	Catalytic Evaluation	95
4	THE EFFECT OF MICROWAVE IRRADIATION AND SOLVENTS ON PHYSICOCHEMICAL PROPERTY OF VANADIUM PHOSPHATE CATALYST	96
4.1	Introduction	96
4.2	Effect of Microwave Irradiation, Isobutyl alcohol and 1-Butanol on the Physicochemical Properties of Vanadium Phosphate Catalysts	99
4.2.1	Phase Identification Using Powder XRD Technique	100



4.2.2	BET Surface Area Measurements and Chemical Analysis	107
4.2.3	Scanning Electron Microscopy	109
4.2.4	Temperature Programmed Reduction (TPR in H ₂)	112
4.3	Effect of Microwave Irradiation, Ethylene glycol and Water Treatment on the Physicochemical Properties of the “Glycolsynthesis Method” Vanadium Phosphate Catalysts	117
4.3.1	Phase Identification Using Power XRD Technique	118
4.3.2	BET Surface Area Measurements and Chemical Analysis	124
4.3.3	Scanning Electron Microscopy	127
4.3.4	Temperature Programmed Reduction (TPR in H ₂)	129
4.4	Conclusions	133
5	EFFECT OF MICROWAVE ON THE PHYSICO-CHEMICAL PROPERTIES OF DOPED-VANADIUM PHOSPHOROUS OXIDE CATALYSTS	138
5.1	Introduction	138
5.2	Phase Identification Using Powder XRD Technique	139
5.2.1	The Doped Precursors	139
5.2.2	The Doped Catalysts	141
5.3	BET Surface Area Measurements and Chemical Analysis	145
5.4	Scanning Electron Microscopy	152
5.5	Temperature Programmed Reduction (TPR in H ₂)	157
5.6	Catalytic Evaluation	163
5.7	Conclusions	165
6	SYNTHESIS OF VANADIUM PHOSPHATE BY NEW ORGANIC ROUTE AND THEIR PHYSICO-CHEMICALS PROPERTIES OF DOPED-CATALYSTS	168
6.1	Introduction	168
6.2	Phase Identification Using Powder XRD Technique	172
6.2.1	The Undoped and Doped New Organic Precursors	172
6.2.2	The Undoped and Doped New Organic Catalysts	174
6.3	BET Surface Area Measurements and Chemical Analysis	179
6.4	Scanning Electron Microscopy	188
6.5	Temperature Programmed Reduction (TPR in H ₂)	194
6.6	Catalytic Evaluation	199
6.7	Conclusions	204
7	SOLVOTHERMAL SYNTHESIS: A NEW METHOD FOR PREPARATION DIRECTLY SOLID SOLUTION OF VANADIUM PYROPHOSPHATE CATALYSTS FOR <i>n</i>-BUTANE OXIDATION	206
7.1	Introduction	206
7.2	Phase Identification Using Powder XRD Technique	208



7.3	BET Surface Area Measurements and Chemical Analysis	224
7.4	Scanning Electron Microscopy	231
7.5	Temperature Programmed Reduction (TPR in H ₂)	239
7.6	Catalytic Evaluation	247
7.7	Conclusions	250
8	SUMMARY AND CONCLUDING REMARKS	253
8.1	Introduction	253
8.2	Summary	254
	REFERENCES	257
	APPENDICES	273
	BIODATA OF THE AUTHOR	280



LIST OF TABLES

Table	Page
2.1 World MA Production Capacity and Effective Production	21
2.2 Application of Maleic Anhydride	22
2.3 Industrial Technologies for Maleic Anhydride Production from <i>n</i> -Butane	28
2.4 Active Sites in Vanadyl Pyrophosphate	51
3.1 Preparation of VOHPO ₄ ·0.5H ₂ O Phase with Reduction of Dihydrate	78
3.2: Preparation of VOHPO ₄ ·0.5H ₂ O Phases using Ethylene Glycol	79
3.3: Preparation of VOHPO ₄ ·0.5H ₂ O Phases using New Organic Route	81
3.4: Preparation Condition of the Doped VOHPO ₄ ·0.5H ₂ O via VPD Route	82
3.5: Preparation of the Doped VOHPO ₄ ·0.5H ₂ O via New Organic Route	83
3.6: Preparation of the VOHPO ₄ ·0.5H ₂ O Phases via Solvothermal Method	85
3.7: Typical Information in a X-ray Diffraction Pattern	86
3.8: Typical Application of Powder X-ray Diffraction Pattern	87
4.1: XRD Data of VPDA1, VPDA2, VPDB1 and VPDB2 Precursors	103
4.2: XRD Data of VODA2, VODA2, VODB1 and VODB2 Catalysts	106
4.3: Total Surface Area and Chemical Analysis for VODA1, VODA2, VODB1 and VODB2 Catalysts	109
4.4: Redaction Activation Energy, Total Amount and Ratio for Oxygen Removed from the Microwave and Conventional Catalysts by Reduction in H ₂ /Ar	116
4.5: XRD Data of VPDC1, VPDD1, VPDC2 and VPDD2 Precursors	119
4.6: XRD Data of VODC1, VODD1, VODC2 and VODD2 Catalysts	124
4.7: Total Surface Area and Chemical Analysis for VODC1, VPDD1, VPDC2 and VODD2 Catalysts	125



4.8:	Redaction Activation Energy, Total Amount and Ratio for Oxygen Removed from the Microwave and Conventional Catalysts by Reduction in H ₂ /Ar	132
5.1	XRD Data of Undoped and Doped Catalysts	144
5.2	Total Surface Area and Chemical Analysis for Undoped and Doped Catalysts	147
5.3	Total Amount of Oxygen Removed, the Values of Reduction Activation Energies and Ratio for Oxygen Removed of V ⁵⁺ /V ⁴⁺ obtained by Temperature Programmed Reduction in H ₂ /Ar	159
5.4	Catalytic Performance of MW and CH for the Oxidation of <i>n</i> -Butane to Maleic Anhydride	164
6.1	XRD Data of Conventional and New Organic Route Catalysts	177
6.2	Table Surface Area and Chemical Analysis for Conventional and New Method Catalysts	180
6.3	Total Amount and Ratio for Oxygen Removed, the Values of Reduction Activation Energies by temperature programmed reduction in H ₂ /Ar	197
6.4	Catalytic Performance of Conventional and New VPO Method for the Oxidation of <i>n</i> -Butane to Maleic Anhydride	200
7.1	XRD Data of Solvothermal and Conventional Precursors	211
7.2	XRD Data of Solvothermal and Conventional Catalysts	218
7.3	Total Surface Area and Chemical Analysis for Solvothermal and Conventional Catalysts	224
7.4	Total Amount of Oxygen and Ratio for Oxygen Removed, the Values of Reduction Activation Energies by Temperature Programmed Reduction in H ₂ /Ar (H ₂ -TPR).	244
7.5	Total Amount of Oxygen and Ratio for Oxygen Removed, the Values of Reduction Activation Energies by Temperature Programmed Reduction in H ₂ /Ar (H ₂ -TPR).	245
7.6	The Catalytic Performances of Organic and Solvothermal method Catalysts in <i>n</i> -Butane Oxidation to Maleic Anhydride	248



LIST OF FIGURES

Figure		Page
1.1	A stainless steel solvothermal autoclave and the component inside the solvothermal autoclave.	13
2.1	Overview of the importance of vanadium in supported metal oxide catalysts	19
2.2	Maleic Anhydride Structure	20
2.3	World Sales of Maleic Anhydride	22
2.4	From Maleic Anhydride and Butadiene to Tetrahydrophalic Anhydride	23
2.5	Structure of Maleic Acid, Fumaric Acid and γ -Butyric Lactone	24
2.6	Selective Oxidation of Benzene to Maleic Anhydride	25
2.7	Process for producing maleic anhydride from benzene	26
2.8	Selective Oxidation of <i>n</i> -Butane to Maleic Anhydride	26
2.9	Flammability Diagram for <i>n</i> -Butane/Oxygen Mixtures	27
2.10	The Layered Structure of $(VO)_2P_2O_7$	29
2.11	Idealized Vanadyl Pyrophosphate Structure in (020) plane	29
2.12	Catalytic oxidation of <i>n</i> -butane on the surface of plate-like $(VO)_2P_2O_7$ crystal. Selective oxidation to maleic anhydride takes place on the (020) basal plane and deep oxidation on the side planes	30
2.13	Selective and unselective oxidation site on crystal faces of $(VO)_2P_2O_7$	30
2.14	<i>n</i> -Butane Conversion as a Function of Amount of Oxygen Removed Associated with V^{5+}	34
2.15	Maleic Anhydride Selectivity as a Function of Amount of Oxygen Removed Associated with V^{5+}	34
2.16	<i>n</i> -Butane Conversion as a Function of Amount of Oxygen Removed Associated with V^{4+}	35
2.17	<i>n</i> -Butane Conversion as a Function of Preparation of Vanadium Phosphate Catalyst	36



2.18	Diagram of Phase Transformations among Vanadium Phosphate Catalysts	41
2.19	Structural Changes and Interconversion of VPO Phases during the activation of $\text{VOHPO}_4 \cdot 0.5\text{H}_2\text{O}$	44
2.20	Scheme of Proposed Evolution of the Vanadium Phosphate Catalyst	45
2.21	Reaction Scheme for the Selective Oxidation of <i>n</i> -Butane	49
2.22	Mechanism of <i>n</i> -Butane Activation on $(\text{VO})_2\text{P}_2\text{O}_7$	52
2.23	Active Sites Present on Vanadyl Pyrophosphate	53
2.24	Schematic of Catalytic Effectiveness as a Function of Metal Oxygen Bond	55
2.25	Oxygen Incorporation in the Oxide Structure from the Gas Phase	57
2.26	Schematic of an Oxygen Sublattice of a Reducible Metal Oxide	57
2.27	Typical Plot for Selectivity to MA as a Function of Conversion	61
2.28	Relationship between Specific Butane Conv. with Catalyst Surface Area	62
2.29	Electromagnetic spectrum	64
2.30	Interaction of microwave with materials	66
2.31	Pressure as a Function of Temperature and Percentage fill of Solvent in a Sealed Vessel	71
2.32	Schematic of a Teflon-lined, Stainless-Steel Autoclave	72
4.1	XRD Diffraction Patterns of VPH1 and VPH2 Vanadyl Phosphate Hydrate	101
4.2	XRD Diffraction Patterns of VPDA1, VPDA2, VPDB1 and VPDB2 Precursors	102
4.3	XRD Diffraction Patterns of VODA1, VODA2, VODB1 and VODB2 Catalysts	104
4.4	SEM Micrograph for VODA1 Catalyst	111
4.5	SEM Micrograph for VODA2 Catalyst	111
4.6	SEM Micrograph for VODB1 Catalyst	111



4.7	SEM Micrograph for VODB2 Catalyst	112
4.8	TPR Profiles for VODA1, VODA2, VODB1 and VODB2 Catalysts	114
4.9	XRD Diffractograms of VPDC1, VPDD1, VPDC2 and VPDD2 Precursors	118
4.10	XRD Diffractograms of VODC1, VODC2, VODD1 and VODD2 catalysts	121
4.11	Relation between the Specific Surface Area and Crystallite Size (020) Phase of $(VO)_2P_2O_7$ Catalysts	126
4.12	SEM Micrograph for VODC1 Catalyst	128
4.13	SEM Micrograph for VODD1 Catalyst	128
4.14	SEM Micrograph for VODC2 Catalyst	129
4.15	SEM Micrograph for VODD2 Catalyst	129
4.16	TPR Profiles for VODC1, VPDD2, VPDC2 and VODD2 Catalysts	131
5.1	XRD Diffraction Patterns for Undoped and Doped VPD Precursors	140
5.2	XRD Diffraction Patterns for Undoped and Doped VOD Catalysts	142
5.3	EDX Microanalysis Spectrum for VODE1	148
5.4	EDX Microanalysis Spectrum for VODE2	149
5.5	EDX Microanalysis Spectrum for VODF1	149
5.6	EDX Microanalysis Spectrum for VODF2	149
5.7	EDX Microanalysis Spectrum for VODG1	150
5.8	EDX Microanalysis Spectrum for VODG2	150
5.9	EDX Microanalysis Spectrum for VODA1	150
5.10	EDX Microanalysis Spectrum for VODA2	151
5.11	SEM Micrograph for Microwave Assisted Nb-doped VOD Catalyst	153
5.12	SEM Micrograph for Conventional Nb-doped VOD Catalyst	153
5.13	SEM Micrograph for Microwave Assisted La-doped VOD Catalyst	154



5.14	SEM Micrograph for Conventional La-doped VOD Catalyst	154
5.15	SEM Micrograph for Microwave Assisted Ce-doped VOD Catalyst	156
5.16	SEM Micrograph for Conventional Ce-doped VOD Catalyst	156
5.17	SEM Micrograph for Microwave Assisted undoped VOD Catalyst	156
5.18	SEM Micrograph for Conventional undoped VOD Catalyst	157
5.19	TPR Profiles for Microwave and Conventional Nb-doped Catalysts	158
5.20	TPR Profiles for Microwave and Conventional La-Doped Catalysts	160
5.21	TPR Profiles for Microwave and Conventional Ce-Doped Catalysts	161
5.22	TPR Profiles for microwave and conventional undoped Catalysts	162
5.23	Changes of <i>n</i> -Butane Conversion and Maleic Anhydride Selectivity with the Microwave Assisted Catalyst Synthesis	164
6.1	XRD Diffraction Patterns for Conventional, Undoped and Doped New Organic VPP Precursors	173
6.2	XRD Diffraction Patterns for Conventional, Undoped and Doped New Organic VPO Catalysts	175
6.3	Relation between Surface Area and I_{020}/I_{204} of the Vanadium Phosphate Catalyst Prepared by New VPO Method	181
6.4	Relation between the Surface Area and Crystallite Size (020) Phase of Vanadium Phosphate Catalyst Prepared by New VPO Method	182
6.5	EDX Microanalysis Spectrum for VPOA1 Catalyst	185
6.6	EDX Microanalysis Spectrum for VPOA2 Catalyst	185
6.7	EDX Microanalysis Spectrum for VPOB1 Catalyst	185
6.8	EDX Microanalysis Spectrum for VPOB2 Catalyst	186
6.9	EDX Microanalysis Spectrum for VPOC1 Catalyst	186
6.10	EDX Microanalysis Spectrum for VPOC2 Catalyst	186
6.11	EDX Microanalysis Spectrum for VPOD1 Catalyst	187



6.12	EDX Microanalysis Spectrum for VPOD2 Catalyst	187
6.13	EDX Microanalysis Spectrum for VPO Catalyst	187
6.14	SEM Micrograph for VPOA1 Catalysts	191
6.15	SEM Micrograph for VPOA2 Catalysts	191
6.16	SEM Micrograph for VPOB1 Catalysts	191
6.17	SEM Micrograph for VPOB2 Catalysts	192
6.18	SEM Micrograph for VPOC1 Catalysts	192
6.19	SEM Micrograph for VPOC2 Catalysts	192
6.20	SEM Micrograph for VPOD1 Catalysts	193
6.21	SEM Micrograph for VPOD2 Catalysts	193
6.22	SEM Micrograph for VPO2 Catalysts	193
6.23	TPR Profiles for VPO Catalysts Prepared by Conventional and New Method	195
6.24	TPR Profiles for Nb Doped VPO Catalysts Prepared by two New Condition Methods	196
6.25	TPR Profiles for La Doped VPO Catalysts Prepared by two New Condition Methods	198
6.26	TPR Profiles for Ce Doped VPO Catalysts Prepared by two New Condition Methods	199
6.27	Changes of <i>n</i> -Butane Conversion and Maleic Anhydride Selectivity with the Microwave Assisted Catalyst Synthesis	201
6.28	Influence of the amount of oxygen atoms removed from the reduction Peaks (TPR) associated with V ⁴⁺ on the <i>n</i> -butane conversion	202
7.1	XRD Diffraction Patterns Recorded from VSPA1, VSPA2, VSPA3 and VPPG Precursors	210
7.2	XRD Diffraction Patterns Recorded from VSPB1, VSPB2, VSPB3 and VPPG Precursors	212
7.3	XRD Diffraction Patterns Recorded from VSPC1, VSPC2, VSPC3 and	213



	VPPG Precursors	
7.4	XRD Diffraction Patterns Recorded from VSPD1, VSPD2, VSPD3 and VPPG Precursors	213
7.5	XRD Diffraction Patterns Recorded from VSPE1, VSPE2, VSPE3 and VPPG Precursors	214
7.6	XRD Diffraction Patterns Recorded from VSPF1, VSPF2, VSPF3 and VPPG Precursors	214
7.7	XRD Diffraction Patterns for VSOA1, VSOA2, VSOA3 and Conventional VPO Catalysts	217
7.8	XRD Diffraction Patterns for VSOB1, VSOB2, VSOB3 and Conventional VPO Catalysts	219
7.9	XRD Diffraction Patterns for VSOC1, VSOC2, VSOC3 and Conventional VPO Catalysts	219
7.10	XRD Diffraction Patterns for VSOD1, VSOD2, VSOD3 and Conventional VPO Catalysts	220
7.11	XRD Diffraction Patterns for VSOE1, VSOE2, VSOE3 and Conventional VPO Catalysts	221
7.12	XRD Diffraction Patterns for VSOF1, VSOF2, VSOF3 and Conventional VPO Catalysts	222
7.13	Relation between Reaction Temperatures and Solvothermal Treatment Crystalline Size	223
7.14	Relation between Surface Area and the Intensity of (020) Phase for the Solvothermal VPO Catalysts	226
7.15	Relation between Surface Area and Reaction Temperature for the Solvothermal VPO Catalysts	227
7.16	EDX Microanalysis Spectrum for VSOA1, VSOA2 and VSOA3 Catalysts	229
7.17	EDX Microanalysis Spectrum for VSOB1, VSOB2 and VSOB3 Catalysts	229
7.18	EDX Microanalysis Spectrum for VSOC1, VSOC2 and VSOC3 Catalysts	229

