UNIVERSITI PUTRA MALAYSIA

GROWTH PERFORMANCE OF ASIAN CATFISH (Clarias batrachus linnaeus) AND WATER QUALITY IN A RECIRCULATING AQUACULTURE SYSTEM

RAHMAT FADHIL

FK 2012 129

GROWTH PERFORMANCE OF ASIAN CATFISH (*Clarias batrachus linnaeus*) AND WATER QUALITY IN A RECIRCULATING AQUACULTURE SYSTEM

RAHMAT FADHIL

MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA

2012

© COPYRIGHT UPM

GROWTH PERFORMANCE OF ASIAN CATFISH *(Clarias batrachus linnaeus)* AND WATER QUALITY IN A RECIRCULATING AQUACULTURE SYSTEM

By

RAHMAT FADHIL

© COPYRIGHT UPM

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

November 2012

DEDICATED TO

My Father,

Drs. H. Syahbuddin Aziz

My Mother,

Hajjah Mardhiah Ibrahim

My Wife,

Drh. Rizki Amelia

My Daugthers,

Rumaisha Khairatun Hisan

Ruhama Althafunnisa

Rufaida Nailal Husna

My Brothers,

Nazaruddin, M.Si

Fitri Yanti (Almarhumah)

Iip Gunawan, ST

And My Nephew and Nieces

Habiburrahman

Izatul Jannah

Nurul 'Aliya

Fathiya Yasmin Putri Gunza

© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
to fulfilment the requirement for the degree of Master of Science

GROWTH PERFORMANCE OF ASIAN CATFISH (*Clarias batrachus linnaeus)* AND WATER QUALITY IN A RECIRCULATING AQUACULTURE SYSTEM

By

RAHMAT FADHIL

November 2012

Chairman : Associate Professor Ir. Johari Endan, Phd

Faculty : Engineering

Strategies to increase fish production are always being done on a continuous basis from time to time to develop a new better system. Water quality management is required in aquaculture systems, as it can affect the production and the organisms that live in the water. The use of Recirculating Aquaculture System/RAS in the fishing industry is the latest challenge to the fishing communities and researchers. The use of multi-media filtrations to maintain healthy water quality for aquaculture remains an important issue to be studied. The main problem is how to maintain the water quality parameters for fish life until the harvest. In this study, the concept of the RAS has been developed to determine the ability of controlling the quality of water for sustainable aquaculture.

© COPYRIGHT UPM

This study aimed to develop catfish breeding in an integrated utility with optimal use of water and soil. The objectives of this study were: (1) to determine the growth rate of catfish with mechanical circulatory system and water quality control through a biological filtration unit, (2) to measure the water quality parameters of fish production tanks, and (3) to study the process and the way water can be mixed up, producing more stable water quality conditions in the unit culture and reducing contaminations.

Asian catfish (*C. batrachus*) was cultured in three separate tanks, each with a size range of 5-8 cm (Tank 1), 8-12 cm (Tank 2), and 12-15 cm (Tank 3) for 30 days. Each tank contained 300 fishes. Water samples were analyzed twice (morning and afternoon) with duplicates during the experiment. Water quality parameters such as temperature, turbidity, dissolved oxygen, pH, and ammonia were measured. The entire tank was connected to water supply through a filter to each tank using PVC pipes. Resulting residues settled in a sedimentation tank. Water was pumped through the filter into a subsequent sedimentation tank and back into the tank through recirculation. Fish was fed at 3% of the total fish weight twice a day (morning and evening).

The RAS and biological filter systems with multi-media and backwash capability have resulted in improved water quality for the intensive culture of catfish. Fish survival rate was above 95% with weight gain of catfish of 11.00$\pm$1.90 gram/fish in Tank 1, 20.45$\pm$0.85 gram/fish in Tank 2, and 25.65$\pm$0.95 gram/fish in Tank 3. Rate of

© COPYRIGHT UPM

increase in length and weight of catfish in Tank 1 was from 6.5±1.5 cm (1.00±0.10 gram) to 12.0±2.0 cm (12.00±1.80 gram), Tank 2 from 10.0±2.0 cm (2.10±0.60 gram) to 16.0±2.0 cm (22.55±1.15 gram), and Tank 3 from 13.5±1.5 cm (5.65±1.25 gram) to 18.5±2.5 cm (32.33±0.30 grams).

Data analysis using Completely Randomized Design/CRD with three treatments and two replicates showed that the total percentage of survival of catfish in Tank 2 was higher than the other tanks. Percentages of survival from the highest to the lowest were 98.33%±0.33 (Tank 2), 95.33%±0.67 (Tank 1), and 95.17%±0.84 (Tank 3). ANOVA test showed no significant difference of water quality and growth performance between one tank and the rest in this trial. The results showed that RAS system can be successfully used and further developed to increase fish production, intensively. There was hardly a huge negative impact on water quality, growth, and survival rate of *catfish* in the tank for the period of observation. This indicates that differences in size of the fish in each tank did not have any great impact on the growth, survival, and quality of water in the catfish culture.

© COPYRIGHT UPM

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

PRESTASI PERTUMBUHAN IKAN KELI *(Clarias batrachus linnaeus)* DAN KUALITI AIR DALAM SEBUAH SISTEM AKUAKULTUR EDARAN SEMULA

Oleh

RAHMAT FADHIL

November 2012

Pengurusi : Profesor Madya Ir. Johari Endan, Phd

Faculti : Kejuruteraan

Strategi untuk meningkatkan pengeluaran ikan sentiasa dilakukan secara berterusan dari semasa ke semasa untuk mendapatkan sistem baru dan yang lebih baik. Pengurusan kualiti air diperlukan dalam sistem akuakultur, kerana ia boleh menjejaskan pengeluaran dan organisma yang hidup di dalam air. Penggunaan Sistem Akuakultur Edaran Semula (*Recirculating Aquaculture System*/RAS) dalam industri perikanan adalah cabaran terbaru untuk komuniti perikanan dan penyelidik. Penggunaan penapisan dengan pelbagai media bagi mengekalkan kualiti air yang sihat untuk akuakultur masih merupakan satu isu yang penting untuk dipelajari. Masalah utama ialah bagaimana mengekalkan parameter kualiti air bagi kehidupan ikan sehingga tempoh tuaian. Dalam kajian ini, konsep RAS telah dibangunkan untuk menentukan keupayaan dalam mengawal kualiti air yang mampan bagi akuakultur.

© COPYRIGHT UPM

Kajian ini bertujuan untuk membangunkan ternakan ikan keli dalam utiliti bersepadu dengan menggunakan air secara cermat dan tanah secara optimum. Objektif kajian ini ialah; (1) untuk menentukan kadar pertumbuhan ikan keli dengan sistem peredaran mekanikal dan pengendalian kualiti air melalui unit penapisan biologi; (2) untuk mengukur parameter kualiti air tangki pengeluaran ikan; dan (3) untuk mengkaji proses dan cara yang ia boleh sebati sehingga menyebabkan lebih banyak keadaan kualiti air yang stabil dalam unit kultur dan juga dalam pengurangan pencemaran.

Ikan keli *(C. batrachus)* dikultur dalam tiga tangki berasingan masing-masing dengan saiz 5-8 cm (Tangki 1), 8-12 cm (Tangki 2) dan 12-15 cm (Tangki 3) selama 30 hari. Setiap tangki mengandungi 300 ekor. Sampel air telah diamati dalam dua waktu (pagi dan petang) dengan masing-masing dua kali ulangan sepanjang tempoh eksperimen. Parameter kualiti air seperti suhu, kekeruhan, oksigen terlarut, pH, dan amonia telah telah diukur. Semua tangki disambungkan ke pembekal air melalui penapis dengan menggunakan paip PVC. Sisa terhasil mendap ke dalam tangki pemendapan. Air yang melalui tangki pemendapan seterusnya dipam ke dalam penapis dan kembali ke tangki kultur ikan secara edaran semula. Ikan diberi makan pada kadar 3% daripada jumlah berat ikan badan dua kali sehari (pagi dan petang).

Sistem RAS dan penapis biologi dengan multimedia dan basuh balik telah berjaya menghasilkan kualiti air untuk kultur ikan keli secara intensif. Kadar kemandirian ikan adalah melebihi 95%, dengan pertambahan berat ikan keli ialah 11.00$\pm$1.90 gram/ikan dalam Tangki 1, 20.45$\pm$0.85 gram/ikan dalam Tangki 2, dan 25.65$\pm$0.95 gram/ikan dalam Tangki 3. Kadar pertambahan panjang dan berat ikan keli dalam

© COPYRIGHT UPM

Tangki 1 dari 6.5±1.5 cm (1.00±0.10 gram) kepada 12.0±2.0 cm (12.00±1.80 gram), Tangki 2 dari 10.0±2.0 cm (2.10±0.60 gram) kepada 16.0±2.0 cm (22.15±0.30 gram), dan Tangki 3 dari 13.5±1.5 cm (5.65±1.25 gram) kepada 18.5±2.5 cm (32.33±0.30 gram).

Analisis data menggunakan reka bentuk rawak lengkap (*Completely Randomized Design*/CRD) dengan tiga rawatan dan dua replikat menunjukkan bahawa jumlah peratusan kemandirian ikan keli pada Tangki 2 bertahan lebih tinggi daripada tangki lainnya. Peratusan kemandirian daripada yang tertinggi kepada yang terendah ialah 98,33%±0,33 (Tangki 2), 95,33±0,67% (Tangki 1) dan 95,17±0,84% (Tangki 3). Ujian ANOVA menunjukkan bahawa tiada perbezaan ketara kualiti air dan prestasi pertumbuhan antara satu tangki dengan yang lain dalam percubaan ini. Keputusan menunjukkan bahawa sistem RAS ini boleh berjaya digunakan dan terus dibangunkan untuk meningkatkan pengeluaran ikan secara intensif. Hampir tidak terdapat kesan negatif yang besar kepada kualiti air, pertumbuhan dan kadar kemandirian ikan keli selama tempoh pemerhatian dalam tangki. Ini bermakna bahawa perbezaan dalam saiz ikan dalam tangki masing-masing tidak memberi kesan yang besar ke atas kadar pertumbuhan, kemandirian dan kualiti air kultur ikan keli.

© COPYRIGHT UPM

ACKNOWLEDGEMENTS

In the name of Allah, Most Gracious and Most Merciful

First of all, I would like to express my sincere gratitude and deep thanks to my supervisor, Assoc. Prof. Ir. Dr. Johari Endan for guidance, suggestions, never-ending patience, kindness and willingness to assist me throughout the whole course of this project. Without his valuable advice and support, it would not have been possible for me to complete my research. I have learnt a lot of useful knowledge from him throughout this research. Thanks is also extended to my co-supervisor, Assoc. Prof. Ir. Muhammad Salih bin Haji Ja'afar and Dr.Farah Saleena Taip for their kind assistance, support, and suggestions for this work and during the preparation of this entire thesis.

Last but not least, to my parents, Drs. Haji Syahbuddin Aziz and Hajjah Mardhiah Ibrahim, my wife, Drh. Rizki Amelia, my brothers, Nazaruddin and Iip gunawan, my sister, Almarhumah Fitri Yanti, and my daughters, Rumaisha Khairatun Hisan, Ruhama Althafunnisa, and Rufaida Nailal Husna for their continuous love, support and encouragement to complete my project thesis.

I convey my thanks to all of my friends, housemate, colleagues and UPM support staff.

© COPYRIGHT UPM

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Ir. Johari Endan, Phd
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Ir. Muhammad Salih bin Hj.Ja'afar
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Farah Saleena Taip, Phd
Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, Phd
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

© COPYRIGHT UPM

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

_____

RAHMAT FADHIL

Date: 9 November 2012

© COPYRIGHT UPM

TABLE OF CONTENTS

		Page
DEDICATION		iii
ABSTRACT		iv
ACKNOWLEDGEMENTS		x
APPROVAL		xi
DECLARATION		xiii
TABLE OF CONTENT		xiv
LIST OF TABLES		xvii
LIST OF FIGURES		xiv
LIST OF APPENDICES		xvi
LIST OF ABBREVIATIONS		xxiii
CHAPTER		
1	**INTRODUCTION**	1
	1.1 Background	1
	1.2 Problem Statement	5
	1.3 Objectives	6
2	**LITERATURE REVIEW**	7
	2.1 Introduction	7
	2.2 Recirculating Aquaculture System	7
	2.3 Biofilter	10
	2.3.1 Nitrification Process	11
	2.3.2 Types of Biofilter	12
	2.3.3 Proper System	14
	2.4 Factors that Affect the Nitrification Process	15
	2.4.1 Dissolved Oxygen	16
	2.4.2 Temperature	17
	2.4.3 pH	18
	2.4.4 Concentration of Substrates (Ammonia, Nitrite, Alkali)	19
	2.4.5 Light	20
	2.5 Filter Unit and Multi Media of Filter	21
	2.5.1 Fiberglass Reinforced Plastic (FRP)	22
	2.5.2 Gravel	23
	2.5.3 Silica Sand (Coarse, Medium and Fine)	25
	2.5.4 Activated Carbon	26
	2.6 Water Quality	28
	2.7 Parameters of Water Quality for Aquaculture	29
	2.7.1 Temperature	32
	2.7.2 Turbidity	34
	2.7.3 pH value	36
	2.7.4 Oxygen	38
	2.7.5 Ammonia	43
	2.8 Catfish	45
	2.8.1 Characteristic of Catfish	46
	2.8.2 Asian Catfish *(Clarias batrachus)*	48

© COPYRIGHT UPM

2.9 Growth Performance 52

3 METHODOLOGY 54
3.1 Description of RAS Component 55
3.1.1 Culture Tank 55
3.1.2 Sump Particulate Tank 55
3.1.3 Biofilter 56
3.1.4 Pump 57
3.1.5 Pipes and Valves 58
3.2 Species Selection 58
3.2.1 Catfish 58
3.2.2 Water Quality Parameter for Catfish 60
3.3 System Layout 60
3.4 Experimental Design 63
3.5 Evaluation of System Performance 63
3.5.1 Water Flow Rate 63
3.5.2 Feed 65
3.5.3 Filtration and Backwash 66
3.6 Measurement of Water Quality 67
3.8.1 Temperature, pH and Dissolved oxygen 68
3.8.2 Turbidity 68
3.8.3 Ammonia 69
3.7 Observation on Fish 70
3.7.1 Survival Rate 70
3.7.2 Total Length 71
3.7.3 Weight Gain 71

4 RESULTS AND DISCUSSION 72
4.1 Operation Performance of Recirculating Aquaculture System 72
4.1.1 Water Exchange Rate 72
4.1.2 Removal of Solids 73
4.1.3 Backwash 74
4.2 Analysis of Water Quality 74
4.2.1 Temperature 76
4.2.2 Turbidity 77
4.2.3 pH 78
4.2.4 Dissolved Oxygen 79
4.2.5 Total Ammonia Nitrogen 80
4.3 Relationship of Water Quality and Feed 81
4.4 Biological Performance of Fish 83
4.4.1 Survival Rate 83
4.4.2 Total Length 85
4.4.3 Weight Gain 85
4.5 Summary 86

5 CONCLUSIONS AND RECOMMENDATIONS 87
5.1 Conclusions 87
5.2 Recommendations 88

© COPYRIGHT UPM

REFERENCES 90
APPENDICES 105
BIODATA OF STUDENT 127
LIST OF PUBLICATIONS 128

© COPYRIGHT UPM

LIST OF TABLE

Table		Page
2.1	The characteristics of nitrifying bacteria	16
2.2	Temperature for growth of nitrifying bacteria	18
2.3	Advantages of FRP products	22
2.4	Standard grade of gravel	24
2.5	Characterstics of gravel	24
2.6	Standard grade of silica sand	26
2.7	Characteristics of silica sand	26
2.8	The physical characteristics and operational specifications of activated carbon	27
2.9	Water quality criteria for aquaculture	31
2.10	General temperature guidelines	33
2.11	Temperature relationship with the dissolved oxygen	33
2.12	Turbidity tolerance levels for aquaculture	35
2.13	The pH value tolerance levels and effect for aquaculture	38
2.14	The influence of dissolved oxygen on a fish	40
2.15	Comparison of pure water temperature and the resulting dissolved oxygen	42
2.16	The relationship between the percentages of total ammonia with pH and Temperature	44
2.17	Factors affecting the toxicity of ammonia to fish	45
2.18	Other name for catfish according to different countries	48
3.1	Water quality parameters for catfish	60
4.1	A system summary of the Asian catfish culture in RAS	72

© COPYRIGHT UPM

4.2 A summary of water quality parameters in RAS culture tank 75

4.3 Relationship between body weights with total feed fed during the 30 days of culture period 82

4.4 Production performance figures of Asian catfish cultures in RAS 84

© COPYRIGHT UPM

LIST OF FIGURE

Figure		Page
2.1	Nitrification process in freshwater	10
2.2	The nitrogen cycle	14
2.3	Gravel	24
2.4	a.Coarse silica sand, b.Medium silica sand, c.Fine silica sand	25
2.5	Activated carbon	27
2.6	Anatomy of catfish *(Clarias* sp*)*	46
2.7	Asian catfish *(Clarias batrachus)*	49
3.1	Location of RAS experiment	54
3.2	Filtration unit made of Fiberglass Reinforced Plastic (FRP)	56
3.3	Water pump	57
3.4	Pipes and valves	58
3.5	The three size of Asian catfish	59
3.6	Layout diagram for Recirculating Aquaculture and Bio-Filtration system	61
3.7	Composition of multi-media filter in filtration unit	62
3.8	Diagram of experimental design	64
3.9	Fish feed	65
3.10	Sample of Mud in the sedimentation tank	66
3.11	Normal Service Run, Backwash and Fast Rinse of Filter	67
3.12	Thermometer/pH meter (HACH SensION2)	68
3.13	DO meter (YSI 55 DO)	68
3.14	The HACH Model 2100P Portable Turbidity meter	69

© COPYRIGHT UPM

3.15 Ammonia meter (HANNA HI 93715) 70

3.16 Common method of fish measurement 71

4.1 Total Feed Fed 82

© COPYRIGHT UPM

LIST OF APPENDIX

Appendix		Page
A.1	Data of water quality	106
A.2	Data of fish	107
B	List of Material and Equipment	108
C	Nutrition Composition and Ingredients of Feed Fish	109
D	Material Safety Data Sheet of Gravel and Silica Sand	110
E.1	Culture Tank	111
E.2	Sumb Particulate Tank	111
E.3	Scooping	112
F.1	Water temperature in tank	113
F.2	ANOVA for temperature	113
F.3	Water turbidity level in tank	113
F.4	ANOVA for turbidity	114
F.5	pH of Water	114
F.6	ANOVA for pH	114
F.7	Level dissolved oxygen in water	115
F.8	ANOVA for dissolved oxygen	115
F.9	TAN Level in water of tank culture	115
F.10	ANOVA for TAN	116
G.1	Survival rate	117
G.2	ANOVA for survival rate	117
G.3	Total length	117
G.4	ANOVA for total length	118

© COPYRIGHT UPM

G.5 Weight gain 118

G.6 ANOVA for weight gain 118

G.7 Initial weight 119

G.8 Final weight 119

H Glossary 120

© COPYRIGHT UPM

LIST OF ABBREVIATIONS

AM or am	- Ante Meridian (morning time)
ANOVA	- Analysis of Variance
ANSI	- American National Standards Institute
APHA	- American Public Health Association
ASTM	- American Society for Testing and Materials
ATC	- Automatically Compensate for Temperature
AWWA	- American Water Works Association
BOD	- Biochemical Oxygen Demand
C	- Celsius
CRD	- Completely Randomized Design
DF	- Degrees of Freedom
DGR	- Daily Growth Rate
DO	- Dissolved Oxygen
E	- Environment
ESC	- Enteric Septicemia of Catfish
F	- Fahrenheit
FE	- Feed Efficiency
FCR	- Feed Conversion Ratio
FRP	- Fiberglass Reinforced Plastic
IA	- Ionized Ammonia
JTU	- Jackson Turbidity Unit
L or l	- Liter

© COPYRIGHT UPM

LSD - Least Significant Different

MS - Mean Square

NSF - National Sanitation Foundation

NTU - Nephelometric Turbidity unit

PM or pm - Post Meridian (afternoon time)

POE - Point of Entry

POV - Point of Use

PVC - Polyvinyl chloride

RAS - Recirculating Aquaculture System

SGR - Specific Growth Rate

SR - Survival Rate

SS - Sum of Squares

T - Tank

TAN - Total Ammonia Nitrogen

TLR - Total Length Ratio

TSS - Total Suspended Solids

UIA - Unionized Ammonia

US EPA - United States Environmental Protection Agency

VOC - Volatile Organic Compound

WG - Weight Gain

WPCF - Water Pollution Control Federation

WQS - Water Quality Standard

Ag - Silver Cathode

CO_2 - Carbon Dioxide

$CaCO_3$ - Calcium Carbonate

© COPYRIGHT UPM

HCl	- Hydrogen Chloride
H_2SO_4	- Sulphuric Acid
H_2O	- Water
$MnCl_2$	- Manganese Chloride
MnO_2	- Manganese Dioxide
NH_3	- Ammonia
NaOH	- Natrium Hydroxide
$Na_2S_2O_3$	- Sodium Thiosulfate
NO_2	- Nitrites
NO_3	- Nitrates
O_2	- Oxygen
I_2	- Iodine
atm	- atmosphere
cm	- centimetre
g	- gram
kg	- kilogram
m	- metre
mg	- milligram
ml	- millilitre
mm	- millimetre
nm	- nanometre
ppm	- part per million
ppt	- part per ton

© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

1. Background

Studies on aquaculture continue to grow in recent years. Its development is very crucial to the sustainability of fish supply for the community, both for direct consumption by the consumer and for further processing by the industry. Aquaculture is an attempt to cultivate aquatic animals or plants, and it became a common term used to describe this specialty. Aquaculture can be developed to produce fish and plants for food, sources of income, and others (Satar, 1984).

Fish, shrimp, seaweed, and other aquatic organisms, depend on water as internal and external media. As an internal medium, water allows reactions of a raw material to take place in the bodies, transports food throughout the body, and carries the metabolic waste to be expelled from the body. As an external medium, water functions as a habitat. The water conditions for aquaculture are very important for the animals to grow and develop optimally. If the environment is in a bad condition, the fish can experience stress, become susceptible to diseases and ultimately die (Hernawati and Suantika, 2007; Boyd, 1990). Therefore, knowing the parameters of water quality alone is not sufficient. A total understanding of the characteristics of water, which is the habitat for water organisms is essential (Kordi and Tancung,

© COPYRIGHT UPM

2007). Parameters of water quality and quantity are crucial for the maintenance of life in the water, so that the results obtained will be optimal. Several important factors that are known are sources of water quality status, the condition of the pond, feed management, water circulation, dense stocking, plankton, dirt, and the environment. These factors greatly affect the overall water quality.

In aquaculture, fish is a general term that covers fish, prawn, oyster, mollusk and others. Metabolism in a fish body secretes compounds that can be toxic to its life (Golz, 1995). In addition, the excessive remains of rotting food can accumulate and become toxic compounds, such as ammonia. In tropical climate conditions, the metabolic processes and decomposition of feed remnants happen very quickly because the environmental temperature is high. Handling these issues, thus, is very important for maintaining healthy fishes. Replacing the water is one good way to tackle this problem, but it requires adequate water supply and can cause fish stress (Sudrajat and Gunawan, 2002). Based on this fact, a recirculating system would be an alternative.

In addition, water quality plays an important role in selecting the types of animal or plant to be cultured. If the freshwater has low oxygen content (below 4-5 ppm), only certain types of fishes can survive (Tucker, 1991). One of the type of fishes that can live in such environmental conditions is the catfish *(Clarias* sp*)* (Singh and Hughes, 1971). The reason is the catfish has an additional respirator known as *arborescent* (air breathing fish), so they can absorb oxygen directly from air (Bruton, 1979; Clay, 1979; Haylor, 1993; Khairuman *et al.*, 2008; Singh and Hughes, 1971).

© COPYRIGHT UPM

The Asian catfish *(Clarias batrachus)* is one of the most potential candidate for aquaculture in Asia, especially in Malaysia, Thailand, India, Indonesia, and Laos. The potentials of these species in aquaculture have been widely reported (Areerat, 1987; Phonekhampheng *et al.* 2008; Sahoo *et al.*, 2008; Sunarma, 2004; Bhat *et al.*, 2008). The availability of stocking material is always considered important for a successful culture of any fish species through collection of natural seeds, but it is not sustainable for an intensive culture. The scarcity of marketable fishes and seeds from the natural ground has been an issue for catfish. So the fish production in hatchery will be the only alternative for obtaining an optimum quantity of fish for the purpose through intensive breeding operation (Sahoo *et al.*, 2008).

Brune (2004) reported that most aquaculture production comes from land-intensive systems, such as catfish farms. Major advantages of pond fish culture are the low capital cost of earthen ponds and the reliability of pond fish production. The disadvantages are the need to continuously manage the pond oxygen concentration and other fluctuating water quality variables, preventing off-flavor, controlling predators and diseases, and providing labor for harvesting. These management difficulties, combined with land, water, and environmental constraints, have driven the search for technological improvements in pond aquaculture. One probable solution is to shift to more energy-intensive systems, either by increasing aeration in low-cost ponds or by abandoning ponds altogether in favor of recirculating tank or raceway processes (Brune *et al.*, 2004).

The traditional aquaculture production in ponds required large quantities of water. Approximately, 1 million gallons of water per acre are required to fill a pond and an

© COPYRIGHT UPM

equivalent volume is required to compensate evaporation and seepage throughout the year. Assuming an annual pond yield of 5,000 pounds of fish per acre, approximately 100 gallons of water are required per pound of fish production (Losordo et al., 1998). This is a large investment in fish farming. Recirculating aquaculture production systems may offer an alternative to reduce the production and investment cost. In the recirculating systems, the water is treated in tanks and reused; thus, substantially less land is required (Losordo *et al.*, 1998; Yudha, 2005). It is expected that the cost needed for a recirculating system is cheaper than that of in a traditional fish production in ponds. The system is also independent of environmental changes.

Recirculating system has several advantages, such as environmentally friendly, and safe from contamination that occurs in aquatic environments. It can minimize the impact of a disease outbreak from the outside environment, and its water quality parameters are more stable. Several weaknesses of this system is the accumulation of organic material, which originated from surplus feed, fish feces, and dead planktons (Yudha, 2005). The management and maintenance of water quality require serious attention because water quality affects fish life.

Research on the concept of recirculating aquaculture production systems is important, especially related to water quality control and growth performance. The investment in the aquaculture farming model is not a problem if it is implemented in a large scale. Considering the needs and benefits of this circulation aquaculture system in the future, the experiment in a small-scale is significant.

© COPYRIGHT UPM

A key to successful recirculating production systems is the use of cost-effective components of the water treatment system. All recirculating production systems remove waste solids, oxidize ammonia and nitrite-nitrogen, remove carbon dioxide, and aerate or oxygenate the water before returning it to the fish tank (Losordo *et al.*, 1998). Applying this concept, the water quality in production ponds can be maintained for the fish survival until the harvesting stage.

1.2 Problem Statement

Strategies to increase fishery's production are constantly evolving to get better results. Water quality management is needed in the aquaculture system, because it can affect the organisms living in the water. The use of Recirculating Aquaculture System in the aquaculture industry is the latest challenge for the fishing communities and researchers. Utilization of filtration with a variety of media to maintain healthy water quality for aquaculture remains an important issue to be investigated. The main problem in fishery aquaculture is the techniques to maintain healthy water quality continuously for growth and survival of fish to the harvesting stage. Recirculating Aquaculture System (RAS) with a bio-filtration unit is developed to study the effectiveness in controlling water quality for sustainable aquaculture.

One of the advantages of this study is the use of domestic biofilter that has been used in the treatment of household water, which is now being applied in a recirculating aquaculture system to produce freshwater fish. Given the fact that domestic biofilter has been commercially available in the market and widely used in

© COPYRIGHT UPM

housing, domestic biofilter is expected to further provide convenience for fishing industry and fish farmers to increase fish production from time to time. Integrating domestic biofilter with RAS is something new, stimulating new research opportunities.

1.3 Objectives

The research aimed at developing a simple integrated system for aquaculture with thrifty water and land. Principally, the fish production under this system can add income value to fish breeders and improve the livelihood of the community. The objectives of this research were:

1. to design a small-scale Recirculation Aquaculture System (RAS).
2. to determine growth performance of various sizes of Asian catfish *(Clarias batrachus)* in RAS.
3. to assess influence of size variation of *Clarias batrachus* on water quality in RAS.

© COPYRIGHT UPM

REFERENCES

Abbas, G. (2000). Length-weight and Condition factor relationship of Anchovy *(Coilia dussumieria)* and Mullet, (Liza carinata) from Bhanbhore backwaters along Sindh coast (Pakistan: Northern Arabian Sea). *Pakistan J. of Zool., 32*(3), 223-228.

Affandi, R., & Tang, U. M. (2002). *Fisiologi Hewan Air (in Indonesian)*. Riau: Universitas Riau Press.

Ahmad, T., Ratnawati, E., & Yakob, M. J. R. (1998). *Budi Daya Bandeng Secara Intensif*. Jakarta, Indonesia: PT.Penebar Swadaya.

Akinwole, A. O., & Faturoti, E. O. (2007). Biological Performance of African Catfish *(Clarias gariepinus)* Cultures in Recirculating System in Ibadan. *Aquacultural Engineering, 36*, 18-23.

Aksnes, A., Izquierdo, M. S., Robaina, L., Vergara, J. M., & Montero, D. (1997). Influence of fish meal quality and feed pellet on growth, feed efficiency and muscle composition in gilthead seabream *(Sparus aurata). Aquaculture, 153*, 251-261.

Amstrong, M. S. (1984). *Environmental Factors Affecting off-flavor in Catfish Production Ponds.* Doctoral, Auburn University, Auburn, Alabama.

Anggoro, S. (2008). Evaluation of Biomass Production, Survival Rate, and Feed Conversion on Culture of *Penaeus monodan* Fab. use Liner Bottom Pond Construction. *Sains Akuatik, 11*, 85-92.

Antoniou, P., Hamilton, J., Koopman, B., Jain, R., Holloway, B., Lyberatos, G., & Svoronos, S. A. (1990). Effect of temperature and ph on the effective maximum specific growth rate of nitrifying bacteria *Water Research, 24*(1), 97-101.

Aquatic Community. (2003). Common Fish Diseases. from Aquatic Community, http://www.aquaticcommunity.com/disease/common.php (accesed February 24, 2011)

Ardiansyah, Hanifah, U., & Setiawan, B. I. (2004, December 3-5, 2004). *Dynamic of Water Balance Analysis in Recirculating System of Fish Hatchery.* Paper presented at the The 3rd International Joint Seminar Toward Harmonization Between Development and Environment Conservation in Biological Production, Banten, Indonesia.

Areerat, S. (1987). *Clarias* Culture in Thailand. *Aquaculture, 63*, 355-360.

Asmawi, S. (1983). *Fish Culture in Floating Cage Pond (in Indonesian)* (1 ed.). Jakarta: PT. Gramedia.

© COPYRIGHT UPM

Atmadja, G. W. (1977). *Bahan Bacaan Akuakultur* (Vol. Jilid I). Bogor,Indonesia: Bagian Akukultur,Dept.Tata Produksi Perikanan,IPB.

Ayanda, O. I. (2008). Comparison of Parasitic Helminth Infection between the Different Age Groups of *Clarias gariepinus* from Asa dam Ilorin, North-Central Nigeria. *African Journal of Environmental Science and Technology, 22*(11), 404-406.

Baber, M. (2005). *Clarias batrachus* (Fish). Retrieved June 2009, from IUCN/SSC Invasive Species Specialist Group (ISSG), http://www.issg.org/database/species/ecology.asp?si=62&fr=1&sts=sss

Bachtiar, Y. (2006). *Complete Guide to African Catfish Culture (in Malay)* (1 ed.). Kuala Lumpur: Synergy Media.

Betlach, M. R., & Tiedje, J. M. (1981). Kinetic Explanation for Accumulation of Nitrite, Nitric Oxide, and Nitrous Oxide During Bacterial Denitrification. *Applied and Environmental Microbiology, 42*(6), 1074-1084.

Bhat, B. V., Corsin, F., Anis, S., Sai, J. P., Biswas, D. K., & Vinod, P. N. (2008). *Present and Future of Pangasius Farming in India.* Paper presented at the International Symposium on Catfish Farming in Asia, Cantho University, Vietnam.

Bio-Systems International. (2010). Frequently Asked Questions, The Nitrification Process & More. Retrieved Mai 3, 2010, 2010, from http://nitrifier.com/FAQNitrificationProcess/tabid/279/Default.aspx

Boreham, S., Ingram, B. A., Jones, P. L., & Collins, R. O. (2004). Water quality and intensive Murray cod aquacuture. In B. A. Ingram & S. S. De Silva (Eds.), *Developments in Intensive Commercial Aquaculture Production Technology for Murray cod. Final report to Fisheries Research and Development Corporation (Project No. 1999/38)*. Melbourne: Department of Primary Industries.

Bovendeur, J., Eding, E. H., & Henken, A. M. (1987). Design and performance of a water recirculating system for high-density culture of the African catfish, *Clarias gariepinus* (Burchell 1822). *Aquaculture, 63*, 329-353.

Boyd, C. E. (1988). *Water Quality in Warmwater Fish Ponds*. Alabama: Auburn University.

Boyd, C. E. (1990). *Water Quality in Ponds for Aquaculture*. Birmingham, Alabama, US: Birmingham Publishing Co.

Boyd, C. E. (2000). *Water Quality an Introduction*. Massachusetts: Kluwer Academic Publishers.

Boyd, C. E., & Lycopper. (1979). *Water Quality Management in Pound Fish Culture Aquacultural Experiment Station*. Alabama,US: Aurbun University.

© COPYRIGHT UPM

Brown, A. L. (1987). *Freshwater Ecology*. London, UK: Heinemann Educational Books.

Brown, M. E. (1957). *The Physiology of Fishes* (Vol. II). New York,USA: Academic Press Inc.Publisher.

Brown, M. E. (1962). *The Physiology of Fishes, Metabolism* (Vol. I). New York: Academic Press Inc.

Brown, S. W. (1981). *Water Quality and off-flavor Problems in Commercial Channel Catfish in West-Central Alabama* Master Thesis, Auburn University, Auburn, Alabama.

Brown, S. W., & Boyd, C. E. (1982). Off-flavor in Channel Catfish from Commercial Ponds. *Transaction of the American Fisheries Society, 111*, 379-383.

Brune, D. E., Schwartz, G., Eversole, A. G., Collier, J. A., & Schwedler, T. E. (2004). Partitioned Aquaculture Systems. from SRAC Publication

Bruton, M. N. (1979). The breeding biology and early development of *Clarias gariepinus* (Pisces claridae) in Lake Sibaya, South Africa, with a review of breeding species of the subgenus *Clarias* (clarices). *Transc. Zool. Soc. London, 35*, 1-45.

Bureau, D. P., Kaushik, S. J., & Cho, C. Y. (2002). Bioenergetics. In John E. Halver & R. W. Hardy (Eds.), *Fish Nutrition* (3rd ed.). California: Academic Press.

Burrows, R.E., and B.D. Combs. 1968. Controlled environments for salmon propagation. *Prog. Fish-Cult*. *30*: 123-136.

Buttner, J. K., Soderberg, R. W., & Terlizzi, D. E. (1993). An Introduction to Water Chemistry in Freshwater Aquaculture (Vol. 170). Nort Dartmouth, Massachusetts: Northeastern Regional Aquaculture Center, University of Massachusetts Dartmouth.

Buwono, I. D. (2000). *Requirement of Essential Amino Acids in Feed Rations*. Jakarta: Agritek YPN.

Chen, S., Ling, J., & Blancheton, J. (2006). Nitrification kinetics of biofilm as affected by water quality factors. *Aquacultural Engineering, 34*(179-197).

Chervinsky, J. (1982). *Environmental Physiology of Tilapia.* Paper presented at the The Biology and Culture of Tilapias. ICLARM, Manila.

Ciudad, G., González, R., Bornhardt, C., & Antileo, C. (2007). Modes of operation and pH control as enhancement factors for partial nitrification with oxygen transport limitation. *Water Research, 41*(20), 4621-4629.

Clay, D. (1979). Population biology, growth and feeding of the African Catfish, *Clarias gariepinus*, with special reference to juveniles and their importance in fish culture. *Arch. Hydrobiol, 87*(4), 453-482.

© COPYRIGHT UPM

Colt, J., & Orwicz, K. (1991). Modeling Production Capacity of Aquatic Culture Systems Under Freswater Conditions. *Aquacultural Engineering, 10*, 1-29.

Colt, J. E., & Amstrong, D. A. (1981). *Nitrogen toxicity to crustaceans, fish, and molluscs.* Paper presented at the Bioengineering Symposium for Fish Culture, Maryland, USA.

D'Silva, A. M., & Maughan, O. E. (1995). Effects of Density and Water Quality on Red Tilapia *(Oreochromis mossambicus X O. urolepis hornorum)* in Pulsed-Flow Culture System. *Journal of Applied Aquaculture, 5*(1), 69-75.

DAFF (2010), Airbreathing or walking catfish, http://www.daff.qld.gov.au/images/Fisheries_SpeciesIdentification/Fish-Pest-Walking-Catfish-600.jpg, Acces on Desember 23, 2010

Darmanto. (2009). *Filtratration (in Indonesian)*. Universitas Gadjah Mada. Yogyakarta.

Darmanto, & Kamulyan, B. (2009). *Pre-treatment in Filtration Process (in Indonesia)*. Universitas Gadjah Mada. Yogyakarta.

De Graaf, G., & Janssen, J. (1996). *Handbook on the Artificial Reproduction and Pond Rearing of the African Catfish Clarias gariepinus in Sub-Saharan Africa - A Handbook* (Vol. 362). Rome: FAO.

De Silva, S. S., & Anderson, A. (1995). *Fish Nutrition in Aquaculture*. London: Chapman and Hall.

Dennis, H. R., & Webster, T. (2008). High ph inhibits nitrifying bacteria. Des Moines, Iowa, USA: Des Moines Water Works (www.dmww.com), American water Works Association (www.awwa.org), www.dmww.com/Laboratory/highpHinhibitsnitrifyingbacteria.pdf, Access on May 22, 2010.

Department of Primary Industries. (2008). Best practice environmental management guidelines for recirculating aquaculture systems. *Victoria Management Report Series, 37*.

Djajadiredja, Rustami, Hatimah, S., & Arifin, Z. (1990). Guidelines for the Introduction of Freshwater Fishery Resources (Fish Species of Economic Importance) (in Indonesian). Jakarta: Direktorat Jendral Perikanan, Departemen Perikanan dan Kelautan Indonesia.

Djajasewaka, H. (1995). *Fish Feed (in Indonesian)*. Jakarta: CV. Yasaguna.

Djatmika, D. H., Farlina, & Sugiharti, E. (1986). *Business of Catfish Culture*. Jakarta: C.V. Simplex.

Djumanto. (2000). Viability of Asian Catfish *(Clarias batrachus)* Eggs Fertilized at Various Ovulation Interval. *J. Fish Science, II*(2), 48-56.

© COPYRIGHT UPM

Dojlido, J. R., & Best, G. A. (1993). *Chemistry of Water and Water Pollution*. Great Britain,UK: Ellis Horwood Limited.

Downing, A. L., Painter, H. A., & Knowles, G. (1964). Nitrification in the activated sludge process *Journal Proc. Inst. Sew Purif., 63*(130-153).

Durborow, M. R., Crosby, M. D., & Brunson, W. M. (1997). Ammonia in Fish Ponds. from Southern Regional Aquaculture Center (SRAC): Mississipi.

Durborow, R. M., Thune, R. L., Hawke, J. P., & Camus, A. C. (1998). Columnaris Disease, A Bacterial Infection Caused by *Flavobacterium columnare*. *SRAC Publication, 479*.

Edhi, W. A. (2001). From Closed Recirculation System to Closed System *Mitra Bahari* (Vol. 2, pp. 51-52). Jakarta: Department of Fisheries and Marine Republic of Indonesia.

Effendi, H. (2003). *Telaah Kualitas Air Bagi Pengelolaan Sumber Daya dan Lingkungan Perairan*. Yogyakarta,Indonesia: Kanisius.

Ellis, M. M., Westfall, B. A., & Ellis, M. D. (1948). Determination of Water Quality *Research Report 9*. Washington D.C: Fish and Wildlife Service, United States Goverment Printing Office.

Environment Canada. (1979). *Water Quality Sourcebook, A Guide to Waters Quality Parameters*. Ottawa: Inland Waters Directorate, Water Quality Branch.

Fdz-Polanco, F., Villaverde, S., & Garcia, P. A. (1994). Temperature effect on nitrifying bacteria activity in biofilters: activation and free ammonia. *Water Science Technology, 30*(11), 121-130.

FFWCC. (2010). How to Measure Freshwater Fish. Retrieved November 24, 2010, from MyFMC.com

Fiberglass Grating Manufacturers Council. (2005). ABC's of FRP. http://www.acmanet.org/fgmc/abc_frp.htm, Acces on November 20, 2010

Filtermedia. (2009). Sand filter media. http://www.statscrop.com/www/ filtermedia.com, Acces on November 20, 2010

Fishbase. (2009). Common names of *Clarias batrachus*. from Fishbase.org, http://www.fishbase.org/comnames/CommonNamesList.php?ID=3054&Genus Name=Clarias&SpeciesName=batrachus&StockCode=3250, Acces on March 20, 2010

Fishing-Khaolak. (2009). Walking Catfish. In Catfish (Ed.), *Fishing-Khaolak.com*. Bangkok, Thailand: http://www.fishing-khaolak.com/fish_species/freshwater_ species/walking_catfish.html.

© COPYRIGHT UPM

Forteath, N., Wee, L., & Frith, M. (1993). Water Quality. In P. Hart, O'Sullivan (Ed.), *Recirculating System: Design, Construction and Management*. Launceston: University of Tasmania, Australia.

Fowler, H. W. (1941). Contributions to the biology of the Philippine archipelago and adjacent regions: fishes of the group Elasmobranchii, Holocephali, Isospondyli and Ostariophysi. *"Albatros" Bulletin 100, 13,* 734-743.

Fritz Industries. (2010). Nitrifying Bacteria Facts. from Bio-Con Labs, Inc, Gainesville, Florida USA, http://www.bioconlabs.com/fritzyme.html , Access on May 3, 2010

Froese, R., & Pauly, D. (2007). Genus *"Clarias"*. Retrieved Jun 2009, from Fish Base, Stockholm, Sweden http://www.fishbase.org/

Froese, R., & Pauly, D. (2009). Common Names of *Clarias batrachus*. from Fish Base, Stockholm, Sweden, http://www.fishbase.org/comnames/CommonNamesList.php?ID=3054&GenusName=*Clarias*&SpeciesName=*batrachus*&StockCode=3250, Acces on March 20, 2010

Gerardi, M. H. (2003). Temperature. In M. H. Gerardi (Ed.), *Nitrification and Denitrification in the Activated Sludge Process*. New York, USA: John Wiley and Sons.Inc.

Gieseke, A., Purkhold, U., Wagner, M., Amann, R., & Schramm, A. (2001). Community Structure and Activity Dynamics of Nitrifying Bacteria in a Phosphate-Removing Biofilm. *Appl. Environ. Microbiol., 67*(3), 1351-1362.

Golz, W. J. (1997). *A Theoretical Model of Nitrification in Floating Bead Filters.* MS Thesis, Louisiana State University, Baton Rouge, LA.

Golz, W. J., Rusch, K. A., & Malone, R. F. (1999). Modeling the Major Limitations on Nitrification in Floating-Bead Filters. *Aquacultural Engineering, 20*, 43-62.

Gomez, A. A., & Gomez, K. A. (1984). *Statistical Procedures for Agricultural Research* (2nd ed.). Canada: John Wiley & Son.

Gregory, J. (2006). *Particles in Water*. London: Taylor & Francis Group.

Gunther, A. (1864). *Catalogue of the Physostomi containing families Siluridae, Characinidae, Haplochitonidae, Sternoptychidae* (Vol. 5). London: Scopelidae in the Collection of the British Museum, London.

Ha, J. H., Ong, S. K., Surampalli, R., & Song, J. (2010). Temperature effects on nitrification in polishing biological aerated filters (BAFs) *Environment Technology, 31*(6), 671-680.

Hall, I. R. (1974). Some studies on nitrification inthe activated-sludge process. *Wat. Pollut. Control, 73*, 538-547.

© COPYRIGHT UPM

Halver, J. E. (1989). *Fish Nutrition*. Washington: Academic Press Inc., University of Washington State

Haylor, G. S. (1993). Aspect of the Biology and Culture of the African Catfish Clarias gariepinus (Burchell,1822) with Particular Reference to Developing African Countries. In J. F. Muir & R. J. Roberts (Eds.), *Recent Advances in Aquaculture IV* (pp. 235-294). New York, USA: Blackwell.

Helfrich, L. A., & Libey G. (2003). *Fish Farming in Recirculating Aquaculture System (RAS)*. Virginia Tech.: Department of Fisheries and Wildlife Sciences.

Hernawati, & Suantika, G. (2007). Recirculating System for Gouramy Fish Seed Culture. *DiSainTek, 1*(1).

Hickling, C. F. (1971). *Fish Culture*. London: Faber and Faber.

Hochheimer, J., & Wheaton, F. (1998). Biological Filters: Trickling and RBC Design. *Proceedings of the Second International Conference on Recirculating Aquaculture*, Virginia Polytechnic Institute and State University Roanoke,Virginia

Holiday, F. G. T. (1969). The Effect of Salinity on The Eggs and Larvae of Teleost. In W. S. Hoar & D. J. Randall (Eds.), *Fish Physiology Volume I* (pp. 293-309). New York: Academic Press.

Hora, S. L. (1936). Siluroid Fishes of India, Burma and Ceylon. *Records of the Indian Mus. XXXVIII,* 347-361.

Horn, H. (1994). Dynamics of a nitrifying bacteria population in a biofilm controlled by an oxygen microelectrode. *Water Science Technology, 29*(10-11), 69-76.

Horn, H., & Hempel, D. C. (2000). Substrate utilization and mass transfer in an autotrophic biofilm system: Experimental results and numerical simulation. *Biotechnology and Bioengineering, 53*(4), 363-371.

Horrigan, S. G., Garlucci, A. F., & Williams, P. M. (1981). Light inhibition nitrification in sea surface films. *Journal of Marine Research, 39*(3), 557-565.

Huisman, E. A., & Richter, C. J. J. (1987). Reproduction, Growth, Health Control and Aquaculture Potential of the African Catfish, *Clarias gariepinus* (Burchell 1822). *Aquaculture, 63*, 1-4.

Hutchinson, W., Jeffrey M. O"Sullivan, D., Casement, D and Clark, S. (2004). Recirculating Aquaculture System Minimum Standard for Design, Construction and Management. South Australia Research and Development Institute.

Jaunkey, K., & Ross, B. (1982). *A Guide to Tilapia Feeds and Feeding*. Stirling, Scotland, UK: Institute of Aquaculture, University of Stirling.

© COPYRIGHT UPM

Jeffries, M., & Mills, D. (1996). *Freshwater Ecology, Principles and Applications* Chickester, UK: John Wiley and Sons.

Khairuman, & Amri, K. (2008). *Intensive Culture of Catfish (in Malay)*. Kuala Lumpur: Synergy Media Books.

Khairuman, Toguan S, & Amri, K. (2008). *African Catfish (C.gariepinus) Culture in Canvas Pond* (1 ed.). Jakarta: AgroMedia Pustaka.

Kır, M. (2009). Nitrification Performance of a Submerged Biofilter in a Laboratory Scale Size of the Recirculating Shrimp System. *Turkish Journal of Fisheries and Aquatic Sciences, 9*, 209-214.

Klontz *et al.* (1979). A Good Deal From Egg Sac to Creel *A Manual for Trout and Salmon Production*. Murray, Utah: Sterling H. Nelson and Sons.

Kompas (2010), Industrialization of catfish, (in Indonesian), *Kompas Newspaper*, 31 Juli 2012, p. 21.

Kordi K., M. G. H. (2004). *Asian Catfish Culture* (1 ed.). Jakarta: Rineka Cipta and Bina Adiaksara.

Kordi, M. G. H. (1996). *Parameter Kualitas Air*. Surabaya, Indonesia: Penerbit Karya Anda.

Kordi, M. G. H. (2004). *Catfish Culture (in Indonesian)*. Jakarta, Indonesia: Rineka Cipta & Bina Adiaksara.

Kordi, M. G. H., & Tancung, A. B. (2007). *Pengelolaan Kualitas Air Dalam Budi Daya Perairan* (1 ed.). Jakarta: PT. Rineka Cipta.

Kottelat, M., & Whitten, A. J. (1996). Freshwater Biodiversity in Asia with Special Reference to Fish *World Bank Technical Paper* (Vol. 343). Washington, USA: Periplus Editions.

Kumari, J., Swain, T., & Sahoo, P. K. (2003). Dietary bovine lactoferrin induces changes in immunity level and disease resistance in Asian catfish *Clarias batrachus*. *Veterinary Immunology and Immunopathology, 94*(1-2), 1-9.

Lawson, T. B. (1995). *Fundamental of Aquaculture Engineering*. New York, USA: Chapman and Hall.

Lelana, I. Y. B., & Triyatmo, B. (2000). African Catfish *(Clarias gariepinus)* Culture with Different Water Exchange Volume. *J. Fish Science, II*(2), 25-30.

Lekang. O-I., (2008). *Aquaculture Engineering*. Hoboken, NJ: Wiley-Blackwell.

Liao, P. B., & Mayo, R. D. (1972). Salmonid hatchery water reuse systems. *Aquaculture, 1*, 317-335.

© COPYRIGHT UPM

Liao, P. B., & Mayo, R. D. (1974). Intensified Fish Culture Combining Water Re-Conditioning with Pollution Abatement. *Aquaculture, 3*, 61–85.

Lim, K. K. P., & NG, H. H. (1999). *Clarias batu,* A New Species of Catfish (Teleostei: Clariidae) from Pulau Tioman, Peninsular Malaysia. *The Raffles Bulletin of Zoology, 6*, 157-167.

Ling, J., & Chen, S. (2005). Impact of organic carbon on nitrification performance of different biofilters. *Aquacultural Engineering, 33*(2), 150-162.

Lloyd, R. (1992). *Pollution and Freshwater Fish*. West Byfleet,USA: Fishing News Books.

Losordo, T. M., Masser, P. M., & Rakocy, J. (1998). Recirculating Aquaculture Tank Production Systems, An Overview of Critical Considerations. from Southern Regional Aquaculture Center (SRAC)

Losordo, T. M., & Westerman, P. W. (1994). An analysis of biological, economic, and engineering factors affecting the cost of fish production in recirculating aquaculture systems. *J. World Aquaculture Society, 24*, 193-203.

Loveless, J. E., & Painter, H. A. (1968). The influence of metal ions and pH on the growth of a *Nitrosomonas* strain isolated from activated sludge. *J. Gen. Microbiol., 52*, 1-14.

Lowe-McConnell, R. H. (1987). *Ecological studies in tropical fish communities*. Cambridge: Cambridge University Press.

Loyless, J. C., & Malone, R. F. (1997). A sodium bicarbonate dosing methodology for pH management in freshwater-recirculating aquaculture systems. *The Progressive Fish Culturist, 59*, 198-205.

Manthe, D. P., Malone, R. F., & Kumar, S. (1998). Submerged rock filter evaluation using an oxygen consumption criterion for closed recirculation systems. *Aquacultural Engineering, 7*, 97-111.

Masser, M. P., Rakocy, J., & Losorso, T. M. (1999). Ricirculating Aquaculture Systems, Management of Recirculating Systems (Vol. 452). Mississippi: The Southern Regional Aquaculture Center.

Menon, A. G. K. (1951). Distribution of Clariid Fishes and Its Significance in Zoogeographical Studies. *Proc. Nat. Inst. Sci. India, 17*(4), 291-299.

Metlaf, & Eddy. (1979). *Waste Water Engineering, Treatment Disposal and Reuse, 2nd. ed.* New Delhi: Tata Mc Graw - Hill Publishing Company Ltd.

Mudjiman, A. (1989). *Fish Feed (in Indonesian)*. Jakarta: PT. Penerbit Swadaya.

Muhajir, Ahmad, F., & Edward. (2004). Variasi Kadar Oksigen Terlarut di Perairan Tanimbar Bagian Utara dan Selatan, Maluku Tengara. *Jurnal Sorihi, III*(1-9).

© COPYRIGHT UPM

National Research Council. (1977). *Nutrient Requirements of Warmwater Fishes and Shell Fishes* Washington D.C: National Academy of Sciences.

Nelson, J. S. (1994). *Fishes of the World*. New York: J. Wiley & Sons.

Nightingale, J. W. (1976). Development of Biological Design Criteria for Intensive Culture of Warm and Cool Water Species. *Technical Report of Kramer*. Retrieved from

Noga, E. J. (1996). *Fish Disease Diagnosis and Treatment*. Weisbaden,Germany: Mosby,St.Louis.

Northern Filter Media (2009). Product of Filtermedia. http://www.northernfiltermedia.com/products.html, Acces on November 25, 2010

Okeyo, D. O., & Montgomery, W. L. (1992). Ingestion, Growth and Conversion Efficiency in the Blue Tilapia, *Oreochromis aureus,* When Fed on Three Aquatic Macrophytes. *Journal of the Arizona-Nevada Academy of Science 24/25*, 1-10.

Olson, R. J. (1981). Differential photo inhibition of marine nitrifying bacteria *Journal of Marine Research, 39*(2), 227-238.

Painter, H. A., & Loveless, J. E. (1983). Effect of temperature and pH value on the grow-rate constant of nitrifying bacteria inactivated sludge process. *Water Research, 17*(3), 237-248.

Pescod, M. B. (1973). *Investigation of Rational Effluent and Stream Standars for Tropical Countries*. Bangkok,Thailand: Asean Institute of Technology.

Phonekhampheng, O., Hung, L. T., & Lindberg, J. E. (2008). Nutritive Value of Potential Feed Resources Used in Laos for African Catfish (*Clarias gariepinus*) Production. *Livestock Research for Rural Development, 20*(12).

Prihatman, K. (2000). Catfish *(Clarias)* Culture (Fishery). Retrieved Jun 3, 2009, from Ristek, Section for Development and Empowerment of Knowledge Science and Technology, www.ristek.go.id

Purwakusuma, W. (2002). *Water Quality*. Ornamental Fish Information Highlights, http://o-fish.com/Profil/Profil.php

Redding, T. A., & Middlen, A. B. (1991). Fish Production in Irrigation Canals. A Review. *FAO Fisheries Technical Papaer, 317,* 111.

Roberts, R. J. (1978). *Fish Pathology*. London: Balliere and Tindall.

Robins, R. H. (2009). Walking Catfish. Retrieved June 2009, from Ichthyology, the Florida Museum of Natural History. US. http://www.flmnh.ufl.edu/fish/Gallery/Descript/WalkingCatfish/WalkingCatfish.html, http://www.flmnh.ufl.edu/fish/Gallery/Descript/WalkingCatfish/Walking Catfish.html

© COPYRIGHT UPM

Rogers, G. L., & Klemetson, S. L. (1985). Ammonia Removal in Selected Aquaculture Water Reuse Biofilters. *Aquacultural Engineering, 4*, 135-154.

Rounsefell, G. A., & Everhart, W. H. (1953). *Fishery Science its Methods and Application*. New York,USA: John Wiley and Sons.

Sahoo, S. K., Giri, S. S., Chandra, S., & Mohapatra, B. C. (2008). Evaluation of Breeding Performance of Asian Catfish *Clarias batrachus* at Different dose of HCG and Latency Period Combinations. *Turkish Journal of Fisheries and Aquatic Sciences, 8*, 249-251.

Saidu, M. M.-G. (2009). *Temperature impact on nitrification and bacterial growth kinetics in acclimating recirculating aquaculture systems biofilters.* Doctor of Philosophy, Louisiana State University, Baton Rouge, LA, USA.

Salvetti, R., Azzellinoa, A., Canziania, R., & Bonomoa, L. (2006). Effects of temperature on tertiary nitrification in moving-bed biofilm reactors *Water Research, 40*(15), 2981-2993

SAS Institute Incorporated. (2004). *SAS for Windows (Version 9.1.3) Service Pack 4.* . North Carolina: SAS Institute Incorporated, Cary, NC

Sastrosupadi, A. (2000). *Practice Experiment Design for Agriculture (in Indonesian)* (6 ed.). Yogyakarta: Kanisius.

Sastry, B. (1996). *A Comparison of Nitrification Capacity in Bead and Tubular Plastic Media.* MS Thesis, Louisiana State University, Baton Rouge, LA.

Satar, M. K. A. (1984). *Fundamentals of Aquaculture (in Malay)*. Serdang, Malaysia: Fakulti Perikanan dan Sains Samudera, Universiti Pertanian Malaysia.

Shammas, N. K. (1986). Interactions o temperature, pH and biomass on the nitrification pocess. *J. Water Pollut. Control Fed., 58*, 52-59.

Shell, E. W. (1987). *Relationship between Rate of Feeding, Rate of Growth and Rate of Conversion on Feeding Trials with Two Species of Tilapia, Tilapia mosasambica (Peters) and Tilapia nilotica (Linneaus).* Paper presented at the The World Symposium on Warm-Water Pond Fish Culture, Rome, Italy.

Shoemaker, C. A., Olivares-Fuster, O., Arias, C. R., & Klesius, P. H. (2008). Flavobacterium columnare genomovar influences mortality in channel catfish *(Ictalurus punctatus)*. *Veterinary Microbiology, 127*, 353-359.

Singh, B. N., & Hughes, G. M. (1971). Respiration of An Air-Breathing Catfish *Clarias batrachus* (Linn.). *J. Exp. Biology, 55*, 421-434.

SITH-ITB, VEDCA, & SEAMOLEC. (2009). *Teknologi Pengelolaan Kualitas Air*. Kualitas biologis dan manipulasi mikroba: bakteri nitrifikasi. Program Alih Jenjang D4 Bidang Akuakultur, Institut Teknologi Bandung (ITB). Bandung, Indonesia.

© COPYRIGHT UPM

Sitio, S. (2008). *Pengaruh Medan Listrik Pada Media Pemeliharaan Bersalinitas 3 ppt Terhadap Tingkat Kelangsungan Hidup dan Pertumbuhan Ikan Gurame Osphronemus gouramy.* Undergraduate, Institut Pertanian Bogor, Bogor.

Smith, H. M. (1934). Contributions to the ichthyology of Siam. *Reprinted from the Jour. of the Siam Soc. Natural History Supp, 9*(8), 287-292.

Smith, L. S. (1980). Digestion and Teleost Fishes. *Fish Feed Technology,* 3-18.

Speece, R.E. (1973), Trout metabolism characteristics and the rational design of nitrification facilities for water reuse in hatcheries. *Trans. Am. Fish. Soc. 2* , 323–334

Steel, R. G. D., & Torie, J. H. (1980). *Principle and Procedure of Statistics*. New York: Mc Graw Hill Inc.

Stickney, R. R. (1979). *Principles of Warmwater Aquaculture*. New York: Jhon Willy and Sons.

Suantika, G. (2001). *Development of a Recirculation System for the Mass Culturing of the Rotifer Brachionus plicatilis* PhD, Universiteit Gent, Belhium Ghent.

Sudarto. (2009). *Systematic Revision and Phylogenetic Relationships among Populations of Clariid Species in Southheast Asia.* Post Graduate Program Summary of Thesis, University of Indonesia, Jakarta, http://www.id.ird.fr/us/ actualities/Sudarto/.

Sudarto, Teugels, G. G., & Pouyaud, L. (2004). Description of A New Clariid Catfish, *Clarias pseudonieuhofii* from West Borneo (Siluriformes:Clariidae). *Zoological Studies, 43*(1), 8-19.

Sudrajat, Y., & Gunawan, B. (2002). Sistem bakteriofiltrasi sebagai sarana pasokan air pada penampungan ikan hidup. *Buletin Teknik Pertanian, 7*(2), 48-50.

Summerfelt, S. T., Davidson, J. W., Waldrop, T. B., Tsukuda, S. M., Bebak-Williams, J., (2004). A partial-reuse system for coldwater aquaculture. *Aquacult. Eng. 31*, 157-181.

Sunarma, A. (2004). *Increase the Business Productivity of Sangkuriang Catfish (Clarias sp.)*. Paper presented at the Conference of Technical Unit and Implementing Business Bandung.

Suresh, A. V., & Lin, C. K. (1992). Effect of Stocking Density on Water Quality and Production of Red Tilapia in Recirculated Water System. *Aquacultural Engineering, 11*, 1-22.

Sutisna, D. H., & Sutarmanto, R. (1995). *Freshwater Fish Hatchery (in Indonesian)*. Yogyakarta,Indonesia: Penerbit Kanisius.

Swann, L. (1992). A Basic Overview of Aquaculture; History, Water Quality, Type of Aquaculture, Production Methods. *Technical Bulletin Series # 102,* 10 p.

© COPYRIGHT UPM

Swingle, H. S. (1963). *Method of Analysis for Water, Organic Matter and Pond Bottom Soils Used in Fisheries Research*. Alabama: Auburn University.

Talbot, C. (1993). *Some Aspects of the Biology of Feeding and Growth in Fish.* Paper presented at the Proceedings of the Nutrition Society, 52, Symposium on 'Fish and Nutrition', A meeting of the Scottish Group of the Nutrition Society, the University of Aberdeen.

Tchobanaglous, G., & Burton, F. (2003). *Wastewater Engineering Treatment, Disposal and Reuse, 4th ed.* (4 ed.). New York, USA: McGraw-Hill.

Tetzlaff and Heidinger, (1990), Basic Principles of Biofiltration and System Design, Southern Illinois University Cooperative (SIUC), *Fisheries Bulletin* No. 9.

Teugels, G. G. (1996). Taxonomy, Phylogeny and Biogeography of Catfishes (Ostariophysi; Siluroidei) : An Overview. *Aquat. Living Resour, 9*, 9-34.

Teugels, G. G., Sudarto, & L, P. (2001). Description of A New *Clarias* Species from Southeast Asia Based on Morphological and Genetical Evidence *(Siluriformes, Clariidae)*. *Cybium, 25*(1), 81-92.

Timmons, M.B, Ebeling, J.M, Wheaton, F.W, Summerfelt, S.T and Vinci, B.J. (2002). Recirculating Aquaculture Systems. 2nd Northeastern Regional Aquaculture Center Publication No. 01-02.

Tucker, C. S. (1991). Water Quantity and Quality Requirements for Channel Catfish Hatcheries. from Southern Regional Aquaculture Center (SRAC)

Twarowska *et. al*. (1997). *Waste Treatment and Waste Characterization Evaluation of an Intensive Recirculating Fish Production System*. North Carolina: Department of Biological and Agricultural Engineering, North Carolina State University.

Unicomposite (2010), *FRP and Pultrusion*, http://www.unicomposite.com/literature_catalog/what%20is%20pultrusion.pdf, Acces on March 20, 2010

US EPA. (1976). *Quality Criteria for Water*. Washington DC: US EPA.

van Rijn, J., & Rivera, G. (1990). Aerobic and Anaerobic Biofiltration in an Aquaculture Unit: Nitrite Accumulation as a Result of Nitrification and Denitrification. *Aquacultural Engineering, 9*, 1-18.

van Rijn, J., & Sich, H. (1992). Nitrite Accumulation by Denitrifying Bacteria Isolated from Fuidized Bed Reactors Operated in an Aquaculture Unit. In B. Moav, V. Hilge & H. Rosenthal (Eds.), *Progress in Aquaculture Research. Special Publication no. 17.* (pp. 39–54). Bredene, Belgium: European Aquaculture Society.

Viveen, W. J. A. R., Richter, C. J. J., van Oordt, P. G. W. J., Janssen, J. A. L., & Huisman, E. A. (1985). *Practical Manual for The Culture of The African Catfish (Clarias gariepinus)* (S. R. Suyanto, Trans.). EB The Hague, The

© COPYRIGHT UPM

Netherlands: The Netherlands Ministry for Development Cooperation, Section for Research and Technology, P.O. Box 20061, 2500.

Watanabe, T. (1988). *Fish Nutrition and Mariculture*. Tokyo: Departement of Aquatic Biosciences. Tokyo University of Fisheries.

Watanabe, T. (2002). Effect of dietary protein levels and feeding period before spawning on chemical components of eggs produced by red sea bream broodstock. *Bull. Jpn. Soc. Sci. Fish, 51*(9), 1501-1509.

Watson, S.W., F.W. Valos, and J.B. Waterbury, 1981. The Family Nitrobacteraceae. In *The Prokaryotes*. Edited by M.P. Starr et al. Berlin: Springer-Verlag.

Weber, M., & de Beaufort, L. F. (1913). *The Fishes of the Indo-Australian Archipelago II. Malacopterygii, Myctophoidea, Ostariophysi: I. Siluroidea*. Leiden: E. Brill, Ltd.

Wedemeyer, G. A. (1996). *Physiology of Fish in Intensive Culture Systems*. New York: Chapman & Hall.

Welch, P. S. (1952). *Limnology*. New York,USA: McGraw-Hill Book Co. Inc.

Wellborn, T. L. (1988). Channel Catfish, Life History and Biology. *SRAC Publication, 180,* 4.

Wheaton, F. (2010). *Recirculating Aquaculture Systems: An Overview of Waste Management*, Department of Biological Resources Engineering, University of Maryland College Park, MD 20742

Wilderer, P. A., Jones, W. L., & Dau, U. (1987). Competition in Denitrification Systems Affecting Reduction Rate and Accumulation of Nitrite. *Water Research 21*(2), 239-245

Wimberly, D. M. (1990). *Development and Evaluation of a Low-Density Media Biofiltration Unit for Use in Recirculating Finfish Culture Systems.* MS Thesis, Louisiana State University, Baton Rouge, LA.

Yacob, N. (2009). Bio-filter in Recirculating Aquaculture System (RAS) (in Malay). *Keli Omega 3, Education And Knowledge About Fish Feed Technology, http://naim-keliomega3.blogspot.com/2009/03/biofilter-di-dalam-recirculating. html, Access on June 29*.

Yudha, I. G. (2005) *Seminar on World Water Day*. Lampung, Indonesia: Universitas Negeri Lampung (UNILA).

Yunus, M., Priyadi, A., & Wahid, A. (2000). Effect of Water Depth in the Rained Pond on the Survival and Growth of Local Catfish (*Clarias batrachus*) (in Indonesian). 98-105.

© COPYRIGHT UPM

Zhu, S., & Chen, S. (1999). An experimental study on nitrification biofilm performances using a series reactor system. *Aquacultural Engineering, 20*, 245-259.

Zhu, S., & Chen, S. (2002). Impact of temperature on nitrification rate in fixed film biofilters. *Aquacultural Engineering, 26*, 221-237.

Zhu, S., Saucier, B., Durfey, J., Chen, S., & Dewey, B. (1999). Waste excretion characteristics of Manila clams *(Tapes philippinarum)* under different temperature conditions. *Aquacultural Engineering, 20*(4), 231-244.

Zonneveld, N., Huisman, E. A., & Boon, J. H. (1991). *Prinsip Prinsip Budidaya Ikan*. Jakarta, Indonesia: PT.Gramedia Pustaka Utama.

Zweig, R. D., Morton, J. D., & Stewart, M. M. (1999). Source Water Quality for Aquaculture, A Guide for Assessment. In R. D. Environmentally and Socially Sustainable Development (Ed.). Washington D.C, USA: The International Bank for Reconstruction and Development, The World Bank.

© COPYRIGHT UPM