



**UNIVERSITI PUTRA MALAYSIA**

***PARENTERAL FORMULATION OF NANOEMULSION LOADED WITH  
CHLORAMPHENICOL FOR MENINGITIS TREATMENT***

**SITI HAJAR MUSA**

**FS 2014 5**



**PARENTERAL FORMULATION OF NANOEMULSION LOADED WITH  
CHLORAMPHENICOL FOR MENINGITIS TREATMENT**

By

**SITI HAJAR MUSA**

**Thesis submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirement for the Degree of Master of  
Science**

**August 2014**

## COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

## **PARENTERAL FORMULATION OF NANOEMULSION LOADED WITH CHLORAMPHENICOL FOR MENINGITIS TREATMENT**

By

**SITI HAJAR MUSA**

**August 2014**

**Chair : Professor Mahiran Basri, PhD**  
**Faculty : Science**

Meningitis is a disease caused by the inflammation of the brain due to bacterial infection. The most common causative agent in bacterial meningitis is *Streptococcus pneumoniae*. The bacterial infection occurs in the brain which is protected by a barrier known as blood-brain barrier (BBB). This barrier consists of a very selective membrane which only allows small particle with lipophilic characteristic to pass through. In the current medication treatment, third generation cephalosporin has been used widely in treating this disease. However, this family of drug (cephalosporin) is no longer effective towards the meningitis bacteria due to the drug resistance problem. Traditional antibiotic such as chloramphenicol has caught medical community's attention back to overcome the problem. Delivery of chloramphenicol could be improved via nanoemulsion system with nano-size droplet range.

From the solubility study, it was found that palm kernel oil esters (PKOEs) and safflower seed oil (1:1 ratio) was the best combination in solubilizing chloramphenicol compared to other oil mixture. Lecithin was best mixed with Tween 80 as co-surfactant in the emulsion system, resulting in smaller particle size (131.5 nm) and low polydispersity index (0.135) reading. Methods of emulsification such as spontaneous method (stirring process), spontaneous with high shear homogenizer, spontaneous with ultrasound homogenizer and spontaneous with high pressure homogenizer were studied. With respect to the particle size and polydispersity index, combination of spontaneous method with high pressure homogenizer gave the best particle size and polydispersity index which was 70.65 nm and 0.154, respectively.

Response Surface Methodology (RSM) was used as a tool in optimization process. Relation between independent variables (oil, lecithin and glycerol amount) and response variables (particle size, zeta potential and osmolality)

were analysed. From three dimension (3D) analysis models, it was observed that particle size, zeta potential and osmolality value of nanoemulsion was largely influenced by the amount of oil, lecithin and glycerol, respectively. RSM has suggested composition with 4% oil, 2.50% lecithin and 2.25% glycerol which resulted with nanoemulsion system with 95.33 nm of particle size, 0.238 of polydispersity index, -36.91 mV of zeta potential and 200 mOsm/kg of osmolality. These actual values show good agreement with the predicted values; 86.96 nm (particle size), 0.234 (polydispersity index), -33.47 mV (zeta potential), 198 mOsm/kg (osmolality).

Dosage optimization showed that stable nano-antibiotic emulsion system was successfully achieved up to 3 mg/ml of chloramphenicol. Three formulations with varied loading dosage (1, 2 and 3 mg/ml) were found to be stable at 4°C. Transmission Electron Microscopy (TEM) of negatively stained nanoemulsion showed that the oil droplets were in a spherical shape and uniform in size. Toxicity analysis shows that nanoemulsion entrapment has successfully reduced the toxicity effect of chloramphenicol towards 3T3 cells. *In-vitro* study showed a good penetration of chloramphenicol through the cellulose acetate membrane which resulted a zero order release mechanism.

Stability evaluation based on the particle size for three months showed that the particle size remain in a nano-size with less than 5% of size changes. Emulsion entrapment study shows no chloramphenicol leakage from the oil droplet. Drug assay through 3 months of storage showed a consistent drug loading in the emulsion with less than 10% of drug lost. In conclusion, the nanoemulsion formulation is a promising vehicle for the delivery of chloramphenicol parenterally (intravenous route).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

## **FORMULASI PARENTERAL BAGI NANOEMULSI YANG MENGANDUNGI CHLORAMPHENICOL UNTUK PENYAKIT RADANG LAPISAN OTAK**

Oleh

**SITI HAJAR MUSA**

**Ogos 2014**

**Pengerusi : Profesor Mahiran Basri, PhD**  
**Fakulti : Sains**

Radang lapisan otak adalah penyakit yang disebabkan oleh jangkitan bakteria di dalam otak. Ejen utama penyebab radang lapisan otak ialah disebabkan oleh jangkitan bakteria *Streptococcus pneumonia*. Jangkitan bakteria terhadap otak berlaku di bahagian yang dilindungi oleh halangan darah-otak (BBB). Halangan ini terdiri daripada membran yang sangat peka dan hanya akan membenarkan zarah lipofilik yang sangat kecil melepaskannya. Kaedah perubatan terkini banyak menggunakan *cephalosporin* generasi ketiga bagi mengubati penyakit ini. Walau bagaimanapun, kumpulan dadah ini tidak lagi efektif terhadap bakteria radang lapisan otak disebabkan oleh masalah rintangan dadah. Perubatan tradisional seperti kloramfenikol telah mendapat tumpuan semula dari komuniti perubatan bagi mengatasi masalah ini. Penghantaran kloramfenikol masih boleh ditingkatkan melalui sistem nanoemulsi yang mempunyai titisan kecil bersaiz-nano.

Daripada kajian kelarutan, didapati ester minyak kernel kelapa sawit (PKOEs) dan minyak biji bunga kesumba (nisbah 1:1) adalah kombinasi yang terbaik bagi melarutkan kloramfenikol berbanding campuran minyak yang lain. Kombinasi lesitin adalah baik bersama dengan Tween 80 sebagai ko-surfaktan di dalam sistem emulsi yang menghasilkan saiz zarah kecil (131.5 nm) dan bacaan indeks polisebaran yang rendah (0.135). Kaedah emulsi seperti kaedah secara spontan (proses pengacauan), kaedah spontan berserta pengacau mericah tinggi, kaedah spontan berserta pengacau ultra-bunyi dan kaedah spontan berserta pengacau bertekanan tinggi telah dikaji. Kombinasi kaedah spontan berserta pengacauan bertekanan tinggi menghasilkan saiz zarah dan indeks polisebaran yang terbaik iaitu masing-masing dengan nilai 70.65 nm dan 0,154.

Kaedah Gerakbalas Permukaan (RSM) digunakan sebagai alat bagi proses pengoptimuman. Hubungan antara pembolehubah bebas (jumlah minyak, lesitin dan gliserol) dengan pembolehubah tindak balas (saiz zarah, keupayaan zeta dan osmolaliti) dikaji. Berdasarkan kepada kajian model tiga dimensi (3D), didapati nilai bacaan saiz zarah, keupayaan zeta dan osmolaliti masing masing adalah sangat besar dipengaruhi oleh jumlah minyak, jumlah lesitin dan jumlah gliserol. RSM mencadangkan komposisi 4% minyak, 2.5% lesitin dan 2.25% gliserol yang menghasilkan nanoemulsi bersaiz zarah 95.33 nm, bacaan indeks polisebaran 0.238, bacaan keupayaan zeta -36.91 mV dan bacaan osmolaliti 200 mOsm/kg. Bacaan sebenar bagi semua nilai ini adalah hampir bersamaan dengan nilai jangkakan; 86.96 nm (saiz zarah), 0.234 (indeks polisebaran), -33.47 mV (keupayaan zeta), 198 mOsm/kg (osmolaliti).

Pengoptimuman dos kloramfenikol menunjukkan sistem emulsi antibiotik-nano berjaya dicapai sehingga 3 mg/mL. Tiga formulasi yang berbeza mengikut dos muatan (1, 2 dan 3 mg/mL) didapati stabil pada suhu 4°C. Penghantaran Elektron Mikroskopi (TEM) nanoemulsi berwarna negatif menunjukkan titik kecil minyak berbentuk sfera dan bersaiz seragam. Analisis toksik menunjukkan perangkap nanoemulsi telah berjaya mengurangkan kesan ketoksikan kloramfenikol terhadap sel 3T3. Kajian secara *in-vitro* menunjukkan penembusan kloramfenikol yang baik melalui membran selulosa acetat yang menghasilkan mekanisme pelepasan tahap sifar.

Penilaian terhadap kestabilan berdasarkan saiz zarah selama tiga bulan menunjukkan saiz zarah masih kekal berada di dalam lingkungan saiz-nano dengan kurang daripada 5% perubahan. Kajian perangkap emulsi menunjukkan tiada kebocoran kloramfenikol daripada butiran minyak. Pemerhatian terhadap muatan dadah sepanjang penyimpanan selama 3 bulan menunjukkan muatan dadah adalah konsisten dengan kehilangan kurang daripada 10%. Kesimpulannya, formulasi nanoemulsi menjanjikan penghantaran kloramfenikol secara parenteral (laluhan intravena) yang baik.

## ACKNOWLEDGEMENTS

First and foremost, I would like to praise Allah S.W.T for the strength and persistence He has given to me throughout my journey in completing this research. The bittersweet journey I have gone through has finally comes to an end with the 'birth' of this thesis.

My deepest gratitude to my excellent supervisor, Prof. Dr. Mahiran Basri for her guidance and endless advice to me throughout the years. No words could describe how thankful am I for being given the opportunity to be a student under her supervision since my undergraduate study. She is and always will be my idol for being a better researcher and a good leader in the future, inshaAllah. I would also like to express my gratitude to all my co-supervisors, Dr. Emilia Abd Malek, Dr. Roghayeh Abedi Karjiban and to Dr. Fuad Shamsuddin for their help in giving me useful opinions and suggestions to improve my research quality and also for all the precious time they spent for my benefit. To Dr. Hamid Reza Fard Masoumi, thank you so much for guiding me on the optimization studies. We finally managed to produce a good publication.

To all members of the EMTECH group, especially to my lab mates in lab 105 and 401, thanks a lot guys. Syarilaida, thank you for lending me your ears, your support, your time and for always being there for me. Woo Fong Yen, thank you for being the best partner in crime I have ever had. To Syafinaz, Siti Norhawani, Ghaidaa, Shazwani, Hasmah, Azren Aida and all lab members, thank you so much for welcoming me to be part of our research group. Thanks for all the sweet memories we have created together. Maybe one day we will all be great researchers and have the opportunity to work together again. Not forgetting, to all UPM staff who helped me in collecting analysis data, thank you.

Special thanks with love for my family, who are my backbone throughout my good and bad times. Ayah, thank you for lending me your shoulders every time I had almost given up. Thanks for always believing in my passion towards chemistry. This is for you. Mak, thank you for never stop supporting me and give me the strength to go through all the challenges I have faced. Along, Angah, Aqim and Adik, I love you and of course, you know that. Wani, Kamal, Izzati and Rayyan, thank you for the non-stop encouragement you guys have shown me throughout this study. Without all of you, I would not have made it this far.



Last but not least, thank you to My Brain 15 (Master), Graduate Research Fellowship (GRF) and Universiti Putra Malaysia for the financial support and facilities.

Thank you.



I certify that a Thesis Examination Committee has met on 29 August 2014 to conduct the final examination of Siti Hajar binti Musa on her thesis entitled "Parenteral Formulation of Nanoemulsion Loaded with Chloramphenicol for Meningitis Treatment" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Mansor bin Hj Ahmad @ Ayob, PhD**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Mohd Zaizi bin Desa, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Nor Azowa binti Ibrahim, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Shahidan Radiman, PhD**

Professor  
Universiti Kebangsaan Malaysia  
Malaysia  
(External Examiner)



---

**NORITAH OMAR, PhD**

Associate Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 19 September 2014

The thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements of the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Mahiran Basri, PhD**

Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Emilia Abd Malek, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Roghayeh Abedi Karjiban, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Ahmad Fuad Shamsuddin, PhD**

Associate Professor  
Faculty of Pharmacy  
Universiti Kebangsaan Malaysia  
(Member)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## DECLARATION

### Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Siti Hajar binti Musa (GS 30331)

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of  
Chairman of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vii
<b>DECLARATION</b>	ix
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xv
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>LIST OF UNITS</b>	xviii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
1.1 Background of Research	1
1.2 Problem statements	1
1.3 Scope of study	2
1.4 Objectives	2
<b>2 LITERATURE REVIEW</b>	<b>3</b>
2.1 Meningitis	3
2.2 Blood-Brain Barrier	3
2.2.1 Structure of Blood-Brain Barrier	4
2.2.2 Function of Blood-Brain Barrier	5
2.3 Antibiotic drug	5
2.3.1 Current antibiotic drugs used in meningitis treatment	6
2.3.2 Chloramphenicol	7
2.4 Intravenous Delivery	8
2.5 Drug-delivery system	9
2.5.1 Emulsion	9
2.5.2 Characteristics of parenteral emulsion	12
2.5.3 Nano-carrier systems	13
2.6 Composition of emulsion	14
2.6.1 Surfactant	14
2.6.2 Palm kernel oil esters	16
2.7 Technique of emulsification	17
2.7.1 Spontaneous emulsification	17
2.7.2 High pressure homogenization	18
2.7.3 Ultrasound homogenization	19
2.7.4 High shear homogenization	20
2.8 Technique of optimization	20
<b>3 MATERIALS AND METHODS</b>	<b>22</b>
3.1 Materials	22
3.2 Selection of nanoemulsion compositions	22
3.2.1 Selection of oil mixture	22
3.2.2 Selection of surfactant mixture	23
3.3 Study on the emulsification technique	23

3.3.1	Selection of emulsification techniques	23
3.3.2	Selection of processing cycles	24
3.4	Optimization of nanoemulsion composition	24
3.5	Modification of nanoemulsion formulation	25
3.5.1	Osmolality alteration	25
3.5.2	Dosage optimization	25
3.5.3	Formulation of nanoemulsion	25
3.6	Characterization of nanoemulsion	26
3.6.1	Particle size, polydispersity index and zeta potential measurement	26
3.6.2	Osmolality measurement	26
3.6.3	pH measurement	26
3.6.4	Viscosity measurement	26
3.6.5	Transmission Electron Microscopy study	27
3.6.6	Toxicity study	27
3.6.7	<i>In-vitro</i> study of chloramphenicol-loaded nanoemulsion release	27
3.6.8	Kinetic study on the chloramphenicol-loaded nanoemulsion release	28
3.7	Stability evaluation of nanoemulsion	28
3.7.1	Nanoemulsion encapsulation efficiency study	29
3.7.2	Chloramphenicol assay study	29
<b>4</b>	<b>RESULTS AND DISCUSSIONS</b>	<b>30</b>
4.1	Selection of nanoemulsion compositions	30
4.1.1	Oil mixture analysis	30
4.1.2	Surfactant mixture analysis	32
4.2	Study on the emulsification technique	33
4.2.1	Emulsification technique analysis	33
4.2.2	Processing cycles analysis	35
4.3	Optimization of nanoemulsion composition	36
4.3.1	Selection of independent variables	36
4.3.2	Fitting the response surface models	37
4.3.3	Response surface analysis	39
4.3.4	Optimized condition using RSM	43
4.4	Modification of nanoemulsion formulation	44
4.4.1	Osmolality value alteration	44
4.4.2	Dosage optimization analysis	44
4.5	Characterization of nanoemulsion	46
4.5.1	Nanoemulsion properties	46
4.5.2	Transmission Electron Microscopy analysis	48
4.5.3	Toxicity analysis	48
4.5.4	<i>In-vitro</i> analysis of chloramphenicol-loaded nanoemulsion release	50
4.5.5	Kinetic analysis of chloramphenicol-loaded nanoemulsion release	51
4.6	Stability evaluation of nanoemulsion	52
4.6.1	Particle size analysis	53
4.6.2	pH analysis	54
4.6.3	Osmolality analysis	55

4.6.4	Nanoemulsion encapsulation efficiency analysis	56
4.6.5	Chloramphenicol assay analysis	57
<b>5</b>	<b>CONCLUSIONS AND RECOMMENDATIONS</b>	<b>58</b>
5.1	Conclusions	58
5.2	Recommendation for further studies	59
	<b>REFERENCES</b>	<b>61</b>
	<b>APPENDICES</b>	<b>74</b>
	<b>BIODATA OF STUDENT</b>	<b>76</b>
	<b>LIST OF PUBLICATION/ PROCEEDINGS</b>	<b>77</b>





## LIST OF TABLES

Table		Page
1	List of commonly used surfactant for parenteral emulsion and recommended concentration	16
2	Physical properties of palm kernel oil esters (PKOEs)	17
3	Coded level for independent variables used in experimental design for nanoemulsion optimization	24
4	The matrix of central composite design (CCD)	37
5	Analysis of variance (ANOVA) for the model	38
6	Individual main effect of independent variables ( $x_i$ ) on response variables ( $Y_i$ )	39
7	Significant ( $P < 0.05$ ) interaction between independent variables ( $x_i$ ) on response variables ( $Y_i$ )	39
8	The predicted and actual values of chloramphenicol-loaded nanoemulsion	43
9	Modification of formulation	45
10	Physicochemical properties of chloramphenicol-loaded nanoemulsion	47
11	Entrapment efficiency study over 3 m storage (F1)	57

## LIST OF FIGURES

Figure		Page
1	Structure of blood-brain barrier	5
2	Nanoemulsion containing antibiotic drug is aimed to penetrate blood-brain barrier (BBB) with the nano-size range of the oil droplets	6
3	Chloramphenicol molecular structure	7
4	Intravenous route	8
5	Illustration of emulsion system	9
6	Instability phenomena of emulsion system	10
7	Picture of nanoemulsion and macroemulsion	11
8	Picture of micelles	15
9	Spontaneous emulsification method	18
10	Illustration of high pressure homogenizer	19
11	Illustration of ultrasound homogenizer	19
12	Illustration of high shear homogenizer	20
13	Solubility study of Chloramphenicol in oil mixture of PKOEs with sesame oil, safflower seed oil, sunflower oil, soybean oil and pine nut oil at ratio 1:1	30
14	The solubility of chloramphenicol in various oil mixtures	32
15	Effect of lecithin with co-surfactant mixture on the particle size and PDI value	33
16	Effect of different types of emulsification techniques on the particle size and PDI value	35
17	Effect of high pressure homogenizer processing cycles on the particle size value	36
18	Significant ( $P < 0.05$ ) interaction effect of lecithin and oil on the particle size ( $Y_1$ ) values	40

19	Significant ( $P < 0.05$ ) interaction effect of lecithin and oil on the zeta potential ( $Y_2$ ) values	41
20	Significant ( $P < 0.05$ ) interaction effect of lecithin and oil on the osmolality ( $Y_3$ ) values	42
21	Phase behaviour of chloramphenicol-loaded nanoemulsion after 3 months of storage at 4 °C	45
22	TEM micrograph of particle size in chloramphenicol-loaded nanoemulsion (F1)	48
23	Effect of chloramphenicol concentration on the percentage of cells viability after 48 h of exposure	49
24	The permeation profile of chloramphenicol-loaded nanoemulsion through cellulose membrane	51
25	Mechanism of chloramphenicol kinetic release for zero order, first order and Higuchi model	52
26	Effect of storage on the particles size value of nanoemulsion	53
27	Effect of storage on the pH value of nanoemulsion	54
28	Effect of storage on the osmolality value of nanoemulsion (F1)	55
29	HPLC-ELSD chromatogram of chloramphenicol in nanoemulsion and aqueous phase of nanoemulsion	56

## LIST OF ABBREVIATIONS

BBB	Blood-brain barrier
CNS	Central nervous system
IV	Intravenous
W/O	Water-in-oil
O/W	Oil-in-water
PDI	Polydispersity index
PKOEs	Palm kernel oil esters
PIT	Phase inversion temperature
PIC	Phase inversion composition
HLB	Hydrophilic-lipophilic balance
RSM	Response surface methodology
CCD	Central composite design
PBS	Phosphate buffer solution
UPLC	Ultra Performance Liquid Chromatography
HPLC	High Performance Liquid Chromatography
ELSD	Evaporative Light Scattering Detection
TEM	Transmission Electron Microscopy

## LIST OF UNITS

nm	Nanometer
mV	Milivolt
mOsm/kg	Miliosmole per kilogram
cP	Centipoise
°C	Degree celcius
%	Percentage
w/w	Weight per weight
mg	Milligram
µg	Microgram
mL	Mililitre
µL	Microlitre
mm	Milimetre
µm	Micrometre
cm	Centimetre
Da	Dalton
rpm	Rotation per minute
m	Month
min	Minute
h	Hour

## CHAPTER 1

### INTRODUCTION

#### 1.1 Background of Research

Human brain is one of the most important organ that play its main role in controlling the whole body system's activity. The brain has more protection compared to other human's organ as a way to make sure that the body systems can function. The feature that encapsulates human's brain is called blood-brain barrier (BBB) which protects the brain from any harmful compounds (Sandoval and Witt, 2008). BBB is recognized as a gate keeper of the brain, an efflux pump, limiting the entrance of many structurally divergent lipophilic molecules, such as peptides and many of the drugs used in psychiatry and neurology (De Klerk *et al.*, 2010). BBB acts as a separator between brain and its blood supply. Thus, this barrier limits many potential therapeutic to reach the central nervous system (Scherrmann, 2002).

Even though BBB is a very selective membrane, but it is still possible for viruses or bacteria to be spread through blood and get into the cerebrospinal fluid, hence causing infection to the brain. One of disease that is caused by the infection of the brain is meningitis. Meningitis is an inflammation of brain caused by either viral meningitis (virus) or bacterial meningitis (bacteria) (Kleine *et al.*, 2003). Statistically, meningitis occurs in 30% of new born and very young infants and 15% to 20% of older children. Between 30%-50% of meningitis survivors will suffer from permanent neurological sequelae (Fitch and Van De Beek, 2007). Despite the availability of effective antibiotic treatment, high mortality rate up to 30% has been observed (Mishal *et al.*, 2008; Weisfelt *et al.*, 2006). Individual that is infected by the meningitis bacteria could also suffer brain damage. Bacterial meningitis also has been recognized as one of the main causes that lead to deafness, seizure disorder and behavioral problems in children (Bonadio, 1997).

#### 1.2 Problem statements

Chloramphenicol is an effective drug towards bacterial infection and is used for meningitis treatment since 1975 (Lewis *et al.*, 1998). However, the use of chloramphenicol has been increasing in the dosage administered in recent years due to the increase incidence in antibiotic resistance. In addition, chloramphenicol is a hydrophobic drug which hardly dissolves in water. Clinically, this kind of hydrophobic drug is used in a salt form as chloramphenicol succinate to be dissolved in water. Due to the less efficient carrier, larger dosage of chloramphenicol is required per injection to ensure it could reach the target cell. Therefore, higher dosages give higher toxicity problem towards cells and also worsen the side effects.

### 1.3 Scope of study

Nanoemulsion loaded with chloramphenicol was formulated using a mixture of oils and surfactants. Screening of these two compositions and selection on the method of emulsification used were carried out. Optimization was carried out by using Response Surface Methodology (RSM) as a tool with respect to the nanoemulsion's composition. The suggested optimized composition was then been modified due to its stability and osmolality value problems. Characterization was carried out by using Transmission Electron Microscopy (TEM), *in-vitro* of chloramphenicol release and the mechanism of kinetic study. Evaluation on the stability of chloramphenicol-loaded nanoemulsion was carried out after 3 months of storage at 4 °C. Stability evaluation was done with respect to the particle size, pH of nanoemulsion, osmolality value, encapsulation efficiency of the droplets and drug assay analysis.

### 1.4 Objectives

The main objective of this study is to develop parenteral nanoemulsion loaded with chloramphenicol for meningitis treatment. In order to successfully achieve the main objective, following specific objectives were carried out:

- i) To formulate and optimize the compositions of chloramphenicol-loaded nanoemulsion suitable for parenteral applications
- ii) To characterize the physicochemical properties of the nanoemulsion.
- iii) To evaluate chloramphenicol-loaded nanoemulsion release potential using *in-vitro* approach
- iv) To evaluate the stability of nanoemulsion.

## REFERENCES

- Ahmed, M., Ramadan, W., Rambhu, D. and Shakeel, F. (2008). Potential of nanoemulsions for intravenous delivery of rifampicin. *Pharmazie*, 63, 806–811.
- Ahn, J.-H., Kim, Y.-P., Lee, Y.-M., Seo, E.-M., Lee, K.-W. and Kim, H.-K. (2008). Optimization of microencapsulation of seed oil by response surface methodology. *Food Chemistry*, 107, 98-105.
- Alam, M. I., Beg, S., Samad, A., Baboota, S., Kohli, K., Ali, J. and Akbar, M. (2010). Strategy for effective brain drug delivery. *European Journal of Pharmaceutical Sciences*, 40(5), 385–403.
- Anarjan, N., Mirhosseini, H., Baharin, B.S. and Tan, C.P. (2010). Effect of processing conditions on physicochemical properties of astaxanthin nanodispersions. *Food Chemistry*, 123, 477–483.
- Anton, N., Benoit, J.-P. and Saulnier, P. (2008). Design and production of nanoparticles formulated from nano-emulsion templates-a review. *Journal of Controlled Release*, 128(3), 185–199.
- Anton, N. and Vandamme, T. F. (2009). The universality of low-energy nano-emulsification. *International Journal of Pharmaceutics*, 377(1-2), 142–7.
- Araújo, F. A., Kelmann, R. G., Araújo, B. V, Finatto, R. B., Teixeira, H. F. and Koester, L. S. (2011). Development and characterization of parenteral nanoemulsions containing thalidomide. *European Journal of Pharmaceutical Sciences*, 42, 238–245.
- Awad, T. and Sato, K. (2002). Acceleration of crystallisation of palm kernel oil in oil-in-water emulsion by hydrophobic emulsifier additives. *Colloids and Surfaces B: Biointerfaces*, 25, 45–53.
- Ballabh, P., Braun, A. and Nedergaard, M. (2004). The blood-brain barrier: an overview. Structure, regulation, and clinical implications. *Neurobiology of Disease*, 16(1), 1–13.
- Banks, W. A. (2011). Drug delivery to the brain in Alzheimer's disease: Consideration of the blood-brain barrier. *Advanced Drug Delivery Reviews*, 64(7), 629–639.
- Banks, W. A. and Erickson, M. A. (2010). The blood-brain barrier and immune function and dysfunction. *Neurobiology of Disease*, 37(1), 26–32.
- Bartels, A. L., Van Berckel, B. N. M., Lubberink, M., Luurtsema, G., Lammertsma, A. A. and Leenders, K. L. (2008). Blood-brain barrier P-



glycoprotein function is not impaired in early Parkinson's disease. *Parkinsonism Related Disorders*, 14(6), 505–508.

Bas, D. and Boyac, I. H. (2007). Modelling and optimization 1: usability of response surface methodology. *Journal of Food Engineering*, 78, 836–845.

Bayraktar, E. (2001). Response surface optimization of the separation of DL-tryptophan using an emulsion liquid membrane. *Process Biochemistry*, 37, 169–175.

Bellac, C. L., Coimbra, R. S., Christen, S. and Leib, S. L. (2006). Pneumococcal meningitis causes accumulation of neurotoxic kynurenine metabolites in brain regions prone to injury. *Neurobiology of Disease*, 24(2), 395–402.

Benita, S. and Levy, M. Y. (1993). Submicron emulsions as colloidal drug carriers for intravenous administration: comprehensive physicochemical characterization. *Journal of Pharmaceutical Sciences*, 82, 1069–1079.

Bezerra, M. A., Santelli, R. E., Oliveira, E. P., Villar, L. S. and Escaleira, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965–77.

Bion, D., Blank, M., Freas, D., Gambogi, C., Rotsides, D., Shahidain, S. and Zhan, B. (2011). *Zero-order kinetic release from capsule reservoirs through semi-permeable membranes* (pp. 1–13).

Bonadio, W. A. (1997). Medical-legal considerations related to symptom duration and patient outcome after bacterial meningitis. *Pediatric Bacterial Meningitis*, 15, 420–423.

Borchardt, J.K. (2004). The history of bacterial meningitis treatment. *Drug News and Perspectives*, 17, 219–224.

Brasnjevic, I., Steinbusch, H. W. M., Schmitz, C. and Martinez-Martinez, P. (2009). Delivery of peptide and protein drugs over the blood-brain barrier. *Progress in Neurobiology*, 87(4), 212–251.

Buszello, K., Harnisch, S., Muller, R.H. and Muller, B.W. (2000). The influence of alkali fatty acids on the properties and the stability of parenteral o/w emulsions modified with solutol HS15. *European Journal of Pharmaceutics and Biopharmaceutics*, 49, 143–149.

Capek, I. (2004). Degradation of kinetically-stable o/w emulsions. *Advances in Colloid and Interface Science*, 107(2-3), 125–55.

Castellanos, I. J., Flores, G. and Griebenow, K. (2005). Effect of the molecular weight of poly (ethylene glycol) used as emulsifier on

$\alpha$  chymotrypsin stability upon encapsulation in PLGA microspheres. *Journal of Pharmacy and Pharmacology*, 57(10), 1261-1269.

Cecchelli, R., Berezowski, V., Lundquist, S., Culot, M., Renftel, M., Dehouck, M.-P. and Fenart, L. (2007). Modelling of the blood-brain barrier in drug discovery and development. *Nature Reviews. Drug Discovery*, 6(8), 650–61.

Costa, P. and Lobo, M. S. (2001). Modeling and comparison of dissolution profiles. *European Journal of Pharmaceutical Sciences*, 13, 123–133.

Cuéllar, I., Bullón, J., Forgarini, A. M., Cárdenas, A. and Briceño, M. I. (2005). More efficient preparation of parenteral emulsions or how to improve a pharmaceutical recipe by formulation engineering. *Chemical Engineering Science*, 60, 2127–2134.

Date, A. A. and Nagarsenker, M. S. (2007). Design and evaluation of self-nanoemulsifying drug delivery systems (SNEDDS) for cefpodoxime proxetil. *International Journal of Pharmaceutics*, 329(1-2), 166–72.

De Jong, I. G., Veening, J.-W. and Kuipers, O. P. (2010). Heterochronic phosphorelay gene expression as a source of heterogeneity in *Bacillus subtilis* spore formation. *Journal of Bacteriology*, 192(8), 2053–67.

De Klerk, O. L., Willemsen, A. T. M., Bosker, F. J., Bartels, A. L., Hendrikse, N. H., Den Boer, J. A. and Dierckx, R. A. (2010). Regional increase in P-glycoprotein function in the blood-brain barrier of patients with chronic schizophrenia: a PET study with [<sup>11</sup>C]verapamil as a probe for P-glycoprotein function. *Psychiatry Research: Neuroimaging*, 183(2), 151–156.

Delmas, T., Piraux, H., Couffin, A., Texier, I., Vinet, F., Poulin, P., Cates, M. E. and Bibette, J. (2011). How to prepare and stabilize very small nanoemulsions. *Langmuir Article*, 27(5): 1683-1692.

Desrumaux, A. and Marcand, J. (2002). Formation of sunflower oil emulsions stabilized by whey proteins with high-pressure homogenization ( up to 350 MPa ): effect of pressure on emulsion characteristics. *International Journal of Food Science and Technology*, 37, 263–269.

Driscoll, D. F., Bistrain, B. R., Demmelmair, H. and Koletzko, B. (2008). Pharmaceutical and clinical aspects of parenteral lipid emulsions in neonatology. *Clinical Nutrition*, 27, 497–503.

Fernandes, C., Soni, U. and Patravale, V. (2010). Nano-interventions for neurodegenerative disorders. *Pharmacological Research*, 62(2), 166–78.

Fitch, M.T. and Van de Beek, D. (2007). Emergency diagnosis and treatment of adult meningitis. *Lancet Infectious Diseases*, 7, 191-200.

- Floyd, A. G. (1999). Top ten considerations in the development of parenteral emulsions. *Pharmaceutical Science and Technology Today*, 4(2), 134–143.
- Floury, J., Desrumaux, A. and Legrand, J. (2003) Effect of high pressure homogenisation on methylcellulose as food emulsifier. *Journal of Food Engineering*, 58, 227-238.
- Frances, Karen, Beek, J. V., Canova, C., Neal, J. W. and Gasque, P. (2003). *The Blood-brain Barrier (BBB)*. Expert Reviews in Molecular Medicine: Cambridge University Press.
- Ganta, S. and Amiji, M. (2009). Coadministration of Paclitaxel and Curcumin in nanoemulsion formulations to overcome multidrug resistance in tumor cells. *Molecular Pharmaceutics*, 6(3), 928–939.
- Ganta, S., Paxton, J. W., Baguley, B. C. and Garg, S. (2008). Pharmacokinetics and pharmacodynamics of chlorambucil delivered in parenteral emulsion. *International Journal of Pharmaceutics*, 360, 115–121.
- Gao, D. Y., Ashworth, E., Kleinhans, F. W., Mazur, P. and Critser, J. K. (1993). Hyperosmotic Tolerance of Human Spermatozoa: Separate Effects of Glycerol , Sodium Chloride , and Sucrose on Spermlolysis. *Biology of Reproduction*, 123, 112–123.
- Garcia-Fuentes, M., Torres, D. and Alonso, M. J. (2002). Design of lipid nanoparticles for the oral delivery of hydrophilic macromolecules. *Colloids and Surfaces B: Biointerfaces*, 27, 159–168.
- García, B. M., Ferrusola, C. O., Aparicio, I. M., Miró-Morán, A., Rodríguez, A. M., Bolaños, J. M. G., Fernández, L. G., Balao da Silva, C. M., Martínez, H. R., Tapia, J. A. and Peña, F. J. (2012). Toxicity of glycerol for the stallion spermatozoa: Effects on membrane integrity and cytoskeleton, lipid peroxidation and mitochondrial membrane potential. *Theriogenology*, 77, 1280–1289.
- Gerber, J., Böttcher, T., Hahn, M., Siemer, A., Bunkowski, S. and Nau, R. (2004). Increased mortality and spatial memory deficits in TNF-alpha-deficient mice in ceftriaxone-treated experimental pneumococcal meningitis. *Neurobiology of Disease*, 16(1), 133–8.
- Główka, E., Wosicka-Frackowiak, H., Hyla, K., Stefanowska, J., Jastrzebska, K., Klapiszewski, L., Jesionowski, T. and Cal, K. (2014). Polymeric nanoparticles-embedded organogel for roxithromycin delivery to hair follicles. *European Journal of Pharmaceutics and Biopharmaceutics*, 88, 75–84.

- Gunawan, E. R. and Suhendra, D. (2008). Synthesis of Wax Esters From Palm Kernel Oil Catalyzed by Lipase. *Jurnal Matematika Dan Sains*, 13, 76–83.
- Han, X., Cheng, L., Zhang, R. and Bi, J. (2009). Extraction of safflower seed oil by supercritical CO<sub>2</sub>. *Journal of Food Engineering*, 92(4), 370–376.
- Hawkins, B. T. and Davis, T. P. (2005). The Blood-Brain Barrier / Neurovascular Unit in Health and Disease. *Pharmacological Reviews*, 57(2), 173–185.
- Herman, C.J. and Groves, M.J. (1992). Hydrolysis kinetics of phospholipids in thermally stressed intravenous lipid emulsion formulations. *Journal of Pharmacy and Pharmacology*, 44, 539–542.
- Hippalgaonkar, K., Majumdar, S. and Kansara, V. (2010). Injectable lipid emulsions-advancements, opportunities and challenges. *AAPS Pharmaceutical Science and Technology*, 11(4), 1526–40.
- Huang, S. H., Stins, M. F. and Kim, K. S. (2000). Bacterial penetration across the blood-brain barrier during the development of neonatal meningitis. *Microbes and Infection*, 2(10), 1237–1244.
- Huynh, G. H., Deen, D. F. and Szoka, F. C. (2006). Barriers to carrier mediated drug and gene delivery to brain tumors. *Journal of Controlled Release*, 110(2), 236–59.
- Jafari, S. M., Assadpoor, E., He, Y. and Bhandari, B. (2008). Re-coalescence of emulsion droplets during high-energy emulsification. *Food Hydrocolloids*, 22(7), 1191–1202.
- Jafari, S. M., He, Y. and Bhandari, B. (2007). Production of sub-micron emulsions by ultrasound and microfluidization techniques. *Journal of Food Engineering*, 82(4), 478–488.
- Jafari, S. M., He, Y. and Bhandari, B. (2006). Nano-emulsion production by sonication and microfluidization - A comparison. *International Journal of Food Properties*, 9, 475-485.
- Jayasooriya, S. D., Bhandari, B. R., Torley, P. and D'Arcy, B. R. (2004). Effect of high power ultrasound waves on properties in meat. *International Journal of Food Properties*, 7(2): 301-319.
- Joshi, M. D. and Müller, R. H. (2009). Lipid nanoparticles for parenteral delivery of actives. *European Journal of Pharmaceutics and Biopharmaceutics*, 71(2), 161–72.
- Jumaa, M. and Müller, B. W. (1999). In vitro investigation of the effect of various isotonic substances in parenteral emulsions on human

erythrocytes. *European Journal of Pharmaceutical Sciences*, 9, 207–212.

Jumaa, M. and Müller, B. W. (1998). The effect of oil components and homogenization conditions on the physicochemical properties and stability of parenteral fat emulsions. *International Journal of Pharmaceutics*, 163, 81–89.

Jumaa, M., Kleinebudde, P. and Müller, B. W. (1998). Mixture experiments with the oil phase of parenteral emulsions. *European Journal of Pharmaceutics and Biopharmaceutics*, 46(2), 161–167.

Keck, C. M. and Müller, R. H. (2006). Drug nanocrystals of poorly soluble drugs produced by high pressure homogenisation. *European Journal of Pharmaceutics and Biopharmaceutics*, 62, 3-16.

Kelmann, R. G., Kuminek, G., Teixeira, H. F. and Koester, L. S. (2007). Carbamazepine parenteral nanoemulsions prepared by spontaneous emulsification process. *International Journal of Pharmaceutics*, 342, 231–239.

Keng, P. S., Basri, M., Ariff, A. B., Abdul Rahman, M. B., Abdul Rahman, R. N. Z. and Salleh, A. B. (2008). Scale-up synthesis of lipase-catalyzed palm esters in stirred-tank reactor. *Bioresource Technology*, 99(14): 6097-6104.

Kibanov, A. (2011). *Controlled Release Technology: Polymeric Delivery Systems for Pharmaceuticals, Proteins, and Other Agents*. Cambridge (MA): Massachusetts Institute of Technology.

Kleine, T. O., Zwerenz, P., Zöfel, P. and Shiratori, K. (2003). New and old diagnostic markers of meningitis in cerebrospinal fluid (CSF). *Brain Research Bulletin*, 61(3), 287–297.

Klement, W. and Arndt, J. O. (1991). Pain on I.V. injection of some anaesthetic agents is evoked by the unphysiological osmolality or pH of their formulations. *British Journal of Anaesthesia*, 66, 189–195.

Krol, S. (2012). Challenges in drug delivery to the brain: Nature is against us. *Journal of Controlled Release*, 164(2), 145–155.

Kwak, S.-Y., Kriven, W. M., Wallig, M. A. and Choy, J.-H. (2004). Inorganic delivery vector for intravenous injection. *Biomaterials*, 25(28), 5995–6001.

Leal-Calderon, F., Thivilliers, F. and Schmitt, V. (2007). Structured emulsions. *Current Opinion in Colloid and Interface Science*, 12(4-5), 206–212.



- Lee, M.-K., Chun, S.-K., Choi, W.-J., Kima, J.-K., Choi, S.-H., Kim, A., Oungbho, K., Park, J.-S., Ahn, W. S. and Kim, C.-K. (2005). The use of chitosan as a condensing agent to enhance emulsion-mediated gene transfer. *Biomaterials*, 26, 2147–2156.
- Leong, T. S. H., Wooster, T. J., Kentish, S. E. and Ashokkumar, M. (2009). Minimising oil droplet size using ultrasonic emulsification. *Ultrasonics Sonochemistry*, 16(6), 721–7.
- Lewis, R. F., Dorlencourt, F. and Pinel, J. (1998). Long-acting oily chloramphenicol for meningococcal meningitis. *The Lancet*, 352, 1998.
- Lopetinsky, R. J. G., Masliyah, J. H. and Xu, Z. (2006). Solids-stabilized emulsions: a review. *Colloidal particles at liquid interfaces*, 186-224.
- Lovell, M. W., Johnson, H. W. and Gupta, P. K. (1995). Stability of a less-painful intravenous emulsion of clarithromycin. *International Journal of Pharmaceutics*, 118, 47–54.
- Lovelyn, C. and Attama, A. A. (2011). Current State of Nanoemulsions in Drug Delivery. *Journal of Biomaterials and Nanobiotechnology*, 2(5), 626–639.
- Lu, Y., Wang, Y. J. and Tang, X. (2008). Formulation and thermal sterile stability of a less painful intravenous clarithromycin emulsion containing vitamin E. *International Journal of Pharmaceutics*, 346, 47–56.
- Lucks, J., Muller, B. and Klutsch, K. (2000). *Parenteral fat emulsions: structure stability and applications. Pharmaceutical Emulsions and Suspensions*. Marcel Dekker, New York.
- Martins, S. M., Sarmiento, B., Nunes, C., Lúcio, M., Reis, S. and Ferreira, D. C. (2013). Brain targeting effect of camptothecin-loaded solid lipid nanoparticles in rat after intravenous administration. *European Journal of Pharmaceutics and Biopharmaceutics*, 85, 488–502.
- McClements, D. J. (2005). *Food emulsions: principles, practice and techniques*. CRC Press.
- McClements, D. J. (2002). Theoretical prediction of emulsion color. *Advances in Colloid and Interface Science*, 97(1-3), 63–89.
- McLean, I. W., Schwab, J. L., Hillegas, A. B. and Schlingman, A. S. (1949). *Susceptibility of micro-organism to chloramphenicol (chloromycetin)* (pp. 953–963).
- Menon, J. U., Ravikumar, P., Pise, A., Gyawali, D., Hsia, C. C. W., Nguyen, K. T. (2014). Polymeric nanoparticles for pulmonary protein and DNA delivery. *Acta Biomaterialia*, 10, 2643–2652.

- Meyer, R. J. and Hussain, A. S. (2005). *Awareness Topic: Mitigating the risks of ethanol induced dose dumping from oral sustained/controlled release dosage form*. Paper presented at the meeting of FDA's ACPS Meeting, USA. October 2005.
- Mirhosseini, H., Ping, C., Hamid, N. S. A. and Yusof, S. (2008). Effect of Arabic gum, xanthan gum and orange oil contents on  $\zeta$ -potential, conductivity, stability, size index and pH of orange beverage emulsion. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 315, 47–56.
- Mirhosseini, H., Tan, C. P., Taherian, A. R. and Boo, H. C. (2009). Modeling the physicochemical properties of orange beverage emulsion as function of main emulsion components using response surface methodology. *Carbohydrate Polymers*, 75(3), 512–520.
- Mishal, J., Embon, A., Darawshe, A., Kidon, M. and Magen, E. (2008). Community acquired acute bacterial meningitis in children and adults: an 11-year survey in a community hospital in Israel. *European Journal of Internal Medicine*, 19(6), 421–426.
- Montgomery, D.C. (2001). *Design and Analysis of Experiments*. 5th Edition, Wiley, New York.
- Müller, R. H., Schmidt, S., Buttle, I., Akkar, A., Schmitt, J. and Brömer, S. (2004). SolEmuls®—novel technology for the formulation of I.V. emulsions with poorly soluble drugs. *International Journal of Pharmaceutics*, 269(2), 293–302.
- Müller, R. H., Radtke, M. and Wissing, S. A. (2002). Nanostructured lipid matrices for improved microencapsulation of drugs. *International Journal of Pharmaceutics*, 242, 121–128.
- Myers, D. (1999). *Surfaces, interfaces and colloids: Principles and applications*. John and Wiley Sons. United States, America: 271.
- Nagarwal, R. C., Kumar, R., Dhanawat, M., Das, N. and Pandit, J.-K. (2011). Nanocrystal technology in the delivery of poorly water soluble drugs: An overview. *Current Drug Delivery*, 8, 398-406.
- Nielloud, F., Marti-Mestres, G., Laget, J.P., Fernandez, C. and Maillols, H. (1996). Emulsion formulations: study of the influence of parameters with experimental designs. *Drug Development and Industrial Pharmacy*. 22, 159–166.
- Pahnke, J., Walker, L. C., Scheffler, K. and Krohn, M. (2009). Alzheimer's disease and blood-brain barrier function-Why have anti-beta-amyloid therapies failed to prevent dementia progression? *Neuroscience and Biobehavioral Reviews*, 33(7), 1099–1108.

- Pal, R. (2011). Rheology of simple and multiple emulsions. *Current Opinion in Colloid and Interface Science*, 16(1), 41–60.
- Peng, L.-C., Liu, C.-H., Kwan, C.-C. and Huang, K.-F. (2010). Optimization of water-in-oil nanoemulsions by mixed surfactants. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 370(1-3), 136–142.
- Penha-Silva, N., Arvelos, L.R., Cunha, C.C., Aversi-Ferreira, T.A., Gouvêa-e-Silva, L.F., Garrote-Filho, M.S., Finotti, C.J., Bernardino-Neto, M. and de Freitas Reis, F.G. (2008). Effects of glycerol and sorbitol on the thermal dependence of the lysis of human erythrocytes by ethanol. *Bioelectrochemistry*, 73, 23–29.
- Pey, C. M., Maestro, A., Solé, I., González, C., Solans, C. and Gutiérrez, J. M. (2006). Optimization of nano-emulsions prepared by low-energy emulsification methods at constant temperature using a factorial design study. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 288(1-3), 144–150.
- Poorgholami-Bejarpasi, N., Hashemianzadeh, M., Mousavi-Khoshdel, S. M. and Sohrabi, B. (2010). Investigation of the mixing behavior of surfactants by lattice Monte Carlo simulation. *Journal of Molecular Modeling*, 16(9), 1499–508.
- Prakash, U. R. T. and Thiagarajan, P. (2011). Nanoemulsions for drug delivery through different routes. *Research in Biotechnology*, 2(3), 1–13.
- Prasad, K., Karlupia, N. and Kumar, A. (2009). Treatment of bacterial meningitis: an overview of Cochrane systematic reviews. *Respiratory Medicine*, 103(7), 945–50.
- Rabinovich-guilatt, L., Couvreur, P., Lambert, G., Goldstein, D., Benita, S. and Dubernet, C. (2004). Extensive surface studies help to analyse zeta potential data : the case of cationic emulsions. *Chemistry and Physics of Lipids*, 131, 1–13.
- Rawat, M., Singh, D., Saraf, S. and Saraf. S. (2006). Nanocarriers: Promising vehicle for bioactive drugs. *Biological and Pharmaceutical Bulletin*, 29(9), 1790–1798.
- Roco, M. C., Williams, R. S. and Alivisatos, P. (2000). *Nanotechnology Research Directions, Biological, medical and health applications* (Eds.), Kluwer Academic Publishers, Boston: Chap 8.
- Roney, C., Kulkarni, P., Arora, V., Antich, P., Bonte, F., Wu, A. and Aminabhavi, T. M. (2005). Targeted nanoparticles for drug delivery through the blood-brain barrier for Alzheimer's disease. *Journal of Controlled Release*, 108(2-3), 193–214.



- Saberi, A. H., Fang, Y. and McClements, D. J. (2013). Fabrication of vitamin E-enriched nanoemulsions: Factors affecting particle size using spontaneous emulsification. *Journal of Colloid and Interface Science*, 391, 95–102.
- Salim, N., Basri, M., Rahman, M. B. A., Abdullah, D. K. and Basri, H. (2012). Modification of palm kernel oil esters nanoemulsions with hydrocolloid gum for enhanced topical delivery of ibuprofen. *International Journal of Nanomedicine*, 7, 4739–4747.
- Sandoval, K. E. and Witt, K. A. (2008). Blood-brain barrier tight junction permeability and ischemic stroke. *Neurobiology of Disease*, 32(2), 200–19.
- Santos-Magalhaes, N. S., Pontes, A., Pereira, V. M. W. and Caetano, M. N. P. (2000). Colloidal carriers for benzathine penicillin G: Nanoemulsions and nanocapsules. *International Journal of Pharmaceutics*, 208, 71–80.
- Scherrmann, J. M. (2002). Drug delivery to brain via the blood-brain barrier. *Vascular Pharmacology*, 38(6), 349–54.
- Scholfield, C. R. (1981). Composition of soybean lecithin. *Journal of the American Oil Chemists' Society*, 58(10), 889–892.
- Seekkuarachchi, I. N., Tanaka, K. and Kumazawa, H. (2006). Formation and characterization of submicrometer oil-in-water (O/W) emulsions, using high-energy emulsification. *Industrial and Engineering Chemistry Research*, 45(1), 372–390.
- Shafiq, S. and Shakeel, F. (2008). Enhanced stability of ramipril in nanoemulsion containing cremophor-EL: a technical note. *AAPS Pharmaceutical Science and Technology*, 9(4), 1097–101.
- Sharma, N., Bansal, M., Visht, S., Sharma, P. K. and Kulkarni, G. T. (2010). Nonemulsion: A new concept of delivery system. *Chronicles of Young Scientists*, 1(2), 2–6.
- Shoab, M. H., Tazeen, J., Merchant, H. A. and Yousof, R. I. (2006). Evaluation of drug release kinetics from ibuprofen matrix tablets using HPMC. *Journal of Pharmaceutical Sciences*, 19(2), 119–124.
- Solans, C., Izquierdo, P., Nolla, J., Azemar, N. and Garcia-Celma, M. J. (2005). Nano-emulsions. *Current Opinion in Colloid and Interface Science*, 10, 102–110.
- Solans, C. and Solé, I. (2012). Nano-emulsions: Formation by low-energy methods. *Current Opinion in Colloid and Interface Science*, 17(5), 246–254.

- Song, J., Shi, F., Zhang, Z., Zhu, F., Xue, J., Tan, X. and Jia, X. (2011). Formulation and Evaluation of Celastrol-Loaded Liposomes. *Molecules*, 16, 7880–7892.
- Stang, M., Schuchmann, H. and Schubert, H. (2001). Emulsification in high-pressure homogenizers. *Engineering in Life Sciences*, 1, 151–157.
- Strickley, R. G. (2004). Solubilizing excipients in oral and injectable formulations. *Pharmaceutical Research*, 21(2), 201–30.
- Swarnalatha, S., Selvi, P. K., Ganesh Kumar, A. and Sekaran, G. (2008). Nanoemulsion drug delivery by ketene based polyester synthesized using electron rich carbon/silica composite surface. *Colloids and Surfaces. B: Biointerfaces*, 65(2), 292–9.
- Tadros, T. (2004). *Application of rheology for assessment and prediction of the long-term physical stability of emulsions* (Vol. 109, pp. 227–258).
- Tadros, T., Izquierdo, P., Esquena, J. and Solans, C. (2004). Formation and stability of nano-emulsions. *Advances in Colloid and Interface Science* 108±109, 303–318.
- Tadros, T.F. and Vincent, B. (1983). *Encyclopedia of Emulsion Technology*, P. Becher (Eds.), Dekker, New York, 1983.
- Taha, E. I., Samy, A. M., Kassem, A. A. and Khan, M. A. (2005). Response surface methodology for the development of self-nanoemulsified drug delivery system (SNEDDS) of all-trans-retinol acetate. *Pharmaceutical Development and Technology*, 10(3), 363-370.
- Tamburini, J., Grimaldi, D., Bricaire, F. and Bossi, P. (2005). Acute bacterial meningitis in a patient receiving ibuprofen. *Journal of Infection*, 51, 336–337.
- Tang, S. Y., Shridharan, P. and Sivakumar, M. (2013). Impact of process parameters in the generation of novel aspirin nanoemulsions - Comparative studies between ultrasound cavitation and microfluidizer. *Ultrasonics Sonochemistry*, 20(1), 485–97.
- Tang, S.Y., Manickam, S., Wei, T.K. and Nashiru, B. (2012). Formulation development and optimization of a novel Cremophore EL-based nanoemulsion using ultrasound cavitation. *Ultrasonics Sonochemistry*, 19, 330–345.
- Theodoridou, K., Vasilopoulou, V. A, Katsiaflaka, A., Theodoridou, M. N., Roka, V., Rachiotis, G. and Hadjichristodoulou, C. S. (2013). Association of treatment for bacterial meningitis with the development of sequelae. *International Journal of Infectious Diseases*, 17(9), 707–13.

- Trapani, G., Altomare, C., Franco, M., Latrofa, A. and Liso, G. (1995). Determination of hydrophile-lipophile balance of some polyethoxylated non-ionic surfactants by reversed-phase thin layer chromatography. *International Journal of Pharmaceutics*, 116(1), 95–99.
- Trias, J. (2001). The role of combichem in antibiotic discovery. *Current Opinion in Microbiology*, 4, 520–525.
- Uskokovi, V. and Drofenik, M. (2005). Synthesis of materials within reverse micelles. *Surface Review and Letters*, 12, 239.
- Vyas, T.K., Shahiwala, A. and Amiji, M.M. (2008). Improved oral bioavailability and brain transport of Saquinavir upon administration in novel nanoemulsion formulations. *International Journal of Pharmaceutics*, 347, 93–101.
- Wang, Y., Zheng, Y., Zhang, L., Wang, Q. and Zhang, D. (2013). Stability of nanosuspensions in drug delivery. *Journal of Controlled Release*, 172, 1126–1141.
- Wang, J.-J., Sung, K. C., Hu, O. Y.-P., Yeh, C.-H. and Fang, J.-Y. (2006). Submicron lipid emulsion as a drug delivery system for nalbuphine and its prodrugs. *Journal of Controlled Release*, 115(2), 140–9.
- Wang, L., Li, X., Zhang, G., Dong, J. and Eastoe, J. (2007). Oil-in-water nanoemulsions for pesticide formulations. *Journal of Colloid and Interface Science*, 314(1), 230–5.
- Weisfelt, M., Van De Beek, D., Spanjaard, L., Reitsma, J. B. and De Gans, J. (2006). Clinical features, complications, and outcome in adults with pneumococcal meningitis: a prospective case series. *The Lancet*, 5(2), 123–9.
- Weiss, J., Coupland, J. N., Brathwaite, D. and McClements, D. J. (1997). Influence of molecular structure of hydrocarbon emulsion droplets on their solubilization in nonionic surfactant micelles. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 121, 53–60.
- Westesen, K., Bunjes, H. and Koch, M.H.J. (1997). Physicochemical characterization of lipid nanoparticles and evaluation of their drug loading capacity and sustained release potential. *Journal of Controlled Release*, 48, 223–236.
- Windhab, E. J., Dressler, M., Feigl, K., Fischer, P. and Megias-Alguacil, D. (2005). Emulsion processing—from single-drop deformation to design of complex processes and products. *Chemical Engineering Science*, 60(8-9), 2101–2113.

Wolburg, H., Noell, S., Mack, A., Wolburg-Buchholz, K. and Fallier-Becker, P. (2009). Brain endothelial cells and the glio-vascular complex. *Cell and Tissue Research*, 335(1), 75–96.

Zaidul, I. S. M., Norulaini, N. A. N., Omar, A. K. M., Sato, Y. and Smith, R. L. (2007). Separation of palm kernel oil from palm kernel with supercritical carbon dioxide using pressure swing technique. *Journal of Food Engineering*, 81(2), 419–428.

Zainol, S., Basri, M., Basri, H., Shamsuddin, A. F., Abdul Ghani, S. S., Karjiban, R. A. and Abdul Malek, E. (2012). Formulation Optimization of a Palm-Based Nanoemulsion System Containing Levodopa. *International Journal of Molecular Sciences*, 13, 13049–13064.

