



UNIVERSITI PUTRA MALAYSIA

***BIOREDUCTION OF HEXAVALENT MOLYBDENUM TO MOLYBDENUM
BLUE USING *Serratia* sp. MIE2 AND PURIFICATION OF
MOLYBDENUM-REDUCING ENZYME***

MOHD IZUAN EFFENDI BIN HALMI

FBSB 2014 35



**BIOREDUCTION OF HEXAVALENT MOLYBDENUM TO MOLYBDENUM
BLUE USING *Serratia* sp. MIE2 AND PURIFICATION OF
MOLYBDENUM-REDUCING ENZYME**

By

MOHD IZUAN EFFENDI BIN HALMI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

November 2014

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

BIOREDUCTION OF HEXAVALENT MOLYBDENUM TO MOLYBDENUM BLUE USING *Serratia* sp. MIE2 AND PURIFICATION OF MOLYBDENUM-REDUCING ENZYME

By

MOHD IZUAN EFFENDI BIN HALMI

November 2014

Chairman: Mohd Yunus Abd Shukor, PhD

Faculty: Biotechnology and Biomolecular Sciences

Molybdenum reduction is an old phenomenon that has received very low attention compare to other well-known and extensively studied metals such as chromium, mercury and lead. Molybdenum has long been known to be toxic to ruminants and not toxic to other organisms. However, more recently it has been increasingly reported that molybdenum shows toxic effects to reproductive organs of fish, mouse and even humans at concentrations between 1 and 100 ppm. Hence its removal from the environment is highly sought after. The isolation of molybdenum reducing bacteria and the elucidation of the reducing mechanism will lead to an efficient bioremediation system. To fulfil this, a new Mo-reducing bacterium was isolated from an agriculture soil plot from Universiti Putra Malaysia. The isolate was tentatively identified as *Serratia* sp. MIE2 based on 16s rDNA molecular phylogeny. *Serratia* sp. MIE2 is a gram negative, oxidase and catalase positive bacterium. The molybdenum blue produced by *Serratia* sp. MIE2 exhibited a unique absorption spectrum with maximum peak at 865 nm and a shoulder at 700 nm. Dialysis tubing experiment showed that molybdate reduction by *Serratia* sp. MIE2 was an enzymatic process and not chemically mediated.

Characterization and optimization of molybdenum blue production by *Serratia* sp. MIE2 was carried out using one factor at a time (OFAT) and Response Surface Methodology (RSM). One factor at a time (OFAT) showed the optimum conditions supporting molybdate reduction occurred at pH 6.0, from 27 to 35 °C and 30-40 g/L sucrose as the carbon source or electron donor. The best nitrogen source was ammonium sulphate with an optimum concentration at 10 g/L. Moreover, the optimum concentrations of phosphate and molybdate were 2 and 10 mM, respectively. Molybdate reduction was maximized and optimized using response surface methodology (RSM) with optimum conditions occurring at 20 mM of molybdate, 25 g/L of sucrose, pH 6.25 and 3.95 mM of phosphate with molybdenum blue production increasing from an OFAT absorbance yield of 10.0 to higher than 20.0 as measured at 865 nm.

Modelling kinetic studies of *Serratia* sp. MIE2 using the optimum conditions obtained from the classical method (OFAT) show that the best model was Teissier followed by Luong, Aiba, Yano and Haldane with correlation coefficient, R^2 values of 0.994, 0.993, 0.992, 0.990 and 0.982, respectively. The calculated values of P_{max} , K_s and K_i of the best model were 0.89 μ mole Molybdenum blue per hour, 5.84 mM and 32.23 mM respectively. Otherwise, modelling kinetics using the optimum condition obtained from RSM showed that the Luong model was the best model followed by Teissier, Aiba, Yano and Haldane with correlation coefficient, R^2 values of 0.999, 0.994, 0.993, 0.992 and 0.965, respectively. However, since Luong exhibited 4 kinetic constants while Teissier has only 3 constants, by default, Teissier model was chosen due to its mathematical simplicity. The calculated values of R^2 , P_{max} , K_s and K_i of the best model, Teissier were 1.97 μ mole Mo-blue per hour, 5.79 mM and 31.48 mM, respectively. Modelling kinetics showed the value of P_{max} was increasing from 0.89 μ mole Molybdenum blue per hour to 1.97 μ mole molybdenum blue per hour indicating that molybdate reduction yield increase several fold after optimization using RSM.

Before purification process, preliminary studies such as effect of storage and chromatographic stabilities, effects of restorative and inhibitive agents were carried out to minimise denaturation and to maximise yield of purified enzyme. The buffer use during storage and purification process was Tris-HCl at pH 7.0. Mo-reducing enzyme was stable when stored at -80°C for both 24 hours and one month followed by storage on ice (0°C). Temperature stability study showed that the enzyme was most stable at 25°C followed by 40°C with complete lost of activity at 60 and 40 minutes of incubation at 54 and 70°C . EGTA or (ethylene glycol tetraacetic acid), EDTA, Triton X-100, DBS and SDS decrease 50% activity of enzyme at concentration 0.1mM, 0.1mM, 0.1%, 0.1%, and 0.1%, respectively. DTT could restore the Mo-reducing enzyme activity of up to 100% at the maximum concentration of 5 mM for DTT and 0.5 mM for β -mercaptoethanol. Effects of cofactor suggest that nickel might be an important cofactor for the enzyme. Heavy metals such as mercury and zinc effect strongly inhibited the Mo-reducing enzyme. The coenzyme such as FMN and FAD were able to restore Mo-reducing enzyme activity. Mo-reducing enzyme was not inhibited by respiratory inhibitors, therefore, the electron transport chain of this bacterium is not the site of molybdate reduction.

Purification of the Mo-reducing enzyme was done using ammonium sulphate precipitation, gel filtration on Zorbax GF-250 and Zorbax GF-450 with a 20.8 purification fold. The molecular mass was estimated to be 100 kDa by SDS-polyacrylamide gel electrophoresis and the enzyme was monomeric. Mo-reducing enzyme showed maximum activity at 35°C and pH 5. The enzyme was assayed using NADH as the electron donor with the maximum initial velocity, V_{max} of 16.18 nmole molybdenum blue/min/mg protein and a Michaelis constant, K_m at 0.89 mM. The optimum concentration of phosphomolybdate (electron acceptor substrate) was 10 mM, with a V_{max} of 6.89 nmole molybdenum blue/min/mg protein (NADH as electron donor at saturated concentrations) and K_m of 6.02 mM. Identification of pure enzyme using MALDI-TOF showed only peptide DNAATRSEAMSLIHGR shows similarity to 35% to nitrile oxidoreductase and GTP cyclohydrolase I. The low similarity value prohibited further analysis to be carried out. Thus, the enzyme is assigned as hypothetical protein.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENURUNAN HEKSAVALEN MOLIBDENUM KEPADA MOLIBDENUM BIRU OLEH *Serratia* sp. MIE2 DAN PENULENAN ENZIM PENURUN MOLIBDENUM

Oleh

MOHD IZUAN EFFENDI BIN HALMI

November 2014

Pengerusi: Mohd Yunus Abd Shukor, PhD
Fakulti: Bioteknologi dan Sains Biomolekul

Penurunan molibdenum adalah suatu fenomena yang kurang mendapat perhatian berbanding dengan logam lain-lain yang terkenal dan dikaji secara meluas seperti kromium, merkuri dan plumbum. Molibdenum telah lama diketahui adalah toksik kepada ruminan dan tidak toksik kepada organisma lain. Walau bagaimanapun, baru-baru ini terdapat laporan yang menunjukkan bahawa molibdenum menunjukkan kesan toksik kepada organ-organ pembiakan ikan, tikus dan juga manusia pada kepekatan antara 1 dan 100 ppm. Oleh itu penyingkirannya daripada alam sekitar mendapat perhatian yang tinggi. Pengasingan bakteria penurun molibdenum dan pengetahuan mengenai mekanisme penurunan akan membawa kepada sistem biopemulihan yang cekap. Satu bakteria Mo-penurun telah diasingkan daripada plot tanah pertanian dari Universiti Putra Malaysia. Isolat ini secara sementara ini dikenalpasti sebagai *Serratia* sp. MIE2 berdasarkan analisa filogenetik molekul 16s rDNA. *Serratia* sp. MIE2 adalah gram negatif, oksidase dan katalase positif. Molibdenum biru dihasilkan oleh *Serratia* sp. MIE2 mempamerkan spektrum penyerapan unik dengan puncak maksimum pada 865 nm dan bahu di 700nm. Eksperimen tiub dialisis menunjukkan bahawa penurunan molibdenum oleh *Serratia* sp. MIE2 adalah suatu proses melibatkan enzim dan bukan secara kimia.

Pencirian dan pengoptimuman pengeluaran molybdenum biru oleh *Serratia* sp. MIE2 telah dijalankan dengan menggunakan satu-faktor-pada-satu-masa (OFAT) dan Kaedah Metodologi Permukaan (RSM). Kaedah satu-faktor-pada-satu-masa (OFAT) menunjukkan keadaan optimum menyokong pengurangan molibdenum berlaku pada pH 6.0, 27-35°C dan 30-40 g/L sukrosa sebagai sumber karbon atau elektron penderma. Sumber terbaik nitrogen adalah ammonium sulfat dengan kepekatan optimum pada 10 g/L. Selain itu, kepekatan optimum fosfat dan molibdat adalah 2 dan 10 mM, masing-masing. Pengurangan molibdenum telah dimaksimumkan dan dioptimumkan menggunakan Kaedah Metodologi Permukaan (RSM) dengan keadaan penghasilan molibdenum biru optimum berlaku pada 20 mM molibdat, 25 g/L sukrosa, pH 6.25 dan 3.95 mM fosfat dengan pengeluaran molibdenum biru meningkatkan dari serapan 10.0 menggunakan kaedah OFAT kepada serapan 20.0 pada jarak gelombang 865 nm menggunakan kaedah RSM.

Model kajian kinetik penurunan oleh *Serratia* sp. MIE2 menggunakan kondisi optimum yang diperolehi daripada kaedah klasik (OFAT) menunjukkan bahawa model yang terbaik ialah Teissier diikuti dengan Luong, Aiba, Yano dan Haldane dengan nilai R^2 0.994, 0.993, 0.992, 0.990 dan 0.982, masing-masing. Nilai P_{max} , K_s dan K_i model yang terbaik adalah 0.89 μmol Mo-biru per jam, 5.84 mM dan 32.23 mM, masing-masing. Kinetik pemodelan menggunakan keadaan optima yang diperolehi daripada kaedah RSM pula menunjukkan bahawa model Luong adalah model terbaik diikuti oleh Teissier, Aiba, Yano dan Haldane dengan, nilai R^2 0.999, 0.994, 0.993, 0.992 dan 0.965, masing-masing. Walau bagaimanapun, sejak Luong menggunakan 4 pemalar kinetik manakala Teissier hanya menggunakan 3 pemalar, maka secara tetapan model Teissier dipilih berdasarkan kesederhanaan matematik. Nilai P_{max} , K_s dan K_i model yang terbaik adalah 1.97 μmole Mo-biru sejam, 5.79 mM dan 31.48 mM, masing-masing. Kinetik pemodelan Teissier menunjukkan nilai P_{max} telah meningkat daripada 0.89 μmol molibdenum biru/jam kepada 1.97 μmole molibdenum biru per jam yang menunjukkan bahawa pengurangan molibdenum meningkatkan hasil beberapa kali ganda selepas pengoptimuman menggunakan kaedah RSM.

Sebelum proses penulenan, kajian awal seperti kesan penyimpanan dan pengstabilan semasa kromatografi, kesan agen pemulihan enzim dan agen perencat telah dijalankan untuk mengurangkan denaturasi dan memaksimumkan hasil enzim yang akan dituliskan. pH optima untuk tujuan penyimpanan dan proses kromatografi adalah Tris-HCl pada pH 7. Mo-penurun enzim adalah stabil apabila disimpan pada -80°C samada pada 24 jam atau satu bulan dan diikuti dengan penyimpanan di dalam ais (0°C). Kajian kestabilan suhu menunjukkan bahawa enzim adalah yang paling stabil pada 25°C diikuti dengan 40°C dengan kehilangan aktiviti berlaku pada 60 dan 40 minit pengeraman pada suhu 54 dan 70°C, masing-masing. EGTA (etilena glikol asid tetraasetik), EDTA (etilena diamina asid tetraasetik), Triton X-100, DBS (dodesil benzena sulfat) dan SDS (sodium dodesil sulfat) mengurangkan aktiviti enzim sebanyak 50% pada kepekatan 0.1mM, 0.1 mM, 0.1%, 0.1%, dan 0.1% masing-masing. DTT boleh memulihkan aktiviti enzim penurun molibdenum sehingga 100% pada kepekatan maksimum 5 mM untuk DTT dan 0.5 mM untuk 2-merkaptotanol. Kesan kofaktor menunjukkan bahawa nikel mungkin menjadi kofaktor penting bagi enzim ini. Logam berat seperti merkuri dan zink merencat aktiviti enzim penurun molibdenum. Koenzim seperti FMN (flavin mononukleotida) dan FAD (flavin adenina dinukleotida) dapat memulihkan aktiviti Mo-penurun enzim. Aktiviti enzim penurun molibdenum tidak direncat oleh perencat respirasi, oleh itu, rantaian pengangkutan elektron daripada bakteria ini bukan merupakan tapak aktiviti penurunan molibdenum.

Penulenan enzim penurun molibdenum telah dilakukan dengan menggunakan fraksinasi ammonium sulfat, gel penapisan menggunakan Zorbax GF-250 dan Zorbax G-450 dengan pekali penulenan sebanyak 20.8. Jisim molekul enzim penurun molibdenum dianggarkan 100 kDa menggunakan kaedah gel elektroforesis-SDS dan enzim penurun molibdenum adalah monomerik. Enzim penurun molibdenum menunjukkan aktiviti maksimum pada suhu 35°C dan pH 5. Enzim ini telah diasai menggunakan NADH sebagai penderma elektron dengan halaju awal maksimum, V_{max} adalah 16.18 nmole molibdenum biru /min/ mg protein dan pemalar Michaelis,

K_m adalah 0.89 mM. Kepekatan optimum fosfomolibdate (elektron penerima substrat) adalah 10 mM , dengan V_{max} adalah 6.89 nmole molibdenum biru/min/mg protein (NADH sebagai penderma elektron pada kepekatan tepu) dan K_m adalah 6.02 mM. Pengenalan enzim tulen menggunakan kaedah MALDI-TOF menunjukkan hanya peptide DNAATRSEAMSLIHGR mempunyai persamaan 35% kepada nitril oksidoreduktase and GTP siklohidrolase 1. Persamaan yang rendah menghalang analisis yang seterusnya. Oleh itu enzim ini dinamakan protein hipotetikal.



ACKNOWLEDGEMENTS

I am very glad and thankful to Allah, for His blessings for me to complete this thesis completely and successfully. In addition, the completion of this project would not have been successful without the assistance, help, and co-operation from a number of people.

I also would like to take this opportunity to special thank to my supervisor, Associate Professor Dr. Mohd. Yunus Abd. Shukor for his guidance, advice and moral support starting from my Bachelor until my PhD. I would like to express my gratitude to my beloved parents, Sam bin Bidin and Sabariah Binti Din and to my late father Halmi bin Hussin who passed away in 1994 for their encouragements and continuous support to keep me going on with this task. My co-supervisor, Dr. Wan Lutfi Bin Wan Johari, Dr. Mohd Shukuri Mohamad Ali and Dr. Noor Azmi Shaharuddin, I appreciate the guidance and suggestions from them during the duration of this project. Thank you to my beloved Norzila binti Kusnin who are always supportive of me and is the source of encouragement and kindness that have encouraged me to finish this thesis.

Thank you very much to my dear labmates, Finally, I would like to thank to all my lecturers especially in the Department of Biochemistry and in the Faculty of Biotechnology and Biomolecular Sciences and UPM in general, my friends and anyone who were involved in assisting me to complete this project. Thank you for your support and understanding.

I certify that a Thesis Examination Committee has met on 17 November 2014 to conduct the final examination of Mohd Izuan Effendi b Halmi on his thesis entitled "Bioreduction of Hexavalent Molybdenum to Molybdenum Blue using *Serratia* sp. MIE2 and Purification of Molybdenum-Reducing Enzyme" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Syahida binti Ahmad, PhD

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Umi Kalsom binti Md Shah, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Shuhaimi bin Mustafa, PhD

Professor

Institut Penyelidikan Produk Halal

Universiti Putra Malaysia

(Internal Examiner)

Amal Kanti Paul, PhD

Professor

Department of Botany

University of Calcutta

(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 23 January 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Mohd Yunus Abd Shukor, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Wan Lutfi Bin Wan Johari, PhD

Senior Lecturer
Faculty of Science and Environmental studies
Universiti Putra Malaysia
(Member)

Mohd Shukuri bin Mohamad Ali, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Noor Azmi Bin Shaharuddin, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Mohd Izuan Effendi Bin Halmi (GS36714)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLES OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xv
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xx
CHAPTER	
1.0 INTRODUCTION	1
1.1 Thesis Objective	2
2.0 LITERATURE REVIEW	3
2.1 The Chemistry of Molybdenum	3
2.2 The Biochemistry of Molybdenum	6
2.2.1 Molybdenum in Mammals	6
2.2.2 Molybdenum in Plant	7
2.3 Molybdenum Toxicity	9
2.4 Molybdenum as a Pollutant	10
2.5 Microbial reduction of metal ions	10
2.6 Enzymatic and Microbial Reduction of Molybdate to Mo-blue	11
2.7 Properties of partial and pure Mo-reducing Enzyme from several Bacteria	16
2.7.1 Mo-reducing enzyme from EC 48 with molybdate as theSubstrate	16
2.7.2 Mo-reducing enzyme from EC 48 with 12-phosphomolybdate as the substrate	17
2.7.3 Using 12-Phosphomolybdate as a substrate	19
2.7.4 Mo-reducing enzyme from EC 48 with Laboratory-prepared phosphomolybdate as the substrate	20
2.7.5 Mo-reducing enzyme from <i>Serratia marcescens</i> . Dr.Y5	20
2.8 Statement of the Problems	21
2.8.1 Isolation and characterization of a more potent Mo-reducing bacterium	21
2.9 Statistical-based optimization compared to one-at-a-time Approach	21
2.10 Kinetics of the Mo-reducing enzyme	22
2.11 Identification of the Mo-reducing enzyme	23

2.12 Significance of the Study	23
--------------------------------	----

3.0 METHODOLOGY	25
3.1 Materials	25
3.1.1 Chemicals and equipments	25
3.2 Preparation of culture medium	25
3.2.1 Low phosphate media (LPM)	25
3.2.2 High phosphate media	25
3.2.3 Low phosphate Agar	25
3.3 Isolation of Mo-Reducing Bacteria	26
3.4 Scanning Spectra of Molybdenum-blue	26
3.5 Dialysis tubing experiment	27
3.6 Identification of Mo-Reducing Bacteria	27
3.6.1 Gram Staining	27
3.6.2 Catalase Test	28
3.6.3 Oxidase test	28
3.6.4 16s rRNA Gene Sequencing	28
3.6.5 Phylogenetic Analysis	28
3.7 Effect of Various Parameter on Molybdate reduction MIE2 using One Factor at a Time (OFAT).	29
3.7.1 Effect of different Electron Donor Sources	29
3.7.2 Effects of electron donor concentrations	30
3.7.3 Screening of nitrogen sources	30
3.7.4 Effects of nitrogen source concentrations	30
3.7.5 Effect of LPM initial pH	31
3.7.6 Effect of Temperature	31
3.7.7 Effect of Phosphate and Molybdate concentration	31
3.8 Optimization of molybdate reduction using Response Surface Methodology (RSM)	32
3.8.1 Screening of significant parameters using Placket-Burman	32
3.8.2 Optimization of significant parameter using Box-Behken Design	33
3.9 Modeling kinetic of molybdate reduction	33
3.9.1 Determination of kinetic parameters for molybdate reduction to molybdenum blue	34
3.10 Purification of Mo-reducing enzyme	35
3.10.1 Preparation of crude enzyme	36
3.10.2 Protein assay	36
3.10.3 Definition of enzyme activity	37
3.10.4 Mo-reducing enzyme stability studies	37
3.10.4.1 Effect of storage pH	39
3.10.4.2 pH stability studies	39
3.10.4.3 Effect of temperature of storage	39
3.10.4.4 Temperature stability studies	39
3.10.4.5 Effect of chelating agent, detergents and organic solvents	40
3.10.4.6 Effect of sulfhydryl agents	40
3.10.4.7 Effect of metal ion cofactors	41
3.10.4.8 Effect of heavy metals ion	41

3.10.4.9	Effect of coenzymes	41
3.10.4.10	Effect of Inhibitors	42
3.10.5	Purification by Ammonium Sulphate Fractionation	42
3.10.6	Purification by Agilent Zorbax™ (G-250) Gel Filtration	43
3.10.7	Purification by Agilent Zorbax™ (GF-450) Gel Filtration	44
3.10.8	SDS-Polyacrylamide Gels Preparation	44
3.10.8.1	Preparation of Samples and Running the Gel	46
3.10.8.2	Gel Staining	46
3.10.9	Characterization of Mo-Reducing enzyme	47
3.10.9.1	Effect of pH on Molybdenum-Reducing Enzyme Activity	47
3.10.9.2	Effect of Temperature on Molybdenum-Reducing Enzyme Activity	47
3.10.10	Molybdenum reducing enzyme kinetics	47
3.10.10.1	K_m and V_{max} NADH as the substrate Electron Donor	48
3.10.10.2	K_m and V_{max} LPPM as the substrate Electron Donor	48
3.10.10.3	Determination of K_{cat}/K_m	48
3.10.11	Protein spots for MALDI-TOF peptide mass	48
4.0	RESULTS & DISCUSSIONS	49
4.1	Isolation and screening molybdenum reducing bacteria	49
4.2	Scanning Spectra of Molybdenum-blue produced by <i>Serratia</i> sp. MIE2	49
4.3	Dialysis Tubing experiment	50
4.4	Identification of Isolate TPU 3	52
4.5	Characterization and optimization of molybdate reduction using one at a time approach	56
4.5.1	Screening of electron donors	56
4.5.2	Effect of sucrose concentration	57
4.5.3	Screening of nitrogen sources	59
4.5.4	Effects of different ammonium sulphate concentrations	60
4.5.5	Effect of initial pH	61
4.5.6	Effect of temperature	63
4.5.7	Effect of phosphate concentration	64
4.5.8	Effect of molybdate concentration	65
4.5.9	Modeling molybdate reduction kinetics after one-factor-at-a-time method (or OFAT)	66
4.6	Optimization of molybdate reduction using Response Surface Methodology (RSM)	69
4.6.1	Screening significant parameter using Plackett-Burman	69
4.6.2	Optimization of molybdate reduction using RSM	72
4.6.3	Determination and validation of optimum conditions	75
4.6.4	Effect of process variables using response surface plot	77
4.6.4.1	Molybdate vs. phosphate concentration	77
4.6.4.2	Molybdate vs. sucrose concentrations	78
4.6.4.3	Molybdate concentration vs. pH	79
4.6.4.4	Phosphate concentration vs. pH	80

4.6.4.5 Phosphate vs. sucrose concentration	82
4.6.4.6 pH vs sucrose concentration	83
4.6.5 Modeling of molybdate reduction kinetics after RSM	84
4.7 Purification of Mo-reducing enzyme from <i>Serratia</i> sp.MIE2	87
4.7.1 Mo-reducing enzyme stability studies	87
4.7.1.1 Effect of storage pH	87
4.7.1.2 pH stability studies.	88
4.7.1.3 Effects of storage temperature	90
4.7.1.4 Temperature stability studies.	91
4.7.1.5 Effect of Chelating Agents, Detergents and Organic Solvents	92
4.7.1.6 Effect of Sulfhydryl Agents	94
4.7.1.7 Effect of Cofactors	95
4.7.1.8 Effect of Heavy metals	95
4.7.1.9 Effect of Coenzymes	96
4.7.1.10 Effect of Inhibitors	97
4.7.2 Purification of the Mo-reducing enzyme	99
4.7.2.1 SDS Polyacrylamide Gel Electrophoresis	102
4.7.3 Characterization and kinetics studies	104
4.7.3.1 Effect of pH on Mo-reducing enzyme activity	104
4.7.3.2 Effect of temperature on Mo-reducing enzyme activity	105
4.7.3.3 Determination of Kinetic Parameter	106
4.7.4 Peptide masses application in databases mining for protein identifications	108
5.0 SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	110
REFERENCES	112
APPENDICES	128
BIODATA OF STUDENT	134
LIST OF PUBLICATION	135

LIST OF TABLES

Table		Page
1:	List of isolated Mo- reducing bacteria.	13
2:	Levels of variables tested in Plackett-Burman Design.	33
3:	Levels of variables tested in Box Behnken Design.	33
4:	Various kinetic models for effect of substrate on Mo-blue production.	35
5:	Solution for preparing 12 % resolving gels for Tris-glycine SDS-polyacrylamide Gel electrophoresis.	45
6	Solution for preparing 5% stacking gels for Tris-glycine SDS-polyacrylamide Gel electrophoresis	46
7:	Parameter estimation for different substrate–inhibition models.	68
8:	Plackett–Burman design to evaluate significant parameters influencing molybdate reduction of <i>Serratia</i> sp. MIE2.	70
9:	ANOVA of Plackett-Burman, Factorial Model and Analysis of variance for molybdate reduction.	72
10:	Box-Behnken matrix for experimental design and predicted response using RSM.	73
11:	Analysis of variance (ANOVA) for the fitted quadratic polynomial model for optimization of molybdate reduction of <i>Serratia</i> sp. MIE2.	75
12:	Predicted and experimental value for the responses at optimum condition using response surface methodology.	76
13:	Summary of optimum condition of <i>Serratia</i> sp. MIE2 using One Factor at a Time (OFAT).	76
14:	Parameter estimates and regression for different substrate–inhibition models.	86
15:	Effects of storage temperature on Mo-reducing enzyme storage stability.	91
16:	Effect of chelating agents, detergents and organic solvents on Mo-reducing enzyme storage stability.	93
17:	Fractionation with ammonium sulphate.	100



© COPYRIGHT UPM

LIST OF FIGURES

Figure		Page
1:	Scanning spectra of Mo-blue from molybdosilicate (A), molybdosulphate (B) and molybdophosphate (C).	5
2:	The chemical structure of 12-phosphomolybdate.	6
3:	The chemical structures of various cofactors of xanthine oxidase (from North, 1998).	8
4:	A schematic presentation of the mechanism of molybdate reduction to Mo-blue by EC 48 (modified from Ghani <i>et al.</i> 1993; Zulhisham, 1993; Danny, 1997).	16
5:	Scanning spectra of molybdenum blue from <i>Serratia marcescens</i> strain Dr.Y10, <i>Acinetobacter</i> sp. strain Dr.Y12, <i>Klebsiella oxytoca</i> strain Dr.Y14 and <i>Serratia</i> sp. strain Dr.Y13 labelled 1, 2, 3 and 4 respectively in comparison with ascorbic acid-reduced phosphomolybdate (uppermost curve) (Shukor <i>et al.</i> , 2006).	19
6:	Scanning spectrum of Mo-blue from <i>Serratia</i> sp. MIE2 after 24 hours of incubation time.	50
7:	Enzymatic reduction of molybdate to Mo-blue in dialysis tube.	52
8:	Phylogram (neighbour-joining method) showing the genetic relationship between <i>Serratia</i> sp. MIE2	54
9:	NCBI deposition 16S rRNA gene sequence of MIE2.	55
10:	Effects of different carbon sources on molybdate reduction of <i>Serratia</i> sp. MIE2.	57
11:	Effects of sucrose concentration on molybdate reduction of <i>Serratia</i> sp. MIE2.	58
12:	Effects of different nitrogen sources on molybdate reduction of <i>Serratia</i> sp. MIE2.	60
13:	Effects of different ammonium sulphate concentrations on molybdate reduction of <i>Serratia</i> sp. MIE2.	61
14:	Effects of different pHs on molybdate reduction of <i>Serratia</i> sp. MIE2.	62
15:	Effects of different temperature on molybdate reduction of <i>Serratia</i> sp. MIE2.	63

16:	Effects of phosphate concentration on molybdate reduction of <i>Serratia</i> sp. MIE2.	64
17:	Effects of molybdenum concentration on molybdate reduction of <i>Serratia</i> sp. MIE2.	65
18:	Effect of different concentration of molybdate on molybdate reduction. Molybdate reduction were measured at wavelength 865nm every for 2 hours until 24 hours.	66
19:	Comparison of molybdate reduction kinetic experimental values with 7 different kinetic models from literature.	68
20:	3D plot (a) and contour plot (b) showing effect of molybdate concentration and phosphate on Mo-blue production.	78
21:	3D plot (a) and contour plot (b) showing effect of molybdate concentration and sucrose concentration on the molybdate reduction.	79
22:	3D plot (a) and contour plot (b) showing effect of molybdate concentration and pH concentration on the molybdate reduction.	80
23:	3D plot (a) and contour plot (b) showing effect of phosphate concentration and pH on the molybdate reduction	81
24:	3D plot (a) and contour plot (b) showing effect of phosphate concentration and sucrose concentration on the molybdate reduction.	82
25:	3D plot (a) and contour plot (b) showing effect of phosphate concentration and sucrose concentration on the molybdate reduction.	83
26:	Effect of different concentration of molybdate on molybdate reduction. Molybdate reduction were measured at wavelength 865nm every for 2 hours until 24 hours.	84
27:	Comparison of molybdate reduction kinetic experimental values with 7 different kinetic models from literature.	86
28:	Effect of pH and types of buffer on Mo-reducing enzyme storage stability.	83
29:	The effects of preincubation pHs of 4 (z), 5 ({), 6 (S), 7 (U), 7.5 ('), 8 (..), 9 (,,) and 10 (;) on the stability of the Mo-reducing enzyme.	89

30:	The effects of preincubation temperatures of 25 °C (,,), 34 °C (...), 40 °C (z), 54 °C (S) and 70 °C ({) on the stability of the enzyme.	92
31:	Effect of the sulfhydryl reagents DTT and 2-mercaptoethanol on the Mo-reducing enzyme storage stability.	94
32:	The effect of potential cofactors to the storage stability of the Mo-reducing enzyme.	95
33:	The effect of heavy metals to the storage stability of the Mo-reducing enzyme.	96
34:	Effect of a cofactor, FAD, on Mo-reducing enzyme storage stability. The error bars represent mean \pm standard deviation for three replicates.	97
35:	Effect of inhibitors to the Mo-reducing enzyme	98
36:	Gel filtration on Zorbax GFC-250. Shaded region indicates Mo-reducing enzyme activity. Each tube number or fraction represents 1 ml.	101
37:	Gel filtration on Zorbax GFC-450. Shaded region indicates Mo-reducing enzyme activity. Each tube number or fraction represents 1 ml.	101
38:	SDS-PAGE electrophoretogram of purified Mo-reducing enzyme stained by Coomassie blue method.	102
39:	Reducing SDS-PAGE electrophoretogram of purified Mo-reducing enzyme stained by Coomassie staining method	103
40:	Effect of pH on enzyme activity using a 25 mM citrate-phosphate buffer system. Error bars are mean \pm standard deviation of triplicates.	105
41:	Effect of temperature on enzyme activity. Error bars are mean \pm standard deviation of triplicates.	106
42:	Michaelis-Menten plot of amount of Molybdenum-blue formed (nmole/min/mg protein) versus the electron donor substrate (NADH).	107
43:	Michaelis-Menten plot of amount of Molybdenum-blue formed (nmole/min/mg protein) versus the electron acceptor substrate (LPPM).	108
44:	List of identified peptide fingerprint	109


LIST OF ABBREVIATIONS

%	percentage
EDTA	ethylene diamine tetraacetic acid
kDa	kiloDalton
M	molarity
min	minute
mM	milimolar
°C	degree Celcius
SDS	sodium dodecyl sulfate
U	unit
µg	microgram
UV	ultraviolet
Abs	Absorbance
et al.	and all
g	gram
HCl	hydrochloric acid
mg	Milligram
mL	milliliter
L	liter

uL	microliter
w	weight
mg/ L	milligram per liter
DNA	Deoxyribonucleic acid
g/cm ³	gram per centimeter cube

g/L	gram per volume
V/V	volume per volume
HPLC	High performance liquid chromatography
MW	molecular weight
AAS	Atomic absorption spectrometry
mg/L	milligram per litre
µg/L	microgram per litre
WHO	World Health Organization
EPA	Environmental Protection Agency
SEM	standard error of the mean
MPL	Maximum Permissible Limit
DOE	Department of Environment
OD	Optical density
n.d	not detected

N/A	not available
ppm	parts per million
OFAT	One Factor at a Time
RSM	Response Surface Methodology
A	Absorbance



Mo

Molybdenum

HP

Hypothetical Protein

AAN

Artificial neural Network

CHAPTER 1

1.0 INTRODUCTION

Heavy metals pollution is a silent threat that has affected water bodies and soils all around the world (Rajkumar et al., 2012; Shukor et al., 2009a; Zakaria et al., 2007; Fadzilah et al., 2014; Sany et al., 2013). There has been a growing concern over public health by heavy metals contamination (Chen et al., 2013; Wee et al., 2014). Molybdenum is one of the essential heavy metals that are required at trace amount and toxic at certain concentration (Othman et al., 2013; Halmi et al., 2013; Yamaguchi et al., 2007; Meeker et al., 2008). Hexavalent molybdenum (Mo^{6+}) exhibit toxic properties due to its solubility in water compared to molybdenum blue which is insoluble in water, thus exhibiting nontoxic properties and limited environmental disruption (Raab & Feldmann, 2003; Lloyd, 2003).

Molybdenum has many important functions in various applications. Molybdenum is a valuable alloying agent that inhibits corrosion in water-base hydraulic systems and automobile engine anti-freeze (Ilevbare & Burstein, 2001). Molybdenum replaces chromium for inhibition of corrosion in mild steel over a wide range of pH (Twite & Bierwagen, 1998). Molybdenum is used due to its low toxicity and is a less aggressive oxidant towards organic additives (Philip, 1992). Another common use of molybdenum is as lubricant in the form of molybdenum disulphide (Lansdown, 1999).

The wide application of molybdenum in industry has resulted in several water pollution cases all around the world such as in the Tokyo Bay and the Black Sea, Japan (Davis, 1991) and Tyrol in Austria (Neunhäuserer et al., 2001), where molybdenum level reaches in the hundreds of ppm. Poland is the latest case where molybdenum reached as high as 10 ppm in soil in Silesian Upland (%XUHü HW DO 2013). In Malaysia, molybdenum is mined as a byproduct of copper and molybdenum mining area in Sabah and there have been episodic cases of pollution in the surrounding area (Yong, 2000).

Molybdenum is very toxic to ruminants with levels as low as several parts per million causing scouring and even deaths (Greenwood and Earnshaw, 1984; Stojek, 2013). It was discovered that molybdenum shows its toxicity by inhibiting spermatogenesis in catfish and mice at levels as low as several parts per million (Yamaguchi et al., 2007; Zhai et al., 2013; Bi et al., 2013; Zhang et al., 2013). This new findings would increase molybdenum exposure as a toxic heavy metals similar to chromium and would increase the number of works on its removal from soil and water bodies. In the past decades researchers have focused on bioremediation as an environmental friendly and low cost method to solve this problem.

Bioremediation is one of the ways to remove toxic metals from the environment (Sar et al., 2013). A variety of molybdenum reducing bacteria has been reported with all

of them required a semi-aerobic condition for maximal production of molybdenum blue (Campbell et al., 1985; Ghani et al., 1993; Shukor et al., 2008; Shukor et al., 2009c; Shukor et al., 2009d; Abo-Shakeer et al., 2013; Shukor et al., 2009a; Shukor et al., 2010a; Ahmad et al., 2013; Shukor et al., 2009e; Shukor et al., 2010b; Othman et al., 2013.). According to Levine, Molybdate reduction was first reported in 1896 by Capaldi and Proskauer (Levine, 1925; Capaldi & Proskauer, 1896). Since then, many more reducers have been isolated (Ghani et al., 1993; Shukor et al., 2008; Shukor et al., 2009a-2009d; Shukor et al., 2010a-2010b). The first successful molybdenum remediation was carried out on an agricultural soil contaminated with molybdenum in Tyrol, Austria. Cows grazing on this soil showed signs of molybdenum toxicity or molybdenosis. The toxicity is actually a Cu deficiency, since Mo decreases Cu uptake in ruminants. The use phytoremediation and microbes from sewage and from the soil itself manages to immobilize the molybdenum into nonsoluble form ultimately reducing its toxicity (Neunhauserer et al., 2001).

Despite this, all of the molybdenum-reducing bacterium isolated so far is not from agricultural soil while molybdenum is particularly very toxic to ruminants. In addition, genetic and strain improvement of the Mo-reducing activity from potent Mo-reducing bacterium using biotechnology would enhance the remediation process. Previously, the first Mo-reducing enzyme was purified from *Serratia* sp. strain DrY5 (Shukor et al., 2014). However the yield of the purified enzyme was very low and prevents identification through sequencing process. To solve this problem, a novel Mo-reducing bacterium isolated from agricultural soil and screened for high Mo-reducing activity is needed. The identification, physiological and biochemical characterization of the isolated bacterium as well as the purification of the Mo-reducing enzyme will be carried out.

1.1 Thesis Objectives

Based on the problem statement and significant of the study, the following objectives are outlined:

1. To isolate and characterize a novel Mo-reducing bacterium from agriculture soil
2. To optimize Mo-blue production through one-factor-at-a-time (OFAT) and Response Surface Methodology (RSM)
3. To determine the kinetics of Mo-blue production in the bacterium before and after RSM
4. To determine the effect of storage pH and temperature, metabolic inhibitor, coenzyme and metal ions on Mo-blue production in the bacterium
5. To purify characterize and identify the Mo-reducing enzyme from the bacterium

REFERENCES

- Abo-Shakeer, L.K.A., Ahmad S.A., Shukor M.Y., Shamaan N.A., and M.A. Syed. 2013. Isolation and characterization of a molybdenum-reducing *Bacillus pumilus* strain lbna. *Journal of Environmental Microbiology and Toxicology* 1(1): 9-14.
- Ahmad, W. A., Zakaria, Z. A., Zakaria, Z., and Surif, S. 2009. Hexavalent Chromium Reduction at Different Growth Phases of *Acinetobacter haemolyticus*. *Environmental Engineering Science* 26(7): 1275-1278.
- Ahmad, S. A., Shukor, M. Y., Shamaan, N. A., Mac Cormack, W. P., & Syed, M. A. 2013. Molybdate Reduction to Molybdenum Blue by an Antarctic Bacterium. *BioMed research international* Article ID 87194.
- Ahmad, W. A., and Mohammed, N. 2010. Application of Response Surface Methodology (RSM) for optimizing removal of Cr (VI) wastewater using Cr (VI)-reducing biofilm systems. *Malaysian Journal of Fundamental and Applied Sciences* 6(1).
- Aiba, S., Shoda, M. and Nagalani, M. 1968. Kinetics of product inhibition in alcohol fermentation. *Biotechnology and Bioengineering* 10, 845±864.
- Annadurai, G., Ling, L. Y., and Lee, J.F. 2008. Statistical optimization of medium components and growth conditions by response surface methodology to enhance phenol degradation by *Pseudomonas putida*. *Journal of hazardous materials* 151(1): 171-178.
- Angelo A J & Kuck J C. 1977, Effects of cyanide on peanut lipoxygenase, *Lipids* 12 682±683.
- Ariff, A. B., Rosfarizan, M., Ghani, B., Sugio, T. and Karim, M. I. A. 1997. Mo-reducing enzyme in *Enterobacter cloacae* strain 48. *World Journal of Microbial Biotechnology* 13: 643-647.
- Balusu, R., Paduru, R. R., Kuravi, S. K., Seenayya, G., and Reddy, G. 2005. Optimization of critical medium components using response surface methodology for ethanol production from cellulosic biomass by *Clostridium thermocellum* SS19. *Process Biochemistry* 40(9): 3025-3030.
- Beyenal. H., Chen S.N. and Lewandowski Z. 2003. The double substrate growth kinetics of *Pseudomonas aeruginosa*. *Enzyme and Microbial Technology* 32; 92- 98.
- Bi, C. M., Zhang, Y. L., Liu, F. J., Zhou, T. Z., Yang, Z. J., Gao, S. Y. And Wang, S. 2013. The effect of molybdenum on the in vitro development of mouse preimplantation embryos. *Systems biology in reproductive medicine* 59(2), 69-73.

- Box, G. E., and Lucas, H. 1959. Design of experiments in non-linear situations. *Biometrika* 46(1/2): 77-90.
- Box, G. E., and Wilson, K. 1951. On the experimental attainment of optimum conditions. *Journal of the Royal Statistical Society. Series B (Methodological)* 13(1): 1-45.
- Bradford, M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-252.
- Braithwaite, E. R. 1981. Molybdenum. R. Thompson, ed., Specialty Inorganic Chemicals. The Royal Society of Chemistry, Burlington House, London W1V 0BN. pp. 350-351.
- %XUHUHZQLDN : -DUR , .XFKDU] yk, J., Narkiewicz, W., & Pasieczna, A. 2013. Investigation of Tin and Molybdenum concentrations in the Soils in the southern part of the Silesian Upland. In *E3S Web of Conferences* (Vol. 1, p. 08005). EDP Sciences.
- Capaldi, A., Proskauer, B., 1896. Beiträge zur Kenntnis der Säurebildung bei Typhusbacillen und Bacterium Coli. *Z. Hyg. Infekt.-Kr.*, 23: 452±474.
- Carpentier. W., Sandra. K., De Smet. I., Brige. A., De Smet. L., Van Beeumen. J., 2003. Microbial reduction and precipitation of vanadium by *Shewanella oneidensis*. *Applied and environmental microbiology* 69: 3636-3639.
- Campbell, M. A., Campbell, A. D. and Villaret, D. B. 1985. Molybdate reduction by *Eschericia coli* K-12 and its *chl* mutants. *Proceeding of the Natural Academy of Science, USA* 82: 227-231.
- Campbell, W. H. and Kinghorn, K. R. 1990. Functional domains of assimilatory of nitrate reductases and nitrite reductases. *Trends in Biochemical Sciences* 15: 315-319.
- Cappuccino, J.G. and Sherman, N. 2005. *Microbiology: A Laboratory Manual* 7th Edition. Pearson Education, Inc, San Francisco. pp 45-47.
- Camargo, F. A. O., Bento, F. M., Okeke, B. C., and Frankenberger, W. T. 2003. Chromate reduction by chromium-resistant bacteria isolated from soils contaminated with dichromate. *Journal of environmental quality* 32(4): 1228-1233.
- Caruso, F., and Schüler, C. 2000. Enzyme multilayers on colloid particles: assembly, stability, and enzymatic activity. *Langmuir* 16(24): 9595-9603.
- Chen, Y., Hu, W., Huang, B., Weindorf, D. C., Rajan, N., Liu, X., and Niedermann, S. 2013. Accumulation and health risk of heavy metals in vegetables from harmless and organic vegetable production systems of China. *Ecotoxicology and environmental safety* 98: 324-330.

- Clesceri, L. S., Greenberg, A. E. and Trussel, R. R. 1989. Standard methods for the examination of wastewater. American Public Health Association, Port City Press, Baltimore, Maryland. 4-166.
- Cotton, F. A., Wilkinson, G., Murillo, C. A., Bochmann, M., and Grimes, R. 1988. Advanced inorganic chemistry (Vol. 5). New York: Wiley.
- Counotte, G. H. M., and Prins, R. A. 1979. Calculation of K_m and V_{max} from substrate concentration versus time plot. *Applied and environmental microbiology* 38(4): 758-760.
- Costa, M., and Klein, C. B. 2006. Toxicity and carcinogenicity of chromium compounds in humans. *CRC Critical Reviews in Toxicology* 36(2): 155-163.
- Cull, M. and McHenry, C.S. 1990. Preparation of extracts from prokaryotes. Deutscher, ed., *Methods in Enzymology, Guide to Protein Purification*. Academic Press, San Diego. 182:425-477.
- Czyryca, E. J. 1993. Advances in high strength steel technology for naval hull construction. *Key Engineering Materials*, 84, 491-520.
- Dancis, A., Roman, D. G., Anderson, G. J., Hinnebusch, A. G., and Klausner, R. D. 1992. Ferric reductase of *Saccharomyces cerevisiae*: molecular characterization, role in iron uptake, and transcriptional control by iron. *Proceedings of the National Academy of Sciences* 89(9): 3869-3873.
- Davis, G. K. 1991. Molybdenum. In: Merian, E. (Ed). *Metals and their compounds in the environment, occurrence, analysis and biological relevance*, VCH Weinheim. New York. 1089±1100.
- Dawson, R.M.C, D.C. Elliot and W.H. Elliot. 1969. 'DWDIRUELRFKHPLFDOUHVHDFK' Clarendon Press, Oxford. pp 66.
- Dewan Bahasa dan Pustaka. 1977. Atlas Kebangsaan Malaysia. Kuala Lumpur. pp 60.
- Dey, S., and Paul, A. K. 2014. Reduction of Hexavalent Chromium by Immobilized Viable Cells of *Arthrobacter* sp. SUK 1201. *Bioremediation Journal* 18(1): 1-11.
- Desai, K. M., Survase, S. A., Saudagar, P. S., Lele, S., and Singhal, R. S. 2008. Comparison of artificial neural network (ANN) and response surface methodology (RSM) in fermentation media optimization: case study of fermentative production of scleroglucan. *Biochemical Engineering Journal*, 41(3): 266-273.
- Devereux, R. and Wilkinson, S. S. 2004. Amplification of ribosomal RNA VHTXHQFHV',Q \$NHUPDQV \$9DQ (OVDV -' DQG 'H %UXLMQ)- HGV Molecular microbial ecology manual. 2nd ed. Kluwer Academic Publishing, Netherland, 1±17.

- Deutscher, M. P. (Ed.). 1990. Guide to protein purification. Gulf Professional Publishing. pp12.
- Deutscher, M. P. 1990. Setting up a laboratory. *Methods in Enzymology: Guide to Protein Purification*, Academic Press, San Diego. pp 19-24
- Diamantino. T.C., L. Guilhermino. L., Almeida. E. and Soares. A.M. 2000. Toxicity of Sodium Molybdate and Sodium Dichromate to *Daphnia magna* Straus Evaluated in Acute, Chronic, and Acetylcholinesterase Inhibition Tests. *Ecotoxicology and environmental Safety* 45: 253-259.
- Dinarvand, M., Rezaee, M., Masomian, M., Jazayeri, S. D., Zareian, M., Abbasi, S., & Ariff, A. B. 2013. Effect of C/N Ratio and Media Optimization through Response Surface Methodology on Simultaneous Productions of Intra-and Extracellular Inulinase and Invertase from *Aspergillus niger* ATCC 20611. *BioMed research international*.
- DOE, Environmental Quality Report. 2010, Department of Environment, Ministry of Science, Technology and the Environment, Malaysia. ISSN 0127-6433.
- Dolin, M. I. 1961. Cytochrome-independent electron transport enzymes of bacteria. I. C. Gunsalus and R. Y. Stainer, eds., *The Bacteria: a treatise of Structure and Function*, Academic Press, New York, pp 425-461
- Fadzilah, M. H. H., Tajam, J., Kamal, M. L., and Daim, N. 2014. Distribution of Heavy Metals, Organic Matter and Mean Size in Sediment at the Perlis River. In *From Sources to Solution*, Springer Singapore, pp. 507-511.
- Freedman. Z., Zhu. C. and Barkay. T. 2012. Mercury resistance and mercuric reductase activities and expression among chemotrophic thermophilic Aquificae. *Applied and Environmental Microbiology*, 78. 6568-6575.
- Felsenstein. J. 1985. Confidence limits on phylogenies. An approach using the bootstrap. *Evolution*, 39:783±791.
- Garbisu. C., Alkorta. I., Llama. M.J. and Serra. J.L. 1998. Aerobic chromate reduction by *Bacillus subtilis*. *Biodegradation*, 9(2): 133±141.
- Ghani, B., Takai, M., Hisham, N.Z, Kishimito, N., M.I.A. Ismail, T. Tano and T. Sugio. 1993. Isolation and characterization of a Mo⁶⁺ -UHGXFLQJ EDFWHULXP' *Applied and Environmental Microbiology*. 59: 1176±1180.
- Glenn J.L and Crane F.L. 1956. Studies on metalloflavoproteins. V. The action of silicomolybdate in the reduction of cytochrome c by aldehyde oxidase. *Biochim. Biophys. Acta*, 22:111±115.
- *XV[F] 3 3HWHUD - DQG /HGDNRZLF] 6 0DWKHPDWLFD0 PRGHOLQJ RI WKH integrated process of mercury bioremediation in the industrial bioreactor. *Bioprocess and Biosystems Engineering* 34(3): 275±285.

- Greenwood, N. N. and Earnshaw, A. 1984. Chemistry of the elements. Pergammon Press, Oxford. pp 1167.
- Gunawan, E. R., Basri, M., Rahman, M. B. A., Salleh, A. B. and Rahman, R. N. Z. A. 2005. Study on response surface methodology (RSM) of lipase-catalyzed synthesis of palm-based wax ester. *Enzyme and Microbial Technology*. 37(7): 739-744.
- Gupta, C. K. 1992. Extractive metallurgy of molybdenum. CRC press, pp 55.
- Halmi, M. I. E., Zuhainis, S. W., Yusof, M. T., Shaharuddin, N. A., Helmi, W., Shukor, M. Y and Ahmad, S. A. 2013. Hexavalent Molybdenum Reduction to Mo-Blue by a Sodium-Dodecyl-Sulfate-Degrading *Klebsiella oxytoca* Strain DRY14. *BioMed research international*.
- Han, K., and Levenspiel, O. 1988. Extended Monod kinetics for substrate, product, and cell inhibition. *Biotechnology and Bioengineering*, 32(4): 430-447.
- Harrison, R. 2002. Structure and function of xanthine oxidoreductase: where are we now?. *Free Radical Biology and Medicine* 33(6):774-797.
- Hem, J. D. 1972. Chemical factors that influence the availability of iron and manganese in aqueous solution. *Geological Society of America Bulletin*, 83: 443±450.
- Holm, R. H., Kennepohl, P. and Solomon, E. I. 1996. Structural and functional aspects of metal sites in biology. *Chemical Review* 2196-2239.
- Hori, T. Sugiyama, M and Himeno, S. 1988. Direct spectrophotometric determination of sulphate ion based on the formation of a blue molybdosulphate complex. *Analyst* 113: 1639 - 1644.
- Ilevbare. G., and Burstein G., 2001, The role of alloyed molybdenum in the inhibition of pitting corrosion in stainless steels. *Corrosion Science* 43:485-513.
- Ilias, M., Rafiqullah, I. M., Debnath, B. C., Mannan, K. S. B., & Hoq, M. M. 2011. Isolation and characterization of chromium (VI)-reducing bacteria from tannery effluents. *Indian journal of microbiology* 51(1): 76-81.
- Jan. A. 1939. La reduction biologique du molybdate d'ammonium par les bactéries du genre *Serratia*. *Bulletin des Sciences Pharmacologiques*, 46 : 336±339.
- Jeffrey, J. 1980. Kinetic aspects of soluble dehydrogenases requiring nicotinamide coenzymes. J. Jeffrey, ed., *Dehydrogenases Requiring Nicotinamide Coenzymes*. Birkhäuser, Verlag. Basel, pp 100-150.
- Johnston, J.J., R.C. Borden and M.A. Barlaz. 1996. Anaerobic biodegradation of alkylbenzenes and trichloroethylene in aquifer sediment down gradient of a sanitary landfill. *Journal of Contamination and Hydrology* 23: 263-283.

- Kazansky, L. P. and Fedotov, M. A. 1980. Phosphorous-31 and oxygen-17 n.m.r. evidence of trapped electrons in reduced 18-molybdodiphosphate (v), $P_2Mo_{18}O_{62}^{8-}$. *Journal of the Chemical Society, Chemical Communication* 13: 644 ± 647.
- Kaiser, B. N., Gridley, K. L., Brady, J. N., Phillips, T., and Tyerman, S. D. 2005. The role of molybdenum in agricultural plant production. *Annals of botany*, 96(5): 745-754.
- Kazansky, L. P. and Fedotov, M. A. 1980. Phosphorous-31 and oxygen-17 n.m.r. evidence of trapped electrons in reduced 18-molybdodiphosphate (v), $P_2Mo_{18}O_{62}^{8-}$. *Journal of the Chemical Society, Chemical Communication* 13: 644 ± 647.
- Khuri, A. I., and Mukhopadhyay, S. 2010. Response surface methodology. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(2): 128-149.
- Kim, J. and Rees, D. C. 1992. Crystallographic structure and functional implications of the nitrogenase molybdenum-iron protein from *Azotobacter Vinelandii*. *Nature* 360: 553-560.
- Kimball, B. A., Nordstrom, D., Runkel, R. L., Vincent, K. R., & Verplanck, P. L. (2006). Questa baseline and pre-mining ground-water quality investigation. 23. Quantification of mass loading from mined and unmined areas along the Red River, New Mexico. U. S. Geological Survey.
- Kwon-Hee Lee, Gyung-Jin Park and Won-Sik Joo. 2005. A Global Robust Optimization Using the Kriging Based Approximation Model. 6th World Congresses of Structural and Multidisciplinary Optimization Rio de Janeiro, 30 May - 03 June 2005, Brazil.
- Lansdown. A. 1999. Molybdenum disulphide lubrication, Access Online via Elsevier.
- LangÅrd, S. and Lison, D. 2012. Chromium, Molybdenum, and Tungsten. *Patty's Toxicology* 106, 176-185.
- Levine, V. E. 1925. The reducing properties of microorganisms with special reference to selenium compounds, *Journal of Bacteriology*, 10, 217±263.
- Lee, J. D. 1977. Concise inorganic chemistry. Van Reinhold Co. pp 325. New York.
- Lepora, N. 2006. *Molybdenum*. Marshall Cavendish. pp: 4-6
- Lim, H. K, M.A. Syed, M.Y. Shukor. 2012. Reduction of molybdate to molybdenum blue by *Klebsiella V S* VWU DL *Qoukud IdfBBSic Microbiology*, 52(3): 296±305,
- Lloyd, J. R. 2003. Microbial reduction of metals and radionuclides. *FEMS microbiology reviews*, 27(2-3): 411-425.
- Lovley, D. R. 1993. Dissimilatory metal reduction. *Annual Reviews in Microbiology* 47: 263-290.

- Luong, J. H. T. 1985. Kinetics of ethanol inhibition in alcohol fermentation. *Biotechnology and bioengineering*, 27(3): 280-285.
- Mamatha, S. S., Ravi, R., and Venkateswaran, G. 2008. Medium optimization of gamma linolenic acid production in *Mucor rouxii* CFR-G15 using RSM. *Food and Bioprocess Technology*, 1(4): 405-409.
- Margush, T. and F.R. McMorris. 1981. Consensus n-WUHH. *Bulletin of Mathematical Biology*, 43: 239±244.
- Mantle, T. J., and Noone, P. 1996. Chromatofocusing. In *Protein Purification Protocols*, Humana Press, pp. 249-254.
- Masood, F., and Malik, A. 2011. Hexavalent chromium reduction by *Bacillus* sp. strain FM1 isolated from heavy-metal contaminated soil. *Bulletin of environmental contamination and toxicology*, 86(1): 114-119.
- Meeker, J. D., Rossano, M. G., Protas, B., Diamond, M. P., Puscheck, E., Daly, D. and Wirth, J. J. 2008. Cadmium, lead, and other metals in relation to semen quality: human evidence for molybdenum as a male reproductive toxicant. *Environmental health perspectives*, 116(11): 1473.
- Mendel, R. R., & Kruse, T., 2012. Cell biology of molybdenum in plants and humans. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* 1823(9), 1568-1579.
- Misra, T. K. 1992. Bacterial resistances to inorganic mercury salts and organomercurials. *Plasmid*, 27(1): 4-16.
- Miyazaki, K., Wintrode, P. L., Grayling, R. A., Rubingh, D. N., and Arnold, F. H. 2000. Directed evolution study of temperature adaptation in a psychrophilic enzyme. *Journal of molecular biology*, 297(4): 1015-1026.
- Molecular Simulations Inc. 2000. Heteropolymolybdate Salts-- A New Structural Model.
- Montgomery, D. C. 1984. *Design and analysis of experiments*. Wiley New York.
- Monod, J. 1949. The Growth of Bacterial Cultures. *Annual Review of Microbiology*. 3: 371±394.
- Mulchandani, A, J. Luong. 1989. Microbial inhibition kinetics revisited. *Enzyme and microbial technology* 11: 66-73.
- Munch, J. C and Ottow, J. C. G. 1983. Reductive transformation mechanism of ferric oxides in hydromorphic soils. *Environmental and Biogeochemical Ecology Bulletin (Stockholm)* 35: 383-394.

Mussatto, S. I., Machado, E. M., Martins, S., and Teixeira, J. A. 2011. Production, composition, and application of coffee and its industrial residues. *Food and Bioprocess Technology* 4(5): 661-672.

Müller, A., Meyer, J., Krickemeyer, E., and Diemann, E. 1996. Molybdenum blue: A 200 year old mystery unveiled. *Angewandte Chemie International Edition in English*, 35(11): 1206-1208.

000HU \$DQG 6HUDLQ & 6ROXEOH PROEGHQXP EOXHV GHV SXGHOV NHUQ
. *Accounts of chemical research*, 33(1): 2-10.

Myers, R. H., and Anderson-Cook, C. M. 2009. *Response surface methodology: process and product optimization using designed experiments*, John Wiley & Sons. 705, 73-153.

Neunhäuserer, C. M., Berreck. and Insam, H. 2001. Remediation of Soils Contaminated with Molybdenum using Soil Amendments and Phytoremediation. *Water Air Soil Pollution* 128: 85-96.

Ng, T. W., Cai, Q., Wong, C.K., Chow, A. T., and Wong, P.K. 2010. Simultaneous chromate reduction and azo dye decolourization by *Brevibacterium casei* Azo dye as electron donor for chromate reduction. *Journal of hazardous materials*, 182(1): 792-800.

North, A. C, 1988. Xanthine oxidase family, [http:// www.Metallo.scripps.edu/promise/xantoxidase.html](http://www.Metallo.scripps.edu/promise/xantoxidase.html).

Opperman. D.J., Piater L.A. and van Heerden. E. (2008). A novel chromate reductase from *Thermus scotoductus* SA-01 related to old yellow enzyme. *Journal of bacteriology*, 190: 3076-3082.

Osborne. F. H. and Ehrlich. H. L. 1976. Oxidation of arsenite by a soil isolate of *Alcaligenes*, *Journal of Applied Bacteriology*. 41(2): 295±305.

Othman, A. R., Bakar, N. A., Halmi, M. I. E., Johari, W. L. W., Ahmad, A., Jirangon, H., & Shukor, M. Y. 2013. Kinetics of Molybdenum Reduction to Molybdenum Blue by *Bacillus* sp. Strain A. rzi, *BioMed research international*.

Pal, A., Datta, S., and Paul, A. K. 2013. Hexavalent chromium reduction by immobilized cells of *Bacillus sphaericus* and 303. *Brazilian Archives of Biology and Technology*, 56(3): 505-512.

Padhiar, A. R., and Modi, H. A. 2013. Optimization Of Lipase Production By *Saccharomonospora Azurea* Ap 11/18 Using Plackett-Burman Design And Response Surface Methodology. *International Journal*, 3(1), 59-66.

Park, C. H., Keyhan, M., Wielinga, B., Fendorf, S., and Matin, A. 2000. Purification to homogeneity and characterization of a novel *Pseudomonas putida* chromate reductase. *Applied and Environmental Microbiology* 66: 1788-1795.

- Pei, Q. H., Shahir, S., Raj, A. S., Zakaria, Z. A., and Ahmad, W. A. 2009. Chromium (VI) resistance and removal by *Acinetobacter haemolyticus*. *World Journal of Microbiology and Biotechnology*, 25(6): 1085-1093.
- Philip. C. 1992. Molecular mechanics study of the interaction of thiophene with a molybdenum disulfide catalyst. *Journal of the Chemical Society, Faraday Transactions*, 88:3225-3232.
- Poole, R. K. 1983. Bacterial Cytochrome Oxidases: A Structurally and functionally diverse group of electron-transfer proteins. *Biochimica et Biophysica Acta* 726: 200-243.
- Privé, G. G. 2007. Detergents for the stabilization and crystallization of membrane proteins. *Methods* 41(4): 388-397.
- Prapulla, S. G., Jacob, Z., Chand, N., Rajalakshmi, D., and Karanth, N. G. 1992. Maximization of lipid production by *Rhodotorula gracilis* CFR-1 using response surface methodology. *Biotechnology and bioengineering*, 40(8): 965-970.
- Rajkumar, B., Sharma, G. D., and Paul, A. K. 2012. Isolation and Characterization of Heavy Metal Resistant Bacteria from Barak River Contaminated with Pulp Paper Mill Effluent, South Assam. *Bulletin of environmental contamination and toxicology*, 89(2): 263-268.
- Raab, A., and Feldmann, J. 2003. Microbial transformation of metals and metalloids. *Science progress*, 86(3): 179-202.
- Rajagopalan, K. V. 1980. Xanthine oxidase and aldehyde oxidase. Pp. 295-306 in W. B. Jakoby, ed., *Enzymatic Basis of Detoxification*. Academic Press, New York.
- Rajwade, J. M., Salunkhe, P. B., and Paknikar, K. M. (1999). Biochemical basis of chromate reduction by *Pseudomonas mendocina*. *Process Metallurgy* 9: 105-114.
- Rüssel, C., & Kämpfer, A. 1998. Electric melting of glass: Influence of cathodic currents on the formation of protective layers on molybdenum electrodes. *Glass science and technology*, 71(1), 6-11.
- Runnells, D.D., D.S. Kaback and E.M. Thurman, 1976. Geochemistry and sampling of molybdenum in Sediments, Soils Plants in Colorado. In: *Molybdenum in the Environment*, W.R. Chappel and K.K. Peterson (eds.). Marcel and Dekker, Inc, New York.
- Sany, S. B. T., Salleh, A., Rezayi, M., Saadati, N., Narimany, L., & Tehrani, G. M. (2013). Distribution and contamination of heavy metal in the coastal sediments of Port Klang, Selangor, Malaysia. *Water, Air, & Soil Pollution* 224(4), 1-18.

- Sar, P., Kazy, S. K., Paul, D., AND Sarkar, A. 2013. Metal Bioremediation by Thermophilic Microorganisms. In *Thermophilic Microbes in Environmental and Industrial Biotechnology*, Springer Netherlands, pp. 171-201
- Sau. G.B., Chatterjee. S. and Mukherjee. S.K. 2010. Chromate reduction by cell-free extract of *Bacillus firmus* KUCr1. *Polish Journal of Microbiology* 59:185±190.
- Saitou. N. And Nei. M. 1987. The neighbour-joining method: A new method for UHFRQVWUXFWLQJSKORJHQWLEFMEUHM. *Molecular Biology and Evolution* 4: 406±425.
- Scopes, R. K. 1988. Protein purification, principles and practice, Springer-Verlag, New York. pp 12.
- Schröder, I., Rech, S., Krafft, T., and Macy, J. M. 1997. Purification and characterization of the selenate reductase from *Thauera selenatis*. *Journal of Biological Chemistry*, 272(38): 23765-23768.
- Schwarz, G., Mendel, R. R., and Ribbe, M. W. 2009. Molybdenum cofactors, enzymes and pathways. *Nature*, 460(7257): 839-847.
- Schindelin, H., Kisker, C., and Rajagopalan, K. V. 2001. Molybdopterin from molybdenum and tungsten enzymes. *Advances in protein chemistry*, 58: 47-94.
- Scopes, R. K. 1994. Protein purification: principles and practice. Springer. pp 10-36.
- Seldén, A. I., Berg, N. P., Söderbergh, A., and Bergström, B. E. 2005. Occupational molybdenum exposure and a gouty electrician. *Occupational Medicine*, 55(2): 145-148.
- Shukor, M. Y. A., Shamaan, N. A., Syed, M. A., Lee, C. H. and Karim, M. I. A. 2000. Characterization and quantification of molybdenum blue production in *Enterobacter cloacae* Strain 48 using 12 molybdophosphate as the reference compound. *Asia Pacific Journal of Molecular Biology & Biotechnology*, 8(2): 167-172.
- Shukor, M. Y., Syed, M. A., Lee, C. H., Karim, M. I. A., and Shamaan, N. A. 2002. A method to distinguish between chemical and enzymatic reduction of molybdenum in *Enterobacter cloacae* strain 48. *Malaysian Journal of Biochemistry*, 7: 71-72.
- Shukor, A., Yunus, M., Lee, C. H., Omar, I., Karim, M. I. A., Syed, M. A., and Shamaan, N. A. 2003. Isolation and Characterization of a Molybdenum Reducing Enzyme in *Enterobacter cloacae* Strain 48. *Pertanika Journal of Science & Technology*, 11(2): 261-272.

- Shukor, M. Y, H. Adam, K. Ithnin, I. Yunus and N.A. Shamaan. 2007. Molybdate reduction to Mo-blue in microbe proceeds via a phosphomolybdate LQWHUPHGLDWHI of *Biological Sciences* 7:1448±1452.
- Shukor M.Y, Habib S.H.M., Rahman M.F.A., Jirangon H., Abdullah M.P.A., Shamaan N.A. and Syed M.A. 2008. Hexavalent molybdenum reduction to molybdenum blue by *S. marcescens* strain Dr.Y6. *Applied Biochemistry and Biotechnology*. 149(1): 33±43.
- Shukor M.Y., Rahman M.F.A., Suhaili Z., Mustafa S., Shamaan N.A., Syed M.A. 2009a. Reduction of Mo(VI) by the bacterium *Serratia* sp. strain DRY5. *Journal of Environmental Biology*, 30(1):65±72.
- Shukor M.Y., Rahman M.F., Shamaan N.A. and Syed M.A. 2009b. Reduction of molybdate to molybdenum blue by *Enterobacter* sp. strain DRY13. *Journal of Basic Microbiology*, 49:1±12.
- Shukor M.Y., Rahman M.F., Suhaili Z., Shamaan N.A. and Syed M.A. 2009c. Bacterial Reduction Of Hexavalent Molybdenum To Molybdenum Blue. *World Journal of Biotechnology and Microbiology* 25:1225±1234.
- Shukor M.Y., Hamdan M.H., Othaman M.A., Shamaan N.A. and Syed M.A. 2009d. Mo(VI) reduction to molybdenum blue by *S. marcescens* strain DRY9. *Polish Journal of Microbiology*. 58(2):141±147.
- Shukor, M. Y., Rahman, M. F., Shamaan, N. A., and Syed, M. A. 2009e. Reduction of molybdate to molybdenum blue by *Enterobacter* sp. strain Dr. Y13. *Journal of basic microbiology*, 49(S1), S43-S54.
- Shukor M.Y., Rahman M.F., Suhaili Z., Shamaan N.A. and Syed M.A 2010a. Hexavalent Molybdenum Reduction to Mo-blue by *Acinetobacter calcoaceticus*. *Folia Microbiology*. 55(2):137±143.
- Shukor MY, Ahmad SA, Abdullah MP, Shamaan NA, Syed MA. 2010b. Molybdate Reduction by *Pseudomonas* sp. Strain DRY2. *Journal of Applied Microbiology*. doi:10.1111/j.1365-2672.2009d.04604.x.
- Shukor, M. Y., and Syed, M. A. 2010C. Microbiological reduction of hexavalent molybdenum to molybdenum blue. *Current Research, Technology And Education Topics in Applied Microbiology and Microbial Biotechnology*, pp 1304-1310.
- Shukor. M.Y. Rahman. M.F., HalmI. M.I.E., Shamaan. N.A., and Syed. M.A., 2014. Molybdenum Reduction to Molybdenum Blue in *Serratia* sp. Strain DRY5 is Catalyzed by a Novel Molybdenum-Reducing Enzyme, *BioMed research international*. vol. 2014, Article ID 853084.

Shen. H. And Wang. Y. Characterization of enzymatic reduction of hexavalent chromium by *Escherichia coli* ATCC 33456. *Applied and Environmental Microbiology* 59: 3171±3777.

Shineldecker, C. L. 1992. Handbook of environmental contaminants: a guide to self assessment. Lewis Publisher, U.S.A. pp 72.

Shen, Hai, and Yi-Tin Wang, 1995. Simultaneous chromium reduction and phenol degradation in a coculture of *Escherichia coli* ATCC 33456 and *Pseudomonas putida* DMP-1. *Applied and environmental microbiology*, 61(7): 2754-2758.

Sharma, P., Singh, L., and Dilbaghi, N. 2009. Optimization of process variables for decolorization of Disperse Yellow 211 by *Bacillus subtilis* using Box± Behnken design. *Journal of hazardous materials*, 164(2): 1024-1029.

Sims, R. P. A. 1961. Formation of heteropoly blue by some reduction procedures used in the micro-determination of phosphorous. *Analyst* 86: 584 - 590.

Sidgwick, N. V. 1984. The chemical elements and their compounds. Clarendon Press, Oxford. pp 21-35.

Sinnakkannu S., Abdullah. A.R, Tahir. N.M and Abas. M.R.. 2004. Degradation of PHWVXOIXURQ PHWKQ LQ VHOHFWHG 0DODVLDQ DJULEXOWXUDO VRLO *Environmental Bulletin*, 13: 258±261.

Slifer, D. 1996. Red river groundwater investigation; Final report. http://www.amigosbravos.org/molywatch/rr_groundwater.html#3.2.2.6. Accessed on 16 May 1999.

Soda, S. O., Yamamura, S., Zhou, H., Ike, M., and Fujita, M. 2006. Reduction kinetics of As (V) to As (III) by a dissimilatory arsenate-reducing bacterium, *Bacillus* sp. SF-1. *Biotechnology and bioengineering* 93(4): 812-815.

Soo, E., Salleh, A. B., Basri, M., Rahman, R. A., and Kamaruddin, K. 2004. Response surface methodological study on lipase-catalyzed synthesis of amino acid surfactants. *Process Biochemistry* 39(11): 1511-1518.

Soni, S. K., Singh, R., Awasthi, A., Singh, M., & Kalra, A. 2013. In vitro Cr (VI) reduction by cell-free extracts of chromate-reducing bacteria isolated from tannery effluent irrigated soil. *Environmental Science and Pollution Research* 20(3), 1661-1674.

Srinath, T., Verma, T., Ramteke, P., and Garg, S. 2002. Chromium (VI) biosorption and bioaccumulation by chromate resistant bacteria. *Chemosphere*, 48(4): 427-435.

Stojek, M. 2013. The concentration of molybdenum and copper in rocks, soils and *SODQWVLQWKH DUHDRI-DEKQNL(DVWHUQ%HVNLGV0WV=DZDUWRüPROL PLHGJL Z VNDFK JOHEDFK L UROLQDFK Z RNROLF\ -DERQHN %HVNLG*

Wschodnie). *Qej tqpc" tqf qykumc" k" \cuqd»y" Pcwwtcnp{ej-Environmental Protection and Natural Resources*, 24(3), 13-17.

Stoll, V. S. and Blanchard, J. S. 1990. Buffers: principles and practice. in M.P. Deutscher, ed., *Methods in Enzymology Vol 182: Guide to Protein Purification*. Academic Press, San Diego, pp 24-37.

Sugio. T., Tsujita. Y., Katagiri. T., Inagaki. K. and Tano. T. 1988. Reduction of Mo^{6+} with elemental sulfur by *Thiobacillus ferrooxidans*. *Journal of Bacteriology* 170: 5956±5959.

Suzuki, T., Miyamata, N., Horitsu, H., Kawai, K., Takamizawa, T., Tai, Y., and Okazaki, M. 1992. NAD(P)H-dependent chromium (VI) reductase of *Pseudomonas ambigua* G-1: A Cr(V) Intermediate is formed during the reduction of Cr(VI) to Cr(III). *Applied and Environmental Microbiology* 174: 5340-5345.

Suhaila, Y. N., Ramanan, R. N., Rosfarizan, M., Latif, I. A., and Ariff, A. B. 2013. Optimization of parameters for improvement of phenol degradation by *Rhodococcus* UKMP-5M using response surface methodology. *Annals of Microbiology* 63(2): 513-521.

Sukumar. M. 2010. Reduction of hexavalent chromium by *Tjk/qrwu" Qt{/cgö*. *African Journal of Environmental Science and Technology* 4(7): 412±418.

Tendo Masayuki, Yutaka Tadokoro, Kazuhiro Suetsugu, and Takanori Nakazawa. 2001. Effects of nitrogen, niobium and molybdenum on strengthening of austenitic stainless steel produced by thermo-mechanical control process. *ISIJ international* 41(3)262-267.

Teissier. G. 1942. Croissance des populations EDFWHULHQHV HW TXDQWLWHGDPOLPH disponible (Growth of bacterial populations and the available substrate concentration), *Revision Science* 80:209,

7KRPSVRQ-' +LJLQV '*DQG *LEVRQ 7- &/867\$:LPSURYLQJ the sensitivity of progressive multiple sequence alignment through sequence weighting, position-VSHFLILFJDSSHQDOWLHVVDQGZHLJKWPDWULFKRFLFH' *Acids Research* 22: 4673±4680.

Thippeswamy. B., Shivakumar C.K., and Krishnappa M. 2012. Bioaccumulation potential of *Aspergillus niger* and *Aspergillus flavus* for removal of heavy metals from paper mill effluent. *Journal of Environmental Biology* 33: 1063±1068.

Tucker, M. D., Barton L. L., and Thomson, B. M. 1997. Reduction and immobilization of molybdenum by *Desulfovibrio desulfuricans*. *Journal of Environmental Quality*. 26: 1146-1152.

- Truex M.J, Peyton B.M, Valentine N.B, Gorby Y.A. 1997. Kinetics of U(VI) reduction by a dissimilatory Fe(III)-reducing bacterium under non-growth conditions. *Biotechnology and Bioengineering* 5;55(3):490-6.
- Twite. R. And Bierwagen G. 1998 Review of alternatives to chromate for corrosion protection of aluminum aerospace alloys. *Progress in organic coatings* 33: 91-100.
- Underwood EJ. 1979. Environmental sources of heavy metals and their toxicity to man and animals. *Programe Water Technology*. 11(4-5):33±45.
- Underwood, E. J. 1966. The mineral nutrition of livestock. Commonwealth Agricultural Bureaux, pp 103.
- Vimalashanmugam, K., and Viruthagiri, T. 2013. Medium optimization for solid state fermentative production of xylanase by *Aspergillus terreus* using central composite design. *Innovative Romanian Food Biotechnology*, 13:18-29.
- Wang, P. C., Mori, T., Komori, K., Sasatsu, M., Toda K. and Ohtake, H. 1989. Isolation and characterization of an *Enterobacter cloacae* strain that reduces hexavalent chromium under anaerobic conditions. *Applied and Environmental Microbiology* 55: 1665-1669.
- Walser, H., and Shields, D. J. 2006. Traditional and emerging applications of molybdenum metal and its alloys. In 18th annual general meeting of IMO, Austria. pp 12-20
- Wee, B. S., Shukor, S. A., Khaidir, A. F., Hamzah, M. S., Rahman, S. A., Elias, M. S., and Hashim, A. 2014. Biomonitoring of Trace Elements Using Epiphytic Lichens Collected in a Suburban Area of Selangor, Malaysia. In *From Sources to Solution*, Springer Singapore, pp 37-41
- Webb, J.L. 1963. *Enzymes and Metabolic Inhibitors*. Boston: Academic Press. Pp 33-34
- Wheeler, P. A., and Kirchman, D. L. 1986. Utilization of inorganic and organic nitrogen by bacteria in marine systems. *Limnology and Oceanography* 31(5): 998-1009.
- Wisniak, J. 2009. Carl Wilhelm Scheele. *Revista CENIC Ciencias Químicas*,40(3). 20-25.
- Williams, R. J. P. 1994. The biochemistry of molybdenum. E. R. Braithwaite and J. Haber, eds., *Molybdenum: an Outline of it's Chemistry and Uses*, Elsevier, Amsterdam, pp-419-450
- World Health Organization. 2003. Molybdenum in drinking-water: background document for development of WHO guidelines for drinking-water quality.

Yano, T., Nakahara, T., Kamiyama, S. & Yamada, K. 1966 Kinetic studies on microbial activities in concentrated solutions. I . Effect of excess sugars on oxygen uptake rate of a cell-free respiratory system. *Agricultural and Biological Chemistry* 30: 42±48.

Yamaguchi, S., Miura, C., Ito, A., Agusa, T., Iwata, H., Tanabe, S. and Miura, T. 2007. Effects of lead, molybdenum, rubidium, arsenic and organochlorines on spermatogenesis in fish: Monitoring at Mekong Delta area and in vitro experiment. *Aquatic toxicology*, 83(1): 43-51.

Yong, F. 2000. Mamut copper mine the untold story, minerals: underpinning seminar on the Malaysian minerals industry, Pacific Sutera Hotel, Kota .LQDEDOX6DEDK0DODVLDSS-24,

Yong, N. K., Oshima M., Blake R. C. and Sugio, T. 1997. Isolation and some properties of an iron-oxidizing bacterium *Thiobacillus ferrooxidans* resistant to molybdenum ion. *Bioscience Biotechnology and Biochemistry* 61: 1523-1526.

Yoshimura K, Ishii M, Tarutani T. 1986. Micro determination of phosphate in water by gel-phase colorimetry with molybdenum blue. *Analytical Chemistry* 58:591±594.

Yuichi Nogi, Hideto Takami and Koki Horikoshi. 2005. Characterization of International Journal of Systematic and Evolutionary Microbiology. 55: 2309±2315.

Yurt. N., Sears. J., and Lewandowski. Z. 2002. Multiple substrate growth kinetics of *Leptothrix disophora* SP- 6. *Biotechnology Progress*, 18: 994- 1002.

Yunus. M. Y. 2014. Revisiting the role of the electron transport chain in molybdate reduction by *Enterobacter cloacae* Strain 48. *Indian Journal of Biotechnology*. IJBT/Auth/2012/2090, accepted.

Zakaria, Z. A., Zakaria, Z., Surif, S., and Ahmad, W. A. 2007. Hexavalent chromium reduction by *Acinetobacter haemolyticus* isolated from heavy-metal contaminated wastewater. *Journal of hazardous materials*. 146(1): 30-38.

Zhai, X. W., Zhang, Y. L., Qi, Q., Bai, Y., Chen, X. L., Jin, L. J. and Liu, F. J. 2013. Effects of molybdenum on sperm quality and testis oxidative stress. *Systems biology in reproductive medicine*, 59(5): 251-255.

Zhang, Y. L., Liu, F. J., Chen, X. L., Zhang, Z. Q., Shu, R. Z., Yu, X. L., and Liu, Z. J. 2013. Dual effects of molybdenum on mouse oocyte quality and ovarian oxidative stress. *Systems biology in reproductive medicine*, 59(6): 312-318.

Zhu, Xueqing, Venosa, A.D., Suidan, M.T., and Lee, K. 2001. Guidelines for the Bioremediation of Marine Shorelines and Freshwater Wetlands. Office of Research and Development, U.S. Environmental Protection Agency.

