



UNIVERSITI PUTRA MALAYSIA

***ANTIDIABETIC ACTIVITIES OF OIL PALM (*Elaeis guineensis* Jacq.)
FRUIT AND PALM OIL MILL EFFLUENT EXTRACTS***

MOHD FAEZ BIN SHARIF

FBSB 2014 36



**ANTIDIABETIC ACTIVITIES OF OIL PALM (*Elaeis guineensis* Jacq.)
FRUIT AND PALM OIL MILL EFFLUENT EXTRACTS**

By

MOHD FAEZ BIN SHARIF

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

October 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**ANTIDIABETIC ACTIVITIES OF OIL PALM (*Elaeis guineensis* Jacq.)
FRUIT AND PALM OIL MILL EFFLUENT EXTRACTS**

By

MOHD FAEZ BIN SHARIF

October 2014

Chairperson: Associate Professor Muhajir Hamid, PhD
Faculty: Biotechnology and Biomolecular Sciences

Diabetes is one of the top ten causes of death in Malaysia. According to the fourth National Health and Morbidity survey conducted in 2011, it is estimated that 15.2% (2.6 million) of Malaysians adults 18 years old and above suffer from diabetes. Most antidiabetic drugs available are associated with several side effects which explain the current prevalence of diabetes. Therefore researches are needed in exploring the new alternative for antidiabetic treatment which are safe, efficient and exert a lesser amount of side effects. Recently the oil palm, *Elaeis guineensis* has been explored in several antidiabetic studies. The oil palm leaves have been found to reduced hyperglycemia in STZ-induced diabetic rats due to the high polyphenolic content. Nevertheless several parts of oil palm which are also contained high amount of polyphenol such as fruit and the effluent from palm oil processing, POME are never been investigated in antidiabetic study. Therefore, this study was conducted to evaluate the antidiabetic properties of oil palm fruit and POME extracts through the *in vivo* antihyperglycemic evaluation. In addition, an initiative was made to study the possible mechanisms involve using *in vitro* models. Raw POME and oil palm fruit were subjected to solvent extraction using ethanol. The extracts collected were further used in the *in vivo* and *in vitro* experiments. To evaluate the antihyperglycemic property of both extracts in diabetic rats, the rats were given the extracts orally using intragastric gavage in the *in vivo* study. The *in vitro* models were design to evaluate the potential antidiabetic mechanisms involve by using the glucose uptake, insulin secretion as well as adiponectin secretion model. The results had shown that acute treatment of POME and oil palm fruit extracts reduced fasting and postprandial hyperglycemia in streptozotocin-induced diabetic rats. Following 28-days treatment, both extracts at concentration of 500 mg kg⁻¹ b.w significantly reduced hyperglycemia, improved the body weight and increased insulin secretion in streptozotocin induced diabetes rats. Through *in vitro* evaluation, the oil palm fruit extract (500µg ml⁻¹) were found to stimulate the insulin secretion from BRIN BD11 cell line the most. Moreover both extracts also enhanced basal and insulin mediated glucose uptake into adipocytes, muscles and liver cells. In the evaluation of adiponectin secreting activity, the POME extracts significantly increased adiponectin secretion in both basal and insulin-stimulated state. However the oil palm fruit extracts significantly increased adiponectin secretion only under the insulin stimulated state. The HPLC analysis had shown the presence of gallic acid and

catechin as part of the bioactive compound for both extracts. In conclusion, from *in vivo* evaluation the treatment of POME and oil palm fruit extracts were shown to reduce hyperglycemia at different prandial state in diabetic rats. Both extracts also did not cause severe hypoglycemia in normal rats. The *in vitro* study suggested that the antihyperglycemic property of POME and oil palm fruits were mediated through insulin secretion from pancreatic β -cells, enhancement of glucose uptake by the muscles, adipocytes and liver cells and amplification of adiponectin secretion from adipocytes cells. Therefore various antihyperglycemic potential of both extracts together with its property that did not cause hypoglycemia make them suitable to be develop as new oral antidiabetic drugs.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

AKTIVITI ANTIDIABETIS EKSTRAK BUAH KELAPA SAWIT (*Eleais guineensis* Jacq.) DAN SISA EFLUEN KILANG MINYAK SAWIT
Oleh

MOHD FAEZ BIN SHARIF

Oktober 2014

Pengerusi: Muhajir Hamid, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Diabetes adalah salah satu penyakit yang berada di tangga teratas yg menyebabkan kematian di Malaysia. Merujuk kepada kaji selidik Kesihatan dan Morbiditi Kebangsaan ke-4 yg dijalankan dalam tahun 2011, dianggarkan sebanyak 15.2% (2.6 juta) penduduk Malaysia dewasa yang berumur 18 tahun dan keatas menghidap penyakit diabetes. Kebanyakan ubat antidiabetik yang terdapat pada masa kini mempunyai kesan sampingan dan ini secara tidak langsung menerangkan tentang fenomena diabetes yang berlaku kini. Oleh itu kajian perlu dilakukan untuk mencari sumber alternatif baru untuk rawatan antidiabetik yang lebih selamat, berkesan dan juga tidak mendatangkan kesan sampingan yang berlebihan. Kini kelapa sawit, *Elais guineensis* telah digunakan dalam beberapa kajian antidiabetik. Daun kelapa sawit telah dibuktikan berjaya menurunkan tahap hyperglisemia tikus aruhan diabetik oleh sebab kandungan polifenol yang tinggi yang terdapat di dalamnya. Walaubagaimanapun kajian masih belum dilakukan terhadap beberapa bahagian lain kelapa sawit yang turut mengandungi kandungan polifenol yang tinggi seperti buah kelapa sawit dan juga sisa dari pemprosesan sawit, POME. Oleh itu kajian ini telah dijalankan untuk menganalisa kesan antidiabetik oleh buah kelapa sawit dan sisa POME melalui penilaian antihyperglisemik secara *in vivo* dan juga untuk mengkaji mekanisme yang mungkin memberi kesan antidiabetik menggunakan model *in vitro*. Sisa POME dan buah kelapa sawit diekstrak menggunakan etanol. Hasil ekstrak seterusnya digunakan di dalam kajian *in vivo* dan *in vitro*. Di dalam kajian *in vivo*, ekstrak diberikan kepada tikus diabetik melalui intragastrik untuk mengkaji sifat antihyperglisemik kedua-dua ekstrak. Model *in vitro* pula direka untuk menghuraikan mekanisme yang mungkin menyebabkan kesan antihyperglisemik menggunakan model pengambilan glukosa, rembesan insulin dan juga model rembesan adiponektin. Keputusan kajian menunjukkan bahawa rawatan secara akut ekstrak sisa POME dan buah kelapa sawit menurunkan hyperglisemia puasa dan posprandial di dalam tikus diabetik aruhan-streptozotocin. Melalui rawatan selama 28 hari, kedua-dua ekstrak pada kepekatan 500 mg kg⁻¹ berat badan menunjukkan penurunan hyperglisemia secara signifikan, menambah jumlah berat dan meningkatkan rembesan insulin di dalam tikus diabetik aruhan-streptozotocin. Melalui kajian *in vitro*, ekstrak buah kelapa sawit (500µg ml⁻¹) didapati paling efektif merangsang perembesan insulin dari sel BRIN BD11. Tambahan lagi, kedua-dua ekstrak juga didapati meningkatkan aktiviti pengambilan glukosa basal dan diperantarakan-insulin ke dalam sel-sel adipos, otot dan hati. Melalui kajian rembesan protein

adiponektin, ekstrak sisa POME didapati merangsang perembesan adiponektin di dalam kedua-dua keadaan basal dan juga diperantarakan-insulin. Walaubagaimanapun ekstrak buah kelapa sawit didapati hanya merangsang perembesan adiponektin secara signifikan di dalam keadaan diperantarakan-insulin sahaja. Analisa HPLC menunjukkan kehadiran asid gallic dan catechin sebagai sebahagian daripada bahan bio aktif kedua-dua ekstrak. Sebagai kesimpulan, rawatan ekstrak sisa POME dan buah kelapa sawit didapati menurunkan hiperglisemia di dalam tikus diabetik. Kedua-dua ekstrak juga didapati tidak mengakibatkan kejadian hipoglisemia di dalam tikus normal. Kajian in vitro yang dijalankan menunjukkan bahawa sifat antihiperglisemik ekstrak sisa POME dan buah kelapa sawit tersebut adalah disebabkan oleh rembesan insulin dari sel β -pankreas, peningkatan dalam pengambilan glukosa ke dalam sel-sel otot, adiposa dan hati dan juga kerana peningkatan rembesan adiponektin dari sel adiposa. Oleh yang demikian, kepelbagaian potensi antihiperglisemia kedua-dua ekstrak dan juga sifat ekstrak yang tidak mengakibatkan hipoglisemia menjadikan ekstrak-ekstrak tersebut sesuai untuk dibangunkan sebagai ubat antidiabetik yang baru.

ACKNOWLEDGEMENTS

First and foremost, all thanks to Allah the Al Mighty, for granting me the chance and ability to successfully complete this study. In preparing this thesis, many people have contributed their advices and knowledge towards my understanding and thoughts. This page is specially designed to express my appreciation to those peoples.

I wish to express my deepest gratitude to my dearest supervisor, Assoc. Prof. Dr. Muhajir Hamid and my co-supervisor Prof. Dr. Amin ismail and Dr. Zainah Adam for the great guidance, encouragements, efforts and valuable advice given throughout the study. Special thanks also to my dearest wife Siti Nurhafizah Binti Abu Samah for all the support and helps during this journey.

Not to be forgotten, my dearest parents, brother and sister whom had support and pray for my success in completing this study. I am also indebted to all lectures of Faculty of Biotechnology and Biomolecular Sciences, lab assistants and fellow friends for their guidance, advice and motivation. Without their continued support and interest, this thesis would not have been the same as presented here.

I certify that a Thesis Examination Committee has met on 13th October 2014 to conduct the final examination of Mohd Faez Bin Sharif on his thesis entitle Antidiabetic activities of oil palm (*Elaeis guineensis* Jacq.) Fruit and Palm Oil Mill Effluent 1971 and the constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Examination Committee were as follows:

Assoc. Prof. Dr. Janna Ong Abdullah, PhD

Associate professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairperson)

Dr. Syahida Ahmad, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Prof. Dr. Md Zuki Abu Bakar, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Dr. Asna Urooj, PhD

Lecturer
Department of Studies in Food Science & Nutrition
Universiti of Mysore
India
(External Examiner)

ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 February 2015

I certify that a Thesis Examination Committee has met on 13 Oktober 2014 to conduct the final examination of Mohd Faez bin Sharif on his thesis entitled "Antidiabetic Activities of Oil Palm (*Elaeis guineensis* Jacq.) Fruit and Palm Oil Mill Effluent Extracts" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Janna Ong binti Abdullah, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Md Zuki bin Abu Bakar @ Zakaria, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Syahida binti Ahmad, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Asna Urooj, PhD

Senior Lecturer
Department of Studies in Food Science & Nutrition
University of Mysore
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 February 2015

Declaration by graduate student

I hereby confirm that:

- x this thesis is my original work;
- x quotations, illustration and citations have been duly referenced;
- x this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- x intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- x written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- x there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date: 20 March 2015

Name and Matric No.: Mohd Faez Bin Sharif, GS 28635

Declaration by Members of Supervisory Committee

This is to confirm that:

- x the research conducted and the writing of this thesis was under our supervision;
- x supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Associate Prof. Dr. Muhajir
Hamid

Signature: _____
Name of
Member of
Supervisory
Committee: Dr. Zainah Adam

Signature: _____
Name of
Member of
Supervisory
Committee: Prof. Dr. Amin Ismail

TABLE OF CONTENTS

	ABSTRACT	Page
	ABSTRAK	i
	ACKNOWLEDGEMENT	iii
	APPROVAL	v
	DECLARATION	vi
	LIST OF TABLES	viii
	LIST OF FIGURES	xiv
	LIST OF ABBREVIATIONS	xv
		xviii
CHAPTER		
1	GENERAL INTRODUCTION	
1.1	Background of study	1
1.2	Statement of problems	1
1.3	Justification of study	2
1.4	Hypothesis of study	3
1.5	Objectives of study	3
2	LITERATURE REVIEW	
2.1	Diabetes mellitus	4
2.2	Type 1 diabetes mellitus	4
2.3	Type 2 diabetes mellitus	5
2.4	Treatment of diabetes mellitus	5
2.4.1	Treatment for type 1 diabetes mellitus	5
2.4.2	Treatment for type 2 diabetes mellitus	6
2.4.2.1	Sulfonylureas (SUs)	6
2.4.2.2	Biguanide	6
2.4.2.3	Thiazolidinediones (TZDs)	7
2.4.2.4	-glucosidase inhibitors	7
2.4.2.5	Meglitinide	8
2.5	Blood glucose level (BGL) regulation in human	8
2.5.1	Stimulation of insulin secretion from pancreatic β -cells	9
2.5.2	Enhancement of glucose uptake into peripheral Cells	9
2.5.3	Augmentation of adiponectin secretion from adipocytes cells	10
2.6	Models for antidiabetic research	11
2.6.1	<i>In vivo</i> models	11
2.6.2	<i>In vitro</i> models	12
2.7	Plants as alternative antidiabetic agents	13
2.8	Oil palm (<i>Elaeis guineensis</i>)	13
2.9	Nutritional properties of palm oil and its component	15
2.10	Palm oil mill effluent (POME)	16

3 EVALUATION OF ANTIHYPERGLYCEMIC ACITVITIES OF *E. GUINEENSIS* FRUITS AND POME EXTRACTS IN STZ-INDUCED DIABETIC RATS

3.1	Introduction	18
3.2	Materials and methods	19
3.2.1	Samples collection	19
3.2.2	Sample preparation and extraction procedures	19
3.2.3	Experimental animals and diabetes induction	19
3.2.4	Hypoglycemic evaluation in normal rats and antihyperglycemic evaluation in fasting and post-prandial diabetic rats	20
3.2.5	Oral glucose tolerance test (OGTT)	20
3.2.6	Evaluation of long term treatment in STZ-induced diabetic rats	20
3.2.7	Statistical analyses	21
3.3	Results	22
3.3.1	The hypoglycemic activity of POME and OPF extract in normal rats	22
3.3.2	Antihyperglycemic activity of POME and OPF extract in STZ-induced diabetic rats at fasting state.	24
3.3.3	Antihyperglycemic activity of POME and OPF extract in STZ-induced diabetic rats at postprandial state.	26
3.3.4	Effect of POME extract on oral glucose tolerance test (OGTT) of STZ-induced diabetic rats	28
3.3.5	Effect of OPF extract on oral glucose tolerance test (OGTT) of STZ-induced diabetic rats	29
3.3.6	Effect of long term four weeks POME treatment	30
3.3.7	Effect of long term four weeks OPF treatment	31
3.3.8	Effect of long term four weeks POME treatment on blood glucose level in normal and STZ-induced diabetic rats	32
3.3.9	Effect of long term four weeks OPF treatment on blood glucose level in normal and STZ-induced diabetic rats	33
3.3.10	Effect of long term four weeks POME and OPF	34
3.4	Discussion	35
3.5	Conclusion	39

4 EVALUATION OF *E. GUINEENSIS* FRUITS AND POME EXTRACTS ON CELL VIABILITY AND ADIPONECTIN SECRETION FROM 3T3F442A ADIPOCYTES

4.1	Introduction	40
4.2	Materials and methods	42
4.2.1	Sample preparation and extraction procedures	42
4.2.2	Cell line	42
4.2.3	Cell culture	42

4.2.4	Cell viability assay	42
4.2.5	Examination of POME and OPF extracts on the secretion of adiponectin from 3T3F442A adipocytes	43
4.2.6	Determination of adiponectin concentration	44
4.2.7	Statistical analyses	44
4.3	Results	45
4.3.1	Effects of POME and OPF extracts on the viability of BRIN BD11 cells	45
4.3.2	Effects of POME and OPF extracts on the viability of Chang liver cells	47
4.3.3	Effects of POME and OPF extracts on the viability of L6 myotubes	49
4.3.4	Effects of POME and OPF extracts on the viability of 3T3F442A adipocytes	51
4.3.5	Effect of OPF and POME extracts on stimulating adiponectin secretion from 3T3F442A adipocytes	53
4.4	Discussions	56
4.5	Conclusion	59
5	EFFECT OF <i>ELAEIS GUINEENSIS</i> FRUITS AND POME EXTRACTS ON GLUCOSE UPTAKE ACTIVITY IN L6 MYOTUBES, 3T3F442A ADIPOCYTES AND CHANG LIVER CELLS	
5.1	Introduction	60
5.2	Materials and methods	61
5.2.1	Sample preparation and extraction procedure	61
5.2.2	Cell line	61
5.2.3	Cell line maintenance	61
5.2.4	Glucose uptake activity of POME and OPF extracts in L6 myotubes	61
5.2.5	Glucose uptake activity of POME and OPF extracts in 3T3F442A adipocytes	62
5.2.6	Glucose uptake activity of POME and OPF extracts in Chang liver cells	63
5.2.7	Statistical analyses	63
5.3	Results	64
5.3.1	Glucose uptake activity of POME and OPF extracts in L6 myotubes	64
5.3.2	Glucose uptake activity of POME and OPF extracts in 3T3F442A adipocytes	66
5.3.3	Glucose uptake activity of POME and OPF extracts in Chang liver cells	69
5.4	Discussions	73
5.5	Conclusion	76

6	EVALUATION OF INSULIN SECRETING ACTIVITY FROM BRIN BD11 CELL AND SCREENING OF POLYPHENOL COMPOUNDS IN OIL PALM FRUIT AND POME EXTRACTS USING HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)	
6.1	Introduction	77
6.2	Materials and methods	79
6.2.1	Sample preparation and extraction procedures	79
6.2.2	Cell line	79
6.2.3	Examination on insulin secreting activity of BRIN BD11 in the presence of POME and OPF extracts	80
6.2.4	Determination of insulin concentration	80
6.2.5	Determination of insulin secretion mechanisms	80
6.2.6	Screening of polyphenolic compounds in POME and OPF extracts using High Performance Liquid Chromatography (HPLC)	80
6.2.7	Statistical analyses	80
6.3	Results	81
6.3.1	Effect of POME and OPF extracts on insulin secreting activity of BRIN BD11 cell line	81
6.3.2	Effect of various insulin modulators on insulin secretion from BRIN BD11 cell line	83
6.3.3	Screening of POME and OPF extracts by HPLC	84
6.4	Discussions	87
6.5	Conclusion	90
7	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	
7.1	Summary and general conclusion	91
7.2	Recommendation for future research	93
	REFERENCES	95
	APPENDICES	108
	BIODATA OF STUDENT	134
	LIST OF PUBLICATION	135

LIST OF TABLES

Table		Page
3.1	Hypoglycemic effect of POME and OPF extracts on fasting blood glucose in normal rats	23
3.2	Antihyperglycemic effect of POME and OPF extracts on fasting blood glucose level in STZ-induced diabetic rats	25
3.3	Antihyperglycemic effect of POME and OPF extracts on postprandial blood glucose level in STZ-induced diabetic rats	27

LIST OF FIGURES

Figure		Page
2.1	Harvested oil palm fruit from a 7 years old oil palm tree	14
2.2	Palm oil mill effluent (POME)	17
3.1	Oral glucose tolerance tests of experimental rats treated with POME extract	28
3.2	Oral glucose tolerance tests of experimental rats treated OPF extract	29
3.3	Ejcpigu"kp"tcvøu"dqf{"y gki jv"fwtkpi"nqpi"vgt o"hqwt"y ggmu" POME treatment	30
3.4	Ejcpigu"kp"tcvøu"dqf{"y gki jv"fwtkpi"long term four weeks OPF treatment	31
3.5	Ejcpigu"kp"tcvøu"D I N"fwtkpi"long term four weeks POME Treatment	32
3.6	Ejcpigu"kp"tcvøu"D I N"fwtkpi"long term four weeks OPF Treatment	33
3.7	Ejcpigu"kp"tcvøu"rncu oc"kpawnkp"ngxgn"fwtkpi"long term four weeks treatment	34
4.1	Cells viability of BRIN BD11 cells after 72 hours exposure to POME extract	45
4.2	Cells viability of BRIN BD11 cells after 72 hours exposure to OPF extract	46
4.3	Cells viability of BRIN BD11 cells after 72 hours exposure to glybenclamide	46
4.4	Cells viability of Chang liver cells after 72 hours exposure to POME extract	47
4.5	Cells viability of Chang liver cells after 72 hours exposure to OPF extract	48
4.6	Cells viability of Chang liver cells after 72 hours exposure to rosiglitazone maleate	48
4.7	Cells viability of L6 myotubes after 72 hours exposure to POME extract	49

4.8	Cells viability of L6 myotubes after 72 hours exposure to OPF extract	50
4.9	Cells viability of L6 myotubes after 72 hours exposure to Metformin	50
4.10	Cells viability of 3T3F442A after 72 hours exposure to POME extract	51
4.11	Cells viability of 3T3F442A after 72 hours exposure to OPF extract	52
4.12	Cells viability of 3T3F442A after 72 hours exposure to rosiglitazone maleate	52
4.13	Effect of OPF extract on basal and insulin-stimulated adiponectin secretion from 3T3F442A adipocytes	53
4.14	Effect of POME extract on basal and insulin-stimulated adiponectin secretion from 3T3F442A adipocytes	54
4.15	Effect of rosiglitazone maleate on basal and insulin-stimulated adiponectin secretion from 3T3F442A adipocytes	55
5.1	Effect of POME extract on basal and insulin-mediated glucose uptake activity of L6 myotubes	64
5.2	Effect of OPF extract on basal and insulin-mediated glucose uptake activity of L6 myotubes	65
5.3	Effect of metformin on basal and insulin-mediated glucose uptake activity of L6 myotubes	66
5.4	Effect of POME extract on basal and insulin-mediated glucose uptake activity of 3T3F442A adipocytes	67
5.5	Effect of OPF extract on basal and insulin-mediated glucose uptake activity of 3T3F442A adipocytes	68
5.6	Effect of rosiglitazone maleate on basal and insulin-mediated glucose uptake activity of 3T3F442A adipocytes	69
5.7	Effect of POME extract on basal and insulin-mediated glucose uptake activity of Chang liver cells	70
5.8	Effect of OPF extract on basal and insulin-mediated glucose uptake activity of Chang liver cells	71
5.9	Effect of rosiglitazone maleate on basal and	

	insulin-mediated glucose uptake activity of Chang liver cells	72
6.1	Effect of POME extract on insulin secretion from BRIN BD11 cells	81
6.2	Effect of OPF extract on insulin secretion from BRIN BD11 cells	82
6.3	Effect of glibenclamide on insulin secretion from BRIN BD11 cells	82
6.4	Effect of various insulin secretion modulators on the insulin secretion activity of 500 µg/ml OPF from BRIN BD11 cells	83
6.5	Chromatogram of POME extract	84
6.6	Chromatogram of OPF extract	85
6.7	Gallic acid	85
6.8	Catechin	86

LIST OF ABBREVIATIONS

ACUC	Animal Care and Use Committee
AUC _{Glucose}	Area under the glucose curve
BGL	Blood glucose level
BSA	Bovine serum albumin
CaCl ₂	Calcium chloride
DMEM	Fw ndgee qøu" o q fkhkg f"gc i ngøu" o g fkw o
DMSO	Dimethylsulfoxide
FBG	Fasting blood glucose
FBS	Fetal bovine serum
FPG	Fasting plasma glucose
HCl	Hydrochloric acid
H ₂ O	Water
H ₂ SO ₄	Sulfuric acid
HEPES	4-(2-hydroxyethyl)piperazine-1-ethanesulfonic acid
IBMX	3-isobutyl-1-methylxanthine
IDDM	Insulin dependent diabetes mellitus
K ⁺ _{ATP} channel	ATP-sensitive potassium channel
K	Potassium
KCl	Potassium chloride
KRB	Krebs Ringer Bicarbonaate buffer
KH ₂ PO ₄	Potassium dihydrogen phosphate
MgSO ₄	Magnesium sulphate
MTT	3-(4,5-Dimethyl-2-thiazol)-2,5-diphenyl-2H-tetrazolium Bromide
NaCl	Sodium chloride
NaHCO ₃	Sodium hydrogen carbonate
NaOH	Sodium hydroxide
NIDDM	Non-insulin dependent diabetes mellitus
OD	Optical density
OPF	Oil palm fruit extract
PBS	Phosphate buffer saline
POME	Palm oil mill effluent
RPMI	Roswell Park Memorial Institute
SDS	Sodium dodecyl sulphate
STZ	Streptozotocin
TNF	Tumor necrosis factor
TZDs	Thiazolidinidiones
TRIS	2-Amino-2-hydroxymethyl-propane-1,3-diol
UPM	Universiti Putra Malaysia
USA	United state of America
WHO	World Health Organization
cm ³	centimeter cubic
µg	microgram
mg	milligram
g	gram
kg	kilogram
µl	microlitre
ml	millilitre

L	Litre
μM	micromolar
mM	millimolar
M	Molar
μm	micrometer
mm	millimeter
m	meter
rpm	round per minute
α	Alpha
β	Beta
γ	Gamma
%	Percentage
$^{\circ}\text{C}$	Degree celcius

CHAPTER 1

GENERAL INTRODUCTION

1.1 Background of Study

Diabetes mellitus which characterized by persistent hyperglycemia in blood is a chronic disease that is relatively common throughout the world. It is resulted from defects of insulin secretion or insulin action and can also be cause by combination of both factors (Alberti and Zimmet, 1998). It is normally accompanied by increased risk to oxidative stress, hypertension and severe atherosclerosis (Reusch, 2003).

Diabetes is one of the top ten causes of death in Malaysia. According to the fourth National Health and Morbidity survey conducted in 2011, it is estimated that 15.2% (2.6 million) of Malaysians adults 18 years old and above suffer from diabetes. The amount has bypassed the assumption made by World Health Organization (WHO) which estimated that in the year of 2030, Malaysia would have a total of 2.48 millions diabetes sufferers. This indicates that Malaysia is having a faster increase rate of diabetic disease. The rapid rising trend in the prevalence of diabetes could possibly due to the urbanization, changes in dietary habits, growth of population and inactive lifestyle (Zanariah *et al.*, 2008).

Poor glycemic controls in diabetes patient will always contribute to the prevalence of diabetes complications such as retinopathy, neuropathy, and albuminuria. The records from the Ministry of Health (MOH) showed that the number of diabetic patients admitted to hospitals had increased from 19, 629 cases in 1991 to 30, 661 in 2001. It shows about 56% increments over 10 years. Diabetes mortality rates also increased from 254 death in 1991 to 380 in 2001 (Ooyub *et al.*, 2004). This indicates that diabetes had become major problem in Malaysia. Therefore a stepped up efforts are required in controlling and preventing this chronic disease.

1.2 Statement of Problems

Diabetes can be classified into two class which are type I and type II diabetes mellitus. In type I diabetes, the maintenance of hyperglycemia is limited to insulin therapy (Jacobson *et al.*, 2009). Different types of insulin analogues are available for this purpose such as Humalog, NovoRapid, Levimir and Lantus. Those analogues were categorized according to their times of action onset and duration. On the other hand the regulation of hyperglycemia in type II diabetes patient were mostly by changes of diet, active lifestyle, regular exercise and the uses of oral antidiabetic drugs. Antidiabetic drugs for the treatment of type II diabetes mellitus vary from sulfonylureas, biguanides, thiazolidinediones, r-glucosidase inhibitor, amylin synthetic derivatives and incretin mimetics. The drugs were categorized into different groups based on its antidiabetic mechanisms and mode of action (Chehade and Mooradian, 2000).

The major problem liaises with the conventional antidiabetic drugs were the limitations of the drugs itself. The drugs can bring up undesirable adverse effects such as vomiting, weight gain, nausea and diarrhoea. Furthermore most conventional antidiabetic drugs were also expensive making them unavailable and unaffordable especially in the third world and developing countries where it is difficult to have access to those modern and high cost drugs (Babu *et al.*, 2007). The clinical trial also were still lacking for most antidiabetic drugs making them less efficient (Kirchheiner *et al.*, 2005). For instance, drugs from sulfonylureas group was reported to cause hypoglycemia and weight gain. There are also reports on occurring of death due to prolonged severe hypoglycemia. Metformin on the other hand always associated also can cause gastrointestinal disturbance, lethal ileus and renal tumors. Meanwhile, drugs from the thiazolidinediones groups can cause fluid retention, liver injury and anemia (Bell, 2002).

Due to the limitations of conventional antidiabetic drugs, searching for a new alternative antidiabetic agent for diabetes treatment is needed. It is crucial to find antidiabetic agents that are safe and efficient to replace the conventional antidiabetic drugs to overcome the limitations brought by those drugs.

1.3 Justification of Study

Palm oil mill effluent (POME) is a waste generated from the oil palm manufacturing process. The POME is often discarded in disposal ponds, resulting in the leaching of contaminants that pollute the groundwater and soil while releasing methane gas to the atmosphere. Tan and colleague reported in 2001 that water soluble phytochemicals from the palm fruit mesocarp which partition into aqueous phase during the oil palm manufacturing process could contains several phenolic compounds. This includes gallic, protocatechuic, gentisic, chlorogenic, coumaric, ferulic, and caffeic acids, as well as hesperidin and catechins (Tan *et al.*, 2001). Recent studies have evaluated the potential of phytochemicals from oil palm in treating various diseases. Oil palm phenolics have been reported previously to inhibit proliferation of estrogen-receptor-positive human breast adenocarcinoma cells, human lung carcinoma cells (Shamala *et al.*, 2010) and promoting vascular relaxation (Mahinda *et al.*, 2002). Furthermore recent study by Rosalina and colleague has revealed the potential of oil palm leaves extract in reducing hyperglycemia and lipid oxidation in STZ-rats (Rosalina *et al.*, 2011). Therefore it is a great effort to evaluate the POME as a new source of oil palm phenolics to treat diabetes. The POME is cheaper and also help reducing environmental pollution as the waste is being utilized.

Phenolic compounds are often found in plants. It comprises of groups such as the phenolic acids, flavonoids and tannins. Several publications on the antioxidative activities of plant derived phenolics have been reported previously (Wang and Ballington, 2007; Materska and Perucka, 2005). Phenolic acids have attracted special attention lately due to its strong inhibitory activity on oxidation induced by peroxyl radicals (Hu and Kitts, 2001). Oil palm fruit is also a rich source of water soluble-phenolics antioxidants like any other fruits. Plants with high phenolics antioxidant

compounds exert high potential as supplements for improving blood glucose control and preventing long-term complications in diabetics (Gallegher *et al.*, 2003). Many research on the antioxidative capacity of the oil palm fruit have been conducted (Neo *et al.*, 2010; Nagendran *et al.*, 2005). Despite the high antioxidant activity of oil palm reported previously however the studies on its potential as alternative antidiabetic agent are never been studied. Therefore this study is conducted to evaluate the potential of oil palm fruit as a new antidiabetic agent to combat diabetes mellitus.

1.4 Hypothesis of Study

- I. Treatment of oil palm fruit and POME extracts reduce hyperglycemia in STZ-induced diabetic rats and has no severe hypoglycemia effect in normal rats
- II. improve insulin secretion activity of BRIN BD11 cell line
- III. improve glucose uptake activity into L6 myotubes, 3T3F442A adipocytes and Chang liver cells and
- IV. improve adiponectin secretion activity of 3T3F442A adipocytes.

1.5 Objectives of study

The main objective of this study was to evaluate the antidiabetic properties of oil palm (*Elaeis guineensis*) fruit and POME extract. The specific objectives of this study were:

1. To evaluate the hypoglycaemic and antihyperglycaemic activity of oil palm fruit and POME extracts in normal and STZ-induced diabetic rats.
2. To evaluate the toxicity of oil palm fruit and POME extracts and their effect on adiponectin secretion from adipocyte cells.
3. To investigate the effect of oil palm fruit and POME extract on glucose uptake into muscle, adipocyte and liver cell.
4. To evaluate the effect of oil palm fruit and POME extract on insulin secretion from pancreatic beta cells and to identify the phenolics compound that exists in both extracts.

REFERENCES

- Abdeljebbar, L. H., Benjouad, A., Morjani, H., Merghoub, N., El Haddar, S., Humam, M., Christen, P., Hostettmann, K., Bekkouche, K. and Amzazi, S. (2009). Antiproliferative effects of withanolides from *Withania adpressa*. *Therapie*, 64 (2): 121-127
- Adam, Z., Khamis, S., Ismail, A. and Hamid, M. (2010). Inhibitory properties of *Ficus deltoidea* on α -glucosidase activity. *Research Journal of Medicinal Plant*, 4 (2): 61-75
- Adam, Z., Hamid, M., Ismail, A. and Khamis, S. (2009). Effect of *Ficus deltoidea* extracts on hepatic basal and insulin-stimulated glucose uptake. *Journal of Biological Sciences*, 9 (8): 796-803
- Ahmad, A., Ismail, S. and Bhatia, S. (2003). Water recycling from palm oil mill effluent (POME) using membrane technology. *Desalination*, 157: 87-95
- Alberti, K. G. and Zimmet, P. Z. (1998). Definition, diagnosis, and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic Medicine*, 15 (7): 539-553
- Allan, V. and Sorens, L. (2012). Insulin initiation in patients with type 2 diabetes mellitus: treatment guidelines, clinical evidence and patterns of use of basal vs premixed insulin analogues. *European Journal of Endocrinology*, 166: 159-170
- Amini, H., Ahmadiani, A. and Gazerani, P. (2005). Determination of metformin in human plasma by high-performance liquid chromatography. *Journal of Chromatography B Analytical Technologies in the Biomedical and Life Sciences*, 824 (1-2): 319-322
- Anandharajan, R., Pathmanathana, K., Shankernarayanan, N. P., Vishwakarmab, R. A. Balakrishna, A. (2005). Upregulation of GLUT4 and PPAR by an isoflavone from *Pterocarpus marsupium* on L6myotubes: A possible mechanism of action. *Journal of Ethnopharmacology*, 97: 253-260
- Anusha, B., Vidhya, V.G. and Ramya, M. (2008). Hypoglycemic effect of *Mucuna pruriens* seed extract on normal and streptozotocin-diabetic rats. *Fitoterapia*, 79: 539-543
- Arner, P. (2003). The adipocyte in insulin resistance: key molecules and the impact of the Thiazolidinediones. *Trends in Endocrinology and Metabolism*, 14 (3): 137-144.
- Ayaki, M. and Iwasawa, A. (2011). Cell viability of four corneoconjunctival cell lines exposed to five preservatives and a surfactant used for infection control in eyedrops. *Biocontrol Science*, 16 (3): 117-121

- Babu, P. S., Prabuseenivasan, S. and Ignacimuthu, S. (2007). Cinnamaldehyde – A potential antidiabetic agent. *Phytomedicine*, 14: 15-22
- Barik, R., Jain, S., Qwatra, D., Joshi, A., Tripathi, G. S. and Goyal, R. (2008). Antidiabetic activity of aqueous root extract of *Ichnocarpus frutescens* in streptozotocin-nicotinamide induced typed II diabetes in rats. *Indian Journal of Pharmacology*, 40 (1): 19-22
- Beatriz, A. G., Luis, A. B., Sergio, G. L., Blanca, G. and Francisca, V. (2009). A general analytical strategy for the characterization of phenolic compounds in fruit juices by high-performance liquid chromatography with diode array detection coupled to electrospray ionization and triple quadrupole mass spectrometry. *Journal of Chromatography A*, 1216: 5398-5415
- Bell, D. S. (2002). Current status of diabetes treatment. *Southern medical Journal*, 95(1): 24-29
- Blumer, R. M., Van Roomen, C. P., Meijer, A. J., Houben-Weerts, J. H., Sauerwein, H. P. and Dubbelhuis, P. F. (2008). Regulation of adiponectin secretion by insulin and amino acids in 3T3-L1 adipocytes. *Metabolism*, 57 (12): 1655-62
- Bolkent, S., Yanardag, R., Tabakoglu-Oguz, A. and Ozsoy-Sacan, O. (2000) Effect of chard (*Beta vulgaris* L. Var. *Cicla*) extract on pancreatic B cells in STZ-diabetic rats: a morphological and biochemical study. *Journal of Ethnopharmacology*. 73: 251-9
- Bosenberg, L. H. and Van Zyl, D. G. (2008). The mechanism of action of oral antidiabetic drugs: A review of recent literature. *Journal of Endocrinology, Metabolism and Diabetes of South Africa*, 13 (3): 80-88
- Cai, Y., Luo, Q., Sun, M. and Corke, H. (2004). Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sciences*, 74: 2157-2184
- Cariello, A. (2005). Perspective in diabetes.postprandial hyperglycaemia and diabetes complications. Is it time to treat? *Diabetes*, 54: 1-7
- Chehade, J. M. and Mooradian, A. D. (2000). A rational approach to drug therapy of type 2 diabetes mellitus. *Drugs*, 60 (1): 95-113
- Cho, S. Y., Park, P. J., Shin, H. J., Kim, Y. K., Shin, D. W., Shin, E. S., Lee, H. H., Lee, B. G., Baik, J. H. and Lee, T. R. (2007). (-)-Catechin suppresses expression of Kruppel-like factor 7 and increases expression and secretion of adiponectin protein in 3T3-L1 cells. *The American Journal of Physiology-Endocrinology and Metabolism*, 292: 1166-1172

- Combs, T. P., Wagner, J. A., Berger, J., Doebber, T., Wang, W. J., Zhang, B. B., Tanen, M., Berg, A. H., O'Rahilly, S., Savage, D. B., Chatterjee, K., Weiss, S., Larson, P. J., Gottesdiener, K. M., Gertz, B. J., Charron, M. J., Scherer, P. E. and Moller, D. E. (2002). Induction of adipocyte complement-related protein of 30 kilodaltons by PPARgamma agonist: a potential mechanism of insulin sensitization. *Endocrinology*, 143 (3): 998-1007
- David, M. K., Matthew, C. R., Julio, R., Dongliang, Z., Dennis, D. K., Mark, S. F. and Alan, D. B. (2005). Effect of exenatide (Exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes care*, 28: 1083-1091
- Elmore, E., Siddiqui, S., Desai, N., Moyer, M. P., Steele, V. E. and Redpath, J. L. (2002). The human epithelial cell cytotoxicity assay for determining tissue specific toxicity: method modifications. *Methods in Cell Science*, 24 (4): 145-53
- Eisenberg, M. L., Maker, A. V., Slezak, L. A., Nathan, J. D., Sritharan, K. C., Jena, B. P., Geibel, J. P. and Anderson, D. K. (2005). Insulin receptor (IR) and glucose transporter 2 (GLUT 2) proteins form a complex on the rat hepatocyte membrane. *Cellular Physiology and Biochemistry*, 15 (1-4): 51-58
- Gallagher, A.M., Flatt, P.R., Duffy, G. and Abdel-Wahab, Y.H.A. (2003). The effects of traditional antidiabetic plants on in vitro glucose diffusion. *Nutrition Research*, 23: 413-424.
- George, P. and McCrimmon, R. (2012). Diazoxide. *Practical Diabetes*, 29(1): 36-37
- Gough, S. C. (2007). A review of human and analogue insulin trials. *Diabetes Research and Clinical Practice*, 77 (1): 1-15
- Grempler, R., Zibrova, D., Schoelch, C., Van Marle, A., Rippmann, J. F. and Redemann, N. (2007). Normalization of prandial blood glucose and improvement of glucose tolerance by liver-specific inhibition of SH2 domain containing inositol phosphatase 2 (SHIP2) in diabetic KKAY mice: SHIP2 inhibition cause insulin-mimetic effects on glycogen metabolism, gluconeogenesis and glycolysis. *Diabetes*, 56 (9): 2235-41
- Guo, L. and Tabrizchi, R. (2006). Peroxisome proliferator-activated receptor gamma as a drug target in the pathogenesis of insulin resistance. *Pharmacology & Therapeutics*, 111 (1): 145-173
- Haverkos, H. W., Battula, N., Drotman, D. P. And Rennert, O. M. (2003). Enteroviruses and type 1 diabetes mellitus. *Biomedicine and Pharmacotherapy*, 57 (9): 379-385
- Hall, A. P. and Davies, M. J. (2008). Assessment and management of diabetes mellitus. *The foundation Years*, 4 (6): 224-229

- Hannan, J., Marenah, L., Ali, L., Rokeya, B., Flatt, P. R. and Abdel-Wahab, Y. H. (2006). Ocimum sanctum leaf extracts stimulate insulin secretion from perfused pancreas, isolated islets and clonal pancreatic beta-cells. *Journal of Endocrinology*, 189(1): 127-136
- Hannan, J., Marenah, L., Ali, L., Rokeya, B., Flatt, P. R. and Abdel-Wahab, Y. H. (2007). Insulin secretory actions of extracts of Asparagus racemosus root in perfused pancreas, isolated islets and clonal pancreatic beta-cells. *Journal of Endocrinology*, 192(1): 159-168
- Hiroki, Y., Mitsuhisa, K., Thomas, S., Toru, A., Tsuyoshi, K., Minoru, N., Geoffrey, W. G. S. and Kiyoshi, H. (1999). cAMP enhances insulin secretion by an action on the ATP –sensitive K⁺ channel-independent pathway of glucose signalling in rats pancreatic islets. *Diabetes*, 48: 1006-1012
- Hu, C. and Kitts, D. D. (2001). Assessing antioxidant and prooxidant activities of phenolics compounds. *Journal of Agriculture and Food Chemistry*, 48: 3597-3604
- Hug, C. and Lodish, H. F. (2005). The role of the adipocyte hormone adiponectin in cardiovascular disease. *Current Opinion in Pharmacology*, 5 (2): 129-134
- Hundal, R. S., Krssak, M., Dufour, S., Laurent, D., Lebon, V., Chandramouli, V., Inzucchi, S. E., Schumann, W. C., Petersen, K. F., Landau, B. R. and Shulman, G. I. (2000). Mechanism by which metformin reduces glucose production in type 2 diabetes. *Diabetes*, 49 (12): 2063-2069
- Hyun, M. K. and Jaetaek, K. (2013). The effect of green tea on obesity and type 2 diabetes. *Diabetes and Metabolism Journal.*, 37 (3): 173-175
- Ian, W. C. (2000). Antidiabetic drugs present and future: Will improving insulin resistance benefit cardiovascular risk in type 2 diabetes mellitus? *Drugs*, 60 (5): 1017-1028
- Iozzo, P., Hallsten, K., Oikonen, V., Virtanen, K. A., Parkkola, R., Kemppainen, J., Solin, O., Lonnqvist, F., Ferrannini, E., Knnuti, J. and Nuutila, P. (2003). Effects of metformin and rosiglitazone monotherapy on insulin-stimulated hepatic glucose uptake and their relation to visceral fat in type 2 diabetes. *Diabetes Care*, 26: 2069-2074
- Irwin, N., McKinney, J. M., Bailey, C. J., Flatt, P. R. and McClenaghan, N. H. (2010). Effect of metformin on BRIN-BD11 beta-cell insulin secretory desensitization induced by prolonged exposure to sulphonylureas. *Diabetes, Obesity and Metabolism*, 12: 1066-1071
- Isela, E. J., Juan, C. D., Jorge L. B., Pedro, H. M., Andres, E. C., Carlos, A. T., Arturo, R. H., Hidemi, A. M., Teresa, R. F. and Deysi, Y. B. (2012). Hypoglycemic effect of *Carica papaya* leaves in streptozotocin induces diabetic rats. *BMC complementary and alternative medicine*, 12: 236

- Islam, M. S. and Choi, H. (2007). Green tea, anti-diabetic or diabetogenic: a dose response study. *Biofactors*, 29 (1): 45-53
- Iwaki, M. M., Matsuda, N., Maeda, T., Funahashi, Y., Matsuzawa, M., Makishima, I. and Shimomura. (2003). Induction of adiponectin, a fat derived antidiabetic and antiatherogenic factor, by nuclear receptors. *Diabetes* 52:1655–1663.
- Jacobson, I. B., Henriksen, J. E., Hother-Nielsen, O., Vach, W. and Beck-Nielsen., H. (2009). Evidence-based insulin treatment in type 1 diabetes mellitus. *Diabetes Research and Clinical Practice*, 86 (1): 1-10
- Jalil, A. M. M., Ismail, A., Chong, P. P., Hamid, M. and Kamaruddin, S. H. S. (2008). Effect of Cocoa extract on glucometabolism, oxidative stress and antioxidant enzymes in obese-diabetic (Ob-db) rats. *Journal of Agricultural and Food Chemistry*, 56: 7877-7884
- Joe, M. C. and Arshag, D. M. (2000). A rational approach to drug therapy of type 2 diabetes mellitus. *Drugs*, 60 (1): 95-113
- Joo, E. B., Jalifah, L., Mohd, P. A., Amin, I. And Muhajir, H. (2010). *Scoparia dulcis* (SDF7) endowed with glucose uptake properties on L6 myotubes compared insulin. *Journal of Ethnopharmacology*, 129: 23-33
- Jung, S. H., Ha, Y. J., Shim, E. K., Choi, S. Y., Jin, J. L., Yun-Choi, H. S. and lee, J. R. (2007). Insulin mimetic and insulin sensitizing activities of tanshinones, diterpine compounds of the root of *Salvia miltiorrhiza* Bunge. *Phytomedicine*, 16: 327-335
- Kadowaki, T. T., Yamauchi, N., Kubota, K., Hara, K., Ueki and Tobe, K. (2006). Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome. *Journal of Clinical Investigation*, 116:1784–1792.
- Kalman, D. S., Schwartz, H. I., Feldman, S. and Krieger, D. I. (2013). Efficacy and safety of *Elaeis guineensis* and *Ficus Deltoidea* leaf extracts in adults with pre-diabetes. *Nutrition Journal*, 12:36
- Kamuhabwa, A., Nshimo, C. and De Witte, P. (2000). Cytotoxicity of some medicinal plant extrats used in Tanzanian traditional medicine. *Journal of Ethnopharmacology*, 70 (2): 143-149
- Kang, Y. and Kim, Y. H. (2004). Glucose uptake-stimulatory activity of *Amomi semen* in 3T3-L1 adipocytes. *Journal of Ethnopharmacology*, 92: 103-105
- Kawamura, Y., Tanaka, Y., Kawamori, R. and Maeda, S. (2006). Overexpression of Kruppel-like factor 7 regulates adipocytokine gene expressions in human adipocytes and inhibits glucose-induce insulin secretion in pancreatic- β cell line. *Molecular endocrinology*, 20: 844-856

- Keckskemati, V., Bagi, Z., Pacher, P., Posa, I., Kocksis, E. and Koltai, M. Z. (2002). New trends in the development of oral antidiabetic drugs. *Current medicinal chemistry*, 9: 53-71
- Kesari, A. N., Gupta, R. K., Singh, S. K., Diwakar, S. and Watal, G. (2006). Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *Journal of Ethnopharmacology*, 107: 374-379
- Kirchheiner, J., I. (2005). Effect of genetic polymorphisms in cytochrome p450 (CYP) 2C9 and CYP2C8 on the pharmacokinetics of oral antidiabetic drugs: clinical relevance. *Clinical Pharmacokinetics*, 44 (12): 1209-1225
- Kim, Y. M., Jeong, Y. K., Wang, M. H., Lee, W. Y. and Rhee, H. I. (2005). Inhibitory effect of pine extract on alpha-glucosidase activity and postprandial hyperglycaemia. *Nutrition*, 211: 756-761
- Ko, B. S., Choi, S. B., Park, S. K., Jang, J. S., Kim, Y. E. and Park, S. (2005). Insulin-sensitizing and insulinotropic action of Berberin from *Cortidis Rizhoma*. *Biological & Pharmaceutical Bulletin*, 28 (8): 1431-1437
- Konrad, R. J., Mikolaenko, I., Tolar, J. F., Liu, K. and Kudlow, J. E. (2001). The potential mechanism of the diabetogenic action of streptozotocin: inhibition of pancreatic beta cell O-GlcNac-selective N-acetyl-beta-D-glucosaminidase. *Biochemical Journal*, 356 (1): 31-41
- Kroemer, G., El-Deiry, W. S., Golstein, P., Peter, M. E., Vaux, D., Vandenabeele, P., Zhivotovsky, B., Blagosklonny, M. V., Malorni, W., Knight, R. A., Piacentini, M., Nagata, S. and Melino, G. (2005). Classification of cell death: recommendations of the Nomenclature committee on Cell Death. *Cell death and differentiation*, 12 (S2): 1463-1467
- Kubota, N., Terauchi, Y., Kubota, T., Kumagai, H., Itoh, S., Satoh, H., Yano, W., Ogata, H., Tokuyama, H., Takamoto, I., Mineyama, T., Ishikawa, M., Moroi, M., Sugi, K., Yamauchi, T., Ueki, K., Tobe, K., Noda, T., Nagai, R. and Kadowaki, T. (2006). Pioglitazone ameliorate insulin resistance and diabetes by both adiponectin-dependent and independent pathways. *The Journal of Biological Chemistry*, 281: 8748-8855
- Li, Y., Wen, S., Kota, B. P., Peng, G., Li, G. Q., Yamaha, J. and Roufogalis, B. D. (2005). Punica granatum flower extract. A potent α -glucosidase inhibitor, improves postprandial hyperglycemia in Zucker diabetic fatty rats. *Journal of Ethnopharmacology*, 99: 239-244
- Li, R., Thorens, B. and Loeken, M. R. (2007). Expression of the gene encoding the high Km glucose transporter 2 by the early postimplantation mouse embryo is essential for neural tube defects associated with diabetic embryopathy. *Diabetologia*, 50(3): 682-689

- Lihn, A. S., Bruun, J. M., He, G., Pederson, S. B., Jensen, P. F. and Richelsen, B. (2004). Lower expression of adiponectin mRNA in visceral adipose tissue in lean and obese subjects. *Molecular and Cellular Endocrinology*, 219 (1-2): 9-15
- Lihn, A. S., Pederson, S. B. and Richelson, B. (2005). Adiponectin: action, regulation and association to insulin sensitivity. *Obesity Reviews*, 6 (1): 13-21
- Limbert, C., Path, G., Jakob, F. and Seufert, J. (2008). Beta-cell replacement and regeneration: Strategies of cell-based therapy for type 1 diabetes mellitus. *Diabetes Research and Clinical Practice*, 79: 389-399
- Liu, B., Asare-Anane, H., Al-Romaiyan, A., Huang, G., Amiel, S. A., Jones, P. M. and Persaud, S. J. (2009). Characterisation of the insulinotropic activity of an aqueous extract of *Gymnema sylvestre* in mouse beta cells and human islet of Langerhans. *Cellular Physiology and Biochemistry*, 23 (1-3): 125-132
- Lotta, W., Billie, P., Ulf, R., Anna, D., Gunilla, J. and Par, W. (2014). Impact of symptomatic hypoglycemia on medication adherence, patient satisfaction with treatment, and glycemic control in patients with type 2 diabetes. *Patient preference and adherence*, 8: 593-601
- Mahinda, A., Irine, R., Mohd, N., Suhaila, M. and Richard, H. (2002). Polyphenol-enriched extract of oil palm fronds (*Elais guineensis*) promotes vascular relaxation via endothelium-dependent mechanisms. *Asia Pacific Journal of Clinical Nutrition*, 11: 467-472
- Malaisse, W. J. (1999). Mechanism of action of a new class of insulin secretagogues. *Experimental and Clinical Endocrinology & Diabetes*, 107: 140-143
- Martineau, L. C., Couture, A., Spoor, D., Ali, B. A., Harris, C., Meddah, B., Leduc, C., Burt, A., Vuong, T., Phuong, M. L., Prentki, M., Bennet, A. S., Arnason, J. T. and Haddad, P. S. (2006). Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. *Phytomedicine*, 13: 612-623
- Materska, M. and Perucka, L. (2005). Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L.) *Journal of Agriculture and food chemistry*, 53: 1750-1756
- Mathews, J. N., Flatt, P. R. and Abdel-Wahab, Y. H. (2006). *Asparagus adscendens* (Shweta musali) stimulates insulin secretion, insulin action and inhibits starch digestion. *British Journal of Nutrition*, 95(3): 576-581
- Matsuzawa, Y. (2005). Adiponectin: Identification, physiology and clinical relevance in metabolic and vascular disease. *Atherosclerosis Supplement*, 6 (2): 7-14
- Mehanna, A. S. (2005). Insulin and oral antidiabetic agents. *American Journal of Pharmaceutical Education*, 69 (5): 1-11

- Merghoub, M., Benbacer, L., Amzazi, S., Morjani, H. and El Mzibiri, M. (2009). Cytotoxic effect of some Moroccan medicinal plant extracts on human cervical cell lines. *Journal of Medicinal Plants Research*, 3 (12): 1045-1050
- Micheal J. Fowler, M. D. (2010). Diabetes treatment: oral agents. *Clinical diabetes*, 28 (3): 132-136
- Motoshima, H., Wu, X., Sinha, M. K., Hardy, V. E., Rosato, E. L., Barbot, D. J., Rosato, F. E. and Goldstein, B. J. (2002). Differential regulation of adiponectin secretion from cultured human omental and subcutaneous adipocytes: effects of insulin and rosiglitazone. *Journal of Clinical Endocrinology & Metabolism*, 87 (12): 5662-5667
- Mustaffa, F., Indurkar, J., Ali, N. I. M., Hanapi, A., Shah, M., Ismail, S. and Mansor, S. M. (2011). A review of Malaysian medicinal plants with potential antidiabetic activity. *Journal of Pharmacy research*, 4 (11): 4217
- Nagendran, B., Tan, Y. A., Ravigadevi, S., Kalyana, S. and Samir, S. (2005). Antioxidant properties of palm fruit extracts. *Asia Pacific Journal of Clinical Nutrition*, 4(4): 319-324
- Nathan, D. M., Buse, J. B., Davidson, M. B., Heine, R. J., Holman, R. R., Sherwin, R. and Zinman, B. (2006). Management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement from American Diabetes Association and European Association for the Study of Diabetes. *Diabetes care*, 29: 759-764
- Neo, Y. P., Aziz, A., Tan, C. P. and Tan Y. A. (2010). Phenolic acid analysis and antioxidant activity assessment of oil palm (*E. Guineensis*) fruit extracts. *Food chemistry*, 122: 353-359
- Nomikos, T., Detopoulou, P., Fragopoulou, E., Pliakis, E. and Antonopoulou, S. (2007). Boiled wild artichoke reduces postprandial glycemic and insulinemic responses in normal subjects but has no effect on metabolic syndrome patients. *Nutrition research*, 27 (12): 741-749
- Nugent, C., Prins, J. B., Whitehead, J. P., Savage, D., Wenworth, J. M., Chatterjee, V. K. and O'Rahilly, S. (2001). Potentiation of glucose uptake in 3T3-L1 adipocytes by PPAR gamma agonists is maintained in cells expressing a PPAR gamma dominant-negative mutant: evidence for selectivity in the downstream responses to PPAR gamma activation. *Molecular Endocrinology*, 15 (10): 1729-1738
- Ooyub, S., Ismail, F. and Daud, N. A. (2004). Diabetes program in Malaysia – current and future. *NCD Malaysia*, 3(2): 6-12

- Ortiz-Andrade, R. R., Garc'ia-Jim' Enez, S., Castillo-Espa~Na, P., Ram'Irez-A'Villa, G., Villalobos-Molina, R. And Estrada-Soto, S. (2007). r-Glucosidase inhibitory activity of the methanolic extract from *Tournefortia hartwegiana*: An anti-hyperglycemic agent. *Journal of Ethnopharmacology*, 109: 48-53
- Parthasarathy, C., Renuka, V. N. And Balasubramaniam, K. (2009). Sex steroids enhance insulin receptors and glucose oxidation in Chang liver cells. *Clinica Chimica Acta*, 399 (1-2): 49-53
- Parveen, F. R., Rajeev, P. S., Ibrahim, M. H. and Norizan, E. (2010). Review of current palm oil mill effluent (POME) treatment methods: vermicomposting as a sustainable practise. *World Applied Science Journal*, 11 (1): 70-81
- Patel, M. B. and Mishra, S. H. (2008). Cell line in diabetic research: A review. *Pharmacognosy Review*, 2 (4): 188-205
- Pieper, A.A., Verma, A., Zhang, J. and Snyder, S.H. (1999). Poly (ADP-ribose) polymerase, nitric oxide and cell death. *Trends in Pharmacological Sciences*, 20:171-181
- Pinent, M., Blay, M., Blade, M. C., Salvado, M. J., Arola, L. and Ardevol, A. (2004). Grape seed derived procyanidins have an antihyperglycemic effect in streptozotocin-induced diabetic rats and insulinomimetic activity in insulin sensitive cell lines. *Endocrinology*, 145 (11): 4985-4990
- Polixeni, A. and Vassilis, J. D. (2010). Medicinal plants used for the treatment of diabetes and its long-term complications. *Plants in Traditional and Modern Medicine: Chemistry and Activity*, 169-175
- Rajavel, V., Munavvar, Z. A. S., Ivy, C., Mahmood, A. A., Normadiah, M. K. and Nor, A. A. (2013). Antioxidant and pro-oxidant effect of oil palm (*Elaeis guineensis*) leaves extract in experimental diabetic nephropathy: a duration-dependent outcome. *BMC complementary and alternative medicine*, 13: 242
- Rees, D. A. and Alcolado, J. C. (2005). Animal models of diabetes mellitus. *Diabetic Medicine*, 22: 359-370
- Rendall, M. (2004). The role of sulfonylureas in the management of type 2 diabetes. *Drugs*, 64: 1339-1358
- Rengarajan, S., Parthasarathy, C., Anitha, M. and Balasubramaniam, K. (2007). Diethyl phthalate impairs insulin binding and glucose oxidation in Chang liver cells. *Toxicology in Vitro*, 21: 99-102
- Reusch, J. E. B. (2003). Diabetes, microvascular complications and cardiovascular complications: What is it about glucose? *Journal of Clinical Investigation*, 112: 986-988

- Romero-Perez, A.I., Lamuela-Raventós, R.M., Andrés-Lacueva, C. and Carmen de la Torre-Boronat, M. (2001). Method for the quantitative extraction of resveratrol and piceid isomers in grape berry skins: effect of powdery mildew on the stilbene content. *Journal of Agricultural and Food Chemistry*, 49, 210–215
- Rosalina, Tan, R. T., Mohamed, S., Samaneh, G. F., Noordin, M. M., Goh, Y. M. and Manap, M. Y. A. (2011). Polyphenol rich oil palm leaves extract reduce hyperglycaemia and lipid oxidation in STZ-rats. *International food research journal*, 18: 179-188
- Roy, M. K., Takenaka, M., Kobori, M., Nakahara, K., Isobe, S. and Tsushida, T. (2006). Apoptosis, necrosis and cell proliferation-inhibition by cyclosporine A in U937 cells (a human monocytic cell line). *Pharmacological Research*, 53 (3): 293-302
- Ryan, C. M. (2002). Hormone-behavior relations of clinical importance: diabetes mellitus. In: *Hormones, Brain and Behaviour*. Edited by Donald W. Pfaff. Elsevier Science. *Academic Press, USA*, pp: 553-592
- Sabino, K. C. C., Gayer, L. C. A., Vaz, L. R. L. and Santos, I. (1999). *In vitro* and *in vivo* toxicological study of the *Pterodon pubescens* seed oil. *Toxicology Letters*, 108: 27-35
- Sachdewa, A. and Khemani, L.D. (2003). Effect of *Hibiscus rosa sinensis* Linn. Ethanol flower extract on blood glucose and lipid profile in streptozotocin induced diabetes in rats. *Journal of Ethnopharmacology*, 89: 61-66
- Sakurai, K., Kawazuma, M., Adachi, T., Harigaya, T., Saito, Y., Hashimoto, N. and Mori, C. (2004). Bisphenol A affects glucose transport in mouse 3T3F442A adipocytes. *British journal of Pharmacology*, 141 (2): 209-214
- Saltiel, A. R. and Pessin, J. E. (2002). Insulin signalling pathways in time and space. *Trends in Cell Biology*, 12 (2): 65-71
- Satake, S., Moore, M. C., Igawa, K., Converse, M., Farmer, B., Neal, D. W. and Cherrington, A. D. (2002). Direct and indirect effects of insulin on glucose uptake and storage by the liver. *Diabetes*, 51 (6): 1663-1671
- Sell, H., Dietze-Schroeder, D. and Eckel, J. (2006). The adipocyte-myocyte axis in insulin resistance. *Trends in Endocrinology & Metabolism*, 17 (10): 416-22
- Shamala, D. S., Soon, S. L., Najwa, A., Kok, K. T., Kalyana, S., Ragivadevi, S. and Mohd, B. W. (2010). Effects of oil palm phenolics on tumor cells in vitro and in vivo. *African journal of food science*, 4 (8): 495-502
- Sharma, V. K., Kumar, S., Patel, H. J. and Hugar, S. (2010). Hypoglycemic activity of *Ficus glomerata* in alloxan induced diabetic rats. *International Journal of Pharmaceutical Sciences Review and Research*. 1 (2): 180-022

- Sharon, W. and Lahiri, M. D. (2012). Management of type 2 diabetes: What is the next step after metformin? *Clinical diabetes*, 30 (2): 72-75
- Snidjer, M. B., Heine, R. J., Seidell, J. C., Bouter, L. M., Stehouwer, C. D., Nijpels, G., Funahashi, T., Matsuzawa, Y., Shimomura, I. And Dekker, J. M. (2006). Associations of adiponectin levels with incident impaired glucose metabolism and type 2 diabetes in older men and women: the hoorn study. *Diabetes care*, 29 (11): 2498-2503
- Spranger, J., Kroke, A., Mohlig, M., Bergmann, M. M., Ristow, M., Boeing, H. and Pfeifer, A. F. (2003). Adiponectin and protection against type 2 diabetes mellitus. *Lancet*, 361 (9553): 226-228
- Sundram, K., Sambarthamurti, R. and Tan, Y. A. (2003). Palm fruit chemistry and nutrition. *Asia Pacific Journal of Clinical Nutrition*, 12 (3): 355-362
- Sundram, K., Thiaharajan, T., Gapor, A. and Basiron, Y. (2002). Palm tocotrienols: new antioxidants for the new millennium. *Inform*, 13: 634-641
- Szkudelski, T. (2001). The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiological Research*, 50: 537-546
- Tan, Y. A., sundram, K. and Sambanthamurthi R. (2001). Water-soluble phenolics from the palm oil industry. In: Pfannhauser W, Fenwick G. R., Khokar S., eds. Biologically-active Phytochemicals in food – analysis, metabolism, bio-availability and function. Cambridge: *The royal society of Chemistry*; 548-551
- Troy, L. M. and Glenn, K. M. (2009). Skeletal muscle glucose uptake during exercise: A focus on reactive oxygen species and nitric oxide signalling. *Life*, 61(5): 479-484
- Trujillo, M. E. and Scherer P. E. (2005). Adiponectin: journey from an adipocyte secretory protein to biomarker of the metabolic syndrome. *Journal of Internal Medicine*, 257:167–175.
- Tschritter, O., Fritsche, A., Shirkavand, F., Machicao, F., Haring, H. and Stumvoll, M. (2003). Assessing the shape of glucose curve during an oral glucose tolerance test. *Diabetes care*, 26 (4): 1026-1033
- Tzeng, Y. M., Chen, K., Rao, Y. K. and Lee, M. J. (2009). Kaempferitin activates the insulin signalling pathway and stimulates secretion of adiponectin in 3T3-L1 adipocytes. *European Journal of Pharmacology*, 607 (1-3): 27-34
- Uebanso, T., Arai, H., Taketani, Y., Fukaya, M., Yamamoto, H., Mizuno, A., Uryu, K., Hada, T. and Takeda, E. (2007). Extracts of *Mimordica charantia* suppress postprandial hyperglycaemia in rats. *Jornal of Nutritional Science and Vitaminology*, 53: 482-488

- Vats, V., Yadav, S. P. and Grover, J. K. (2004). Ethanolic extract of *ocimum sanctum* leaves partially attenuates streptozotocin-induced alterations in glycogen content and carbohydrate metabolism in rats. *Jornal of Ethnopharmacology*, 90: 155-160
- Vijayakumar, M. V., Singh, S., Chhipa, R. R. and Bhat, M. K. (2005). The hypoglycaemic activity of fenugreek seed extract is mediated through the stimulation of an insulin signalling pathway. *British Journal of Pharmacology*, 146: 41-48
- Viljanen, A. P., Virtanen, K. A., Jarvisalo, M. J., Hallsten, K., Parkkola, R., Ronnema, T., Lonnqvist, F., Iozzo, P., Ferrannini, E. and Nuutila, P. (2005). Rosiglitazone treatment increases subcutaneous adipose tissue glucose uptake in parallel with perfusion in patients with type 2 diabetes: a double-blind, randomized study with metformin. *Journal of Clinical Endocrinology & Metabolism*, 90(12): 6523-6528
- Wang, S. Y. and Ballington, J. R. (2007). Free radical scavenging capacity and antioxidant enzyme activity in deer berry (*Vaccinium stamineum* L.) *LWT-Food Science and Technology*, 40: 1352-1361
- Wang, H. & Helliwell, K. (2001). Determination of flavonols in green and black tea leaves and green tea infusion by high performance liquid chromatography. *Food Research International*, 34, 223-227
- Watson, R. T. and Pessin, J. E. (2006). Bridging the GAP between insulin signalling and GLUT 4 translocation. *Trends in Biochemical Sciences*, 31 (4): 215-222
- Wiecek, A., Adamczak, M. and Chudek, J. (2007). Adiponectin- an adipokine with unique metabolic pproperties. *Nephrology Dialysis Transplantation*, 22 (4): 981-988
- Wild, S., Roglic, G., Green, A., Sicree, R. and King, H. (2004). Global prevalence of Diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes care*, 27: 1047-1053
- Xu, A., Wang, H., Hoo, R. L., Sweeney, G., Vanhoutte, P. M., Wang, Y., Wu., D., Chu, W., Qin, G. and Lam, K. S. (2009). Selective elevation of adiponectin production by the natural compounds derived from a medicinal herb alleviates insulin resistance and glucose tolerance in obese mice. *Endocrinology*, 150 (2): 625-633
- Yamauchi, T., kamon, J., Waki, H., Terauchi, Y., Kubota, N., Hara, K., Mori, Y., Ide, T., Murakami, K., Tsuboyama-Kasaoka, N, Ezaki, O, Akanuma, Y., Gavrilova, O., Vinson, C., Reitman, M. L., Kagechika, H., Shudo, K., Yoda, M., Nakano, Y., Tobe, K., Nagai, R., Kimura, S., Tomita, M., Froguel, P. and Kadowaki, T. (2001). The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoathropy and obesity. *Nature Medicine*, 7 (8): 941-946

- Yousef, W. M., Omar, A. H., Morsy, M. D., El-Wahed, M. M. A. and Ghanayem, N. M. (2005). The mechanism of action of calcium channel blockers in the treatment of diabetic nephropathy. *International Journal of Diabetes & Metabolism*, 13: 76-82
- Zanariah, H. Chandran, L. R., Wan Mohamad, W. B., Wan Nazaimoon, W. M., Letchuman, G. R., Jamaiah, H. Fatanah, I. Nurain, M. N. Helen Tee, G. H. and Mohd Rodi, I. (2008). DWP1-3 Prevalence of diabetes mellitus in Malaysia in 2006 - Results of the 3rd National Health and Morbidity Survey (NHMS III). *Diabetes Research and Clinical Practice*, 79 (S1): S21
- Zangeneh, F., Kudva, Y. C. and Basu, A. (2003). Insulin sensitizers. *Mayo Clinic Proceedings*, 78 (4): 471-479
- Ziyou, C., Xiulan, C., Bingwen, L., Sung, K. P., Tao, X., Zhensheng, X., Peng, X., Junjie, H., Haiying, H., John, R. Y. and Fuquan, Y. (2009). Preliminary quantitative profile of differential expression between rat L6 myoblast and myotubes by stable isotope labelling by amino acids in cell culture. *Proteomics*, 9(5): 1274-1292
- Zou, C. and Shen, Z. (2007). An optimized in vitro assay for screening compounds that stimulate liver cell glucose utilization with low cytotoxicity. *Journal of Pharmacological and Toxicological Methods*, 56 (1): 58-62
- Zygmunt, K., Faubert, B., MacNeil, J. and Tsiani, E. (2010). Naringenin, a citrus flavonoid, increases muscle cell glucose uptake via AMPK. *Biochemical and Biophysical Research Communications*, 398 (2): 178-183