



UNIVERSITI PUTRA MALAYSIA

***CHARACTERISATION AND PATHOGENICITY OF *Neofusicoccum ribis*
AND GENE EXPRESSION STUDY OF PATHOGENESIS –RELATED GENES
IN RUBBER (*Hevea brasiliensis* Muell.Arg)***

NYAKA NGOBISA AURELIE IRENE CLAIRE

FP 2014 22



**CHARACTERISATION AND PATHOGENICITY OF *Neofusicoccum ribis*
AND GENE EXPRESSION STUDY OF PATHOGENESIS –RELATED
GENES IN RUBBER (*Hevea brasiliensis* Muell.Arg)**

By

NYAKA NGOBISA AURELIE IRENE CLAIRE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

February 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of University Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of University Putra Malaysia.

Copyright © Universiti Putra Malaysia

© COPYRIGHT UPM

DEDICATION

The work in this thesis is dedicated to the memory of my loving father Nyaka Emmanuel (28/12/1939-19/05/2009) and to the entire Nyaka family. To my husband Mandengue Louis Lucien Seurard for his boundless love, understanding, encouragement and patience throughout my study

Abstract of thesis presented to the senate of Universiti Putra Malaysia in fulfilment of the requirement of the degree of Doctor of Philosophy

**CHARACTERISATION AND PATHOGENICITY OF *Neofusicoccum ribis*
AND GENE EXPRESSION STUDY OF PATHOGENESIS-RELATED GENES
IN RUBBER (*Hevea brasiliensis* Muell.Arg)**

By

NYAKA NGOBISA AURELIE IRENE CLAIRE

February 2014

**Chairman: Associate Professor Zainal Abidin Mior Ahmad, PhD
Faculty: Agriculture**

Rubber tree (*Hevea brasiliensis*) is well known as the tree of life for its by-products such as latex and wood. However, the growth and performances of *H. brasiliensis* are reduced substantially by Fusicoccum Leaf Blight disease, a re-emerging malady first detected in Malaysia in 1987. This study provides a foundation for understanding the infection of healthy rubber plant tissues by *Fusicoccum* spp. in selected regions in Malaysia, with emphasis on pathogenicity and cross pathogenicity evaluation of associated hosts. The study provides a leaf colonisation pattern since the resistance mechanism provided by pathogenesis-related proteins (PR) and phenylalanine ammonia lyase (PAL) enzymes, both induced in infected rubber, are fairly well studied. Fungal isolates were obtained from leaf samples collected in several rubber plantations in Selangor, Perak and Johor in 2010. All isolates were identified based on cultural characteristics such as conidial production in various cultural media (PDA, MEA, CDA, CMA) and sequence data generated from the internal transcribed spacers (ITS1 and ITS2), the 5.8S rRNA gene and an unknown locus (BotF15) containing microsatellite repeats. Pathogenicity tests were conducted using representative isolates from three locations (SB30, SK10 and SJ20) for confirmation by detached leaf method and seedlings in glasshouse, and cross pathogenicity tests on detached leaves. The roles of β -1, 3-glucanase and chitinase, the two most widely studied groups of pathogenesis-related (PR) proteins (PR3 and PR2) and phenylalanine ammonia lyase (PAL) enzymes present during pathogenesis were investigated.

Thirty three isolates were derived from this study. In different culture media and pH tested, significant differences were observed in spore production, dimensions of the conidia and growth rates among isolates. On PDA, pH 6 supported the best growth compared to others and optimum temperatures of growth were between 25 °C to 30 °C. The growth rate was 21.2 mm/day on PDA, 17.9 mm/day on CDA, 17.1 mm / day on MEA and 14.2 mm/day on CMA. The highest spore production of 161.33×10^6 spores / ml was obtained on PDA. The optimum temperature for spore germination in distilled water was 25 °C – 35 °C within a period of twenty- four hours. The fungus colony color was white to grey; dark greenish grey or greyish.

Conidia were either septate or aseptate, hyaline when young and turning brown with age, oval, elliptic to ovaloid or subglobose in shape. Conidia size ranged 9.0-25.8 μm in length and 1.9-8.5 μm width with the L: B ratio of 3.4.

Molecular characterisation revealed fungal sequences generated from ITS and BotF15 genes, indicating therefore that they formed a supported clade with *Neofusicoccum. ribis*. Tested isolates (SB30, SK10 and SJ20) were pathogenic to *H. brasiliensis* but varied in symptoms. Pathogenicity tests showed high susceptibility of percentage of disease severity on detached rubber leaves from clones RRIM 2024 and PB 350 (39 - 56%) and low susceptibility on clones RRIM 2002 and RRIM 2007 (6.6 - 18.8%). Some alternate hosts such as mango and guava exhibited high susceptibility (50 - 80%) while others (rambutan and papaya) were low (3 - 10%).

Induction of pathogenesis-related genes in rubber leaves in the presence of fungal pathogen showed that patterns of expressions of these enzymes were up-regulated in both tolerant (RRIM 2002) and susceptible (PB 350) rubber clones over time although their expression was much higher in the latter. Generally, pathogenesis-related proteins from *H. brasiliensis* could be used as molecular probes to monitor defence-response activation while the mechanism in *N. ribis* isolate SK10 could serve as an inducer of systemic resistance in rubber.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENCIRIAN DAN PATOGENISITI *Neofusicoccum ribis* DAN KAJIAN
EKSPRESI GEN PERKAITAN-PATOGENESIS DALAM GETAH
(*Hevea brasiliensis* Muell.Arg)**

Oleh

NYAKA NGOBISA AURELIE IRENE CLAIRE

Februari 2014

Pengerusi: Profesor Madya Zainal Abidin Mior Ahmad, PhD

Fakulti: Pertanian

Pokok getah (*Hevea brasiliensis*) lebih dikenali sebagai pokok hayat bagi produk sampingan seperti getah dan kayu. Walau bagaimanapun, pertumbuhan dan perkembangan *H. brasiliensis* terjejas dengan ketara disebabkan oleh penyakit daun hawar *Fusicoccum*, penyakit yang dikesan muncul semula di Malaysia pada tahun 1987. Kajian ini memberi asas pemahaman jangkitan *Fusicoccum* spp. terhadap tisu tumbuhan getah yang sihat di kawasan terpilih di Malaysia, dengan penekanan kepada patogenisiti dan penilaian patogenisiti silang pada hos yang berkaitan. Kajian ini adalah wajar dalam menerangkan corak kolonisasi daun apabila mekanisme rintangan terhasil dari protein yang berkaitan dengan patogenesis (PR) dan enzim ammonia phenylalanine lyase (PAL) yang mana kedua-duanya disebabkan oleh getah yang telah dijangkiti.

Pengasingan kulat diperolehi daripada sampel daun yang dikutip di beberapa ladang getah di Selangor, Perak dan Johor pada tahun 2010. Semua pengasingan telah dikenal pasti berdasarkan ciri-ciri kultur seperti pengeluaran conidial dalam pelbagai medium kultur (PDA, MEA, CDA, CMA) dan urutan data yang terhasil daripada ruang salinan dalaman (ITS1 dan ITS2), gen rRNA 5.8S dan lokus yang tidak diketahui (BotF15) yang mengandungi mengulangi mikrosatelit. Ujian patogenisiti telah dijalankan menggunakan pengasingan mewakili tiga lokasi (SB30, SK10 dan SJ20) untuk pengesahan dengan kaedah helaian daun dan benih di rumah kaca, dan ujian silang patogenisiti pada daun berkembar. Fungsi β -1, 3-glucanase dan chitinase, dua kumpulan ezmin yang dikaji paling meluas yang berkaitan dengan patogenesis (PR) protein (PR3 dan PR2) dan penyediaan enzim lyase ammonia phenylalanine (PAL) dalam patogenesis telah dikaji.

Tiga puluh tiga pengasingan telah diperolehi daripada kajian ini. Dalam medium kultur berbeza dan pengujian pH, perbezaan ketara telah diperhatikan dalam pengeluaran spora, dimensi konidia dan kadar pertumbuhan di kalangan pengasingan. Pada PDA, pH 6 menyokong pertumbuhan yang terbaik berbanding

dengan yang lain dan suhu optimum pertumbuhan adalah di antara 25 °C hingga 30 °C. Kadar pertumbuhan adalah 21.2 mm / hari untuk PDA, 17.9 mm / hari untuk CDA, 17.1 mm / hari MEA dan 14.2 mm / hari untuk AKM. Pengeluaran tertinggi spora 161.33×10^6 spora / ml telah diperoleh untuk PDA. Suhu optimum untuk percambahan spora di dalam air suling adalah antara 25 °C - 35 °C dalam tempoh dua puluh empat jam. Warna koloni kulat adalah putih kelabu, kelabu kehijauan gelap atau kekelabuan. Konidia sama ada septate atau aseptate, hialin ketika muda dan bertukar perang pada usia, bujur, eliptik ke ovaloid atau berbentuk subglobose. Saiz panjang konidia antara 9.0-25.8 μm dan lebar antara 1.9-8.5 μm dengan L: B nisbah 3.4.

Pencirian molekul mendedahkan urutan kulat terhasil daripada ITS dan gen BotF15, ini menunjukkan bahawa kulat tersebut membentuk klad yang baik yang dibantu oleh *Neofusicoccum. ribis*. Hasil ujian penasingan (SB30, SK10 dan SJ20) adalah patogenik pada *H. brasiliensis* tetapi berbeza pada simptom. Ujian Patogenisiti menunjukkan kecenderungan yang tinggi pada daun getah berkembar dari klon RRIM 2024 dan PB 350 (39 - 56%) dan kecenderungan rendah pada klon RRIM 2002 dan RRIM 2007 (6.6-18.8%). Seseengah alternatif hos rumah seperti mangga dan jambu menunjukkan kecenderungan yang tinggi (50 - 80%) manakala yang lain (rambutan dan betik) adalah rendah (3 - 10%). Induksi gen yang berkaitan dengan patogenesis dalam getah daun tanpa kulat patogen menunjukkan bahawa corak ungkapan enzim ini adalah up-selia kedua-dua toleran (RRIM 2002) dan mudah (PB 350) klon getah dari semasa ke semasa walaupun ungkapan mereka adalah jauh lebih tinggi dalam kedua. Secara umumnya, protein yang berkaitan dengan patogenesis daripada *H. brasiliensis* boleh digunakan sebagai probe molekul untuk mengawal pengaktifan tindak balas pertahanan manakala mekanisme untuk ketegangan *N. ribis* SK10 yang mampu bertindak sebagai pencetus penentangan sistemik dalam getah.

ACKNOWLEDGEMENTS

Thanks to the almighty GOD for his continuous guidance and for giving me the courage and determination to complete this study.

This thesis represents the fruit of endless effort by many people and the support of several institutions to which I'd like to extend my gratitude. I wish to express my gratitude to my supervisor Assoc. Prof. Dr. Zainal Abidin Mior Ahmad who was abundantly helpful and offered invaluable assistance, patience, understanding, support and guidance. His scientific intuition, idea for my project and passion for science has inspired me and enriched my growth as a student and a researcher. I am looking forward to future collaboration.

To my co-supervisors Assoc. Prof. Dr. Wong Mui Yun and Assoc. Prof. Dr. Wan Mohamed Noordin Wan Daud without whose knowledge, advice and crucial assistance this would not have been successful. They have triggered and nourished my intellectual maturity, from which I will benefit for a long time to come.

I am very indebted to the Malaysian Rubber Board. I also would like to record our appreciation to Dr. Mohd Nasarudin bin Mohd Aris, the Head of Crop Improvement and Protection Unit of the Malaysian Rubber Board and, Mrs Murnita Masahuling for their permission, and assistance to conduct this research.

Deepest gratitude are due to Datuk Dr. Abdul Aziz S.A. Kadir, the Secretary General of the International Rubber Research and Development Board (IRRDB) and Dr. Ismail Hashim who carried out all necessary formalities and requirements for me to conduct this study.

My appreciation goes to Camerooninan Government, specifically the Institute of Agricultural Research for Development (IRAD) who made possible my coming to Malaysia and provided me logistic support each time it was needed during field work. Especially warm thanks to Dr Zok Simon the former General Manager of IRAD, Dr Woin Noe the General manger of IRAD, Dr Mouen Bedimo Joseph Deputy General manger of IRAD, Dr Ehabe Eugene Ejolle, principal researcher and Head of Latex Plants latex programme IRAD; Dr Begoude Boyogueno Aime-Didier, principal researcher from the Regional Biological Control and Applied Microbiology Laboratory of IRAD.

During my study, I had a number of excellent colleagues and friends. Special mention to Dr Boubakar Barry, Coordinator of Association of African Universities, Ghana; Dr Abubakar Muhammad Salisu, lecturer at Usmanu Danfodiyo University, Sokoto-Nigeria; Dr Mfegue Virginie research officer at IRAD, Cameroon; Beyegue Djonko Honore lecturer at the University of Dschang, Cameroon and, PhD fellow at Universiti Sains Islam Malaysia; Michael Mbenoun PhD fellow at Forestry and Agricultural Biotechnology Institute at University of Pretoria, South Africa, Eunice

Adgidzi PhD fellow at Univerity of Nigeria Nsuk ka for their constant presence, care and moral support.

Special thanks are also given to staff at the Department of Plant Protection, Faculty of Agriculture, Universiti Putra Malaysia and at the International Research Rubber Developpment Board, especially to Mr Shamsudin Bojang, Mr Mohamed Nazri Abdul Rahman, Mrs Asmalina Abu Bakar and Mr Johari Mohd Sarikat, Mrs Norani Saad, Mr Aizat Hazman Mohamed Khairi who have assisted me in various ways.

I am tempted to individually thank all my lab mates at UPM, and RRIM, but as the list is long and for fear I might omit someone, I will simply and genuinely say: Thank you to you all for your positive input, cheerful attitude, care and trust. Particular I would like to cite Ms Salha Elmahdi, Mr Abbas Nashi, Mrs Farnaz Abed, Ms Dzarifah Zulperi, Mr Mamma Sawaneh, Ms. Azimah Izhar, Mrs Khairita Khairuddin, Mr Soni Othman, Mr Sega, Mr Guna and Mrs. Erusama.

I am greatly indebted to my family and in-laws for their inseparable support and prayers. Their dedication, love and persistent confidence in me, has taken the load off my shoulder. Words fail me to express my appreciation to my beloved husband, Mandengue Louis Lucien Heurard, as well as to my lovely kids (Louis-Simon and Charles-Emmanuel) who are for me, more than water in the desert.

Finally, thank to all those who contributed to the successful realization of this thesis and my apologies to those that I could not mention here.

I certify that a Thesis Examination Committee has met on 17 February 2014 to conduct the final examination of Nyaka Ngobisa Aurelie Irene Claire on her thesis entitled "Characterization and Pathogenicity of *Neofusicoccum ribis* and Gene Expression Study of Pathogenesis-Related Genes in rubber (*Hevea brasiliensis* Muell.Arg.)" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Kamaruzaman bin Sijam, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Ganesan a/l Vadamalai, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Ho Chai Ling, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Paul W J Taylor, PhD

Associate Professor
University of Melbourne
Australia
(External Examiner)



NORITAH OMAR, PhD

Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 10 March 2014

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Zainal Abidin Mior Ahmad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Wong Mui Yun, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Wan Mohamed Noordin Wan Daud, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

Declaration by the student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature:..... Date: 17th February 2014

Name and Matric No: NYAKA NGOBISA AURELIE IRENE CLAIRE, (GS26788)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as slated in Rule 41 in Rules 2003 (Revision 2012-2013) were adhered to.

Signature -----

Zainal Abidin Mior Ahmad, PhD

(Chairman)

Signature -----

Wong Mui Yun, PhD

(Member)

Signature -----

Wan Mohamed Noordin Wan Daud, PhD

(Member)

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 The Rubber Tree Profile	4
2.2 Diseases of rubber	5
2.2.1 Fusicoccum Leaf Blight (FLB) disease on rubber	7
2.3 The Botryosphaeriaceae Fungi	8
2.3.1 Taxonomy and nomenclature of <i>Neofusicoccum</i> spp.	9
2.3.1.1 Morphology of <i>Neofusicoccum</i> spp.	10
2.3.1.2 Molecular methods	12
2.3.1.3 Microsatellite markers	13
2.3.1.4 Use of other molecular tools	13
2.4 Biology of <i>Neofusicoccum</i> spp.	14
2.4.1 Distribution and host range	14
2.4.2 Disease symptoms caused by <i>Neofusicoccum</i> spp.	15
2.4.3 Spore development, dispersal and epidemiology	16
2.5 Plant defence responses to pathogens	17
2.5.1 Induced resistance in plants	18
2.5.1.1 Systemic acquired resistance (SAR)	19
2.5.1.2 Induced systemic resistance (ISR)	20
2.5.2 Pathogenesis-related (PR) proteins in induced resistance	21
2.5.3 Phenylalanine ammonia-lyase (PAL) gene	23
2.5.4 Defence enzymes in rubber	24
3 ISOLATION AND MORPHOLOGICAL CHARACTERISATION OF <i>NEOFUSICOCCUM</i> SPP.	26
3.1 Introduction	26
3.2 Materials and Methods	27
3.2.1 Collection of samples	27
3.2.2 Isolation of leaf blight fungus	28

	3.2.2.1	Cultures morphology	29
3.2.3		Morphological characteristics of cultured fungal colonies	29
	3.2.3.1	Effect of media on colony growth	29
	3.2.3.2	Effect of media on pH sporulation	30
	3.2.3.3	Effect of temperature on germination of spores	30
	3.2.4	Experimental design and data analysis	30
3.3		Results and Discussion	31
	3.3.1	Samples symptoms	31
	3.3.2	Isolation of leaf blight fungus	31
	3.3.2.1	Colony and conidial morphology	33
	3.3.3	Morphological characteristics of cultured fungal colonies	37
	3.3.3.1	Effect of media on colony growth	37
	3.3.3.2	Effect of pH on colony sporulation	38
	3.3.3.3	Effect of temperature on germination of spores	40
3.4		Conclusions	42
4		MOLECULAR CHARACTERISATION AND PHYLOGENETIC ANALYSIS OF <i>NEOFUSICOCCUM</i> SPP.	43
	4.1	Introduction	43
	4.2	Materials and Methods	44
	4.2.1	DNA extraction	44
	4.2.2	Amplification of Target DNA	44
	4.2.2.1	Polymerase chain reaction amplification using Internal Transcribed Spacers (rDNA-ITS) and Microsatellite BotF5 regions	44
	4.2.2.2	Gel Electrophoresis and Staining	45
	4.2.3	Purification, sequences and analyses	45
	4.2.4	Phylogenetic Analysis	45
	4.3	Results and discussion	46
	4.3.1	Genomic DNA extraction	46
	4.3.2	Amplification of target DNA	46
	4.3.2.1	Polymerase chain reaction amplification	46
	4.3.3	DNA sequencing and Analysis	50
	4.4	Conclusions	59
5		PATHOGENICITY OF ISOLATES OF <i>Neofusicoccum ribis</i> TO RUBBER AND EVALUATION OF CROSS PATHOGENICITY ON HOST PLANTS	60
	5.1	Introduction	60
	5.2	Materials and Methods	61
	5.2.1	Fungal Isolates	61
	5.2.2	Pathogenicity tests	61

5.2.2.1	Pathogenicity tests on detached rubber leaves in laboratory	61
5.2.2.2	Pathogenicity tests on rubber seedlings in glasshouse	62
5.2.2.3	Cross-inoculation on detached leaves of host plants in laboratory	63
5.2.3	Statistical Analysis	63
5.3	Results and Discussion	64
5.3.1	Pathogenicity tests	64
5.3.1.1	Pathogenicity tests on detached rubber leaves in laboratory	64
5.3.1.2	Pathogenicity test on rubber seedlings in glasshouse	68
5.3.1.3	Cross-inoculation on detached leaves of host plants in laboratory	70
5.4	Conclusions	74
6	DETECTION OF PATHOGENESIS-RELATED GENES INDUCED BY <i>Neofusicoccum ribis</i> ISOLATE ON RUBBER	75
6.1	Introduction	75
6.2	Materials and Methods	76
6.2.1	Plant materials and treatments	76
6.2.2	Total RNA extraction and purification	76
6.2.3	Synthesis of cDNA	77
6.2.4	Amplification of the cDNA Template	77
6.2.4.1	Analysis of Total RNA, cDNA and PCR products with Gel electrophoresis	78
6.2.4.2	Nucleotide Sequence Analysis	78
6.3	Results and Discussion	79
6.3.1	Fungal infection	79
6.3.2	Total RNA Extraction, Purification and cDNA synthesis	80
6.3.3	Gene Expression Analysis	81
6.3.4	DNA Sequence Analysis	85
6.3.4.1	Sequences BLAST Results	85
6.4	Conclusions	86
7	SUMMARY, GENERAL CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH	87
7.1	Summary and Conclusions	87
7.2	Recommendations for future research	88
	REFERENCES	90
	APPENDICES	114
	BIODATA OF STUDENT	144
	LIST OF PUBLICATIONS	145

LIST OF TABLES

Table		Page
2.1	The morphological distinction on <i>Neofusicoccum</i> species	11
2.2	Different families of pathogenesis-related proteins	22
3.1	Source of 33 <i>Neofusicoccum</i> spp. derived from LB infected rubber leaves for research	28
3.2	Cultural Characteristics of 8-day Old Fungal Isolates on PDA.	34
3.3	Measurements of conidia of 10 to 15 day old Fungal Isolates on PDA (n =30).	36
3.4	Radial growth rates (mm/day) of isolates of leaf blight from rubber on culture media	38
3.5	Mean germination rates of three isolates of <i>Neofusicoccum</i> spp. at different incubation periods at 25° C.	41
4.1	BLAST results of 33 isolates for ITS regions related to <i>N. ribis</i> .	49
4.2	BLAST results of 6 isolates for BotF15 region related to <i>N. ribis</i> .	50
4.3	Sources of <i>Neofusicoccum</i> species used for phylogenetic analyses	52
4.4	Morphological character of <i>Neofusicoccum</i> species used for phylogenetic analyses.	58
5.1	Isolates of <i>N. ribis</i> used in pathogenicity tests	61
5.2	Pathogenicity of representative <i>N.ribis</i> isolates on different rubber clones using detached leaf assay after 15 days inoculation.	67
5.3	Means rating (R, M, S) of isolates resulting from inoculation trials	68
5.4	Pathogenicity of the representative <i>N.ribis</i> isolates on hosts plant using detached leaf assay at 15 day	70
5.5	Pathogenicity of representative <i>N. ribis</i> isolates on rubber and fruit plants using detached leaf assay after 15 days inoculation. Means rating (R, M, S) of isolates resulting from inoculation trials.	73
6.1	Primers for gene-specific amplification and expected lengths for cDNA given in base pair (bp)	78
6.2	Blast results of <i>Hevea brasiliensis</i> chitinase (Chit), β -1-3 glucanase (Glu) Phenylalanine lyase (PAL) and Glyceraldehyde phosphate dehydrogenase (GAPHD) genes	86

LIST OF FIGURES

Figure		Page
2.1	Rubber tree with cup of latex (a), leaves (b) fruits (c) flower (d), and seeds (e) of <i>Hevea brasiliensis</i>	5
2.2	Conidia morphology of anamorph and teleomorph stage of <i>Fusicoccum</i> . Asci of <i>Botryosphaeria dothidea</i> , teleomorph of <i>Fusicoccum aesculi</i> (a), <i>Neofusicoccum</i> conidia (b), aged conidia of <i>Neofusicoccum</i> with brown wall and one to two septa (c). Scale bar =10 μ m.	10
2.3	Overview of local and systemic signalling in Arabidopsis disease resistance.	19
3.1	Infected leaflets of <i>H. brasiliensis</i> by <i>Neofusicoccum</i> sp.	31
3.2	Twelve day- old culture of 33 isolates grown on PDA: SB from Selangor, SK from perak, and SJ from Johor.	32
3.3	Culture and conidial morphology of <i>Neofusicoccum</i> spp. isolated from rubber leaf: a. one week old culture on PDA producing pycnidia, b conidia observed under the light microscope, (c) germinating conidia in distilled water after 24 h of incubation. Scale bar = 10 μ m	35
3.4	Effect of four types of media on colony growth of <i>Neofusicoccum</i> spp. at 4 days after incubation	37
3.5	Sporulation of three isolates of <i>Neofusicoccum</i> spp. on CDA at different pH after 12 days of incubation	39
3.6	Sporulation of three isolates <i>Neofusicoccum</i> spp on PDA at different pH after 12 days of incubation	40
3.7	Effect of temperature on spore germination of three isolates of <i>Neofusicoccum</i> spp on water agar (WA) after 24 hours incubation	41
4.1	Total genomic DNA extracted from representative <i>Neofusicoccum</i> spp.	46
4.2	Gel electrophoresis showing representative bands of PCR product from ITS region. The amplification fragments were approximately 550 bp. (A: SB isolate from Selangor, B: SK isolate from Perak and C: SJ isolate from Johor)	48
4.3	Gel electrophoresis showing representative band of PCR product from BotF15 region. The amplification fragment was approximately 350 bp	50
4.4	Phylogenetic tree based on ITS region. One of the most parsimonious trees obtained from Maximum Parsimony analyses of the ITS sequence data of the Botryosphaeriaceae. The scale 1 shows 1 change and the Bootstrap support (%) from 1000 replications is given on the branch. Isolates marked in bold represent those from	55

	rubber	
4.5	Phylogenetic tree based on the unknown locus (Bot F15). One of the most parsimonious trees obtained from Maximum Parsimony analyses of the BotF15 sequence data of the Botryosphaeriaceae. The scale 10 shows 10 changes and the Bootstrap support (%) from 1000 replications is given on the branch. Isolates marked in bold represent those from rubber.	56
4.6	Phylogenetic tree based on combine sequences ITS and BotF15. Most parsimonious unrooted tree inferred from Maximum Parsimony analyses of the combined ITS and BotF15 sequence data of the representative taxa of the Botryosphaeriaceae. The scale 10 shows 10 changes and the Bootstrap support (%) from 1000 replications is given on the branch. Isolates marked in bold represent those from rubber.	57
5.1	Level of infection of the representative <i>N.ribis</i> on detached leaves of different rubber clones.	65
5.2	Pictures showing different responses on seedlings of different rubber clones inoculated with one selected <i>N.ribis</i> isolate (SB30) after 30 days	69
5.3	Mean percentage disease severity of representative <i>N.ribis</i> on different seedlings after 30 days of inoculation	69
5.4	Level of infection of the representative isolates on detached leaves of different host plants	71
5.5	Effect of rubber isolates on fruits plants. Mean lesion (mm) on different detached leaves of fruit plants post inoculated with different isolates (A: SB30); (B: SJ20), (C: SK10).	72
6.1	Developments of infection on rubber leaves after inoculation by <i>N.ribis</i> isolate SK10.	80
6.2	Distinct rRNA bands (28S and 18S) on 1% TAE agarose.	81
6.3	Semi-quantitative reverse transcription polymerase chain reaction analysis of β -1,3 glucanase (Glu), chitinase(Chit) genes and phenylalanine ammoniase lyase (PAL) expression in response to clone RRIM 2002 leaf inoculation with <i>N.ribis</i> strain SK10(JX035744) and sterile distilled water (control) over day. The GAPDH (glyceraldehyde phosphate dehydrogenase) was used as control for normalising mRNA quantity	82
6.4	Semi-quantitative reverse transcription polymerase chain reaction analysis of β -1,3 glucanase (Glu), chitinase(Chit) genes and phenylalanine ammoniase lyase (PAL) expression in response to clone PB 350 leaf inoculation with <i>N.ribis</i> strain SK10(JX035744) and sterile distilled water (control) over day. The GAPDH (glyceraldehyde phosphate dehydrogenase) was used as control for normalising mRNA quantity	84

LIST OF ABBREVIATIONS

ABCT	Gene encoding ATP-binding cassette transporters
ACO2	Aminocyclopropanecarboxylate oxidase 2
ANPC	Association of Natural Rubber Producing Countries
ANOVA	Analysis of variance
Avr gene	Avirulence gene
BLAST	Basic Local Alignment Search Tool
Bot F15	Microsatellites
Bp	Base pair
°C	Degree Celsius
cDNA	Complementary deoxyribonucleic acid
CRD	Completed randomised design
CTAB	Hexadecyltrimethyl-Ammonium Bromide
Chit	Chitinase
DNA	Deoxyribonucleic acid
dNTPs	Deoxynucleotides
DAI	Day after inoculation
DEPC	Diethylprocarbonate
DMS	Degrees Minutes Seconds
DNase	Deoxyribonuclease
DI	Disease incidence
DSI	Disease severity index
EDTA	Ethylenediaminetetraacetic acid
ETR 1	Ethylene receptor protein 1
FLB	Fusicoccum Leaf Blight
GAPDH	Glyceraldehyde phosphate dehydrogenase

Glu	Glucanase
GPX	Gluthation peroxidase
ITS	Internal Transcribed Spacer
IRSG	International Rubber Study Group
ISR	Induction of systemic resistance
Kb	Kilo base –pair
LiCl	Lithium Chloride
mL	Millilitre
Mm	Millimetre
mM	Milimolar
MW	Molecular weight
Ng	Nanogram
Nm	Nanometre
NCBI	National Centre for Biotechnology Information
OD	Optical density
PAL	Phenylalanine ammonia lyase
PCR	Polymerase chain reaction
PDB	Potato dextrose broth
PDI	Percent disease intensity
pH	Potential of hydrogen
PR	Pathogenesis related protein
PVP	Polyvinyl pyrrolidone
RNA	Ribonucleic Acid
RNase	Ribonuclease
rRNA	Ribosomal Ribonucleic Acid
Rpm	Revolution Per Minute
RRIM	Rubber Research Institute Malaysia
SDS	Sodium dodecyl sulphate

SA	Salicylic acid
SAR	Systemic acquired resistance
SE	Standard error
SSR	Simple Sequence repeats
TAE	Tris acetate EDTA
µg	Microgram
µM	Micromole
µl	Microlitre
U	Unit
UV	Ultra violet
V	Volts
V/V	Volume per volume
W/V	Weight per volume

CHAPTER 1

INTRODUCTION

The para rubber tree (*Hevea brasiliensis*) is well known as the tree of life for its latex and wood (Hayashi, 2009). It belongs to the genus *Hevea* (Euphorbiaceae, Malpighiales) which accommodates three species that yield usable rubber: *H. brasiliensis*, *H. guianensis* and *H. benthamiana*. Although the *H. brasiliensis* is native to the South American Amazonian rainforest, the plant gained popularity in Asia and became increasingly important in some African and then Latin American countries where it was commercially planted as the primary source of natural rubber, the main structural component of which is the poly (*cis* -1,4 isoprene) (Thomas et al., 2011).

Rubber plantation has a critical social implication worldwide as over 20 million people are dependent on the rubber industry. Owing to demand following the phenomenal growth of the tyre and automobile industries (IRSG, 2013), natural rubber production is expected to increase considerably in the years ahead and to even double by the year 2020 (Burger and Smit, 2004; IRGS, 2007). Indeed, the Association of Natural Rubber Producing Countries (ANRPC, 2011) estimated land use for rubber production in Malaysia at 1.25 million hectares in 2007 with a total production of 1.20 million tonnes contributing to about a third of the world's rubber production (MRB, 2009) next to Thailand and Indonesia (ANRPC, 2010; 2011). However, the planted area has been continuously declining since 1982, a situation that is worsened by the concurrent drop in rubber cultivated in all West Malaysia owing to some constraints: aging of trees, less and less availability of suitable land, abiotic and biotic stresses (Jayasinghe, 2009; Thomas et al., 2011).

That notwithstanding, biotic stress, climate change and introduction of new clones in Malaysian rubber plantations could contribute to a change in the diseases scenario. A new disease dubbed Fusicoccum Leaf Blight (FLB) was detected in Johore, Malaysia (Radziah and Chee, 1989) that was caused by a fungal pathogen that infected leaves of mature and immature rubber and inflicted defoliation and premature leaf fall (Hashim et al., 2010) resulting in significant reduction in growth performance, yield and survival of the four –year-old trees in a mixed planting of rubber clones: RRIM 600 (55%), PB260 (15%), PB255 (15%), PB261 (15%), and PB217 (15%), with GT1 as supplies (Radziah and Chee, 1989).

The causal agent was first identified based on the morphology of the vegetative and reproductive structures on infected tissues (Radziah and Chee, 1989). However, although many identification methods for pathogenic fungi associated with leaf blight disease have been developed, the clear taxonomic position of this fungus remains uncertain (Phillips et al., 2008). Hence, the use of a combination of morphological and molecular techniques for the identification of FLB disease would be the choice to implement the appropriate control strategies in the future.

Current knowledge of the taxonomy of *Fusicoccum* through morphological characterisation is unclear. Crous et al. (2006) hence introduced a single generic name for the anamorph which occurs with a Dichomera-like synanamorph, which is the more informative morphological state: *Neofusicoccum* (Crous, Slippers and A.J.L. Phillips, gen. nov. MycoBank MB500870). Interest in this fungal group emerged from the fact that it can lead an endophytic life style within plant organs (Slippers and Wingfield, 2007), and ability to become pathogenic under favourable conditions.

Despite its seriousness to the rubber industry, the mechanism(s) by which the disease develops on rubber tree, the identity of the causal fungus based on morphology and molecular characteristic; and pathogenicity in Malaysia are not known. Indeed, a better understanding of the plant's response to fungal infections could lead to improvement in identification of disease and determination of resistant genes involved as the induction of systemic resistance (ISR) in host plants results in unremitting protection against many pathogens (Kloepper et al., 1992). The ISR is phenotypically similar to systemic acquired resistance (SAR) that develops when plants successfully activate their defence mechanism in response to primary infection by a pathogen (Compant et al., 2005). Pathogen infections can trigger the expression of several classes of pathogenesis –related (PR) genes in a wide range of plant species (Shaji, 2006). However, very little is known about pathogen-induced defence responses in rubber. To gain more insight into the molecular mechanisms of attack by *Neofusicoccum* spp, the role of chitinase, β -1, 3-glucanase; the most widely studied groups of pathogenesis-related (PR) proteins (PR3, PR2), that have been previously demonstrated as reliable markers for induced defense responses in other plant species (Dixon and Harrison, 1990; Van Loon et al., 2006) and phenylalanine ammonia lyase (PAL), a key enzyme in the phenylpropanoid pathway that could perform defence-related functions (Wen et al., 2005), during pathogenesis would be useful.

This research was therefore initiated with the following objectives:

- 1) To isolate fungus from FLB infected leaves from rubber plantations in West Malaysia.
- 2) To determine the characteristics of the *Neofusicoccum* spp. based on morphology and molecular characterisations.
- 3) To conduct pathogenicity tests of *Neofusicoccum* spp. on rubber clones and cross inoculations on alternative host plants (guava, mango, papaya and rambutan).
- 4) To study gene expression of pathogenesis –related (PR) genes during plant – pathogen interaction.

The hypotheses of the study were as follows:

- 1) There is diversity among the fungal isolates from FLB infected rubber in terms of morphological and molecular characteristics.

- 2) Resistant and susceptible rubber clones and other alternative host plants will express different symptoms of disease when inoculated with different isolates of the fungus.
- 3) The fungus induces pathogenesis –related (PR) genes in rubber.

© COPYRIGHT UPM

REFERENCES

- Adesemoye, A. O. and Eskalen, A. (2011). First report of *Spencermartinsia viticola*, *Neofusicoccum australe* and *N. parvum* causing branch canker of citrus in California. *Plant Disease* 95 (6): 770-770.
- Agrios, G.N. (2005). Plant pathology. fifth edition. Academic Press. United States of America. pp. 208-210.
- Agrios, G.N. (1997). How plants defend themselves against pathogens. In: Smith MD, editor. Plant pathology. San Diego: Academic; 1997. pp. 93–114.
- Ahimera, N., Driever, G.F. and Michailides, T.J. (2003). Relationships among propagule numbers of *Botryosphaeria dothidea*, latent infections and severity of panicle and shoot blight in pistachio orchards. *Plant Disease* 87 (7): 846-853.
- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Research* 25:3389-3302.
- Alves, A., Philips, A.J.P., Henriques, I. and Corrieia, A. (2007). Rapid differentiation of species of Botryosphaeriaceae by PCR fingerprinting. *Research in Microbiology* 158 (2): 112-121.
- Alves, A., Crous, P.W., Corrieia, A. and Philips, A.J.P. (2008). Morphological and molecular data reveal cryptic speciation in *Lasiodiplodia theobromae*. *Fungal Diversity* 28 (8):1-13.
- Amponsah, N.T., Jones, E.E., Ridgway, H.J. and Jaspers, M.V. (2011). Susceptibility of grapevine tissues to *Neofusicoccum luteum* conidial infection. *Plant Pathology* 61(4):719-729.
- Anand, A., Zhou, T., Trick, H.N., Gill, B., Bockus, W.W. and Muthukrishanan, S. (2003). Greenhouse and field testing of transgenic wheat plants stably expressing genes for thaumatin-like protein, chitinase and Glucanase against *Fusarium graminearum*. *Journal of Experimental Botany* 54 (384):1101-1111.
- Annakutty, J. (2006). Enzymes in host-pathogen interactions. In: Corynespora leaf Disease of *Hevea brasiliensis*. Strategies for management. Rubber research Institute of India, Kottayam, Kerala, India pp. 54-62.
- Antoniw, J.F., Ritter, C.E., Pierpoint, W.S. and Van Loon, L.C. (1980). Comparison of three pathogenesis-related proteins from plants of two cultivars of tobacco infected with TMV. *Journal of General Virology* 47 (1):79-87.
- ANRPC (Association of Natural Rubber Producing Countries), (2010). Natural Rubber Trends and Statistics. 3(6), pp 29. June 2010.
- ANRPC (Association of Natural Rubber Producing Countries), (2011). Natural Rubber Trends and Statistics. 3, (9), pp. 49. September 2011.

- Arauz, L.F and Sutton, T.B. (1989). Influence of temperature and moisture on germination of Ascospore and conidia of *Botryosphaeria obusta*. *Phytopathology* 79 (6):667-674.
- Bagyalakshami, R., Therese, K.L., Prasanna, S. and Madhavan, H.N. (2008). Newer emerging pathogens of ocular non-sporulating molds (NSM) identified by polymerase chain reaction (PCR)-based DNA sequencing technique targeting internal transcribed spacer (ITS) region. *Current Eye Research*. 33 (2):139-147.
- Bakker, P.A.H.M., Pieterse, C.M.J. and Van Loon. (2007). Induced systemic resistance by inflorescent *Pseudomonas spp.* *Phytopathology* 97 (2):239-243.
- Barber, P.A., Burgess, T., Hardy, G.S.J., Slippers, B., Keane, P.J. and Wingfield, M.J. (2005). *Botryosphaeria* species from *Eucalyptus* in Australia are pleoanamorphic, producing *Dichomera* synanamorphs in culture. *Mycological Research* 109 (12):1347–1363.
- Begoude, B.A.D., Slippers, B., Wingfield, M.J. and Roux, J. (2010). Botryosphaeriaceae associated with *Terminalia catappa* in Cameroon, South Africa and Madagascar. *Mycological Progress*. 9 (1):101-123.
- Begoude, B.A.D., Slippers, B., Wingfield, M.J. and Roux, J. (2011). Identity and pathogenicity of botryosphaeriaceous fungi on *Terminalia* species in Cameroon. *Forest Pathology* 41 (4): 281-292.
- Begoude, B.A.D., Slippers, B., Perez, G., Wingfield, M.J. and Roux, J. (2012). High gene flow and outcrossing within populations of two cryptic fungal pathogens on a native and non-native host in Cameroon. *Fungal Biology* 116 (3):343-353.
- Biagioni, M., Nali, C., Heimler D. and Lorenzini, G. (1997). PAL activity and differential ozone sensitivity in tobacco, bean and poplar. *Journal of Phytopathology* 145 (11-12):533-539.
- Breton, F., D'Auzac, J., Garcia, D., Sanier, C and Eschbach, J.M. (1996). Recent researches on *Corynespora cassicola/Hevea brasiliensis* interaction. In: Proceedings, Workshop on Corynespora Leaf Fall Diseases of *Hevea* Rubber, 16-17 December 1996. Medan, Indonesia, pp. 49-78.
- Breton, F., Sanier, C. and D'Auzac, J. (1997). Scopoletin production and degradation in relation to resistance of *Hevea brasiliensis* to *Corynespora cassicola*. *Journal of Plant Physiology*. 151 (5):595-602.
- Bridge, P.D., Spooner, B.M. and Roberts, P.J. (2005). The impact of molecular data in fungal systematics. *Advances In Botanical Research* 42:33-67.

- Brogie, K., Chet, I., Holliday, M., Cressman, R., Biddle, P., Knowlton, S., Mauvais, C.J. and Brogie, R. (1991). Transgenic Plants with Enhanced Resistance to the Fungal Pathogen *Rhizoctonia solani*. *Science* 254 (5035):1194-1197.
- Brooks, F.E. and Ferrin, D.M. (1994). Branch Dieback of Southern California Chaparral Vegetation caused by *Botryosphaeria dothidea*. *Phytopathology* 84 (1):78-83.
- Brunings, A.M., Datnoff, L.E., Ma, J.F., Mitani, N., Nagamura, Y., Rathinasabapathi, B. and Kirst, M. (2009). Differential gene expression of rice in response to silicon and rice blast fungus *Magnaporthe oryzae*. *Annals of Applied Biology* 155(2):161-170.
- Burger, K and Smit, H. P. (2004). "Natural Rubber Planting Policies and the Outlook for Prices and Consumption." In Full Texts, International Rubber Conference 2004, Chiang Mai, 8-9 April 2004. Bangkok: J Charoen Printing Co.
- Burgess, T.I., Wingfield, M. J. and Wingfield, B.D. (2001). Simple sequence repeat (SSR) markers distinguish between morphotypes of *Sphaeropsis sapinea*. *Applied Environmental Microbiology* 67 (1):354-362.
- Burgess, T.I., Barber, P.A. and Hardy, G.E.St.J. (2005). *Botryosphaeria* spp. associated with eucalypts in Western Australia, including the description of *Fusicoccum macroclavatum* sp. nov. *Australasian Plant Pathology* 34 (4): 557–567.
- Burgess, T.I., Sakalidis, M.L. and Hardy, G.E.St.J. (2006). Gene flow of the canker pathogen *Botryosphaeria australis* between *Eucalyptus globulus* plantations and native eucalypt forests in Western Australia. *Austral Ecology* 31 (5): 559-566.
- Butland, S.L., Chow, M.L. and Ellis, B.E. (1988). A diverse family of phenylalanine ammonia-lyase genes expressed in pine trees and cell cultures. *Plant Molecular Biology* 37 (1):15-24.
- Camm, E.L. and Towers, G.H.N. (1973). Review article: Phenylalanine ammonia lysae. *Phytochemistry* 12(1):962-973.
- Chacko, N., Philip, S., Zacharia, C.A and Jacob, C.K (2005). Artificial induction of chitinase and phenylalanine ammonia lyase activity in *Hevea brasiliensis*. In: International Natural Rubber Conference, India 2005. (Eds. N.M. Mathew *et al.*). Rubber Research Institute of India, Kottayam, Kerala, India, pp. 523-526.
- Chaendaekattu, N. and Kavitha K. M. (2011). Breeding for Disease Resistance in *Hevea* spp.—Status, Potential Threats, and Possible Strategies. Proceedings of the 4th International Workshop on Genetics of Host-Parasite Interactions

in Forestry: Disease and Insect Resistance in Forest Trees. July 31 – August 5, 2011 — Eugene, Oregon, USA.

- Chee, K.H. (1976). Micro-organisms associated with rubber (*Hevea brasiliensis* Mull Arg) Rubber Research Institute of Malaysia, Kuala Lumpur, Malaysia. pp 78.
- Chee, K.H. (1990). Present status of rubber diseases and their control. *Review of Plant pathology* 69 (7): 430-443.
- Chen, S.F., Pavlic, D., Roux J., Slippers, B., Xie, Y.J., Wingfield, M. and Zhou, X.D. (2011). Characterization of Botryosphaeriaceae from plantation-grown *Eucalyptus* species in South China. *Plant Pathology* 60 (1): 739-751.
- Choi, I., Sharm, P.K. and Cheong, S. (2012). First Report of *Neofusicoccum parvum* associated with Bark Dieback of Blueberry in Korea. *Plant Pathology Journal*. 28 (2): 217 -217.
- Choudhary, D.K., Prakash, A. and Johri, B.N. (2007). Induced Systemic Resistance (ISR) in plants: Mechanism of action. *Indian Journal of Microbiology* 47(4): 289-297.
- Churngchow, N. and Rattarasarn. (2001). Biosynthesis of scopoletin in *Hevea brasiliensis* leaves inoculated with *Phytophthora palmivora*. *Journal of Plant Physiology* 158 (7): 875-882.
- Cohn, J., Sessa, G. and Martin, G.B. (2001). Innate immunity in plants. *Current Opinion in Immunology*. 13 (1):55-62.
- Compant, D., Duffy, B., Nowak, J., Clement, C. and Barka, E.A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: Principles, mechanisms of action, and future prospects. *Applied Environmental Microbiology* 71 (9):4951-4959.
- Crous, P.W. and Palm, M.E (1999). Reassessment of the anamorph genera *Botryodiplodia*, *Dothiorella* and *Fusicoccum*. *Sydowia* 52:167–175.
- Crous, P.W., Slippers, B., Wingfield, M.J., Rheeder, J., Marasas, W.F.O., Philips, A.J.L., Alves, A., Burgess, T., Barber, P. and Groenewald, J.Z. (2006). Phylogenetic lineages in the Botryosphaeriaceae. *Studies In mycology* 55 (1): 235-253.
- Chye, M.-L. and Cheung K.Y.(1995). β -1,3-glucanase is highly expressed in the laticifers of *Hevea brasiliensis*. *Plant Molecular Biology* 29:397– 402.
- Davidson, E.M. and Tay, C.S. (1983). Twig, branch, and upper trunk cankers of *Eucalyptus marginata*. *Plant Disease* 67:1285-1287.

- Delaney, T.P., Uknes, S., Vernooij, B., Friedrich, L., Weymann, K., Negrotto, D., Gaffney, T., Gur-Rella, M., Kessmann, H., Ward, E. and Ryals, J. (1994). A Central Role of Salicylic Acid in Plant Disease Resistance. *Science* 266 (5188): 1247-1250.
- Delledonne, M., Zeier, J., Marocco, A. and Lamb, C. (2001). Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease-resistance response. *Proceedings of National Academy of Sciences USA* 98:13454-13459.
- Denman, S., Crous, P.W., Taylor, J.E., Kang, J-C., Pascoe, I. and Wingfield, M.J. (2000). An overview of the taxonomic history of *Botryosphaeria*, and a re-evaluation of its anamorphs based on morphology and ITS rDNA phylogeny. *Studies in Mycology* 45:129–140.
- Denman, S., Crous, P.W., Groenewald, J.Z., Slippers, B., Wingfield, B.D. and Wingfield, M.J. (2003). Circumscription of *Botryosphaeria* species associated with *Proteacea* based on morphology and DNA sequence data. *Mycologia* 95 (2):294-307.
- Deon, M., Scomparin, A., Tixier, A., Mattos, C.R.R., Leroy, T., Seguin, M., Roeket-Drevet, P. and Pujade-renaud, V. (2012). First characterization of endophytic *Corynespora cassiicola* isolates with variant cassiicolin genes recovered from rubber trees in Brazil. *Fungal Diversity* 54 (1):87-99
- Dhingra, O.D. and Sinclair, J.B. (1995). Basic Plant Pathology Methods. (2nd Edition) CRC Press.Inc. Boca Raton, Florida: Lewis Publishers. pp 434.
- Dixon, R. A. and Harrison, M. (1990). Activation, structure and organization of genes involves in microbial defence in plants. *Advances in Genetics* 28:165-234.
- Dornelas, C.M. and Rodriguez, M.A.P. (2005). The rubber tree (*Hevea brasiliensis* Muell. Arg.) homologue of the LEAFY/FLORICAULA gene is preferentially expressed in both male and female floral meristems. *Journal of Experimental Botany* 56 (417):1965–1974.
- Durner, J., Wendehenne, D. and Klessig, D.F (1998). Defense gene induction in tobacco by nitric oxide, cyclic GMP, and cyclic ADP-ribose. *Proceedings of National Academy of Sciences USA*. 103228-10333.
- Durrant, W.E and Dong, X. (2004). Systemic acquired resistance. *Annual Review of Phytopathology* 42:185-209.
- Egea, C., Dickinson, M.J., Candela, M. and Candela, M.E. (1999). β -1,3-glucanase isozymes and genes in resistant and susceptible pepper (*Capsicum annuum*) cultivars infected with *Phytophthora capsici*. *Physiology Plant* 107:312–318.

- Elmsly, T.A. and Dixon, J. (2008). Growth rate of ripe rot fungi at different temperatures. *New Zealand Avocado Grower's Association. Annual Research Report* 18:77-84.
- Espinoza, J. G., Briceño, E. X., Chávez, E. R., Urbez-Torres, J. R. and Latorre, B. A. (2009). *Neofusicoccum* spp. associated with stem canker and dieback of blueberry in Chile. *Plant Disease* 93 (11):1187-1194.
- Everett K.R and Pak H.A (2002). Infection criteria for pathogens causing body rots in avocados. *New Zealand Avocado Growers Annual Research Report*. 2:1-7.
- FAO (Food and Agriculture Organization of the United Nations), (2005). Selected Indicators of Food and Agriculture Development in Asia Pacific Region 1994–2004. Food and Agriculture Organization of the United Nations, Regional Office for Asia and the Pacific, Bangkok.
- Farr, F.D., Elliot, M. and Rossman A.Y. (2005). *Fusicoccum arbuti* sp .nov. causing cankers on Pacific madrone in western Nord America with notes on *Fusicoccum dimidiatum*, the correct name for *Scytalidium dimidiatum* and *Nattrassia mangiferae*. *Mycologia* 97 (3):730-741.
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783-791
- Feys, B.J and Parker, E.J. (2000). Interplay of signalling pathways in plant disease resistance. *Trends in genetics* 16 (10):449-455.
- Findlender, M., Inbar, I. and Chet, L. (1991). Biological control of soil-borne plant pathogens by a β -1, 3-glucanase producing *Pseudomonas cepacia*. *Soil Biology and Biochemistry* 25 (9):1211-1221.
- Fontaine, T., Simenel, C., Dubreucq, G., Adam, O., Delepierre, M., Lemoine, J., Vorgias, C.E., Diaquin, M. and Latge, J.P. (2000). Molecular organization of the alkali-insoluble fraction of *Aspergillus fumigatus* cell wall. *Journal of Biology and Chemistry* 275(36):27594–27607.
- Friedrich, L., Lawton, K., Rues, W., Masner. P., Specker, N., GutRella, M., Meier, B., Dincher, S., Staub, T., Uknes, S., Metaux, J.P., Kessman, H. and Ryals J. (1996). A benzothiadiazole derivative induces systemic acquired resistance in tobacco. *Plant* 10 (1):61-70.
- Frick, U.B. and Schaller, A. (2002). cDNA microarray analysis of fusicoccin-induced changes in gene expression in tomato plants. *Planta* 216 (1):83-94.
- Garcia, D.C., Sanier, J.J., Macheix. And D' Auzac., J. (1995). Accumulation of scopoletin in *Hevea brasiliensis* infected by *Microcyclus ulei* (P.Henn) V. ARX and evaluation of its fungitoxicity for three leaf pathogen of rubber tree. *Physiological and Molecular Plant Pathology* 47 (4):213-223.

- Gazis, R. and Chaverri, P. (2010). Diversity of fungal endophytes in leaves and stems of wild rubber trees (*Hevea brasiliensis*) in Peru. *Fungal Ecology* 3 (3): 240-254.
- Gilbert, G.S. and De Steven, D. (1996). A canker disease of seedlings and saplings of *Teragastria panamensis* (Burseraceae) cause by *Botryosphaeria dothidea* in lowland tropical forest. *Plant Disease* 80 (6):684-687.
- Golzar, H. and Burgess, T.I (2011). *Neofusicoccum parvum*, a causal agent associated with cankers and decline of Norfolk Island pine in Australia. *Australasian Plant Pathology* 40 (5):484-489.
- Gramaje, D., Agustí-Brisach, C., Pérez-Sierra, A., Moralejo, E., Olmo, D., Mostert, L., Damm, U. and Armengol, J. (2012). Fungal trunk pathogens associated with wood decay of almond trees on Mallorca (Spain). *Persoonia: Molecular Phylogeny and Evolution of Fungi* 28:1–13.
- Grossenbacher, J. G. and Duggar, B.M. (1911). A contribution to the life history, parasitism and biology of *Botryosphaeria ribis*. *New York State, Agricultural Experiment Station Technical Bulletin*. 18:113–190.
- Grove, W.B. M.A. (1935). *British Stem and Leaf Fungi. (Coelomycetes)*. Verlag Von J. Cramer, Lehre. Germany. 1. pp. 3301.
- Gu, Y.Q., Wildermuth, M.C., Chakravarthy, S., Loh, Y.T., Yang, C., He, X., Han, Y. and Martin, G.B. (2002). Tomato transcription factor Pti4, Pti5, and Pti6 activate defense response when expressed in *Arabidopsis*. *Plant cell* 14 (4): 817-831.
- Guarro, J., Gene, J. and Stchigel, A.M. (1999). Developpments in fungal taxonomy. *Clinical Microbiology Reviews* 12 (3):454-500.
- Hahlbrock, K. and Scheel, D. (1989). Physiology and molecular biology of phenyl propanoid metabolism. *Annual Review of physiology and Plant Molecular Biology* 40:347-369.
- Hammerschmidt, R. and Kuc, J. (1995). *Induced Resistance to Disease in Plants. (Development in plant pathology, book 4)*. Springer, 1995 edition. pp 183.
- Hammerschmidt, R. (1999). Induced disease resistance: How do induce plants stop pathogens? *Physiological and Molecular Plant pathology* 55 (2):77-84.
- Hashim, H., Haron, M.A., Osman, F.N. and Jung, S.A.M. (2010). Classification of Rubber tree leaf Disease using Spectrometer. 4th Asia International Conference on Mathematical / Analytical Modelling and Computer Simulation. *IEEE Computer Society* pp. 302-306.

- Ismail, H. and Jeyanayagi, I. 1999. Occurrence and identification of physiological races of *Corynespora cassiicola* of *Hevea*. In 'Proceedings of the IRRDB Symposium 1999' (Eds QB Chen, JN Zhou). (Hainnan Publishing House: Haikou. pp. 263-272.
- Iturrizxae, E., Slippers, B. and Mesanza, N. (2011). First Report of *Neofusicoccum parvum* causing canker and die-back of *Eucalyptus* in Spain. *Australasian Plant Disease* 6 (1):57-59.
- Jacobs, K.A, and Rehner, S.A. (1998). Comparison of cultural and morphological characters and ITS sequences in anamorphs of *Botryosphaeria* and related taxa. *Mycologia* 90 (4):601–610.
- Jacobs, R. (2002). Characterisation of *Botryosphaeria* species from mango in South Africa. Msc.Thesis, University of Pretoria, Republic of South Africa. pp. 162.
- Javier-Alva, J., Gramaje, D., Alvarez, L.A. and Armengol, J. (2009). First Report of *Neofusicoccum parvum* associated with dieback of mango trees in Peru. *Plant Disease* 93 (4):426. <http://dx.doi.org/10.1094/PDIS-93-4-0426B>
- Jayasinghe, C.K. and Silva, W.P.K. (1996). Current status of *Corynespora* Leaf Fall in Sri Lanka. In: Proceedings Workshop on *Corynespora* Leaf Fall Disease of *Hevea* Rubber, 16-17 December 1996, Medan, Indonesia. Indonesia Rubber Research Institute. pp. 15-9.
- Jayasinghe, C.K. (1999). Pest and Diseases of *Hevea* Rubber and their geographical distribution. *Bulletin of the Rubber research Institute of Sri Lanka*. 40:1-8.
- Jayasinghe, C.K. (2000). *Corynespora* leaf fall of rubber in Sri Lanka. Diversity of the pathogen and pathogenesis. IRRDB Workshop on *Corynespora* Leaf Fall Disease, Kuala Lumpur and Medan, 6-14 June, 2000.
- Jayasinghe, C.K. (2007). Disease scenario of the rubber tree: history and current status in: Preprint of papers, IRRDB, International Natural Rubber Conference, 11-13 November 2007, Siem Reap, Cambodia, Cambodia Rubber Research Institute. pp. 265-73.
- Jayasinghe, C.K. (2009). The Most Threatening Diseases of the Rubber Tree. MRB-IRRBD Workshop on Agronomy and TOT on Latex harvesting, Kuala Lumpur, Malaysia, 10-15 August 2009.
- Jayasinghe, C.K. (2011). White root disease. The most devastating root disease of rubber tree. First edition 2011. International Rubber Research and Development Board. pp. 24.
- Jayakumar, V., Rajalakshmi, S. and Amaresan, N. (2011). Leaf spot caused by *Neofusicoccum parvum* reported on nutmeg in India. *New Disease Reports* 23:19. [doi:10.5197/j.2044-0588.2011.023.019].

- Jebakumar, R.S., Anandaraj, M. and Sarma, Y.R. (2001). Induction of PR-proteins and defense related enzymes in black pepper due to inoculation with *Phytophthora capsici*. *India Phytopathology* 54 (1):23-28.
- Joosten, M.H.A.J. and De Wit, J.D.M. (1989). Identification of several pathogenesis-related proteins in tomato leaves inoculated with *Cladosporium fulvum* (Syn. *Fulvia fulva*) as β -1,3-glucanases and chitinases. *Plant Physiology* 89 (3):945-951.
- Kanokratana, P., Chantasigh, D., Champredra, V., Tanapongpipat, S., Pootanakit, K. and Eurwilaichitr, L. (2008). Identification and expression of cellobiohydrolase (CBHI) gene from an endophytic fungus, *Fusicoccum* sp. (BCC4121) in *Pichia pastoris*. *Protein Expression and Purification* 58 (1): 148-155.
- Kapsprewska, A. (2003). Plant Chitinases- Regulation and Function. *Cellular and Molecular Biology Letters* 8:809-824.
- Keefe, D., Hinz, U. and Meins, F. Jr. (1990). The effect of ethylene on the cell-type specific and intracellular localization of β -1, 3- glucanases and chitinase in tobacco leaves. *Planter* 182:34-51.
- Kemp, G., Botha, A.M., Kloppers, F.J. and Pretorius, Z.A. (1999). Disease development and β -1,3-glucanase expression following leaf rust infection in resistant and susceptible near-isogenic wheat seedlings. *Physiological and Molecular Plant Pathology* 55 (1):45-52
- Khin, A.A., Chiew, E.F, C., Shamsudin, M.N. and Zainal, A.M. (2008). Natural rubber price forecasting in the world market. AGREX08. International Conference on Agricultural Extension 2008, 16-19 June, Di Bangi-Putrajaya, Kuala Lumpur Malaysia. pp. 13.
- Kirk, P.M., Cannon, P.F., Minter, D.W. and Stalpers, J.A. (2008). Dictionary of the Fungi (10th ed.). Wallingford, UK: CABI. p. 340. ISBN 978-0-85199-826-8.
- Kloepper, J.W., Tuzun, S. and KuD, J.A (1992). Proposed definitions related to induced disease resistance. *Biocontrol Science and Technology* 2:349-351.
- Kombrink, E., M. Schroder and K. Hahlbrock. (1988). Several “pathogenesis-related” proteins in potato are 1,3-glucanases and chitinases. *Proceedings of National Academy of Science USA*. 85:782-786.
- Kombrink, E. and Somssich, I.E. (1995). Defence responses of plants to pathogens. In JH Andrews, IC Tommerup, eds, *Advance in Botanical Research* (incorporating *Advances in Plant Pathology*), Academic Press, London, 21: 1-34.

- Koukol, J. and Conn E.E. (1961). The metabolism of aromatic and properties of the Phenylalanine deaminase of *Hordeum vulgare*. *Journal of Biological and Chemistry* 236: 2692-2698.
- Knogge, W. (1996). Fungal Infection of Plants. *The plant Cell* 8 (10): 1711-1722.
- Kragh, K.M., Jacobsen, S. and Mikkelsen, J.D. (1990). Induction, purification and characterization of barley leaf chitinase. *Plant Science* 71:55-68.
- Kriel, W.M., Swart, W.J. and Crous, P.W. (2000). Foliar endophytes and their interactions with host plants, with specific reference to Gymnospermae. *Advance in Botanical Research* 33:1-29
- Krishnaveni, S., Muthukrishnan, S., Liang, G.H., Wilde, G. and Manickam, A. (1999). Induction of chitinases and β -1,3-glucanases in resistant and susceptible cultivars of sorghum in response to insect attack, fungal infection and wounding. *Plant Science* 144:9–16.
- Kumar, K.K. and Jacob, C.K. (2006). Post-infectious changes of enzymes and proteins in *Hevea brasiliensis* and their role in resistance to *Corynespora cassiicola* In: *Preprints of papers. International natural Rubber Conference*, 13-15 November 2006, Ho Chi Minh City, Vietnam. Rubber Research Institute of Vietnam, pp. 440-449.
- Kuruvilla, J.C. (2006). *Corynespora* leaf disease of *Hevea brasiliensis*. Strategies for management. Rubber Research Institute of India, kottoyam-686009, Kerala, India. pp.188.
- Lazzizzera, C., Frisullo, S., Alves, A. and Phillips, A.J.L. (2008). Morphology, phylogeny and pathogenicity of *Botryosphaeria* and *Neofusicoccum* species associated with drupe rot of olives in Southern Italy. *Plant Pathology* 57 (5): 948–956.
- Lee, S.W., Nazar, R.N., Powell, D.A and Robb, J. (1992). Reduced PAL gene suppression in *Verticillium*-infected resistant tomatoes. *Plant Molecular Biology* 18:345-352
- Leubner-Metzger, G. and Meins, F.J. (1999). Function and regulation of plant β -1, 3-glucanases. In: S.K. Datta and S.Muthukrishnan (eds), *Pathogenesis-related Proteins in Plants*. Boca raton, FL: CRC press. pp. 49-76.
- Leubner-Metzger, G. (2003). Class I β -1, 3-glucanase in the endosperm of tobacco during germination. *Plant Physiology* 109:751-759.
- Leong, S.K. (1976). Propagation and establishment methods in *Hevea*. In Samsudin.T. (ed RRIM training manual on rubber planting and nursery techniques, Rubber Research Institute of Malaysia, Kuala Lumpur. PP. 15-21.

- Linaldeddu, B.T., Francheschini, A., Luque, J. and Phillips, A.J.L. (2007). First report of canker disease caused by *Botryosphaeria parva* on cork trees in Italy. *Plant Disease* 91(3):324.-324.
- Linaldeddu, B.T., Scanu, B., Schiaffino, A. and Serra, S. (2010). First report of *Neofusicoccum australe* associated with grapevine cordon dieback in Italy. *Phytopathologia Mediterranea* 49 (3):417-420.
- Linthrost, H.J., Melchers, L.S., Mayer, A., Van Rockel, J.S.C., Cornelisen, B.J.C. and Bol, J.F. (1991). Analysis of gene families encoding acidic and basic β -1,3-glucanases of tobacco. *Proceedings of National Academy of science USA*, 87:8756-8760.
- Liyanage, A.D.S. and Jacob, C.K. (1992). Disease of economic importance in rubber. In Sethuraj, M.R., Mathew, N.M. eds. Natural rubber: biology, cultivation and technology. The Netherland: Elsevier Science Publishers: *Developments in Crop Science* 23:324-369.
- Lois, R., Dietrich, A., Hahlbrock, K. and Schulz, W. (1989). A phenylalanine ammonia-lyase gene from parsley: structure, regulation and identification of elicitor and light responsive cis-acting elements. *The EMBO* 8:1641-1648.
- Lotan, T., Ori, N. and Fluhr, R. (1989). Pathogenesis-related proteins are developmentally regulated in tobacco flowers. *Plant Cell* 1 (9):881-887.
- Ma, Z. and Michialides, T.J. (2002a). Characterization of *Botryosphaeria dothidea* Isolates Collected from Pistachio and Other Plant Hosts in California. *Phytopathology* 92 (5):519-526.
- Ma, Z. and Michialides, T. J. (2002b). A PCR- based Technique for Identification of *Fusicoccum* sp. from pistachio and various other hosts in California. *Plant Disease* 86 (5):515-520.
- Macedo, D.M. and Barreto, R.W. (2008). First record of *Botryosphaeria ribis* associated with leaf spots on *Magnolia aff. Candollei* in Brazil. *Brazilian Journal of Microbiology* 9 (3):321-324.
- Maclean, D.J., Braithwaite, K.S., Manners, J.M. and Irwin, J.A.G. (1993). How do we identify and classify fungal plant pathogens in the era of DNA analysis? *Advances in Plant Pathology* 10:207-244.
- McDonald, V., Lynch, S. and Eskalen, A. (2009). First report of *Neofusicoccum australe*, *N. luteum*, and *N. parvum* associated with avocado branch canker in California. *Plant Disease* 93 (9):967-967.
- Mahe, A and Dron, M. (1992). Fungal and plant specific gene markers to follow the bean anthracnose infection process and normalize a bean chitinase mRNA induction. *Molecular Plant Membrane Interaction* 5:242-248.

- Maher, E.A Bate, N/J., Ni, W., Elkind, Y., Dixon, R.A and Lamb, C.J. (1994). Increased disease susceptibility of transgenic tobacco plants with suppressed levels of performed phenylpropanoid products. *Proceedings of the National Academic of Sciences USA* 91:7802-7806
- Manju, M. J., Vinod, K.K., Idicula S.P., Kuruvilla, J. C., Nazeer, M.A and Benagi, V.I. (2010). Susceptibility of *Hevea brasiliensis* clones to *Corynespora* Leaf Fall disease. *Mycology Plant Pathology* 40 (40):603-609.
- Mauch, J., Mauch-Mani, B. and Boller, T. (1988). Artificial hydrolyses in pea tissue II. Inhibition of fungal growth by combinations of chitinases and β -1, 3-glucanases. *Plant physiology* 88:936-942.
- Mauch-Mani, B. and Slusarenko, A.J. (1996). Production of salicylic acid precursors is a major function of phenylalanine ammonia -lyase in the resistance of *Arabidopsis* to *Peronospora Parasitica*. *Plant Cell* 8:203-212.
- Marianne, B., Sela-Buurlage, Anne, S., Ponstein, Sandra, A., Bres-Vloemans, S.Leo, S., Melchers, Peter, J.M. van den Elzen, and Ben J. C. Cornelissen, B.J.C (1993). Only Specific Tobacco (*Nicotiana tabacum*) Chitinases and, β -1, 3-Clucanases Exhibit Antifungal Activity. *Plant Physiology* 101:857-863.
- Matheus, R.J.S., Pereira, C.S., Rodrigues, F. A., Luiz Antônio Zañão, J, Renildes L.F. F. and Oliveira, M.G.A. (2012). Silicon and manganese on the activity of enzymes involved in rice resistance against brown spot. *Tropical Plant Pathology* 37(5):339-345.
- Meins, F.J., Neuhaus, J.M., Sperisen, C. and Ryals, J. (1992). The primary structure of plant pathogenesis-related glucanohydrolases and their genes. In T.Boller and F. Meins (eds), *Genes Involved in Plant Defense*. Berlin: Springer-Verlag, pp. 245-282.
- Melo, D.F. and Mello, S.C.M. de C. (2009). Ideal culture conditions for *Dicyma pulvinata* conidia mass production. *Presquis Agropecuaria. Brasileira* 44 (10) :1232- 1238.
- Metraux, J.P., Signer, H., Ryals, J., Ward, E., Wyss-Benz, M., Gaudin, J., Raschdorf, K., Shmid, E., Blum, W. and Inverardi, B. (1990). Increase in salicylic acid at the onset of systemic acquired resistance in cucumber. *Science* 250 (4983):1004-1006.
- Metraux, J.P. (2001). Systemic acquired resistance and salicylic acid: current state of knowledge. *European Journal of Plant pathology* 107:13-18.
- Michailides, T.J. (1991). Pathogenicity, distribution, sources of inoculum and infection courts on *Botryosphaeria dothidea* on pistachio. *Phytopathology* 81:566-573.

- Mohali, S., Slippers, B. and Wingfield, M. J. (2006). Two new *Fusicoccum* species from *Acacia* and *Eucalyptus* in Venezuela based on morphology and DNA sequence data. *Mycological Research* 110 (4):405-415.
- Mohali, S., Slippers, B. and Wingfield, M.J. (2007). Identification of Botryosphaeriaceae from *Eucalyptus*, *Acacia* and *Pinus* in Venezuela. *Fungal Diversity* 25:103-125.
- Mohali, S.R., Slippers, B. and Wingfield, M.J. (2009). Pathogenicity of seven species of the Botryosphaeriaceae on *Eucalyptus* clones in Venezuela. *Australasian Plant Pathology* 38 (2):135-140.
- Moller, E.M., Bahnweg, G., Sandermann, H. and Geiger., H. H. (1992). A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Research* 20:6115-6116.
- Munch-Garthoff, S., Neuhaus, J.M., Boller, T., Kemmerling, B. and Kogel, K.H. (1997). Expression of beta-1, 3-glucanase and chitinase in healthy, stem –rust –affected and elicitor –treated near –isogenic wheat lines showing Sr5-or Sr-24- specified race-specific rust resistance. *Planta* 201: 235-244.
- MRB (Malaysian Rubber Board), (2009). Rubber Plantation and Processing Technologies. Malaysian Rubber Board, Kuala Lumpur. pp. 404.
- Murnita, M.M., Adam Malik., A.Z. and Masahuling, B. (2008). A study on morphology and characteristics of *Fusicoccum* isolates a causal of leaf blight on rubber. Poster presented during the International Conference of Plant Protection in the Tropics, Kuala Lumpur. 27th –29th August 2008.
- Narayanasamy, P. (2010). Microbial Plant Pathogens-Detection and Disease Diagnosis: *Fungal Pathogens* 1:20-26.
- Nei, C., Hull, J., Jenkins, D., Steinberger, D. and Bent, N. (2005). ‘SPSS: statistical package for the social sciences 11.0.’ (McGraw-Hill: Chicago, IL).
- Neuhaus, J.M., Ahl-Goy, P., Hinz, U., Flores, S. and Meins, F. Jr. (1991) High-level expression of a tobacco chitinase gene in *Nicotiana sylvestris*. Susceptibility of transgenic plants to *Cercospora nicotianae* infection. *Plant Molecular Biology* 16:141-151.
- Nghai, N.A., Kadir, J., Sunderasan, E., Abdullah, M.P., Malik, A. and Napsis, S. (2008). Morphological and Inter SimpleSequence Repeat (ISSR) Markers Analyses of *Corynespora cassiicola* Isolates from Rubber Plantation in Malaysia. *Mycopathologia* 166 (4):189-201.

- Ni, H.F., Yang, H.R., Chen, R.S., Liou, R.F. and Hung, T.H. (2012). New Botryosphaeriaceae fruit rot of mango in Taiwan: identification and pathogenicity. *Botanical Studies* 53:467-478.
- Nilsson, T.H., Kristiansson, E., Ryberg, M., Hallenberg, N. and Larsson, K-H. (2008). Intraspecific ITS Variability in the Kingdom Fungi as Expressed in the International Sequence Databases and Its Implications for Molecular Species Identification. *Evolutionary Bioinformatics* 4:193-201.
- Noordin, W.D., Shafar, J.M and Che, F.I. (2012). Assessment of selected *Hevea brasiliensis* (RRIM 2000 Series) seeds for rootstocks production. *African Journal of Agricultural Research* 7 (21):3209-3216.
- Ntahimpera N., Drieve G.F., Felts D., Morgan D. P. and Michailides, T. J. (2002). Dynamics and patterns of latent infection caused by *Botryosphaeria dothidea* on Pistachio buds. *Plant Disease* 86:282-287.
- Odjakova, M. and Hadjiivanova, C. (2001). The complexity of pathogen defense in plants. *Review. Bulgarian Journal of Plant Physiology* 27(1-2):101-109.
- Ogbebor, O.N. (2010). The status of three common leaf disease of para rubber in Nigeria. *Journal of Animal and plant Sciences* 6 (1):567-570.
- Oliveira Costa, D.V.S., Michereff, S.J., Martins, R.S., Gava, C, A. T., Mizubuti, E.G.S., and Camara, M.P.S. (2010). Species of Botryosphaeriaceae associated on mango in Brazil. *European Journal of Plant Pathology* 127(4):509-519.
- Omo-Ikerodah, E.E., Omokhafa, K.O., Akpobome, F.A., and Mokwunye, M.U. (2009). An overview of the potentials of natural rubber (*Hevea brasiliensis*) engineering for the production of valuable proteins. *African Journal of Biotechnology* 8 (25):7303-7307.
- Ooi, C.B. (1978). Propagation and planting materials in *Hevea*. In Samsudin.T. (ed) RRIM short course on rubber planting and nursery techniques. Rubber research Institute of Malaysia. Kuala Lumpur. pp.8-16.
- Pavlic, D., Slippers, B., Coutinho, T.A. and Wingfield, M.J. (2007). The Botryosphaeriaceae occurring on *Syzygium cordatum* in South Africa, and their potential threat to *Eucalyptus*. *Plant Pathology*. 56(4):624-636.
- Pavlic, D., Wingfield, M.J., Barber, P., Slippers, B., Hardy, G.E.St.J. and Burgess, T.L. (2008). Seven new species of the Botryosphaeriaceae from baobab and other native trees in Western Australia. *Mycologia*. 100 (6):851-866.
- Pavlic, D., Slippers, B., Coutinho, T.A. and Wingfield, M.J. (2009a). Multiple gene genealogies and phenotypic species of the Botryosphaeriaceae: A case study on the *Neofusicoccum parvum*/ *N.ribis* complex. *Molecular Phylogenetics and Evolution* 51(2):259-268.

- Pavlic, D., Slippers B., Coutinho TA. and Wingfield, M.J (2009b). Molecular and phenotypic characterization of the tree phylogenetic species discovered within the *Neofusicoccum parvum*/ *N.ribis* complex. *Mycologia* 101(5): 635-647
- Pennycook, S.R. and Samuels, G.J (1985). *Botryosphaeria* and *Fusicoccum* species associated with ripe fruit rot of *Actinidia deliciosa* (kiwifruit) in New Zealand. *Mycotaxon* 24: 445–458.
- Pieterse, C.M.J., Van Wees, S.C.M., Hoffland, E., Van Pelt, J.A. and van Loon, L.C (1998). A novel signalling pathway controlling induced systemic resistance in *Arabidopsis*. *Plant cell* 10 (9): 1571-1580.
- Pieterse, C.M.J. and Van Loon., L.C. (2004). NPR1: The spider in the web of induced resistance signalling pathways. *Current Opinion in Plant Biology* 7: 456-464.
- Philip, S., Joseph, A., Kumar, A., Jacob, C.K., and Kothandaraman., R. (2001). Detection of β -1, 3- glucanase isoforms against *Corynespora* leaf disease of rubber (*Hevea brasiliensis*). *Indian Journal of Natural Rubber Research* 14(1):1-6.
- Phillips, A.J.L. (1998). *Botryosphaeria dothidea* and other Fungi Associated with Excoriose and Dieback of Grapevines in Portugal. *Journal Phytopathology* 146 (7):327–332.
- Phillips, A.J.L., Rumbos, I.C., Alves, A. and Correia., A. (2005). Morphology and phylogeny of *Botryosphaeria dothidea* causing fruit rot of olives. *Mycopathologia* 159 (5):433-439.
- Phillips, A.J.L., Crous, P.W. and Alves, A. (2007). *Diplodia seriata*, the anamorph of “*Botryosphaeria*” obtusa. *Fungal Diversity* 25:141-155.
- Phillips, A.J.L., Alves, A., Pennycook, S.R., Johnston, P.R., Ramaley, A., Akulov, A. and Crous, P.W. (2008). Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the Botryosphaeriaceae. *Persoonia*. 21: 29-55.
- Phillips, A.J.L., Alves, A., Abdollahzadeh, J., Slippers, B., Wingfield, M.J., Groenewald, J.Z. and Crous, P.W. (2013). The Botryosphaeriaceae: genera and species known from culture. *Studies in Mycology*. 76 (1): 51
- Ploetz, R.C., Pérez-Martínez, J.M., Palmateer, A.J., and Tarnowski, T. (2009). Influence of temperature, light intensity, and isolate on the development of *Neofusicoccum parvum*–induced dieback of *Eugenia*, *Syzygium paniculatum*. *Plant Disease* 93 (8):804-808.

- Punithalingam, E. and Holliday, P. (1973). *Botryosphaeria ribis*. Commonwealth Mycological Institute, CMI Descriptions of Pathogenic Fungi and bacteria. Kew, Surrey, England, No.395.
- Punithalingam, E. (1980). Plant diseases attributed to *Botryodiplodia theobromae* Pat. J. Cramer, Vaduz. Germany. pp. 42-43.
- Pusey, P.L. (1989). Influence of water stress on susceptibility of non-wounded peach bark to *Botryosphaeria dothidea*. *Plant Disease* 73:1000-1003.
- Priyadarshan, P.M. (2003). Contributions of weather variables for specific adaptation of rubber tree (*Hevea brasiliensis* Muell. - Arg) clones. *Genetics and Molecular Biology* 26 (4):435-440.
- Radziah, N.Z. and Chee, K.H. (1989). A new foliar disease of rubber. *Plant Pathology* 38:293-296.
- Radziah, N.Z. and Ismail, H. (1990). Major leaf diseases of rubber and their management. *Rubber Board Bulletin* 26 (2):20-25.
- Radziah, N. Z. (1992). Fusicocum Leaf Blight of Rubber. *Malaysia Plant Protection Society* 16 (1): 3-4.
- Ramos, L.J., Lara, S.P., McMillan, R.T.Jr. and Narayanan, K. R. (1991). Tip dieback of mango (*Mangifera indica*) caused by *Botryosphaeria ribis*. *Plant Disease* 75 (3):315-318.
- Rao, B.S. (1975). *Maladies of Hevea in Malaysia*, Rubber research Institute of Malaysia, Kuala Lumpur, Malaysia. pp. 108.
- Rayachhetry, M. B., Blakelslee, G. M., Webb, R.S. and Kimbrough, J.W. (1996). Characteristics of the Fusicocum anamorph of *Botryosphaeria ribis*, a potential biological control agent for *Malaleuca quinquenervia* in South Florida. *Mycologia* 88 (2):239-248. *ent. Rubber Board Bulletin* 26: 20-25.
- Rayner, R.W. (1970). A mycological color chart. Commonwealth Mycological Institute and British Mycological Society, U.K, pp. 34.
- Rodas, C.A., Slippers, B., Gryzenhout, M. and Wingfield, M. J. (2009). Botryosphaeriaceae associated with *Eucalyptus* canker diseases in Colombia. *Forest Pathology* 39:110-23.
- Rodesuchit, A. and Kajornchaiyakul, P. (1996). Screening *Corynespora* resistant clones of rubber in Thailand. In Darussamin, A., Pawirosoemardjo, S., Basuki, Azwar, R. and Sadarudin, eds, Proceedings of the workshop on *Corynespora* Leaf Fall Disease of *hevea* Rubber. Indonesian Rubber Research Institute, Indonesia, pp.163-176.

- Ruiz, J.M., Bretones, G., Baghour, M., Ragala, L., Belakbir, A. and Romero, L. (1998). Relationship between boron and phenolic metabolism in tobacco leaves. *Phytochemistry* 48:269-272.
- Ryals, J.A., Neuenschwander, U.H., Willits, M.G., Molina, A., Steiner, H.Y., Hunt, M.D. (1996) Systemic acquired resistance. *The Plant Cell* 8 (10): 1809–1819.
- Saha, A., Mandal, P., Dasgupta, S., and Saha, D. (2008). Influence of culture media and environmental factors on mycelial growth and sporulation of *Lasiodiplodia theobromae* (Pat.) Griffon and Maubl. *Journal of Environmental Biology* 29(3): 407-410.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). *Molecular Cloning: a laboratory manual*. 2nd ed. N.Y., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, 1659 p. ISBN 0-87969-309-6
- Sánchez, M.E., Venegas, J., Romero, M.A., Phillips, A.J.L and Trapero, A. (2003). *Botryosphaeria* and related taxa causing oak canker in southwestern Spain. *Plant Disease* 87:1515–1521.
- Saskia van Wees, C.M., De Swart, E.A.M., Van Pelt, J.A., Van Loon, L.C. and Pieterse, C. M.J. (2002). Enhancement of induced disease resistance by simultaneous activation of salicylate- and jasmonate- dependent defense by pathways in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America* 97:8711-8716
- Sathik, M.M.B., Kuruvilla, L., Thomas, M., Luke, L.P., Satheesh, P.R., Annamalainathan and Jacob, J. (2012). Quantitative expression analysis of stress responsive genes under cold stress in *Hevea brasiliensis*. *Rubber Science* 25(2):199-213.
- Sathyapriya, H., Sariah, M., Siti Nor Akmar, A. and Wong, M. (2011). Root colonisation of *Pseudomonas aeruginosa* strain UPMP3 and induction of defence related genes in oil palm (*Elaeis guineensis*). *Annals of applied Biology* 160:137-144
- Schlumbaum, A., Mauch, F., Vogelli, U. and Boller, T. (1986). Plant Chitinase are Potent Inhibitor of Fungal Growth. *Nature* 324:365-367.
- Schreiber, L.R. (1964). Stem canker and Dieback of Rhododendron caused by *Botryosphaeria ribis*. Grossenbacher and Duggar. *Plant Disease Reporter* 48: 207-210.
- Sekeli, R., Tan Yik Wan, Y., Kulaveerasingam, H., Napis, S. and Hee, T.S. (2003). The Isolation and Expression Analysis of a Class I Chitinase from Developing Winged Bean Seed (*Psophocarpus tetragonolobus*). *Asia Pacific Journal of Molecular Biology and Biotechnology* 11 (2):63-69.

- Sergeeva, V., Alves, A. and Phillips A.J.L. (2009). *Neofusicoccum luteum* associated with leaf necrosis and fruit rot of olives in New South Wales, Australia. *Phytopathologia Mediterranea* 48 (2):294–298.
- Shaji, P. (2006). Pathogenesis-related proteins, phytoalexins their role in resistance of *Hevea brasiliensis* to leaf diseases In: *Corynespora* leaf Disease of *Hevea brasiliensis*. Strategies for management. Rubber research Institute of India, Kottayam, Kerala, India pp. 63-69.
- Shen, Y. M., Chao, C. H. and Liu, H. L. (2010). First Report of *Neofusicoccum parvum* Associated with Stem Canker and Dieback of Asian Pear Trees in Taiwan. *Plant Disease* 94 (8):1062-1062.
- Sibounnavoung, P., Kalaw, S.P., Divina, C.C. and Soyong, K. (2009). Mycelial growth and sporulation of *Emericella nidulans*, a new fungal antagonist on two culture media. *Journal of Agricultural Technology* 5 (2): 317-324.
- Silva, W.P.K., Deraniyagala, S.A., Wijesundera, R.L.C., Karaunanayake, E.H. and Priyanka, U.M. S. (2001). Isolation of scopoletin from leaves of *Hevea brasiliensis* and the effect of scopoletin to pathogens of *H.brasiliensis*. *Mycopathologia* 153 (4):199-202
- Simmons, R. (1994). The physiology and molecular biology of plant β -1, 3-glucanase and 1, 3: 1, 4- β -D- glucanases. *Critical Reviews in Plant Sciences* 13:325.-325.
- Slippers, B., Crous, P.W., Denman, S., Coutinho, T.A., Wingfield, B.D., and Wingfield, M.J. (2004a). Combined multiple gene genealogies and phenotypic characters differentiate several species previously identified as *Botryosphaeria dothidea*. *Mycologia* 96 (1):83-101.
- Slippers, B., Burgess, T., Crous, P.W., Coutinho, T.A., Wingfield, B.D. and Wingfield, M. J. (2004b). Development of Simple Sequence Repeat markers for *Botryosphaeria* spp. with *Fusicoccum* anamorphs. *Molecular Ecology Resources* 4 (4):675-677.
- Slippers, B., Fourie, G., Crous, P.W., Coutinho, T.A., Wingfield, B.D., Carnegie, A.J and Wingfield, M.J. (2004c). Species and distribution of *Botryosphaeria* sp. on native and introduced *Eucalyptus* trees in South Africa and Australia. *Studies in Mycology* 50:343-358.
- Slippers, B., Fourie, G., Crous, P.W., Coutinho, T.A., Wingfield, B.D. and Wingfield, M.J. (2004d). Multiple gene sequences delimit *Botryosphaeria australis* sp.nov from *B. lutea*. *Mycologia* 96 (5): 1028-1039.

- Slippers, B., Johnson, G.I., Crous, P.W., Coutinho, T.A., Wingfield, B.D., and Wingfield, M.J. (2005). Phylogenetic and morphological re-evaluation of the *Botryosphaeria* species causing diseases of *Mangifera indica*. *Mycologia* 97(1):99-110.
- Slippers, B. and Wingfield, M.J. (2007). Botryosphaeriaceae as endophytes and latent pathogens of woody plants: diversity, ecology and impact. *Fungal Biology Reviews* (21):90-106.
- Slippers, B., Smit, W.A., Crous, P.W., Coutinho, T.A., Wingfield, B.D. and Wingfield, M.J. (2007). Taxonomy, phylogeny and identification of Botryosphaeriaceae associated with pome and stone fruit trees in South Africa and other regions of the world. *Plant pathology* 56:128-139.
- Slippers, B., Burgess, T., Pavlic, D., Ahumada, R., Maleme, H., Mohali, S., Rodas, C. and Wingfield, M.J. (2009). A diverse assemblage of Botryosphaeriaceae infects *Eucalyptus* in native and non-native environments. *Southern Forests* 71:101-110.
- Smith, H., Wingfield M.J, Crous, P.W, and Coutinho, T.A (1996). *Sphaeropsis sapinea* and *Botryosphaeria dothidea* endophytic in *Pinus* spp. and *Eucalyptus* spp. in South Africa. *South Africa Journal of Botany* 62:86-88.
- Smith, H. (2001). Biology of *Botryosphaeria dothiodes* and *Sphaeropsis sapinea* as endophytes of Eucalypts and Pines in South Africa. PhD thesis. University of Pretoria, Pretoria, South Africa pp.236
- Smith, H., Crous, P.W., Wingfield M. J., Coutinho, T.A. and Wingfield M.J. (2001). *Botryosphaeria eucalyptorum* sp.nov., a new species in the *B.dothidea*-complex on *Eucalyptus* in South Africa . *Mycologia* 93 (2):277-284.
- Smith, D.R and Stanosz, G.R. (2001). Molecular and morphological differentiation of *Botryosphaeria dothidea* (anamorph *Fusicoccum aesculi*) from some other fungi with *Fusicoccum* anamorphs. *Mycologia* 93 (3):505-515.
- Sutton, B.C. (1980). The *Coelomycetes*. Fungi Imperfect with pycnidia, acervuli and stroma. Commonwealth Mycological Institute, Kew, Surrey, England. pp. 32-32.
- Sutton, B.C. (1981). Production and dispersal of ascospore and conidia by *Physalospora obtuse* and *Botryosphaeria dothidea* in apple orchards. *Phytopathology* 71:584-589.
- Sticher, L., Mauch-Mani, B. and Mettraux, J.P. (1997). Systemic acquired resistance. *Annual Review of Phytopathology* 35 (1):235-270.
- Swofford, D.L. (2002). PAUP. Phylogenetic Analysis Using Parsimony (and other methods). Version 4.0b10. Sunderland, MA, USA: Sinauer Associates.

- Taira, T., Toma, N. and Ishihara, M. (2005). Purification, characterization, and antifungal activity of chitinases from pineapple (*Ananas comosus*) leaf. *Bioscience, Biotechnology and Biochemistry* 69 (1):189-196.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24:1596-1599.
- Tan, A.M., Loo, T.P., Vadivel, G., Bachik, M.R. and Yoon, K.K. (1992). Survey of major leaf disease of rubber in Peninsular Malaysia. *Planters Bulletin*. 211: 51-62.
- Tan, A.M., Loo, T.P., Vadivel, P.R., Malik, A., Badarudin, B. E., Yeang, H.Y. and Nathan, S. (2011). *Hevea* genetic transformation: A molecular approach to enhance resistance to *Corynespora* leaf disease. Paper presented at IRRDB-CIRAD plant breeders seminar, 4-7 April 2011. Bahia. Brazil.
- Taylor, J.W., Jacobson, D.J., Kroken, S., Kasuga, T., Geiser, D.M., Hibbett, D.S. and Fisher, M.C. (2000). Phylogenetic Species Recognition and Species Concepts in Fungi. *Fungal Genetics and Biology* 31 (1):21–32.
- Taylor, A., Hardy, G.E.S., Wood, P. and Burgess, T. (2005). Identification and pathogenicity of *Botryosphaeria* species associated with grapevine decline in Western Australia. *Australasian Plant Pathology* 34 (2):187-195.
- Thanseem, I., Venkatachalam, P. and Thulaseedharan, A. (2003). Sequence characterization of *Hevea*, β -1, 3-glucanase genes through genomic and cDNA cloning. *Indian Journal of Natural Rubber Research* 16 (1-2):106-14.
- Thanseem, A., Joseph and Thulaseedharan, A. (2005). Induction and differential expression of β -1, 3-glucanase mRNAs in tolerant and susceptible *Hevea* clones in response to infection by *Phytophthora meadii*. *Tree Physiology* 25: 1361-1368.
- Thanseem and Thulaseedharan, A. (2006). Optimization of RQRT-PCR protocols to measure β -1,3-glucanase mRNA levels in infected tissues of rubber tree (*Hevea brasiliensis*). *Indian Journal of Experimental Biology* 44: 492-498.
- Thomidis, T., Michailides, T.J., and Exadaktylou, E. (2011). *Neofusicoccum parvum* associated with fruit rot and shoot blight of peaches in Greece. *European Journal of Plant Pathology* 131:661–668.
- Thomas, K.U., Minimol, R., Bin, K., and Saha, K. (2011). Divergent DNA Methylation patterns Associated with Abiotic stress in *Hevea brasiliensis*. *Molecular plant* 4 (6):996-1013.
- Turner, N.C. (1969). Fusicoccin: a Fungal Toxin that opens Stomata. *Nature* 223 (5210):1070-1071.

- Uknes, S. Mauch-Mani, B., Moyer, M., Potter, S., Williams, S., Dincher, S., Chandler, D., Slusarenko, A., Ward, E., and Ryals, J. (1992). Acquired resistance in *Arabidopsis*. *Plant Cell* 4 (6): 645-656.
- Úrbez-Torres, J.R. and Gubler, W.D. (2009). Pathogenicity of Botryosphaeriaceae species isolated from Grapevines in California. *Plant Disease* 93(3):584-592.
- Úrbez-Torres, J.R., Battany, M., Bettiga, L.J., Gispert, C., McGourty, G., Roncoroni, J., Smith, R.J., Verdegaal, P., and Gubler, W.D. (2010). Botryosphaeriaceae species spore-trapping studies in California vineyards. *Plant Disease* 94 (6):717-724.
- Úrbez-Torres, J.R., Peduto, F., Striegler, R.K., Urrea-Romero, K.E., Rupe, J.C., Cartwright, R.D. and Gubler, W.R. (2012). Characterization of fungal pathogens associated with grapevine trunk disease in Arkansas and Missouri. *Fungal Diversity* 25(1):169-189.
- Valencia-Botin, A.J., Sandoval-Islas, J.S., Cardenas-Soriano, E., Michailides, T.J., and Rendon Sanchez, G (2004). A new stem spot disease of pitahaya (*Hylocereus undatus* H.) caused by *Fusicoccum*-like anamorph of *Botryosphaeria dothidea* in Mexico. *Revista Mexicana de Fitopatologia* 22. (1):40-42.
- Van Loon, L.C. (1997). Induced resistance in plants and the role of pathogenesis-related proteins. *European Journal of Plant Pathology* 103:753-765
- Van Loon, L.C., Bakker, P.A.H.M. and Pieterse, C.M.J. (1998). Systemic resistance induced by rhizosphere bacteria. *Annual Review of phytopathology* 36 (1): 453-483.
- Van Loon and Van Strien, (1999). The families of pathogenesis related proteins, their activities, and comparative analysis of PR-1 type proteins. *Physiological and Molecular Plant Pathology* 55:85-97.
- Van Loon, L.C. (1999). Occurrence and properties of plant pathogenesis-related proteins. In: S.K. Datta and S. Muthukrishnan (eds), *Pathogenesis –related proteins in Plants*. Boca raton, FL: CRC Press. pp. 1-9.
- Van Loon, L.C., Rep, M. and Pieterse, C.M.J. (2006). Significance of inducible defense-related proteins in infected plants. *Annual review of Phytopathology* 44:135-162.
- Van Niekerk, J.M., Crous, P.W., Groenewald, J.Z., Fourie, P.H. and Halleen, F. (2004). DNA phylogeny, morphology and pathogenicity of *Botryosphaeria* species on grapevines. *Mycologia* 96 (4):781-798.
- Vernooij, B., Friedrich, L., Morse, A., Reist, R., Kolditz-Jawhar, R., Ward, E., Ukness, S., Kessman, H., and Ryals, J. (1994). Salicylic acid is not the

translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *Plant Cell* 6 (7):959-965.

Vogeli-Lange, R., A. Hansen-Gehri, T. Boller and F. Meins, Jr. (1988). Induction of the defense-related glucanohydrolases, β -1,3-glucanase and chitinase by tobacco mosaic virus infection of tobacco leaves. *Plant Science* 54:171–176.

Von Arx, J. A. (1987). Plant Pathogenic Fungi. *Behefte zur Nova Hedwigia*. 87: 288.

Walters, D. and Heil, M. (2007). Costs and trade-offs associated with induced resistance. *Physiological and Molecular Plant Pathology* 71(1-3):3-17.

Wang, C.L. and Hsieh, H.Y. (2006). Occurrence and Pathogenicity of Stem Canker of guava in Taiwan Caused by *Botryosphaeria rhodina*. *Plant Pathology Bulletin* 15:219-230.

Wang, H.L., P.H. Chen, P.H., Ni, H.F. and Chen, R.S. (2007). Physiological characterization and screen of control chemicals for *Lasiodiplodia theobromae* of papaya. *Plant Pathology Bulletin* 16:71-77.

Wei, G., Kloepper, J. W. and Tuzun, S. (1991). Induction systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology* 81(12):1508-1512.

Wen, P-F., Chen, J-Y., Kong, W-F., Pan, Q-H., Wan, S-B. and Huang, W-D. (2005). Salicylic acid induced the expression of phenylalanine ammonia-lyase gene in grape berry. *Plant Science* 169: 928–934

White, T. J., Bruns, T., Lee, S. and Talor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics In: *PCR protocols, a guide to methods and applications*. ed. by Innis M. A., Gelfand, D. H., Snisky, J. J. and White, T. J., pp. 315-322. Academic press, New York, USA

Williams, W.V., Rosenbaum, H. and Weiner, D.B.(1992). Effect of RNA concentration on cDNA synthesis for DNA amplification. *Genome Research* 2:86-88.

Xi, Z., Ruhfel, B.R., Schaefer, H., Amorim, A.M., Sugumaran, M., Wurdack, K.J., Endress, P.K., Matthews, M.L., Stevens, P.F., Mathews, S. and Davis, CC. (2012) Phylogenomics and a posteriori data partitioning resolve the Cretaceous angiosperm radiation Malpighiales. *Proceedings of the National Academy of Sciences of the United States of America*, 109(43):17519-17524.

Xue, L., Charest, P.M. and Jabaji-Hare, S.H. (1998). Systemic induction of peroxidases, β -1,3-glucanases, chitinases and resistance in bean plants by binucleate *Rhizoctonia* species. *Phytopathology* 88(4):359–365.

- Yano K., Kawada Y., and Sato T. (2003). Stem blight of eggplant caused by *Fusicoccum aesculi* Corda. *Japan. Journal Phytopathology* 69 (2):132-135.
- Yamada, T., Hashimoto, H., Shiraishi, T. and Oku, H. (1989). Suppression of Pisatin, Phenylalanine ammonia-lyase mRNA, and chalcone synthase mRNA accumulation by a putative pathogenicity factor from the fungus *Mycosphaerella pinodes*. *Molecular plant-Microbe Interactions* 2 (5):256-261.
- Yi, S.Y. and Hwang, K.B. (1996). Differential induction and accumulation of β -1,3-glucanase and chitinase isoforms in soybean hypocotyls and leaves after compatible and incompatible infection with *Phytophthora megasperma* f. sp. glycinea. *Physiological and Molecular Plant Pathology* 48:179–192.
- Yu, D., Tang, H., Zhang, Y., Du, Z., Yu, H., and Chen, Q. (2012). Comparison and Improvement of Different Methods of RNA Isolation from Strawberry (*Fragria X ananassa*). *Journal of Agricultural Science* 4 (6):51-56.
- Yusof A and Ismail B.H (1974). Artificial Defoliation by Aerial Spraying for Avoiding Secondary Leaf Fall of *Hevea brasiliensis*. Rubber Research Institute of Malaysia, Kuala Lumpur.
- Zane, L., Bargelloni, L. and Patarnello, T. (2002). Strategies for microsatellite isolation: a review. *Molecular Ecology* 11 (1):1-16.
- Zemanek, A.B., Ko, T-S., Thimmapuram, J., Hammerschlag, F.A. and Korban, S.S. (2002). Changes in β -1,3-glucanase mRNA levels in peach in response to treatment with pathogen culture filtrates, wounding and other elicitors. *Journal of Plant Physiology* 159 (8):877–889.
- Zhou, S. and Stanosz, G.R. (2001). Relationships among Botryosphaeria species and associated anamorphic fungi inferred from the analyses of ITS and 5.8S rDNA sequences. *Mycologia* 93 (3): 516-527.