



UNIVERSITI PUTRA MALAYSIA

**PREVALENCE AND CHARACTERISATION OF LISTERIA
MONOCYTOGENES ISOLATED FROM CHICKEN AND BEEF**

SAMUEL LIHAN

FSTM 2007 15



**PREVALENCE AND CHARACTERISATION OF *LISTERIA*
MONOCYTOGENES ISOLATED FROM CHICKEN AND BEEF**

By

SAMUEL LIHAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

May 2007

DEDICATED TO MY:

*MOTHER
LATE FATHER
BROTHERS
LATE BROTHER
SISTERS
WIFE
SON*

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**PREVALENCE AND CHARACTERISATION OF *LISTERIA*
MONOCYTOGENES ISOLATED FROM CHICKEN AND BEEF**

By

SAMUEL LIHAN

May 2007

Chairman: Professor Son Radu, PhD

Faculty : Science and Food Technology

Listeria monocytogenes is an opportunistic haemolytic pathogen of humans and animals involved in several outbreaks and sporadic cases of listeriosis associated with the consumption of contaminated food. Growing antibiotic resistance demands the constant reassessment of antimicrobial efficacy, particularly in countries with wide antibiotic used in veterinary science, where higher resistance prevalence is often found and it has been reported that antibiotic resistant is associated with plasmid in bacteria. The purpose of this study is to characterize *L. monocytogenes* isolated from chicken and beef meat by antibiotic resistance test, plasmid DNA profile and arbitrarily primed (AP)- and repetitive sequence (RS)-PCR analysis and to determine whether there is any correlation between the antibiotic resistant and the incidence of plasmid. A total of 112 samples of chicken meat and 101 samples of beef were collected from different wet markets and night markets in Malaysia. *Listeria* spp. were detected in 39 (34.8%) of the chicken meat and 21 (20.8%) of the

beef samples, respectively. *L. monocytogenes* were detected in 27 (24.1%) and 17 (16.8%) of the chicken meat and beef, respectively. Out of all the *Listeria* spp., 42 and 20 isolates from the chicken meat and beef, respectively, were confirmed as *L. monocytogenes*. The antibiotic susceptibility test with sixteen different types of antibiotics revealed that the highest prevalence of resistance among the *L. monocytogenes* isolates was observed against nalidixic acid (100%). None (0%) of the isolates were resistant to vancomycin. The *L. monocytogenes* strains showed resistance to at least two or more of the antibiotics tested with their multiple antibiotic resistance index (MARI) ranging between 0.13 to 0.63. Thirty-three different patterns of resistance were observed among all the *L. monocytogenes* isolates. Plasmid analysis of the *L. monocytogenes* strains revealed that 24 (38.7%) of the strains carried plasmid DNA ranging in sizes from 1.75 to 104.0 megadalton (MDa). Spearman's rho correlation analysis was utilised to determine the correlation coefficient between MARI and the incidence of plasmid in the *L. monocytogenes* isolates. The result shows that there were no significant correlation between the MARI and the incidence of the plasmid ($r= 0.143$, $p= 0.269$). The AP- and RS-PCR analyses generated diverse PCR patterns with multiple DNA fragments in sizes ranging between 250 and 3000 bp. Dendrograms generated from the PCR patterns clustered the isolates into several groups and subgroups. PCR analyses with primers GEN1-50-02, GEN1-50-10 and repetitive primer clustered the chicken isolates into 3, 13 and 6 main groups; and the beef isolates into 2, 4 and 5 main groups; respectively. These main groups were further clustered into several subgroups. Isolation of *L. monocytogenes* from the beef and chicken meat suggests that there is a risk of acquiring listeriosis through these popular meat sources in Malaysia. *L. monocytogenes* will be killed by cooking and raw or semi raw meat are not

consumed in Malaysia. However, *L. monocytogenes* in raw beef and chicken meat may pose a health hazard in kitchen if contaminating cooked food or other ready to eat food. Considering outbreaks of listeriosis associated with different foods, avoidance of consumption of insufficiently cooked meat by at-risk populations is recommended. Hygiene quality control of meat and its products must be recommended during the slaughtering, transportation carriage, other used devices and stuff carriers. Diligent enforcement of sanitary conditions of food contact surface and handling areas, and personal hygiene practices should reduce the potential contamination of meat products by *L. monocytogenes* at the retail level. Resistance to two or more antibiotics among these isolates was common. It is suggested that incorrect use of these antimicrobial agents for therapeutical purposes in veterinary science may lead to the development of antibiotic resistance. The high MARI value indicated that the *L. monocytogenes* strains were originated from high risk sources of contamination in the geographical area. *L. monocytogenes* can no longer be thought to be uniformly susceptible to antibiotics active against Gram-positive bacteria. Continued surveillance of emerging antimicrobial resistance among *L. monocytogenes* is important to ensure effective treatment of human listeriosis. The non-significant correlation between the antibiotic resistance and the incidence of plasmid suggest that the plasmids could be cryptic plasmids which have no apparent effect on the phenotype, especially the antibiotic resistance of the host. The presence of hemolysin gene in the beef and chicken meat isolates is of public health concern, as this virulence gene is associated with pathogenicity of the bacteria in human listeriosis. Both AP-PCR and RS-PCR, having high discriminatory power, have revealed the high diversity of food related *L. monocytogenes* isolates and their suitability to track down the contamination sources. The results suggest how

complex the epidemiology of the *L. monocytogenes* in the study area, as a result of several strains as opposed to the widespread transmission of a single type. The results do not support that certain genetically related strains are better adapted to a particular food source. The genomic heterogeneity of the *L. monocytogenes* found in this study confirms the usefulness of the AP- and RS-PCR analyses for the rapid differentiation and grouping of this organism.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KEKERAPAN DAN PENCIRIAN *LISTERIA MONOCYTOGENES* YANG
DIPENCILKAN DARI DAGING AYAM DAN DAGING LEMBU**

Oleh

SAMUEL LIHAN

Mei 2007

Pengerusi: Profesor Son Radu, PhD

Fakulti : Sains dan Teknologi Makanan

Listeria monocytogenes adalah sejenis patogen hemolitik oportunistik bagi manusia dan haiwan yang terlibat dalam beberapa kes wabak dan kes terpencil listeriosis yang berkaitan dengan memakan makanan yang tercemar. Kerentanan antibiotik yang semakin meningkat memerlukan penilaian semula yang berterusan terhadap keberkesanan antimikrobial, terutamanya di negara yang menggunakan antibiotik dengan meluas dalam sains veterinar dimana kekekerapan kerentanan tinggi selalunya ditemui dan terdapat laporan yang mengatakan bahawa kerentanan antibiotik berkaitan dengan plasmid dalam bakteria. Kajian ini adalah bertujuan untuk mencirikan *L. monocytogenes* yang dipencilkan dari daging ayam dan daging lembu dengan menggunakan ujian kerentanan antibiotik, analisis profil DNA plasmid dan analisis primer arbitrari (PA)- dan jujukan berulang (JB)-PCR dan menentukan samada terdapat sebarang korelasi antara kerentanan antibiotik dengan kehadiran plasmid. Sebanyak 112 sampel daging ayam dan 101 sampel daging lembu

telah diperolehi dari pelbagai pasar basah dan pasar malam di Malaysia. *Listeria* spp. telah dikesan dalam 39 (34.8%) sampel daging ayam dan 21 (20.8%) sampel daging lembu tersebut. *L. monocytogenes* telah dikesan dalam 27 (24.1%) and 17 (16.8%) sampel daging ayam dan daging lembu, masing-masing. Dari kesemua pencilan *Listeria* spp. ini, 42 dan 20 pencilan dari daging ayam dan daging lembu, masing-masing, telah dikenalpasti sebagai *L. monocytogenes*. Ujian kerentanan antibiotik terhadap 16 jenis antibiotik yang berbeza menunjukkan bahawa kekerapan kerentanan tertinggi dalam pencilan *L. monocytogenes* tersebut diperhatikan terhadap nalidixic acid (100%). Tiada (0%) pencilan yang rentan terhadap vancomycin. *L. monocytogenes* menunjukkan kerentanan terhadap sekurang-kurangnya 2 atau lebih antibiotik dengan indeks kerentanan pelbagai (IKP) dari 0.13 hingga 0.63. Tiga puluh tiga corak kerentanan pelbagai diperhatikan dalam kesemua pencilan tersebut. Analisis plasmid menunjukkan 24 (38.7%) strain membawa plasmid yang bersaiz 1.75 hingga 104.0 megadalton (MDa). Analisis korelasi Spearman's rho telah digunakan untuk menentukan korelasi koefisien antara indeks kerintangan pelbagai dan insiden plasmid dalam pencilan *L. monocytogenes*. Keputusan menunjukkan bahawa tiada perkaitan antara IKP dengan insiden plasmid ($r=0.143$, $p=0.269$). Analisis PA- dan JB-PCR menghasilkan pelbagai corak PCR dengan penghasilan fragmen-fragmen DNA yang bersaiz dari 250 hingga 3000 bp. Dendrogram yang dihasilkan dari corak PCR mengumpulkan pencilan kepada beberapa kumpulan dan subkumpulan. Analisis PCR dengan menggunakan primer GEN1-50-02, GEN1-50-10 dan primer jujukan berulang mengumpulkan pencilan daging ayam kepada 3, 13 dan 6 kumpulan utama manakala pencilan daging lembu dikumpulkan kepada 2, 4 dan 5 kumpulan utama, masing-masing, dengan menggunakan primer-primer yang sama. Kumpulan-kumpulan utama ini seterusnya dibahagikan kepada subkumpulan.

Pemencilan *L. monocytogenes* dari daging lembu dan daging ayam tersebut mencadangkan bahawa terdapat risiko untuk mendapat penyakit listeriosis melalui kedua-dua sumber daging popular tersebut di Malaysia. *L. monocytogenes* akan dibasmi bila makanan dimasak dan daging mentah atau separa masak tidak dimakan oleh penduduk Malaysia. Walaubagaimanapun, *L. monocytogenes* dalam daging lembu dan daging ayam mungkin mendatangkan bahaya kesihatan di dapur kalau makanan yang sudah masak atau makanan yang sedia untuk dimakan dicemari. Dengan mengambilkira wabak penyakit listeriosis berkaitan dengan pelbagai makanan, mengelak dari memakan daging yang tidak dimasak sepenuhnya oleh kumpulan berisiko adalah dicadangkan. Kawalan mutu daging dan hasilan daging adalah dicadangkan semasa penyembelihan, pengangkutan atau peralatan yang dipakai dan pembawa barangan. Penguatkuasaan secara diligen tentang keadaan kebersihan permukaan yang bersentuhan dengan makanan dan kawasan sekitar kerja dan amalan kebersihan individu sepatutnya mengurangkan kemungkinan pencemaran daging dengan *L. monocytogenes* pada peringkat jualan runcit. Kerentanan terhadap dua atau lebih antibiotik dalam pencilan-pencilan ini adalah kerap. Adalah dicadangkan bahawa penyalahgunaan antibiotik untuk terapeutik dalam sains veteriner mungkin menjadi pendorong kepada peningkatan kerentanan antibiotik. Nilai IKP yang tinggi menunjukkan bahawa *L. monocytogenes* tersebut berasal dari sumber kontaminasi yang berisiko tinggi dalam kawasan geografi berkenaan. *L. monocytogenes* tidak lagi difikirkan kesemua “susceptible” terhadap antibiotik yang berkesan terhadap bakteria Gram positif. Pemantauan yang berterusan terhadap kemunculan kerentanan antibiotik dalam *L. monocytogenes* adalah perlu untuk memastikan rawatan yang efektif terhadap listeriosis manusia. Korelasi yang tidak bererti antara kerentanan antibiotik dan kehadiran plasmid mencadangkan plasmid

yang dibawa oleh *L. monocytogenes* adalah plasmid kriptik yang mana ianya tidak memberi kesan terhadap fenotip, terutamanya kerentanan antibiotik perumah. Kehadiran gen hemolisin dalam pencilan *L. monocytogenes* dari daging ayam dan daging lembu memerlukan perhatian darisegi kesihatan awam kerana gen virulen ini terlibat dalam kepatogenikan bakteria ini dalam listeriosis manusia. Kedua-dua PA- dan JB-PCR, mempunyai kuasa pengasingan yang tinggi, telah menunjukkan kepelbagaian *L. monocytogenes* yang tinggi dalam makanan dan kesesuaian kedua-duanya digunakan untuk mengesan sumber pencemaran. Keputusan ini mencadangkan bahawa betapa rumitnya epidemiologi *L. monocytogenes* dalam kawasan berkenaan, hasil dari pelbagai strain bertentangan dengan perebakan satu jenis strain. Keputusan ini tidak menyokong bahawa setengah-setengah strain yang rapat dari segi genetik adalah lebih sesuai mengadaptasi terhadap satu jenis sumber makanan. Kepelbagaian genetik *L. monocytogenes* yang dijumpai dalam kajian ini mengesahkan kepentingan analisis PA- dan JB-PCR untuk pengasingan dan pengumpulan organisma ini dengan cepat.

ACKNOWLEDGEMENTS

The author would like to express his whole hearted indebtedness to his main supervisor, Professor Dr. Son Radu from Department of Food Science, Faculty of Science and Food Technology, Universiti Putra Malaysia for his endless motivation and guidance throughout the course of research and transcription of this thesis. His meticulous nature of evaluating the manuscript has been instrumental in making this thesis a valuable one.

The author owe his heartfelt thanks to all his supervisory committees, Associate Professor Dr. Zaiton Hassan and Professor Dr. Raha Abdul Rahim for their invaluable guidance, encouragement and assistance throughout this research project.

The author heartily thank his friends from Center for Research Laboratory, Department of Biotechnology, Universiti Putra Malaysia; Suhaimi Mustafa, Christina Virmala, Ernie Eileen Rose, Varma Virmala, Tan Li Lung, Yanti bt. Toosa, Cheng Leong for their excellent technical expert assistance in all the molecular work.

The author also would like to acknowledge all postgraduates and under graduates students in the Foodborne Pathogen Laboratory, Department of Food Science, Faculty of Science and Food Technology, Universiti Putra Malaysia; Dr. Sahilah Abd. Mutalib, Dr. Yuherman, Dr. Nasreldin El Hadi, Dr. Endang Purwati, Dr. Ahmad Zainuri Mohd. Dzomir, Dr. Ooi Wai Ling, Dr. Cheah Yoke Queen, Dr. Norzaleha Awang Salleh, Dr. Woo Kwan Kit, Lee Yoke Ann, Lim Yin Sze, Apinya Vanichpun, Dr. Lesley Maurice, Dr. Yousr Abd. Hadi, Shushil Kaur, Sitizuliha

Hanapi, Mohd Nizam Lani, Noorlis Ahmad and all the others that are not mentioned here, for their help, assistance and critical comment on the methodology and the outcomes of this research.

The author heartily thanks the Ministry of Science and Technology, and Graduate School of Universiti Putra Malaysia for their sponsorship through the Graduate School Scholarship (GSS) and also the Government of Malaysia for funding this research project through the IRPA grant under vot no. 51538 and 50522.

Last but not least, the author wishes to thank his beloved family for their strong moral and financial support from the beginning until completion of this research project.

I certify that an Examination Committee met on 15 May 2007 to conduct the final examination of Samuel Lihan on his Doctor of Philosophy thesis entitled “Prevalence and Characterisation of *Listeria monocytogenes* Isolated from Chicken and Beef ” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee were as follows:

Khatijah Mohd. Yusoff, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Norihan Mohd. Salleh, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal examiner)

Raja Noor Zaliha Raja Abdul Rahman, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal examiner)

Susumu Kumagai, PhD

Professor
Graduate School of Agriculture and Life Sciences
University of Tokyo
(External examiner)

Hasanah Mohd Ghazali, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Son Radu, PhD

Professor
Faculty of Science and Food Technology
Universiti Putra Malaysia
(Chairman)

Zaiton Hassan, PhD

Associate Professor
Faculty of Science and Technology
Universiti Sains Islam Malaysia
(Member)

Raha Abdul Rahim, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:16/10/08

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

.....
SAMUEL LIHAN

Date:

TABLES OF CONTENTS

	PAGE
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF TABLES	xix
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxii

CHAPTER

I	INTRODUCTION	1
II	LITERATURE REVIEW	7
	Introduction	7
	Taxonomy	7
	Morphology and Colonial Characteristics	9
	Biochemical Properties	11
	Ecology	13
	Natural Environment	13
	Human and Animal Carrier	14
	Risk Assessment of Acquiring Listeriosis	19
	Foodborne Listeriosis	20
	Typing Methods	24
	Serological Typing	24
	Bacteriophage Typing	26
	Molecular Typing Techniques	27
	Arbitrarily Primed-PCR	28
	Repetitive Element-PCR	29
	Multilocus Enzyme Electrophoresis (MEE)	32
	Restriction Enzymes Method	33
	Pulse Field Gel Electrophoresis (PFGE)	34
	Ribotyping	35
	Fatty Acid Analysis	36
	Quantitative Analysis for Epidemiological Investigation of <i>Listeria</i> spp.	37
	Occurrence in Foods	37
	Poultry	38
	Meat	39
	Milk and Dairy products	41
	Seafood	43
	Vegetables	44
	Methodology for Isolation of <i>Listeria</i>	44
	Enrichment Procedures	45
	Selective Enrichment	47
	Plating Media	49
	Primer for species-specific identification	51
	Antibiotic Resistance	53

Nature of Antibiotic Resistance Gene	53
Transfer of Antibiotic Resistant	54
Treatment of Listeriosis	56
Antibiotic Resistance Mechanism	56
Resistance to β -lactam Antibiotic	57
Resistance to Quinolones	58
Resistance to Tetracycline	58
Resistance to Aminoglycosides	59
Resistance to Glycopeptides Antibiotics	60
Resistance to Macrolides and Lincoasamide	60
Resistance to Trimethoprim and Sulfonamides	61
Antibiotic Residues and Antibiotic Resistant Organisms	61
Plasmids in <i>Listeria</i>	62
Disc Diffusion Assay	65
III INCIDENCE OF <i>LISTERIA MONOCYTOGENES</i> AND OTHER <i>LISTERIA</i> SPP.	67
Introduction	67
Materials and Methods	71
Samples Collections	71
Isolation of <i>L. monocytogenes</i> and other <i>Listeria</i> spp.	71
Identification of <i>L. monocytogenes</i> and other <i>Listeria</i> spp.	72
Biochemical Tests	72
Microbact™ <i>Listeria</i> Identification System	72
Detection of Haemolysin Gene by PCR	73
Statistical Analysis	74
Results	76
Isolation and Identification of <i>L. monocytogenes</i> and Other <i>Listeria</i> spp.	76
Occurrence of <i>L. monocytogenes</i> and Other <i>Listeria</i> spp. in chicken and beef meat	80
Statistical Analysis	83
Discussion	85
IV ANTIMICROBIAL DRUG RESISTANCE AND PLASMID DNA AMONG <i>LISTERIA MONOCYTOGENES</i>	95
Introduction	95
Materials and Methods	99
Antimicrobial Drug Resistance Testing	99
Bacterial Strains, Cultivation and Media	99
Antimicrobial Drug Agents and Susceptibility Testing	100
Reading the Antibiogram	100
Multiple Antibiotic Resistance Index (MARI)	101
Plasmid DNA Analysis	101
Determination of Molecular Weight of Plasmid DNA	102
McFarland Turbidity Standard	103
Statistical Analysis	105
Testing for Normality	105
Results	106
Antimicrobial Drug Resistance	106
Plasmid DNA Analysis	106

Statistical Analysis	109
Discussion	110
V GENETIC DIVERSITY OF <i>LISTERIA MONOCYTOGENES</i> AS DETECTED BY ARBITRARILY PRIMED- AND REPETITIVE SEQUENCE-POLYMERASE CHAIN REACTION	119
Introduction	119
Materials and Methods	122
<i>L. monocytogenes</i> Strains and DNA Extraction	122
Arbitrarily Primed (AP)-PCR analysis	122
Repetitive Sequence-PCR Analysis	123
Agarose Gel Electrophoresis Procedure	124
Arbitrarily Primed (AP)- and Repetitive Sequence-PCR Pattern Analysis	125
Results	126
Arbitrarily Primed (AP)-PCR Analysis	126
Repetitive Sequence-PCR Analysis	126
Discussion	137
VI GENERAL DISCUSSION AND CONCLUSION	146
REFERENCES	163
APPENDICES	202
A: GENERAL MEDIA AND SOLUTIONS	202
B: SOLUTION FOR PLASMID EXTRACTION	207
C: REAGENTS FOR PCR	208
D: CONVENTIONAL BIOCHEMICAL TEST RESULTS	209
E: MICROBACT BIOCHEMICAL IDENTIFICATION	212
F: STATISTICAL ANALYSIS RESULTS	250
G: HEMOLYSIN GENE DETECTED AMONG <i>L. MONOCYTOGENES</i>	254
H: PLASMID PROFILE OF <i>L. MONOCYTOGENES</i>	257
BIODATA OF THE AUTHOR	259

LIST OF TABLES

Table		Page
2.1	Biochemical characteristics of the genus <i>Listeria</i>	11
2.2	The biochemical tests used for differentiation of the <i>Listeria</i> genus	12
3.1	Occurrence of <i>L. monocytogenes</i> and other <i>Listeria</i> spp. in chicken meat	82
3.2	Occurrence of <i>L. monocytogenes</i> and other <i>Listeria</i> spp. in beef meat	83
3.3	Prevalence of <i>Listeria</i> spp. in chicken meat	84
3.4	Prevalence of <i>Listeria</i> spp. in beef meat	84
4.1	Molecular weight of plasmid DNA of <i>E. coli</i> V517	103
4.2	Guilford's rule of thumb for interpreting degree, size, strength or magnitude of relationship	105
4.3	Antibiotic resistance patterns and possession of plasmid DNA among <i>L. monocytogenes</i> examined	107
4.4	The Spearman's rho correlation coefficient (r) between antibiotic resistance and the incidence of plasmid in <i>L. monocytogenes</i>	109
5.1	Arbitrary primers screened for arbitrarily primed-PCR analysis	123

LIST OF FIGURES

Figure		Page
3.1	Flow diagram for the isolation and identification of <i>L. monocytogenes</i> and other <i>Listeria</i> spp. from chicken and beef meat	75
3.2	Growth of <i>Listeria</i> spp. colonies on PALCAM agar	77
3.3	Growth of <i>L. monocytogenes</i> colonies on blood agar	78
3.4	Agarose gel electrophoresis of PCR product indicating the presence of haemolysin gene	79
4.1	The graphical method of relating the logarithm of the molecular weight of a DNA molecule (log C) to its electrophoretic mobility (m)	104
4.2	Representative profile of plasmid DNA possessed by the <i>L. monocytogenes</i> strains	109
5.1	Representative profile of arbitrarily primed-PCR patterns obtained with primer GEN1-50-01 to GEN1-50-10 of the <i>L. monocytogenes</i> isolates	127
5.2	Representative profile of arbitrarily primed-PCR patterns obtained with primer GEN1-50-02 of the chicken <i>L. monocytogenes</i> isolates	128
5.3	Representative profile of arbitrarily primed-PCR patterns obtained with primer GEN1-50-10 of the chicken <i>L. monocytogenes</i> isolates	128
5.4	Representative profile of arbitrarily primed-PCR patterns obtained with primer GEN1-50-02 of the beef <i>L. monocytogenes</i> isolates	129
5.5	Representative profile of arbitrarily primed-PCR patterns obtained with primer GEN1-50-10 of the beef <i>L. monocytogenes</i> isolates	129
5.6	Representative profile of repetitive sequence-PCR patterns obtained with repetitive primer of the chicken <i>L. monocytogenes</i> isolates	130
5.7	Representative profile of repetitive sequence-PCR patterns obtained with repetitive primer of the beef <i>L. monocytogenes</i> isolates	130
5.8	Dendrogram generated from the arbitrarily primed-PCR patterns with primer GEN1-50-02 of the <i>L. monocytogenes</i> isolates from chicken	131
5.9	Dendrogram generated from the arbitrarily primed-PCR patterns with primer GEN1-50-10 of the <i>L. monocytogenes</i> isolates from chicken.	132
5.10	Dendrogram generated from the arbitrarily primed-PCR patterns with primer GEN-1-50-02 of the <i>L. monocytogenes</i> isolates from beef	133
5.11	Dendrogram generated from the arbitrarily primed-PCR patterns with primer GEN-1-50-10 of the <i>L. monocytogenes</i> isolates from beef	134

5.12	Dendrogram generated from the repetitive sequence-PCR patterns with repetitive primer of the <i>L. monocytogenes</i> isolates from chicken	135
5.13	Dendrogram generated from the repetitive sequence-PCR patterns with repetitive sequence primer of the <i>L. monocytogenes</i> isolates from beef.	136

LIST OF ABBREVIATIONS

%	Percentage
β	Beta
μl	Microliter
μm	Micrometer
16S rRNA	16 subunit ribosomal ribose nucleic acid
A	Adenine or adenosine
Amp	Ampicillin
AP-PCR	Arbitrarily primed-polymerase chain reaction
ATCC	American type culture collection
ATP	Adenosine triphosphate
B	Bacitracin
bp	Base pair
CAMP Test	Christie-Atkins-Munch-Peterson Test
Car	Carbenicillin
CCC	Covalently close circular
CC	Close circular
C	Chloramphenicol or Cytosine
CDC	Centre for Disease Control
cDNA	Complementary DNA
Cfs	Cefsulodin
cm	Centimeter
Cn	Gentamicin
CO ₂	Carbon dioxide
dCTP	Deoxycytosine triphosphate
DNA	Deoxyribonucleic acid
dNTP	Deoxyribonucleoside triphosphate
dTTP	Deoxythymidine triphosphate
DW	Distilled water
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	Ethylenediaminetetra-acetic acid
E	Erythromycine
g	Gram

G	Guanine
H ₂ SO ₄	Sulfuric acid
HACCP	Hazard analysis critical control point
HCl	Hydrochloric acid
HIV	Human immunodeficiency virus
HLA	Haemolytic ceftazidime lithium chloride agar
I. U.	International unit
K	Kanamycin
Kb	Kilobase
KDa	Kilodalton
Kg	Kilogram
KH ₂ PO ₄	Potassium phosphate
<i>L. grayii</i>	<i>Listeria grayii</i>
<i>L. innocua</i>	<i>Listeria innocua</i>
<i>L. ivanovii</i>	<i>Listeria ivanovii</i>
<i>L. monocytogenes</i>	<i>Listeria monocytogenes</i>
<i>L. seelegeri</i>	<i>Listeria seeligeri</i>
<i>L. welshimeri</i>	<i>Listeria welshimeri</i>
LB	Luria-Bertani
LEB	<i>Listeria</i> enrichment broth
LMBA	<i>Listeria monocytogenes</i> basic agar
LPM	Lithium chloride phenylethanol moxlactam medium
Ltd.	Limited
MARI	Multiple antibiotic resistance index
MDa	Megadalton
MgCl ₂	Magnesium chloride
mg	Miligram
min	Minute
ml	Mililiter
mM	milimolar
MMWR	Morbidity and Mortality Weekly Report
Mox	Latamoxef
MR	Methyl red
mRNA	Messenger RNA

mr	Relative molecular weight
Na ₂ HPO ₄	Di-sodium hydrogen phosphate
NaCl	Sodium chloride
Na	Nalidixic acid
NaOH	Sodium hydroxide
NCCLS	National committee for clinical laboratory standard
Nm	Night market
nm	Nanometer
nt	Nucleotide
°C	Degree celcius
OD	Optical density
PCR	Polymerase chain reaction
PFGE	Pulsed-field gel electrophoresis
P	Penicillin G
<i>R. equi</i>	<i>Rhodococcus equi</i>
r.p.m.	Revolutions per minute
RAPD	Random amplified polymorphic DNA
RNA	Ribonucleic acid
<i>S. agalactiae</i>	<i>Streptococcus agalactiae</i>
<i>S. aureus</i>	<i>Staphylococcus aureus</i>
SDS	Sodium dodecyl sulphate
spp.	Species
ssDNA	Single stranded DNA
S	Streptomycin
Taq	<i>Thermus aquaticus</i>
TBE	Tris-Boric acid-EDTA
Te	Tetracycline
TE	Tris-EDTA
TSA	Trypticase soy agar
TSA-YE	Trypticase soya agar-yeast extract
TSB	Trypticase soy broth
TSI	Triple sugar ion
UK	United Kingdom
UPM	Universiti Putra Malaysia