

Evaluation of commercial soy sauce koji strains of *Aspergillus oryzae* for γ -aminobutyric acid (GABA) production

ABSTRACT

In this study, four selected commercial strains of *Aspergillus oryzae* were collected from soy sauce koji. These *A. oryzae* strains designated as NSK, NSZ, NSJ and NST shared similar morphological characteristics with the reference strain (*A. oryzae* FRR 1675) which confirmed them as *A. oryzae* species. They were further evaluated for their ability to produce γ -aminobutyric acid (GABA) by cultivating the spore suspension in a broth medium containing 0.4 % (w/v) of glutamic acid as a substrate for GABA production. The results showed that these strains were capable of producing GABA; however, the concentrations differed significantly ($P < 0.05$) among themselves. Based on the *A. oryzae* strains, highest GABA concentration was obtained from NSK (194 mg/L) followed by NSZ (63 mg/L), NSJ (51.53 mg/L) and NST (31.66 mg/L). Therefore, *A. oryzae* NSK was characterized and the sequence was found to be similar to *A. oryzae* and *A. flavus* with 99 % similarity. The evolutionary distance (K_{nuc}) between sequences of identical fungal species was calculated and a phylogenetic tree prepared from the K_{nuc} data showed that the isolate belonged to the *A. oryzae* species. This finding may allow the development of GABA-rich ingredients using *A. oryzae* NSK as a starter culture for soy sauce production.

Keyword: *Aspergillus oryzae*; Glutamic acid; γ -Aminobutyric acid; Fermentation; Soy sauce