



UNIVERSITI PUTRA MALAYSIA

***UTILIZATION OF SPENT MUSHROOM WASTE AND PEAT AS
SOILLESS GROWING MEDIUM FOR *Brassica oleracea* var. *Alboglabra****

HATARI SENDI

FP 2014 47



**UTILIZATION OF SPENT MUSHROOM WASTE AND PEAT AS
SOILLESS GROWING MEDIUM FOR *Brassica oleracea* var. Alboglabra**

By

HATARI SENDI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Agricultural Science**

December 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright@ Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Agricultural Science

UTILIZATION OF SPENT MUSHROOM WASTE AND PEAT AS SOILLESS GROWING MEDIUM FOR *Brassica oleracea* var. Alboglabra

By

HATARI SENDI

December 2014

Chairman: Mahmud Tengku Muda Mohamed, PhD

Faculty: Agriculture

Peat moss (PM) is the most widely used growing substrate for the pot culture. Due to diminishing availability and increasing price of PM, researchers are looking for viable alternatives for peat as a growth media component for potted plants. A pot study was conducted with a view to investigate the possibility of using spent mushroom waste (SMW) for kailan (*Brassica oleracea* var. Alboglabra) production replacing PM in growth media. The treatments evaluated were 100% SMW, 100% PM, and mixtures of SMW and PM in different ratios like 1:1, 1:2 and 2:1 (v/v) with or without NPK amendment. The experiment was arranged in a completely randomized design (CRD) with five replications per treatment. Chemical properties like pH and salinity level (EC) of SMW were within the acceptable range of crop production but, nutrient content, especially nitrogen content was not enough to provide sufficient nutrition to plant for normal growth. Only PM (100%) and SMW and PM mixture in 1:2 ratio with NPK amendment performed equally in terms of kailan growth. This study confirms the feasibility of replacing PM by SMW up to maximum 50% in the growth media, and suggests NPK supplementation from inorganic sources to ensure higher productivity of kailan.

Further testing was conducted to study the effects of various rates of SMW with virgin peat (VP) and non-virgin peat (NVP) at different location on the growth and yield of kailan. The peat sampling for VP and NVP was collected at Malaysian Agricultural Research and Development Institute (MARDI), Pontian Johor. The treatments evaluated were 100% SMW, 100% VP, 100% NVP and mixtures of SMW and VP or NVP in different ratios like 1:1, 1:2 and 2:1 (v/v) with NPK amendment. The experiment was arranged in a completely randomized design (CRD) with five replications per treatment. Result was obtained from peat soil in Peninsular Malaysia (Johor) indicated that no significant differences occurred between treatments that contained the portion of SMW with VP and SMW with NVP when amended with NPK. Plant height, leaf number, leaf fresh weights, leaf dry weight, total leaf area and chlorophyll content increased significantly when kailan were planted with amended NPK especially on NVP. The media with 100%

NVP also showed a better plant growth because of the existing of fertilizer in the soil which is came from previous agriculture activities.

For experiment three, the peat sampling was collected at Malaysian Agricultural Research and Development Institute (MARDI), Saratok Sarawak. Result obtained from observation in Sarawak also showed the media with amended NPK, fertilized soilless, there is not much differences among VP and NVP. This is due to the pH reading for all treatments gave almost similar result in between pH 3.63 to pH 5.04.

Plants grown in soilless media that contained portions of SMW and NVP gave better result compared to plants grown in media that contained SMW and VP. Further studies should be conducted under various conditions with plant species or other horticultural crops. Since imported PM is expensive and not economical to use as part of the soilless media, more studies need to be done to determine the modifications on soil fertility, and fertilizer requirements should be taken into account to enable SMW and peat compound to become a soilless media.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains Pertanian

PENGGUNAAN SISA BUANGAN CENDAWAN DAN GAMBUT SEBAGAI MEDIA PERTUMBUHAN KURANG TANAH UNTUK *Brassica oleracea* var. *Alboglabra*

Oleh

HATARI SENDI

Disember 2014

Pengerusi: Mahmud Tengku Muda Mohamed, PhD

Fakulti: Pertanian

Peat moss (PM) adalah media tanaman yang paling meluas digunakan untuk kultur pot. Disebabkan ketersediaan pengurangan dan kenaikan harga PM, para penyelidik melihat gambut sebagai alternatif media pertumbuhan bagi tanaman berpasu. Kajian ini dilaksanakan untuk menilai kemungkinan menggunakan bahan buangan cendawan (SMW) untuk pengeluaran kailan (*Brassica oleracea* var. *Alboglabra*) menggantikan PM dalam media tanaman. Rawatan yang telah dibuat adalah 100% SMW, 100% PM pada nisbah berbeza seperti 1:1, 1:2, 2:1 dengan pengubahan NPK. Rekabentuk kajian yang digunakan ialah rekabentuk penuh rawak lengkap (CRD) dengan lima replikasi setiap rawatan. Sifat kimia seperti pH dan tahap kemasinan bahan buangan cendawan adalah di dalam lingkungan yang boleh diterima untuk pengeluaran tanaman tetapi kandungan nutrien khususnya kandungan nitrogen adalah tidak mencukupi untuk dijadikan zat makanan supaya tanaman ini tumbuh dalam keadaan biasa. Hanya PM (100%) dan campuran SMW dan PM pada nisbah 1:2 dengan NPK menunjukkan pertumbuhan sebenar kailan. Kajian ini mengesahkan SMW dapat menggantikan PM sehingga 50% dalam media tanaman dan mencadangkan gantian NPK dari sumber bukan organik untuk memastikan produktiviti kailan meningkat.

Kajian selanjutnya dijalankan untuk menilai keberkesanan kepelbagaian kadar SMW dengan tanah gambut dara (VP) dan bukan tanah gambut dara (NVP) di kawasan yang berbeza terhadap pertumbuhan dan hasil kailan. Persampelan gambut untuk VP dan NVP telah diambil di Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI), Pontian Johor. Rawatan yang telah dinilai adalah 100% SMW, 100% VP dan 100% NVP pada nisbah berbeza seperti 1:1, 1:2, 2:1 dengan pengubahan NPK. Rekabentuk kajian yang digunakan ialah rekabentuk penuh rawak lengkap (CRD) dengan lima replikasi setiap kawatan. Keputusan yang diperolehi di Semenanjung Malaysia (Johor) menunjukkan perbezaan yang tidak ketara pada rawatan yang mengandungi SMW dengan VP dan SMW dengan NVP apabila adanya pengubahan NPK. Ketinggian tanaman, jumlah bilangan daun, berat basah pucuk, berat kering pucuk, jumlah keluasan daun dan

kandungan klorofil meningkat dengan ketara apabila di tanam dengan campuran baja NPK terutamanya pada media NVP. Media yang mengandungi 100% NVP juga menunjukkan pertumbuhan tanaman yang lebih baik berbanding dengan rawatan kawalan disebabkan oleh kewujudan baja di dalam kandungan tanah yang terhasil daripada aktiviti pertanian sebelum ini.

Untuk kajian ketiga, persampelan gambut untuk VP dan NVP telah diambil di Institut Penyelidikan dan Kemajuan Pertanian Malaysia (MARDI), Saratok Sarawak. Keputusan yang diperolehi daripada pemerhatian di Sarawak menunjukkan media yang dibaja NPK, ianya tidak menunjukkan perbezaan yang ketara di antara VP dan NVP. Ini berikutan bacaan pH tanah untuk kesemua rawatan memberikan keputusan yang hampir sama di antara pH 3.63 hingga pH 5.04.

Pertumbuhan pokok dalam media kurang tanah yang mengandungi SMW dan NVP memberikan keputusan yang baik berbanding dengan tanaman pada media yang mengandungi SMW dan VP. Kajian pada masa akan datang haruslah dibuat dengan mengambil kira pelbagai keadaan dengan spesis tanaman atau tanaman hortikultur yang lain. Memandangkan PM import adalah mahal dan tidak ekonomi untuk digunakan sebagai sebahagian daripada media kurang tanah, kajian lanjut haruslah dibuat untuk menentukan pengubahsuaian pada kesuburan tanah, dan keperluan pembajaan perlulah di ambil kira untuk membolehkan SMW dan kandungan gambut boleh menjadi media kurang tanah.

ACKNOWLEDGEMENTS

In the name of Allah the Beneficial and the Compassionate

First of all, I thank Allah for giving me the strength and ability to complete this study.

I also wish to express my sincere thank and profound appreciation to the chairman of my supervisory committee, Professor Dr. Mahmud Tengku Muda Mohamed, Department of Crop Science, Faculty of Agriculture for his supervision, advice, encouragement and friendship. Special thanks to Associate Professor Dr. Halimi Mohd Saud as a member of my supervisory for his ideas and suggestion during the course of this project. Furthermore, I would also like to acknowledge with much appreciation to Dr. Anwar Perves for the useful comments on my thesis writing.

My special appreciation and gratitude to my beloved parents and my wife for their continuous support and encouragement to achieve higher level of education carry out this research.

Last but not least, most profound thanks to MARDI for giving me the opportunity to further my study at UPM. ALHAMDULILLAH...

I certify that a Thesis Examination Committee has met on 19 December 2014 to conduct the final examination of Hatari bin Sendi on his thesis entitled "Utilization of Spent Mushroom Waste and Peat as Soilless Growing Medium for *Brassica oleracea* var. *Alboglabra*" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Agricultural Science.

Members of the Thesis Examination Committee were as follows:

Mohd Rafii bin Yusop, PhD

Professor
Institute of Tropical Agriculture
Universiti Putra Malaysia
(Chairman)

Siti Hajar binti Ahmad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Datin Rosenani binti Abu Bakar, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Mohd Yusoff Abdullah, PhD

Senior Lecturer
Malaysian Agricultural Research and Development Institute (MARDI)
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 13 May 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Agricultural Science. The members of the Supervisory Committee were as follows:

Mahmud Tengku Muda Mohamed, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Halimi Mohd Saud, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: HATARI SENDI (GS22556)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of

Supervisory Committee: Mahmud Tengku Muda Mohamed, PhD

Signature: _____

Name of Member of

Supervisory Committee: Halimi Mohd Saud, PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xix
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	3
2.1 Mushroom substrates	3
2.2 Soilless culture	3
2.3 Spent mushroom waste	3
2.4 Kailan (<i>Brassica oleracea</i> var. Alboglabra)	4
2.5 Peat in Malaysia	6
2.6 Characteristics of peat soil in Malaysia	6
3 EFFECTS OF DIFFERENT RATIOS OF SPENT MUSHROOM WASTE AND PEAT MOSS AS SOILLESS MEDIA ON GROWTH AND YIELD OF KAILAN (<i>Brassica oleracea</i> var. Alboglabra)	11
3.1 Introduction	11
3.2 Materials and methods	13
3.2.1 Experimental site	13
3.2.2 Seedling preparation	13
3.2.3 Experimental treatments and design	13
3.2.4 Media preparation and potting	14
3.2.5 Data collection	15
3.2.6 Statistical analysis	15
3.3 Results	16
3.3.1 Acidity and salinity of media	16
3.3.2 Nutrient status of media	17
3.3.3 Growth and yield responses	18
3.3.3.1 Plant height of kailan	18
3.3.3.2 Leaf number of kailan	19
3.3.3.3 Shoot fresh weight of kailan	20
3.3.3.4 Shoot dry weight of kailan	21
3.3.3.5 Root length of kailan	22
3.3.3.6 Leaf area of kailan	23
3.3.3.7 SPAD value of kailan	24

3.4	Discussion	25
3.5	Conclusion	28
4	EFFECTS OF DIFFERENT RATIOS OF SPENT MUSHROOM WASTE MIXED WITH VIRGIN AND NON-VIRGIN PEAT AS SOILLESS MEDIA ON GROWTH AND YIELD OF KAILAN (<i>Brassica oleracea</i> var. <i>Alboglabra</i>)	
4.1	Introduction	29
4.2	Materials and methods	31
4.2.1	Experimental site	31
4.2.2	Seedling preparation	31
4.2.3	Experimental treatments and design	31
4.2.4	Media preparation and potting	33
4.2.5	Data collection	34
4.2.6	Statistical analysis	34
4.3	Results	35
4.3.1	Acidity and salinity of media	35
4.3.2	Nutrient status of media	36
4.3.3	Growth and yield responses	37
4.3.3.1	Plant height of kailan	37
4.3.3.2	Leaf number of kailan	38
4.3.3.3	Shoot fresh weight of kailan	39
4.3.3.4	Shoot dry weight of kailan	40
4.3.3.5	Root length of kailan	41
4.3.3.6	Leaf area of kailan	42
4.3.3.7	SPAD value of kailan	43
4.4	Discussion	44
4.5	Conclusion	46
5	EFFECTS OF DIFFERENT RATIOS OF SPENT MUSHROOM WASTE MIXED WITH SARAWAK'S VIRGIN AND NON-VIRGIN PEAT AS SOILLESS MEDIA ON GROWTH AND YIELD OF KAILAN (<i>Brassica oleracea</i> var. <i>alboglabra</i>)	
5.1	Introduction	47
5.2	Materials and methods	50
5.2.1	Experimental site	50
5.2.2	Seedling preparation	50
5.2.3	Experimental treatments and design	50
5.2.4	Media preparation and potting	52
5.2.5	Data collection	52
5.2.6	Statistical analysis	53
5.3	Results	54
5.3.1	Acidity and salinity of media	54
5.3.2	Nutrient status of media	55
5.3.3	Growth and yield responses	56
5.3.3.1	Plant height of kailan	56
5.3.3.2	Leaf number of kailan	57
5.3.3.3	Shoot fresh weight of kailan	58

5.3.3.4	Shoot dry weight of kailan	59
5.3.3.5	Root length of kailan	60
5.3.3.6	Leaf area of kailan	61
5.3.3.7	SPAD value of kailan	62
5.4	Discussion	63
5.5	Conclusion	65
6	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	66
	REFERENCES	69
	APPENDICES	80
	BIODATA OF STUDENT	89
	PUBLICATIONS	90

LIST OF TABLES

Table	Page
2.1 Nutrient composition of kailan (Per 100 gram edible portion)	5
2.2 Division of peat area distinguished by countries	8
2.3 The area (ha) of peatlands in Peninsular Malaysia by state	9
2.4 Physical properties of peat soil in Peninsular Malaysia	9
2.5 Geotechnical properties of Sarawak peat	10
3.1 Soilless media at different ratios of spent mushroom waste (SMW) and peat moss (PM)	14
3.2 Chemical composition of study media	16
4.1 The soilless media at different ratios of spent mushroom waste (SMW) with virgin peat (VP) and non-virgin peat (NVP)	32
4.2 Chemical composition of study media	35
5.1 Estimates of the peatland area in Sarawak	49
5.2 The soilless media at different ratios of spent mushroom waste (SMW) with virgin peat (VP) and non-virgin peat (NVP) at Sarawak, West Malaysia	51
5.3 Chemical composition of study media	54

LIST OF FIGURES

Figure	Page
3.1 Media of different ratios of spent mushroom waste (SMW) and peat moss (PM).	14
3.2 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on plant height of <i>Brassica alboglabra</i> at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$.	18
3.3 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on number of leaves of <i>Brassica alboglabra</i> at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$.	19
3.4 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on shoot fresh weight of <i>Brassica alboglabra</i> at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$.	20
3.5 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on shoot dry weight of <i>Brassica alboglabra</i> at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$.	21

- 3.6 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on root length of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$. 22
- 3.7 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on total leaf area of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$. 23
- 3.8 Effect of different ratios of spent mushroom waste (SMW) and peat moss (PM) on chlorophyll contents of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter were not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ PM; $T_3=1(\text{SMW}):1(\text{PM})$; $T_4=1(\text{SMW}):2(\text{PM})$; $T_5=2(\text{SMW}):1(\text{PM})$; $T_6=1(\text{SMW}):1(\text{PM}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{PM}) + \text{NPK}$ and $T_8=2(\text{SMW}):1(\text{PM}) + \text{NPK}$. 24
- 4.1 Media of different ratios of spent mushroom waste (SMW) and virgin peat (VP). 32
- 4.2 Media of different ratios of spent mushroom waste (SMW) and non-virgin peat (NVP). 33
- 4.3 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on plant height of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 37

4.4 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on number of leaves of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. T₁=100% SMW; T₂=100% VP; T₃=100% NVP; T₄=1(SMW):1(VP) + NPK; T₅=1(SMW):1(NVP) + NPK; T₆=1(SMW):2(VP) + NPK; T₇=1(SMW):2(NVP) + NPK; T₈=2(SMW):1(VP) + NPK; T₉=2(SMW):1(NVP) + NPK. 38

4.5 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on shoot fresh weight of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. T₁=100% SMW; T₂=100% VP; T₃=100% NVP; T₄=1(SMW):1(VP) + NPK; T₅=1(SMW):1(NVP) + NPK; T₆=1(SMW):2(VP) + NPK; T₇=1(SMW):2(NVP) + NPK; T₈=2(SMW):1(VP) + NPK; T₉=2(SMW):1(NVP) + NPK. 39

4.6 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on shoot dry weight of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. T₁=100% SMW; T₂=100% VP; T₃=100% NVP; T₄=1(SMW):1(VP) + NPK; T₅=1(SMW):1(NVP) + NPK; T₆=1(SMW):2(VP) + NPK; T₇=1(SMW):2(NVP) + NPK; T₈=2(SMW):1(VP) + NPK; T₉=2(SMW):1(NVP) + NPK. 40

4.7 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on root length of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. T₁=100% SMW; T₂=100% VP; T₃=100% NVP; T₄=1(SMW):1(VP) + NPK; T₅=1(SMW):1(NVP) + NPK; T₆=1(SMW):2(VP) + NPK; T₇=1(SMW):2(NVP) + NPK; T₈=2(SMW):1(VP) + NPK; T₉=2(SMW):1(NVP) + NPK. 41

4.8 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on total leaf area of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. T₁=100% SMW; T₂=100% VP; T₃=100% NVP; T₄=1(SMW):1(VP) + NPK; T₅=1(SMW):1(NVP) + NPK; T₆=1(SMW):2(VP) + NPK; T₇=1(SMW):2(NVP) + NPK; T₈=2(SMW):1(VP) + NPK; T₉=2(SMW):1(NVP) + NPK. 42

- 4.9 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on chlorophyll contents of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 43
- 5.1 Media of different ratios of spent mushroom waste (SMW) and virgin peat (VP). 51
- 5.2 Media of different ratios of spent mushroom waste (SMW) and non-virgin peat (NVP). 52
- 5.3 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on plant height of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 56
- 5.4 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on number of leaves of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 57
- 5.5 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on shoot fresh weight of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 58

- 5.6 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on shoot dry weight of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 59
- 5.7 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on root length of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 60
- 5.8 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on total leaf area of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 61
- 5.9 Effect of different ratios of spent mushroom waste (SMW) mixed with virgin peat (VP) or non-virgin peat (NVP) with NPK amended on chlorophyll contents of *Brassica alboglabra* at 5 weeks after transplanting (WAT). Means with the same letter are not significantly different ($p < 0.05$) according to least significant different (LSD) test. $T_1=100\%$ SMW; $T_2=100\%$ VP; $T_3=100\%$ NVP; $T_4=1(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_5=1(\text{SMW}):1(\text{NVP}) + \text{NPK}$; $T_6=1(\text{SMW}):2(\text{VP}) + \text{NPK}$; $T_7=1(\text{SMW}):2(\text{NVP}) + \text{NPK}$; $T_8=2(\text{SMW}):1(\text{VP}) + \text{NPK}$; $T_9=2(\text{SMW}):1(\text{NVP}) + \text{NPK}$. 62

LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
CRD	Completely randomized design
EC	Electrical conductivity
LSD	Least significant difference
NAP	National agrofood policy
NVP	Non-virgin peat
PM	Peat moss
SMC	Spent mushroom compost
SMS	Spent mushroom substrate
SMW	Spent mushroom waste
VP	Virgin peat
WAT	Week after transplanting

© COPYRIGHT UPM



CHAPTER 1

INTRODUCTION

Mushroom cultivation is a profitable agribusiness. In 2007, global mushroom production amounted to 3.4 million tons (USITC, 2010). China, European Union and the United States were the largest global producers of mushrooms. Other important global producers in year 2007 included Canada, Japan, India, Australia and Indonesia. Global mushroom consumption amounted to 3.3 million tons and China, the European Union and the United States were the leading consumers of mushrooms. Other major consumers included Canada, Japan, Russia, Australia and India. All consumption in China, European Union and India was supplied from domestic production. Global exports of canned mushrooms amounted to 458 thousand tons with China accounting for 87% of total volume in 2008. However, global imports of canned mushroom amounted to 292 thousand tons with the United States and Russia accounting for the largest of total import volume in 2008 (USITC, 2010). For the global exports of fresh mushroom, Canada and the United States were the largest global exporters in 2008 together accounting almost 80% of the total volume. Other major exporters of fresh mushrooms in 2008 included Malaysia and Mexico. For the global imports of fresh mushroom, Russia and the United States together were the most important global import markets (USITC, 2010).

In Malaysia, the cultivation technique of the mushroom has been introduced into Malaysia from Canton and has been grown at Penang since 1934 (Baker, 1934). In 2008, there were 436 mushroom growers in Malaysia (Khairul *et al.*, 2012). In year 2007, total import for mushroom is 16,243 tonnes with a value of RM 50.40 million compared to total quantity of exported mushroom which was only 3,506 tonnes with a value of RM 25.1 million (DOSM, 2005). Among the types of mushrooms grown in Malaysia, grey oyster mushroom (*Pleurotus sojar-caju*) is widely cultivated. Oyster mushroom is harvested four to seven times from each production cycle. Normally, at the end of the mushroom fruiting cycle, “spent” (used) mushroom substrate is left abandoned or discarded (Danny and Seung, 2004). If not properly handled, these spent substrates may eventually cause environmental hazard.

Worldwide, mushrooms are produced on natural materials obtained from agriculture, woodlands, animal husbandry and manufacturing industries. Some of these wastes include banana leaves, peanut hull and corn leaves, mango fruits and seeds, sugarcane leaves, wheat and rice straw (Cangy and Peerally, 1995). The widely used substrate for cultivation of the oyster mushroom in Asia is rice straw (Thomas *et al.*, 1998). Rice straw considered as the best substrate in terms of yield and protein content of mushroom. In Europe, wheat straw is used as a growing medium for mushroom, while in the South East Asian countries sawdust is more common. Adding of soil, peat moss or both components to the media will increase the nutrient contents. Hence, growth of plants in media that was formulated with peat, soil or both amended with fertilizers is better than without amendment.

In recent years, plant growing in a soilless culture has a great importance worldwide. Lohr *et al.* (1984) and Maher (1991) reported that spent mushroom waste (SMW) also was used as plant growing medium. Mixtures of SMW with high quality growing medium such as peat, vermiculite and bark were given a significant result than used alone in studies on SMW (Calvin *et al.*, 1994). SMW from *Agaricus bisporus* production is used in horticulture as a component of potting soil mixes, soil amendment to improve turf-grass, in wetlands for remediation of contaminated water, in stabilizing severely disturbed soils, as bedding for animals and to control plant diseases (Danny, 2002). SMW from *Agaricus bisporus* has found acceptance as feed or animals, ingredients in the cultivation of other mushroom species, fuel, medium for vermiculture, in agriculture or landscape to enrich soils and as a matrix for bio-remediation of contaminated soils (Danny, 2002).

Shi *et al.* (2008) reported that the electronic conductivity value of SMW is high but low for peat. Thus, the pH value of SMW can be lowered by mixing with peat. If SMW is used in proper proportion, it can act as peat substitute in soilless culture (Shi *et al.*, 2008). According to Wang *et al.* (1984) increased plant growth and yield were obtained with the addition of 20 to 30% of SMW with sandy loam. However, vegetables grown at 50% of SMW with sandy loam exhibited some stunting effect.

In Europe, farmers are using SMW, supplemented with other materials to cultivate plants but, in Malaysia there are not much scientific research done to use it in combination with other agricultural by-products. Study on the use of SMW as a part of soilless media is still scarce in Malaysia, thus this study was deemed necessary. This study was undertaken with an ultimate aim to determine the effects of SMW alone and in combination with peat moss (PM) and peat on growth and quality of kailan, *Brassica oleracea* var. Alboglabra. This variety is a popular variety, easy to grow and contain high nutritional value. The specific objectives were:

- i. To evaluate the effects of spent mushroom waste and different ratios of spent mushroom waste and peat moss with and without fertilizer on growth and yield of *Brassica oleracea* var. Alboglabra at an open field.
- ii. To evaluate the effects of different ratios of spent mushroom waste, virgin peat and non-virgin peat in soilless media on growth and yield of *Brassica oleracea* var. Alboglabra at an open field.
- iii. To evaluate the effects of different ratios of spent mushroom waste, virgin peat and non-virgin peat in soilless media on growth and yield of *Brassica oleracea* var. Alboglabra at Sarawak, East Malaysia under netted structure.

REFERENCES

- Abad, M., Noguera, P., & Bures, S. (2001). National inventory of organic wastes for use as growth media for ornamental potted plant production: case study in Spain. *Bioresource Technology*, 77, 197-200.
- Akanbi, W.B., Togun, A.O., Adedrian, J.A., & Ilupeju, A.O. (2010). Growth dry matter and fruit yields component of okra under organic and inorganic sources of nutrients. *American Eurasian Journal of Sustainable Agriculture*, 49(1), 1-13.
- Akande, M.O., Oluwatoyinbo, F.I., Makinde, E.A., Adepujo, E.S., & Adepujo, I.S. (2010). Response of okra to organic and inorganic fertilization. *Nature and Science*, 8(11), 261-266.
- Altland, J.E., Gilliam, C.H., Edwards, J.H., Keever, G.J., Fare, D.C., & Sibley, J.L. (2002). Rapid determination of nitrogen status in annual vinca. *Journal of Environment and Horticulture*, 20, 189-194.
- Altland, J.E. (2006). Substrate pH, a tricky topic. *Digger*, 50, 42-47.
- Altland, J.E., & Krause, C. (2010). Modification of switchgrass substrate pH using compost, peatmoss, and elemental sulphur. *HortTechnology*, 20(6), 950-956.
- Ambak, K., & Melling, L. (2000). *Management practices for sustainable cultivation of crop plants on tropical peatland*. Paper presented at the International Symposium on Tropical Peatlands, Bogor, Indonesia.
- Anderson, J.A.R. (1964). The structure and development of the peat swamps on Sarawak and Brunei. *The Journal of Tropical Geography*, 18, 7-16.
- Anderson, J.A.R. (1972). *Trees of peat swamp forests of Sarawak*. Forest Department, Sarawak.
- Appanah, S. (1999). *Flora survey of North Selangor Peat Swamp Forest*. Sustainable Management of Peat Swamp Forest in Peninsular Malaysia. Volume I, Resource and Environment. Forestry Department, Kuala Lumpur, Peninsular Malaysia.
- Awang, Y., Shaharom, A.S., Mohamad, R.B., & Selamat, A. (2010). Growth dynamics of *Celosia cristata* grown in cocopeat, burnt rice hull and kenaf core fibre mixtures. *American Journal of Agricultural and Biological Sciences*, 5, 70-76.
- Baker, J.A. (1934). Mushroom growing in Wellesley and Penang Provinces. *Malay Agriculture Journal*, 22, 25-28.

- Bennett, E.L., & Gombek, F. (1991). *Wildlife and conservation in Sarawak's peat swamp forests*. Paper presented at the International Symposium on Tropical Peatlands, Kuching, Sarawak, Malaysia.
- Bunt, A.C. (1988). *Media and mixes for container-grown plants*. Second ed. of modern potting composts. Unwin Hyman Ltd., London, UK.
- Burger, B.W., Hartz, T.K., & Forister, G.W. (1997). Composted green waste as a container medium amendment for the production of ornamental plants. *HortScience*, 32(1), 57-60.
- Bustamante, M.A., Paredes, C., Moral, R., Agullo, E., Perez-Murcia, M.D., & Abad, M. (2008). Composts from distillery wastes as peat substitutes for transplant production. *Elsevier*, 52(5), 792-799.
- Calvin, C., Cline, R.A., & Rinker, D.L. (1994). Bark and peat amended spent mushroom compost for containerized culture of shrubs. *HortScience*, 29, 781-784.
- Cangy, C., & Peerally, A. (1995). Studies of *Pleurotus* production on sugarcane bagasse. *African Journal of Mycology and Biotechnology*, 3, 67-79.
- Chang, S.T. (1982). Cultivation of *Volvariella* mushrooms in Southeast Asia. *Tropical mushrooms: biological nature and cultivation methods*. 12, 221-252.
- Chang, S.T., & Miles, P.G. (1982). Introduction to mushroom science. *Tropical mushrooms: biological nature and cultivation methods*. 1, 1-10.
- Chew, T.A., Abu Hasssan, I., & Mohayidin, M.G. (2000). Estimating the environmental benefits of sago cultivation. *Repositioning of the Agriculture Industry in the Next Millenium*. Eds A Radan and FM Arshad. Universiti Putra Malaysia, Serdang, Selangor. 235-251.
- Chiu, S.W., Ching, M.L., Fong, K.L., & Moore, D. (1998). Spent oyster mushroom substrate perfoms better than many mushroom mycelia in removing the biocide pentachlorophenol. *Cambridge Journals*, 102(12), 1553-1562.
- Chong, C., Cline, R.A., Rinker, D.L., & Allen, O.B. (1991a). Growth and mineral nutrient status of containerized woody species in media amended with spent mushroom compost. *Journal of the American Society for Horticultural Science*, 116, 242-247.
- Chong, C., Cline, R.A., & Rinker, D.L. (1991b). Rapid leaching of salts- Key to successful plant growth in spent mushroom compost potting mixes. *Mushroom News*, 39(10). 12-13.
- Chong, C., Rinker, D.L., & Cline, R.A. (1991c). A comparison of live spent mushroom composts for container culture of ornamental shrubs. *Mushroom Science*, 13(2), 637-644.

- Chong, C., & Rinker, D.L. (1994). Use of spent mushroom substrate for growing containerized woody ornamentals. An overview. *Journal of the American Society for Horticultural Science*, 2(3), 45-53.
- Chong, C., Cline, R.A., & Rinker, D.L. (1994). Bark- and peat-amended spent mushroom compost for containerized culture of shrubs. *HortScience*, 29(7), 781-784.
- Chong, C. (2005). Experiences with wastes and composts in nursery substrates. *HortTechnology*. 15, 739-747.
- Danny, L.R. (2002). Handling and using “spent” mushroom substrate around the world. *Mushroom Biology and Mushroom Products*. Sánchez et al. (Eds).UAEM. ISBN 968-878-105.
- Danny, L.R., & Seung, W.K. (2004). *Recycling of Spent Mushroom Substrate*. Oyster Mushroom Cultivation. 187-191.
- Davis, J.G., & Wilson C.R. (2005). *Choosing a Soil Amendment* online from <http://www.ext.colostate.edu/pubs/garden/07235.html> (Retrieved 23 February 2011)
- DOA. (2012). Department of Agriculture. Hectareage, production and value of production of vegetables, Malaysia. *Vegetables and Cash Crops Statistic Malaysia*.
- Eklind, Y., Ramert, B., & Wivstad, M. (2001). Evaluation of growing media containing farmyard manure compost, household waste compost or chicken manure for the propagation of lettuce (*Lactuca sativa* L.) transplants. *Biological Agriculture & Horticulture. Journal for Sustainable Production Systems*, 19(2), 157-181.
- Ersin, P., Ibrahim, U.H., Bulent, T., Kubilay, O., Naci, O.A., & Mehmet, K. (2009). Effects of spent mushroom compost on quality and productivity of cucumber (*Cucumis sativus* L.) grown in greenhouses. *African Journal of Biotechnology*, 8(2), 176-180.
- Eudoxie, G.D., & Alexander, I.A. (2011). Spent mushroom substrate as a transplant media replacement for commercial peat in tomato seedling production. *Journal of Agricultural Science*, 3, 41-49.
- FAO. (1990). Soilless culture for horticultural crop production. *FAO Plant Production and Protection Paper*, ROME, 188.
- FAO. (2012). Food and Agriculture Organization. *World mushroom production quantity* online from [http:// faostat.fao.org/site/567/Desktop Default.aspx](http://faostat.fao.org/site/567/DesktopDefault.aspx) (Retrieved 3 August 2013)
- Fasidi, I.O., Kadiri, M., Jonathan, S.G., Adenipekun, C.O., & Kuforiji, O.O. (2008). *Cultivation of tropical mushrooms*. Ibadan University Press, Ibadan.

- Geoffrey, I.M. (2001). *Making Use of Sawdust* online from http://www.suite101.com/pages/article_old.cfm/organic/68287 (Retrieved 23 February 2011)
- Graham, K.M. (1975). Studies on the padi mushroom (*Volvariella volvacea*). V. Performance of single-basidiospore isolates. *Malaysian Agricultural Research Journal*, 5, 31-36.
- Griffin, D.H. (1994). *Fungal Physiology*, 2nd edn. Wiley Liss, New York.
- Grigatti, M. (2008). Growth and nutritional status of bedding plants on compost-based growth media. *Acta Horticulturae*, 77(9), 607-614.
- Gui-xiao, L., Ping, F., Yi-bo, T., & Ya-juan, L. (2009). Effect of CO₂ enrichment on the glucosinolate contents under different nitrogen levels in bolting stem of Chinese kale (*Brassica alboglabra* L.). *Journal of Zhejiang University Science B.*, 10(6), 454-464.
- Hammond, R.F. (1975). The origin, formation and distribution of peatland resources. *Peat in horticulture*. Academic press inc. (London) Ltd. 1,1-22.
- Hashim, R., & Islam, S. (2008). Engineering properties of peat soils in Peninsular, Malaysia. *Journal of Applied Sciences*, 8(22), 4215-4219.
- Handreck, K.A., & Black, N. (1991). *Growing media for ornamental plants and turf*. New South Wales University Press. Kensington. Australia.
- Idowu, O.O., & Kadiri, M. (2013). Growth and yield response of okra to spent mushroom compost from the cultivation of *Pleurotus ostreatus* an edible mushroom. *Academic Journal of Agricultural Research*, 1(1), 39-44.
- Idris, A., & Abdull Ghaffar, O. (2010). Tanaman Cendawan. *Nota Ringkas Tanaman Makanan*. 53-57.
- Irshad, F., & Saleem, Q.T (2011). Designing a Low Cost Super Salinity Measuring Device for Soil and Water. *International Journal of Chemical and Environmental Engineering*, 2, 56-61.
- Ismail, A.B. (1984). *Characterization of lowland organic soils in Peninsular Malaysia*. Paper presented at the Workshop Classification and Management of Peat, Kuala Lumpur, Malaysia.
- Ismail, A.B., Asing, J., & Zulkifli, M. (2007). Residual impact of various land clearing techniques on peat chemical characteristics. In *Impact of Land Clearing on Peat Ecosystems. A Case Study at MARDI Peat Research Station, Sessang, Sarawak, Malaysia*, 33-61.

- Issarakraisila, M., Ma, Q., & Turner, D.W. (2007). Photosynthetic and growth responses of juvenile Chinese kale (*Brassica oleracea* var. *alboglabra*) and Caisin (*Brassica rapa* subsp. *parachinensis*) to waterlogging and water deficit. *Scientia Horticulturae*, 111(2), 107-113.
- Iwase, Y., Umezawa, K., & Masuda, K. (2000). Cultivation of *Pleurotus ostreatus* with beer spent grain and utilization. *Mushroom Science*, 15, 819-826.
- Jayasinghe, G.Y., Arachichi, I.D.L., & Yoshihiro, T. (2010). Evaluation of containerized substrates developed from cattle manure compost and synthetic aggregates for ornamental plant production as a peat alternative. *Resource Conservation and Recycling*, 54(12), 1412-1418.
- Jonathan, S.G. (2002). *Vegetative growth requirements and antimicrobial activities of some higher fungi in Nigeria*. PhD dissertation. University of Ibadan, Ibadan.
- Jonathan, S.G., Lawal, M.M., & Oyetunji, O.J. (2011). Effect of spent mushroom compost of *Pleurotus pulmonaris* on growth performance of four Nigerian vegetables. *Mycobiology*, 39, 164-169.
- Jordan, S.N., Mullen, G.J., & Murphy, M.C. (2008). Composition of variability of spent mushroom compost in Ireland. *Bioresource Technology*, 99, 411-418.
- Kadiri, M., & Mustapha, Y. (2010). The use of spent mushroom substrate of *lentinus subnudus* (Berk) as a soil conditioner for vegetables. *Bayero Journal of Pure and Applied Science*, 3, 16-19.
- Kanapathy, K. (1968). *Peat soils in diversification of Malaysia Agriculture*. West Malaysia.
- Khairul Asfamawi, K., Yaseer Suhaimi, M., Mohamad, A.M., Aminuddin, H., & Mohd Hassan, W. (2012). *Manual Teknologi Penanaman Cendawan Tiram Kelabu*. MARDI.
- Khan, Z., Tiyagi, S.A., Mahmood, I., & Rizvi, R. (2012). Effects of N fertilization, organic matter, and biofertilizers on the growth and yield of chilly in relation to management of plant-parasitic nematodes. *Turkish Journal of Botany*, 36(81), 73-81.
- Lee, B.S., & Noor Auni, H. (1990). *Panduan Penanaman Cendawan. Siri Panduan Tanaman*, MARDI.
- Lee, H.S. (1991). *Utilization and conservation of peat swamp forest in Sarawak*. Paper presented at the International Symposium on Tropical Peatland, Kuching, Sarawak, Malaysia.
- Lemaire, F., Dartigues, A., & Riviere, L.M. (1985). Properties of substrate made with spent mushroom compost. *Acta Horticulturae*, 172, 13-29.

- Lemaire, F. (1997). The problem of the biostability in organic substrates. In *Proceedings of the ISHS International Symposium on Growing Media and Plant Nutrition*. Roeber, Freising, Germany. *Acta Horticulturae*, 450, 63-70.
- Lennartsson, M. (1997). *The peat conservation issue and the need for alternatives*. Paper presented at the International Peat Conference on Peat in Horticulture, Schmilewski, Amsterdam.
- Lette, R. (2006). *Malaysia's peat swamp forest conservation and sustainable use*. Published by United Nations Development Programme Malaysia. Ministry of Natural Resources and Environment, Malaysia.
- Lim, C.K., Chin, Y.K., & Bolle-Jones, E.W. (1973). Crop indicators of nutrient status of peat soil. *Malaysian Agricultural Research Journal*, 49, 198-207.
- Liong, T.Y., & Siong, K.H. (1979). A review of lowland organic soils of Sarawak. *Technical paper no. 4*. Research Branch Department of Agriculture Sarawak.
- Lohr, I.V., Wang, S.H., & Wolt, J.D. (1984). Physical and chemical characteristics of fresh and aged spent mushroom compost. *HortScience*, 19, 681-683.
- Lucas, R.E., Rieke, P.E., Shickluna, J.C., & Cole, A. (1975). Lime and fertilizer requirements for peats. *Peat in horticulture*. Academic press inc. (London) Ltd. 4, 51-70.
- Maher, M.J. (1988). Spent mushroom compost as an organic manure and potting compost component. *Compost processes in waste management*. Brussels., 71-82.
- Maher, M.J. (1991). Spent mushroom compost as a nutrient source in peat based potting substrates. *Science and Cultivation of Edible Fungi*. Balkema, Rotterdam, 645-650.
- Maher, M.J. (1994). The use of spent mushroom substrate (SMS) as an organic manure and plant substrate component. *Compost Science and Utilization*, 2, 37-44.
- MARDI. (1996). *Master plan for MARDI Sessang Peat Research Station*. MARDI, Serdang.
- MARDI. (2005). *Anggaran Kos Pengeluaran dan Pendapatan untuk Sayuran dan Rempah*. 64-67.
- Medina, E., Paredes, C., Perez-Murcia, M.D., Bustamante, M.A., & Moral, R. (2009). Spent mushroom substrates as component of growing media for germination and growth of horticultural plants. *Elsevier*. 10(18), 4227-4232.

- Melling, L., Ambak, K., Osman, J., & Husni, A. (1999). *Water management for the sustainable utilisation of peat soils for agriculture*. Paper presented at the International Conference & Workshop on Tropical Peat Swamps, Penang, Malaysia.
- Melling, L. (2000). *Dalat and Mukah sago plantation peat soil study*. DOA, Sarawak, 37.
- Mohd Zahari, A.B., Abd Wahab, N., Ting, C.C., & Abd Rahim, N. (1981). Distribution, characteristics and utilities of problem soils in Malaysia. *A country report. Symp. on Distribution, Characteristics and Utilization of Problem Soils*. Tsukuba Ibaraki, Japan.
- Munita, J. (2001). Characteristics and classification of soils, in *Chemical and Mining Company of Chile*, 11th edn., Santiago, Chile. p. 1515.
- Mutalib, A.A., Lim, J.S., Wong, M.H., & Koonvai, L. (1991). *Characterization, distribution and utilization of peat in Malaysia*. Paper presented at the International Symposium on Tropical Peatlands, Kuching, Sarawak, Malaysia.
- Naidu, N.R. (1971). Cultivation of padi straw mushroom *Volvariella volvacea* (Fr.) Sing. Using oil palm bunch waste as a medium. *The Planter*, 47, 190-193.
- Natural Standard Professional Monograph. (2012). <http://www.naturalstandard.com/index-abstract.asp?create-abstract=brassica.asp&title=Cabbage,%20Broccoli,%20Cauliflower,%20Collard,%20Kale,%20Brussels%20sprouts,%20Kohlrabi> (Retrieved 5 May 2012)
- Okazaki, M., Yonebayashi, K., & Zahari, A.B. (1989). *Characteristic of soil and water in Wedland along the western coast of Johor, Malaysia*. Paper presented at the Workshop on Tropical Peat Ecosystem, Serdang, Selangor, Malaysia.
- Okazaki, M., Watanabe, C., Yoshikawa, M., Yamaguchi, C., & Yoshimura, N. (1999). *Chemical compounds in gas emitted from tropical peat soil with burning*. Paper presented at the International Symposium on Tropical Peatlands, Bogor, Indonesia.
- OMAF. (2003). Ontario Ministry of Agriculture and Food. *Nursery and landscape plant production and IPM, Publ. 383*. Queen's Printer for Ontario, Toronto, ON.
- Ostos, J.C., Lopez-Garrido, R., Murillo, J.M., & Lopez, R. (2008). Substitution of peat for municipal solid waste and sewage sludge-based composts in nursery growing media. Effects on growth and nutrition of the native shrub *Pistacia lentiscus* L. *Bioresource Technology*, 99, 1793-1800.

- Paramananthan, S., Zauyah, S., Lim, L., Chan, C.P., & Boaklan, Y.K., (1984). *Proposal for united classification of organic soil in Malaysia*. Paper presented at the Workshop on classification and Management of Peat in Malaysia, Kuala Lumpur, Malaysia.
- Paramananthan, S., & Omar, W. (2008). *Proposals for the classification of organic soils of Malaysia*. Paper presented at the meeting and seminar, Malaysia-Netherlands joint committee on carbon emissions. Bangi, Selangor, Malaysia.
- Perez-Murcia, M.D., Moral, R., Caselles, J.M., Espinosa, A.P., & Paredes, C. (2006). Use of composted sewage sludge in growth media for broccoli. *Bioresource Technology*, 97, 123-130.
- Peter, S. (2003). *Environmental issues of peatland development in Sarawak*. Paper presented at the workshop on management and conservation of peatland, Sibul, Sarawak, Malaysia
- Puustjarvi, V., & Robertson, R.A. (1975). Physical and Chemical properties. *Peat in horticulture*. Academic press inc. (London) Ltd. 2, 23-38.
- Radjagukguk, B. (1991). *Utilization and management of peat land in Indonesia for agriculture and forestry*. Paper presented at the International Symposium on Tropical Peatland, Kuching, Sarawak, Malaysia.
- Rainbow, A., & Wilson, N. (1998). The transformation of composted organic residues into effective growing media. *SHS Acta Horticulturae 469: International Symposium on Composting & Use of Composted Material in Horticulture*, 469, 79-88.
- Rashidah, A., Ismail, B., Devapriya, C.W., & Adnan, Z. (2012). Overview of the sustainable uses of peat soil in Malaysia with some relevant geotechnical assessments. *International Journal of Integrated Engineering*. Special Issue on ICONCEES, 4(3), 38-46.
- Ribas, L.C.C., De Mendonca, M.M., Camellini, C.M., & Soares, C.H.L. (2009). Use of spent mushroom substrates from *Agaricus subrufescens* (syn. *A. blazei*, *A. brasiliensis*) and *Lentinula edodes* productions in the enrichment of a soil-based potting media for lettuce (*Lactuca sativa*) cultivation: Growth promotion and soil bioremediation. *Bioresource Technology*, 100, 4750-4757.
- Ribeiro, H.M., Romero, A.M., Pereira, H., Borges, P., Cabral, F., & Vaconcelos, E. (2007). Evaluation of a compost obtained from forestry wastes and solid phase of pig slurry as a substrate for seedlings production. *Bioresource Technology*, 98, 3294-3297.
- Romaine, C.P., & Holcomb, E.J. (2001). Spent mushroom substrate: a novel multifunctional constituent of potting medium for plants. *Mushroom News*, 49, 4-15.

- Sa'don, N.M., Abdul Karim, A.R., Jaol, W., & Wan Lili, W.H. (2015). Sarawak peat characteristics and heat treatment. *Journal of Civil Engineering*. UNIMAS. 6-12.
- SAS. (2003). Statistical Analysis System. *The SAS system for Windows, Version 9.2*. SAS Inst. Inc., Cary, NC, USA.
- Schisler, L.C. (1982). Biochemical and mycological aspects of mushroom composting. In P. Wuest (Eds.). *Penn State handbook for commercial mushroom growers*. Pa. State University, University Park., 3-10.
- Sheard, G.F. (1975). Loamless substrates for use in containers and as unit products. *Peat in horticulture*. Academic press inc. (London) Ltd. 9, 119-132.
- Sherry, H.W., Virginia, I.L., & David, L.C. (1984). Growth response of selected vegetable crops to spent mushroom compost application in a controlled environment. *Plant and Soil*, 82, 31-40.
- Shi, L., Zhang, Z., Liu, D., Li, W., & Bao, R. (2008). Comparison of physiochemical properties between spent mushroom compost and peat substrate and adjustment. *Nongye Gongcheng Xuebao/Transactions of the Chinese Society of Agricultural Engineering*. 24(4), 199-203.
- Stewart, D.P.C., Cameron, K.C., Cornforth, I.S., & Sedcok, J.R. (1998). Effects of spent mushroom substrate on soil physical conditions and plant growth in an intensive horticultural system. *Australian Journal of Soil Research*, 36(6), 899-912.
- Tariq, U., Rehman, S.U., Khan, M.A., & Yunus, A. (2012). Agricultural and municipal waste as potting media components for the growth and flowering of *Dahlia hortensis* Figaro. *Turkish Journal of Botany*, 36(4), 378-385.
- Tay, T.H., & Wee, Y.C. (1972). Liming of peat soils and its effects on pineapple. Malay Pineapple Industry Board. *Malays Pineapple Journal*, 2, 53-59.
- Teng, C.S. (2003). *Peatland and agriculture in Sarawak. Research and development support and challenges. Sarawak DOA's experience*. Paper presented at the workshop on management and conservation of peatland, Sibu, Sarawak, Malaysia.
- Thomas, G.V., Prabhu, S.R., Reeny, M.Z., & Bopaiah, B.M. (1998). Evaluation of lignocellulosic biomass from coconut palm as substrate for cultivation of *Pleurotus sajor-caju* (Fr.) Singer. *World Journal of Microbiology and Biotechnology*, 14, 879-882.
- Tie, Y.L. (1979). A review of lowland organic soils of Sarawak. *Technical paper 4*. Department of Agriculture, Sarawak. 11-16.

- Tie, Y.L. (1991). Report on detailed soil surveys of the peat experimental stations. *Report No. 271*, Soil Division, Research Branch, Department of Agriculture, Sarawak.
- Tie, Y.L., and Esterle, J.S. (1991). *Formation of lowland peat domes in Sarawak, Malaysia*. Paper presented at the International Symposium on Tropical Peatlands, Kuching, Sarawak, Malaysia.
- USDA. (1975). *Soil Taxonomy*. A basic system of soil classification for making and interpreting soil surveys. Soil conservation services, USDA Agri. Handbook No. 436, U.S. Government Printing Office, Washington D.C.
- USDA. (2010). *National Nutrient Database for Standard Reference* online from <http://ndb.nal.usda.gov/ndb/foods/show/3642?fg=&man=&lfacet=&count=&max=25&sort=&qlookup=Broccoli%2C+chinese%2C+cooked&offset=&format=Abridged&new=1> (Retrieved 20 February 2012)
- USITC. (2010). *Mushrooms. Industry & Trade Summary*. United States International Trade Commission. 28-31 online from http://www.usitc.gov/publications/332/ITS_7.pdf (Retrieved 23 August 2013)
- Van Poppel, G., Verhoeven, D.T., Verhagen, H., & Goldbohm, R.A. (1999). Brassica vegetables and cancer prevention. Epidemiology and mechanisms. *Adv. Exp. Med. Biol.*, 472, 159-168.
- Vimala, P., & Chan, S.K. (1990). Tanah dan pembajaan. *Panduan pengeluaran sayur-sayuran*. MARDI. 5, 29-40.
- Wang, H.L.S. (1983). *Spent mushroom compost as a soil amendment for vegetables*. Ph.D. theses. The University of Tennessee.
- Wang, H.L.S., Virginia, I. L., & David, L.C. (1984). Growth response of selected vegetable crops to spent mushroom compost application in a controlled environment. *Plant and Soil*, 82, 31-41.
- Watson, M.E. (2003). *Testing compost*. Ohio State University Fact Sheet. Agriculture and Natural Resources, Ohio.
- Wetlands International Malaysia. (2010). *A quick scan of peatlands in Malaysia*. Wetlands International-Malaysia. Petaling Jaya, Malaysia. 50 pp.
- Wong, L.J. (2003). *Status of Peat Swamp Forest of Sarawak*. Sarawak Forestry Corporation/Alterra Wageningen UR: Kuching, Sarawak/Wageningen, Netherlands.
- Williams, B.C., McMullan, J.T., & McCahey, S. (2001). An initial assessment of spent mushroom compost as potential energy feedstock. *Bioresource Technology*, 79, 227-230.

- Yong, Y.C., and Graham, K.M. (1973). Studies on the mushroom (*Volvariella volvacea*). Use of oil palm pericarp waste as an alternative substrate. *Malaysian Agricultural Research Journal*, 2, 15-22.
- Yong, Y.C., and Graham, K.M. (1974). Studies on the mushroom (*Volvariella volvacea*). Effect of urea on mycelial development and upon yield in palm oil pericarp waste. *Malay Agriculture, Malaysian Agricultural Research Journal*, 3, 7-11.
- Young, J.R., Holcomb, E.J., & Heuser, C.W. (2002). Greenhouse growth of marigolds in three leached sources of spent mushroom compost over a 3-year period. *HortTechnology*, 12(4), 701-705.
- Zhang, R.H., Li, X.J., & Fadel, J.G. (2002). Oyster mushroom cultivation with rice and wheat straw. *Bioresource Technology*, 82, 277-284.
- Zhang, R.H., Duan, Z.Q., & Li, Z.G. (2012). Use of spent mushroom substrate as growing media for tomato and cucumber seedlings. *Pedosphere*, 22(3), 333-342.
- Zuraifah, A.M., Daranee, C., Kisana, B., Monique, P., Jisnuson, S., Pattana, S.H., & Chantragan, S. (2011). Proteomic analysis of Chinese kale (*B. alboglabra*) leaves during growth. *Journal of Integrated Omics*. 102-107.