



UNIVERSITI PUTRA MALAYSIA

***OCCURRENCE OF MULTIDRUG RESISTANT *E.coli* AND *Campylobacter*
IN CHICKEN AND CHICKEN MEAT AND THEIR ASSOCIATED
RISK FACTORS***

IBRAHIM JALO MUHAMMAD

FPV 2015 9



**OCCURRENCE OF MULTIDRUG RESISTANT *E.coli* AND *Campylobacter*
IN CHICKEN AND CHICKEN MEAT AND THEIR ASSOCIATED
RISK FACTORS**

By

IBRAHIM, JALO MUHAMMAD

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

May 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This thesis is specially dedicated to:

My beloved parents,

ALHAJI IBRAHIM JALO MUHAMMAD

and

HAJIYA FATSUMA IBRAHIM JALO

My beloved wife and children,

HUSSAINA ABUBAKAR LAWAN

FATIMA MUHAMMAD JALO (AMNA)

ABUBAKAR MUHAMMAD JALO (AMMAR)

Who always pray, support and encourage me to do the best

And finally,

In memory of my late sister

HAUWA IBRAHIM JALO

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Master of Science

**OCCURRENCE OF MULTIDRUG RESISTANT *E.coli* AND *Campylobacter*
IN CHICKEN AND CHICKEN MEAT AND THEIR ASSOCIATED
RISK FACTORS**

By

IBRAHIM JALO MUHAMMAD

May 2015

Chairperson: Professor Saleha Abdul Aziz, PhD

Faculty: Veterinary Medicine

Antibiotic usage is an important factor that have been widely reported to bring about the emergence and spread of antibiotic resistant microorganisms in both animal and human health. Chickens may be infected with antibiotic resistant *Campylobacter* and *E.coli* which may be resistant to one or more antibiotics. There is an alarming increase of multidrug resistance among bacterial pathogens affecting human and animal populations globally. There is limited data on the occurrence of multidrug resistance (MDR) *E. coli* and *Campylobacter* in poultry and poultry meat in Malaysia. This study was aimed at investigating the occurrence of multidrug resistant *E. coli* and *Campylobacter* in chicken and chicken meat and their associated risk factors. The specific objectives were: to determine the occurrence of *E.coli* and *Campylobacter* in chickens and farm environment and chicken meat retailed in the markets and their associated risk factors and to determine the antibiotic resistance and MDR profiles of the isolates from chicken and chicken meat. Two hundred and ten (210) samples were collected from five farms and seventy (70) samples from seven markets. Five (5) farms were visited three (3) times each during the study period. The first visit was to sample chicken at market age (35-42 days) prior to harvesting (considered as first population) and farm environment (which is included water, feed, flies and chicken house floor). The second visit was to sample farm environment at two (2) weeks after the chicken farms were emptied (all chicken were sold), cleaned and disinfected and the third visit was carried out on a new batch of chickens at market age also prior to harvesting (as second population) and the environment. In each farm, thirty (30) cloacal swab samples and twelve (12) environmental samples which consisted of three (3) samples each of water from the drinker, feed from the feeder, chicken house floor and pooled samples of flies each consisting 5-7 per pooled sample were collected (3). Chicken meat which consisted of breast, thigh and wings including the skin were randomly chosen from stalls in each market. The number of stalls visited in seven markets varied from 4 to 10 per market depending on the number of poultry meat retailers in that market. Cutting boards and weighing scales were also sampled. Fifty five point seven percent (60.0% and 51.3%) of the chickens in the first and second population were colonized with *Campylobacter*. The proportion of *Campylobacter* isolated from environmental samples in the first and third visits was 7.5% (1.7% and 13.3%). Only one water

sample (6.7%) was positive (first visit) and two (13.3%) in the third visit. Flies were found positive for *Campylobacter* only in the third visit (33.3%). Among the chicken house floor samples, only one was positive (6.7%). All feed samples were negative (0%). The most frequently isolated species was *C. jejuni* at 74.9% (82.2% and 67.5%), followed by *C. coli* 25.2% (17.8% and 32.5%). At the markets, the occurrence of *Campylobacter* was 20.0% for chicken meat, and 5.7% for weighing scales and cutting boards. The species identified were 78.6%, 17.9% and 3.6% for *C.jejuni*, *C.coli* and *C.upsaliensis* respectively. The occurrence of *E.coli* in chickens is 53.0% (56.0% and 50.0%). Similarly, the occurrence of *E.coli* in environmental samples at first and third visits were 33.4% (26.7%, 0% and 40.0%). Among the environmental samples, the isolation of *E.coli* in feed was 16.7%, floor 40.0%, water 20.0% and flies 56.7%. The occurrence of *E.coli* in chicken meat at markets was 45.0%. In the farms, the occurrence of the *Campylobacter* and *E.coli* in the first and second population were almost similar; this was probably due to the farm practices were similar and the biosecurity measures practised in the farm may not be sufficient. The presence of flies and birds around the farm and/or the use of contaminated water may transfer these bacteria to the chickens or the workers brought them into the chicken houses from the farm environment. At the markets, cross contamination was a possible factor because during meat handling and cutting, the surfaces of poultry carcasses could become contaminated with *Campylobacter* and *E.coli* from the intestinal content due to accidental rupture of the gut or from contaminated equipment or water. The absence of *Campylobacter* in feed samples may probably be due to poor resistance of *Campylobacter* to atmospheric condition and other environmental pressures during storage that would have converted to viable-but-non-culturable (VBNC) form. It was reported that most flocks were negative until two (2) weeks of age, and once *Campylobacter* colonized a broiler flock, the spread is very rapid and up to 100% of birds within a flock can become colonized within three days. The antibiotic resistance of these *Campylobacter* (n=208) and *E.coli* (n=269) isolates was done using disc diffusion method against 12 different antibiotics. *Campylobacter* isolates from first and second population showed high resistance to penicillins at 75.5% (74.4% and 76.6%) and the least resistance was to amoxicillin_clavulanic acid at 33.7% (32.2% and 35.1%). Sixty two point five percent (62.5%) *Campylobacter* isolated from chicken meat was resistant to ampicillin and the least was to streptomycin, 3.1%. *E.coli* isolated from chicken in first population showed high resistance to penicillin, erythromycin, ampicillin and tetracycline at 98.8%, 96.4%, 94.0% and 92.8% respectively and the least resistance was to amoxicillin_clavulanic at 22.9%. Those isolated from the second population were 100% resistant to erythromycin and tetracycline and the least resistance was to cefotaxime at 20.0%. In chicken meat, the highest resistance was to erythromycin and penicillins at 100% each and the least was to gentamicin at 40%. The high resistance may likely be as a result of imprudent use of these agents for growth promotion and for prophylactic purposes at farm level. Multidrug resistance (MDR) (resistance to 3 or more classes of antibiotics) was high in *Campylobacter* isolated in chicken farms, at 87.5%, chicken meat 71.9%, while in *E.coli* it was 99.5% for chicken, and for chicken meat was 100%. The most common multidrug resistance profile for *Campylobacter* isolates in chicken was to 7 (EPNaCtxSCipAmp) and 10 (EPNaCipTeAmcSxtCnEnrAmp) classes of antibiotics at 13.2% each in the first population. In the second population, the common MDR pattern was to 9 (EPNaCtxSCipAmcEnrAmp) classes at 12.9%. In chicken meat it was to 4 (TeAmcEnrAmp) and 5 (EPTeCtxAmp) classes at 15.6% each. The most

common MDR profile for *E. coli* was to 9 (EPNaSCipTeAmcSxtAmp) and 10 (EPNaSCipTeAmcSxtEnrAmp) antibiotic classes at 16.0% each in the first population and in the second population, the most common profile was to 8 (EPNaSCipTeAmcSxtEnrAmp) classes at 19.2%. It was to 8 (EPNaSCipTeEnrAmp) classes of antibiotics at 21.4% in chicken meat. The findings suggested the possible persistence of the resistant organisms in the farm environment even after cleaning and disinfection of the chicken houses. Flies may have played a role in introducing the resistant bacteria to the chicken farm and environment. Imprudent use of antibiotics as growth promoters or for therapeutic purposes not administered professionally could lead to development of resistance by these bacteria. However, health records were not available in the farms and information on antibiotics used could not be obtained making it impossible to make sound deduction except to make assumptions. Risk factors that were significantly ($p < 0.05$) associated with *Campylobacter* contamination in chickens meat included; no working attire (OR 2.7, CI 1.144-6.374, $p = 0.033$), fair and poor usage of protective equipment (OR 12.6, CI 1.186-133.899, $p = 0.036$) and (OR 38.50, CI 2.915-508.463, $p = 0.006$) respectively, poor stall hygiene (OR 44.00, CI 2.193-882.66, $p = 0.013$) and use of wood counter surface (OR 6.1, CI 1.198-31.164, $p = 0.029$). The relevant but not significant factor was poor working hygiene (OR 5.250, CI 0.988-27.895, $p = 0.05$). This study identified five risk factors for *Campylobacter* contamination that, if taken together, might account for most sporadic cases. The presence of high MDR *Campylobacter* and *E. coli* species could compromise treatment in humans and in particular, if the bacteria is resistant to the drugs of choice and alternative drugs for treatment and therefore poses a significant public health risk. Certainly, at farm level, the prevalence of MDR *Campylobacter* and *E. coli* in broiler flocks should be monitored. From this study it was observed that there is a need to stress the awareness among farmers to observe good hygienic practices and prudent use of antibiotics to reduce the menace of antibiotic resistance.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Sains Veterinar

KEHADIRAN *E.coli* DAN *Campylobacter* RINTANG PELBAGAI DRUG PADA AYAM DAN DAGING AYAM SERTA FAKTOR RISIKO YANG BERKAITAN

Oleh

IBRAHIM JALO MUHAMMAD

Mei 2015

Pengerusi: Professor Saleha Abdul Aziz, PhD

Fakulti: Perubatan Veterinar

Penggunaan antibiotik adalah faktor penting yang membawa kepada kemunculan dan penyebaran mikroorganisma rintang antibiotik pada kesihatan veterinar dan manusia. Ayam boleh dijangkiti *Campylobacter* dan *E. coli* yang rintang terhadap satu atau lebih antibiotik. Terdapat peningkatan yang membimbangkan terhadap kerintangan antibiotik pada patogen bakteria manusia dan haiwan secara global. Terdapat data yang terhad berkaitan kejadian *E.coli* dan *Campylobacter* rintang pelbagai drug (MDR) pada ayam dan daging ayam di Malaysia. Kajian ini dijalankan untuk mengkaji *E.coli* dan *Campylobacter* yang rintang pelbagai drug pada ayam dan daging ayam dan faktor risiko yang berkaitan. Objektif khusus adalah: untuk menentukan kehadiran *E.coli* dan *Campylobacter* pada ayam dan persekitaran ladang dan daging ayam yang dijual di pasar dan faktor risiko yang berkaitan dan untuk menentukan profil rintangan antibiotik bakteria yang diasingkan daripada ayam dan daging ayam. Dua ratus sepuluh (210) ayam daging disampel di lima ladang dan 70 daging ayam di gerai jualan ayam di tujuh pasar. Lima (5) ladang tersebut telah dilawati sebanyak tiga kali dalam tempoh kajian. Lawatan pertama adalah untuk mengambil sampel daripada ayam pada umur pasaran (35-42 hari) sebelum dijual (populasi pertama) dan persekitaran ladang, lawatan kedua adalah untuk mengambil sampel persekitaran ladang dua (2) minggu selepas ladang ayam dikosongkan (semua ayam telah dijual), dibersihkan dan dibasmi kuman dan lawatan ketiga telah dijalankan dengan mengambil sampel ke atas kumpulan baru ayam pada umur pasaran juga sebelum dijual (populasi kedua) dan kawasan sekitarnya. Dalam setiap ladang, tiga puluh (30) sampel swab kloaka ayam dan dua belas (12) sampel kawasan sekitar, terdiri daripada tiga (3) sampel setiap jenis sampel iaitu air dari bekas minuman (3), sampel makanan daripada bekas makanan (3) sampel lantai rumah ayam (3) dan sampel lalat (disatukan 5-7 lalat dalam satu botol sebagai satu sampel) (3) dikumpulkan. Daging ayam telah dipilih secara rawak yang mengandungi bahagian dada, paha dan sayap, yang disaluti kulit. Bilangan sampel daripada setiap pasar adalah berbeza iaitu daripada 4 hingga 10 gerai jualan ayam yang ada di setiap pasar bergantung kepada jumlah gerai di pasar tersebut. Papan pemotong dan alat timbang juga disampel. Lima puluh lima perpeuluhan tujuh (60.0% and 51.3%) ayam pada populasi satu dan dua dijangkiti *Campylobacter*. *Campylobacter* yang diasingkan daripada persekitaran ladang adalah 7.5% (1.7% dan 13.3%). Hanya satu sampel air yang positif *Campylobacter* (6.7%) pada lawatan pertama dan dua

(13.3%) positif pada lawatan ketiga. Lalat yang positif *Campylobacter* hanya pada lawatan ketiga (33.3%) dan hanya satu sampel lantai (6.7%) yang positif. Semua sampel makanan adalah negatif *Campylobacter*. Spesies yang paling kerap dipencilkan pada populasi pertama dan kedua adalah *C.jejuni* (74.9%), diikuti oleh *C.coli* (25.2%). Di pasar, kehadiran *Campylobacter* adalah 20.0% pada daging ayam, dan 5.7% daripada alat timbang dan papan pemotong dan spesies *Campylobacter* yang dikenalpasti adalah *C. jejuni*, *C.coli* dan *C.upsaliensis* masing-masing 78.6%, 17.9% dan 3.6%. Kehadiran *E. coli* pada ayam adalah 53.0%. Begitu juga, kehadiran *E.coli* pada sampel persekitaran pada lawatan pertama dan ketiga masing-masing 26.7% dan 40.0%, dan semua sampel adalah negatif pada lawatan kedua. Di antara sampel persekitaran, kadar pencilan pada makanan 13.3%, lantai 60.0%, air 13.3% dan lalat 73.3%. Kehadiran *E.coli* pada daging ayam di pasar adalah 45.0%. Di ladang, kehadiran *Campylobacter* dan *E. coli* pada lawatan populasi pertama dan kedua adalah hampir sama; ini mungkin kerana langkah-langkah amalan ladang adalah sama, biosekuriti yang diamalkan di ladang mungkin tidak mencukupi. Kehadiran lalat dan burung di sekitar ladang atau penggunaan air yang tercemar boleh memindahkan bakteria ini kepada ayam atau pekerja membawa bakteria tersebut ke dalam rumah ayam daripada persekitaran ladang. Di gerai jualan ayam di pasar, kontaminasi silang mungkin menjadi faktor semasa pengendalian dan memotong daging dan permukaan daging ayam boleh tercemar dengan *Campylobacter* dan *E.coli* daripada kandungan usus kerana usus terpecah atau daripada peralatan yang tercemar atau daripada air. Ketiadaan *Campylobacter* dalam sampel makanan mungkin kerana ketahanan *Campylobacter* yang lemah kepada keadaan dan tekanan persekitaran yang lain semasa penyimpanan sehingga mereka bertukar kepada bentuk *viable-but-not culturable*. Kebanyakan kelompok ayam adalah negatif sehingga berumur dua (2) minggu, dan apabila *Campylobacter* menjangkiti kelompok ayam daging, penyebaran bakteria adalah sangat cepat dan sehingga 100% daripada ayam dalam kelompok boleh dijangkiti dalam tempoh tiga (3) hari . Ujian kerintangan antibiotik telah dijalankan untuk menentukan corak rintangan antibiotik ke atas pencilan *Campylobacter* (n = 208) dan *E.coli* (n = 269) dengan menggunakan kaedah “disc diffusion” dan diuji terhadap 12 antibiotik yang berbeza. Pencilan *Campylobacter* menunjukkan rintangan yang tinggi terhadap *penicillin* adalah 75.5% (74.4% dan 76.6%) dan rintangan yang paling rendah adalah terhadap *amoxicillin_clavulanic acid* 33.7% (32.2% dan 35.1%). Pada daging ayam, *Campylobacter* menunjukkan rintangan yang tinggi terhadap *ampicillin* (62.5%) dan yang paling rendah adalah kepada *streptomycin* 3.1%. Didapati *E.coli* yang dipencil daripada ayam pada populasi pertama rintang terhadap *penicillin*, *erythromycin*, *ampicillin* dan *tetracycline* yang masing-masing pada 98.8%, 96.4%, 94.0% dan 92.8% dan rintangan yang paling rendah adalah terhadap *amoxicillin_cluvanic acid* 22.9% pada lawatan pertama. Manakala pada populasi kedua, rintangan yang paling tinggi adalah terhadap *erythromycin* dan *tetracycline* iaitu 100%, dan rintangan yang paling rendah adalah terhadap *cefotaxime* 20.0%. Bagi pencilan daripada daging ayam, rintangan yang paling tinggi adalah kepada *erythromycin* dan *penicillin* iaitu masing-masing 100% dan rintangan yang paling rendah adalah kepada *gentamicin* pada 40%. Rintangan yang tinggi mungkin terjadi hasil daripada penggunaan antibiotik yang kurang bijak untuk menggalakkan pertumbuhan dan untuk tujuan profilaksis di ladang dan dengan itu bakteria akan terbawa daripada ayam kepada daging ayam. Rintangan pelbagai drug (MDR) (rintangan kepada 3 atau lebih kelas antibiotik) adalah tinggi pada *Campylobacter* yang di pencilkan daripada ladang ayam iaitu 87.5% dan daging ayam 71.9%, manakala bagi *E.coli* adalah 99.5% dan

untuk daging ayam 100%. Rintangan pelbagai drug paling kerap diperolehi untuk *Campylobacter* yang diasingkan dari ayam adalah kepada 7 (EPNaCtxSCipAmp) dan 10 (EPNaCipTeAmcSxtCnEnrAmp) kelas antibiotik, 13.2% setiap satu pada populasi pertama dan pada populasi kedua corak MDR yang kerap adalah kepada 9 (EPNaCtxSCipAmcEnrAmp) kelas iaitu 12.9%. Pada daging ayam, MDR ialah kepada 4 (TeAmcEnrAmp) dan 5 (EPTeCtxAmp) kelas iaitu pada 15.6% setiap satu. *E.coli* pada ayam, menunjukkan MDR paling kerap kepada 9 (EPNaSCipTeAmcSxtAmp) dan 10 (EPNaSCipTeAmcSxtEnrAmp) kelas antibiotik iaitu 16.0% setiap satu dalam populasi pertama dan populasi kedua. Corak yang paling kerap adalah kepada 8 (EPNaSCipTeAmcSxtEnrAmp) kelas antibiotik iaitu pada 19.2%. Pada daging ayam MDR adalah terhadap 8 (EPNaSCipTeEnrAmp) kelas antibiotik pada 21.4%. Hasil kajian memperlihatkan kehadiran organisma rintang antibiotik dalam persekitaran yang berpanjangan walaupun selepas pembersihan ladang. Lalat juga boleh memainkan peranan dalam membawa masuk bakteria rintang antibiotik ke ladang. Penggunaan antibiotik yang tidak bijak sebagai penggalak pertumbuhan atau untuk tujuan terapeutik yang digunakan secara tidak profesional boleh membawa kepada pembentukan rintangan oleh bakteria ini. Rekod kesihatan tidak terdapat di ladang-ladang dan maklumat tentang antibiotik yang digunakan tidak dapat diperolehi menyukarkan untuk membuat sesuatu kesimpulan dan hanya dapat membuat andaian sahaja. Faktor risiko ketara ($p < 0.05$) yang berkaitan dengan pencemaran *Campylobacter* pada ayam daging termasuklah: tiada pakaian kerja (OR 2.7, CI 1,144-6,374, $p = 0.033$), kekurangan penggunaan peralatan perlindungan yang baik (OR 12.6, CI 1,186-133,899, $p = 0.036$) dan (OR 38.50, CI 2,915-508,463, $p = 0.006$), kandang yang kurang basih (OR 44.00, CI 2,193-882,66, $p = 0.013$) dan penggunaan permukaan pemotong daripada kayu (OR 6.1, CI 1,198-31,164, $p = 0.029$). Faktor yang berkaitan tetapi tidak ketara adalah kerja yang kurang bersih (OR 5.250, CI 0,988-27,895, $p = 0.05$). Kajian ini mengenal pasti lima (5) faktor risiko pencemaran oleh *Campylobacter* dan jika diambil kira semuanya, mungkin boleh menerangkan sebab bagi kebanyakan kes-kes sporadik. Kehadiran MDR *Campylobacter* dan *E.coli* yang tinggi boleh menjejaskan rawatan pada manusia terutamanya jika antibiotik diperlukan dan jika organisma tersebut rintang kepada ubat dan ubat alternatif untuk rawatan dan dengan itu boleh menimbulkan risiko kesihatan yang ketara. Sudah tentu, di peringkat ladang, kehadiran *Campylobacter* dan *E.coli* pada ayam daging perlu dipantau untuk memastikan pelaksanaan langkah-langkah amalan penternakan, veterinar dan biosekuriti yang baik. Daripada kajian ini juga didapati bahawa terdapat keperluan untuk mewujudkan kesedaran dalam kalangan penternak untuk mematuhi amalan kebersihan yang baik dan penggunaan antibiotik secara berhemah untuk mengurangkan ancaman rintangan antibiotik.

ACKNOWLEDGEMENTS

All praises be to Allah (SWT), the creator, nourisher, cherisher, sustainer and provider of one and all who bestowed the ability in me making my dreams a reality. I would like to express my sincere appreciation and profound gratitude to the Chairman of my supervisory committee Prof. Dr. Saleha Abdul Aziz for her unwavering support, scholarly criticisms throughout the research and the program as a whole. Her thorough scrutiny and suggestions made this reality. I am grateful and indebted to supervisory committee members, Assoc. Prof. Dr. Jalila Abu and Assoc. Prof. Dr. Siti Khairani-Bejo for their valuable suggestions throughout this study. I also thank Hj. Abd Razak bin Jaafar for his valuable assistance during my sampling.

Many thanks to Puan Fauziah Nordin of Veterinary Public Health Lab, Krishnama and all staff of Bacteriology Lab of the Faculty of Veterinary Medicine for their kind support, assistance and cooperation throughout my laboratory work. I am indeed very grateful to my lab mates Dauda Goni, Rasheed, Teguh, Yousif, Wint Wint, Emelia, Abdelrahman, Yusuf Yaqoub and Shah for their friendly support and cooperation which I cherish a lot. I am very thankful to my friends and house mates Alhassan, MD Usman, Abdulaziz Ibrahim, Sadiq Nesta, Dr. Konto, Sani, Abdullahi Adamu, Mohammed Ngab, Syafiqah, Zubaida, Abdullahi Adamu and many others whom space would not permit me to mention. All are good friends especially Lawan Lawu, Butali, Ibrahim Ayya, Samba, Daddy, La'ari and supportive brothers and sisters such as Idi, Dr. Idiris, Nura, Sadiq, Usman and Umami, Mami, Hadiza, Larabiyyo. My special gratitude and thanks go to rest of my family, relatives and all well wishers for their prayers and support.

I certify that a Thesis Examination Committee has met on 11 May 2015 to conduct the final examination of Ibrahim Jalo Muhammad on his thesis entitled "Occurrence of Multidrug Resistant *E. coli* and *Campylobacter* in Chicken and Chicken Meat and their Associated Risk Factors" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Abdul Aziz Saharee, PhD
Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Zunita Zakaria, PhD
Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Sharifah Syed Hassan, PhD
Associate Professor
Monash University Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 22 September 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory committee were as follows:

Saleha Abdul Aziz, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Jalila Abu, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Siti Khairani-Bejo, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of graduate studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work
- quotations, illustrations and citations have been duly referenced
- the thesis has not been submitted previously or concurrently for any other degree at any institutions
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be owned from supervisor and deputy vice –chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Ibrahim, Jalo Muhammad GS35558

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iv
ACKNOWLEDGEMENTS		vii
APPROVAL		viii
DECLARATION		x
LIST OF TABLES		xiv
LIST OF FIGURES		xv
LIST OF ABBREVIATIONS		xvi
CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	3
2.1	The genus <i>Campylobacter</i>	3
2.1.1	History, taxonomy and microbiological overview of <i>Campylobacter</i>	3
2.1.2	Epidemiology of <i>Campylobacter</i>	5
2.1.3	Physical and Biochemical Properties of <i>Campylobacter</i>	6
2.1.4	Sources of infection and mode of transmission of <i>Campylobacter</i> in human.	6
2.1.5	<i>Campylobacter</i> in poultry and other animals	8
2.1.6	Risk factors associated with colonization of <i>Campylobacter</i> in chicken	9
2.1.7	Occurrence of <i>Campylobacter</i> species in Malaysia	11
2.2	The genus <i>Escherichia coli</i>	11
2.2.1	History, taxonomy and microbiology of <i>E.coli</i>	11
2.2.2	Commensal <i>E.coli</i>	12
2.2.3	Pathogenic <i>E.coli</i>	12
2.2.4	Public health significance of <i>E.coli</i>	13
2.2.5	Occurrence of <i>E.coli</i> in Malaysia	13
2.3	Antibiotic Resistance	14
2.3.1	Public Health significance of antibiotic resistance	15
2.3.2	Antibiotic resistance in <i>Campylobacter</i>	16
2.3.3	Antibiotic resistance in <i>E.coli</i>	17
2.3.4	Multidrug Resistance (MDR)	18
2.3.5	Surveillance and monitoring of antibiotic resistance	18
3	OCCURRENCE OF <i>CAMPYLOBACTER</i> IN CHICKEN, FARM ENVIRONMENTS AND CHICKEN MEAT AT WET MARKETS	20
3.1	Introduction	20
3.2	Materials and Methods	21
3.2.1	Collection of samples	25
3.2.2	Isolation of <i>Campylobacter</i> species	26
3.2.3	Phenotypic identification of <i>Campylobacter</i>	27
3.2.4	Confirmation of <i>Campylobacter</i> isolates by Multiplex Polymerase Chain Reaction (m-PCR) assay	28
3.2.5	Agarose Gel Electrophoresis	29
3.3	Statistical Analysis	30
3.4	Results	30

3.5	Discussion	39
3.6	Conclusion	42
4	OCCURRENCE OF <i>E.COLI</i> IN CHICKEN, FARM ENVIRONMENT AND CHICKEN MEAT IN WET MARKETS	43
4.1	Introduction	43
4.2	Materials and Methods	44
4.2.1	Collection of samples	44
4.2.2	Isolation of <i>E.coli</i> species	44
4.2.3	Phenotypic identification of <i>E.coli</i>	45
4.2.4	Confirmation of <i>E.coli</i> isolates by Polymerase Chain Reaction (PCR) assay	45
4.2.5	Agarose Gel Electrophoresis	46
4.2.6	Statistical Analysis	46
4.3	Results	46
4.4	Discussion	51
4.5	Conclusion	53
5	ANTIBIOTIC RESISTANT PROFILES OF <i>CAMPYLOBACTER</i> AND <i>E.COLI</i> ISOLATES	54
5.1	Introduction	54
5.2	Materials and Methods	55
5.2.1	Bacterial isolates and growth condition	55
5.2.2	Antibiotic Susceptibility Testing	55
5.2.3	Data Analysis	57
5.3	Results	57
5.3.1	Multidrug resistance (MDR)	72
5.3.2	Antibiotic resistance pattern of <i>Campylobacter</i> and <i>E.coli</i>	74
5.4	Discussion	84
5.5	Conclusion	86
6	RISK FACTORS ASSOCIATED WITH THE OCCURRENCE OF <i>CAMPYLOBACTER</i> IN CHICKEN AND CHICKEN MEAT	87
6.1	Introduction	87
6.2	Materials and Method	88
6.2.1	Data Analysis	89
6.3	Results	89
6.4	Discussion	93
6.5	Conclusion	95
7	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCH	96
7.1	Summary and general conclusion	96
7.2	Recommendations for future research	98
	REFERENCES	99
	APPENDICES	136
	BIODATA OF STUDENT	155
	LIST OF PUBLICATIONS AND CONFERENCES	156

LIST OF TABLES

Table	Page
3.1 Location of broiler chicken farms and number of samples collected	21
3.2 Location of wet markets and number of samples collected	24
3.3 Number of samples taken and the period of farm visits	25
3.4 Biochemical tests for identification of <i>Campylobacter</i> species (Lastovica & Allos, 2008)	28
3.5 Primers for the amplification of <i>Campylobacter</i> genes (Yamazaki-Matsune et al (2007)	29
3.6 Description and management system of the farms	30
3.7 Occurrence of <i>Campylobacter</i> in broiler chickens in five (5) farms	33
3.8 Occurrence of <i>Campylobacter</i> species in five (5) broiler chicken farms	35
3.9 Occurrence of <i>Campylobacter</i> in chicken house environment in five different farms	37
3.10 Occurrence of <i>Campylobacter</i> isolated from chicken meat in wet markets	38
3.11 Occurrence of <i>Campylobacter</i> isolates from cutting boards and weighing scales	39
4.1 Primers for <i>E.coli</i> confirmation (Wang et al., 2002)	46
4.2 Occurrence of <i>E.coli</i> isolates from broiler chicken in five (5) farms	49
4.3 Proportion of <i>E.coli</i> isolated from chicken house environment in five farms	50
4.4 Proportion of <i>E.coli</i> isolated from chicken meat, cutting board and weighing scales in wet markets	51
5.1 Breakpoint of the disc diffusion method to determine antimicrobia susceptibility of <i>Campylobacter</i> and <i>E.coli</i> isolates	56
5.2 Antibiogram of <i>Campylobacter</i> isolates from broiler chicken and farm environment	75
5.3 Antibiogram of <i>Campylobacter</i> isolates from chicken meat, weighing scales and cutting boards	78
5.4 Antibiogram of <i>E.coli</i> isolates from broiler chicken and farm environment	79
5.5 Antibiogram of <i>E.coli</i> isolates from broiler chicken meat, cutting boards and weighing scales	82
6.1 Definition and description scores of exposure variables of meat handling practices associated with <i>Campylobacter</i> in selected markets	90
6.2 Univariate logistic regression for risk factors associated with <i>Campylobacter</i> isolates in markets	92
6.3 Fisher's exact tests for risk factors associated with the occurrence of <i>Campylobacter</i> isolated from chicken and farm environments	93

LIST OF FIGURES

Figure	Page
3.1 Location of broiler farms sampled	22
3.2 Farm environment two (2) weeks after the chicken houses were emptied (all chickens were sold), cleaned and disinfected	22
3.3 Chickens at market age (35-42 days) prior to harvesting and were raised under closed house intensive management system. Chicken feed and water are given automatically	23
3.4 Location of wet markets sampled	24
3.5 Colonies of <i>Campylobacter</i> on mCCDA	31
3.6 Cellular morphology of <i>Campylobacter</i> species are gram negative, spiral, curve or rod shaped	31
3.7 Colonies of <i>Campylobacter</i> on CBA	32
3.8 Representative m-PCR assay to confirm <i>Campylobacter</i> isolates from chicken and farm environment	32
3.9 Representative mPCR assay for confirmation of <i>Campylobacter</i> species isolated from chicken meat, weighing scales and cutting boards	33
4.1 <i>E.coli</i> colonies on Chromocult Chromogenic agar plates	46
4.2 Colonies of <i>E.coli</i> on EMB agar plates	47
4.3 Representative PCR assay for confirmation of <i>E.coli</i> species isolated from chicken and farm environments.	48
4.4 Representative PCR assay for confirmation of <i>E.coli</i> species isolated from chicken meat, weighing scales and cutting boards	48
5.1 Percentage of <i>Campylobacter</i> isolates in chicken	58
5.2 Percentage of <i>Campylobacter</i> isolates from water	60
5.3 Percentage of <i>Campylobacter</i> isolates from flies	61
5.4 Percentage of <i>Campylobacter</i> isolates from chicken meat, weighing scales and cutting board	62
5.5 Prevalence of <i>E.coli</i> isolates from chicken	64
5.6 Prevalence of <i>E.coli</i> isolates from water	66
5.7 Occurrence of <i>E.coli</i> isolates from flies	67
5.8 Percentage of <i>E.coli</i> isolated from feed	69
5.9 Prevalence of <i>E.coli</i> isolates from chicken house floor	70
5.10 Percentage of <i>E.coli</i> isolates from meat, weighing scale and cutting board	71
5.11 Percentage of MDR <i>Campylobacter</i> in chicken (first population) and farm environment (first visits)	72
5.12 Percentage of MDR <i>Campylobacter</i> in chicken (second population) and farm environment (third visits)	73
5.13 Percentage of MDR <i>Campylobacter</i> in chicken meat, weighing scales and cutting boards	73

LIST OF ABBREVIATIONS

ATCC	American Type Culture Collection
bp	Base pairs
CCUG	Culture Collection of the University of Goteborg
cueE	Siderophoreenterochelin
CLSI	Clinical Laboratory Standard Institute
°C	Degree celcius
DNA	Deoxyribonucleic acid
EDTA	Ethylenediaminetetraacetic
flaA	Flagellin A gene
glyA	Serine hydroxyl methyl transferase gene
g	Gram(s)
h	Hour(s)
hip	Hippuricase gene
Kb	Kilobase
ml	Mililitre
mg	Milligram
mg	Milligram(s)
min	Minute(s)
MDR	Multidrug resistance
mPCR	Multiplex Polymerase Chain Reaction
NaCl	Sodium chloride
NCCLS	National Committee for Clinical Laboratory Standards
PFGE	Pulsed Field Gel Electrophoresis
PBS	Phosphate buffered saline
PCR	Polymerase Chain Reaction
RNA	Ribonucleic acid

rRNA	Ribosomal RNA
s	Second(s)
Spp.	Species
TAE	Tris-acetate-EDTA
TBE	Tris-borate EDTA
UV	Ultraviolet
V	Volt
Taq	Thermophilus aquaticus
WHO	World Health Organization
μL	Micro liter
μg	Micro Gram
μM	Micro molar
EMB	Eosin Methylene Blue
SIM	Sulfide-Indole-Motility medium
MR-VP	Methyl Red- Voges-Proskauer
BPW	Buffered Peptone Water
NA	Nutrients Agar
\$	Chi-square Tests
<i>P</i>	Probability

CHAPTER 1

INTRODUCTION

In livestock production, the use of antibiotics for disease treatment and prevention is important. It is recognised that the use of these agents in food animals creates the development of antibiotic resistant bacteria, which disseminate via the food chain. Globally, the increase and spread of antibiotic resistance especially to foodborne zoonotic bacteria with its reservoirs in healthy food animals such as poultry, pigs and cattle has become public health concern (EFSA, 2008a). In developed countries, there is increasing scientific reports regarding the widespread of antibiotic usage in food animal production that leads to the development of resistant pathogenic organisms in the food chain (Marshall & Levy, 2011; Philips et al., 2004).

The resistance elements reduce the efficiency of antibiotic therapy, which results in an increased morbidity and mortality associated with disease outbreaks (Da Costa et al., 2009). It has been reported by Schwarz and Chaslus-Dancla (2001) that an antibiotic therapy with a specific agent has been either accompanied or followed shortly by the occurrence of resistant bacteria, and for the past ten years the effect of antibiotics on the development of resistance have gained much attention (Smith et al., 2007). In developed and developing countries, *Campylobacter* species are among the leading causes of zoonotic infections (EFSA, 2013; Anonymous, 2008) that the prevalence rate is increasing. In Europe, campylobacteriosis is the most reported zoonotic enteric disease surpassing salmonellosis (EFSA 2012; Fosse et al., 2007).

Poultry meat is regarded as the major source of campylobacteriosis in human, apart from chicken; other animals such as pigs and cattle also serve as reservoir hosts (Stanley & Jones, 2003). *Campylobacter* species is also found in various other domestic and wild animals, including goats, horses, cats, rodents and dogs (Man, 2011); they are also isolated from marine animals such as shellfish and dolphins. Poultry is considered as the main reservoir of *Campylobacter*. Among *Campylobacter*, *Campylobacter jejuni* and *Campylobacter coli* are the frequently cause of foodborne diseases in human (Uaboi-Egbenni et al., 2012). The risk to *Campylobacter* infection in human include consumption of undercooked poultry meat, handling of raw poultry carcasses, drinking untreated water, drinking unpasteurized milk or ingestion of dairy products made from raw unpasteurized milk and international travel (Danis et al., 2009). The majority of the cases occur intermittently. The consumption of poultry meat, contact with infected animals, drinking contaminated water or travel are the highest risk associated with the infection (Everest and Ketley, 2002). In poultry, the occurrence of *Campylobacter* varies according to age and the type of farm; they are seldom detected in broiler chicken less than 2-3 weeks of age and in those managed under close house system (Sahin et al., 2002).

Campylobacter infection is normally self-limiting, but it may be associated with complications such as Guillain-Barre Syndrome (neurological) and Reiter's Syndrome (reactive arthritis) (Yan et al., 2005). Once a broiler chicken becomes infected, *Campylobacter* spread rapidly to other broiler chickens in that flock, and up to slaughter age, or at thinning the chickens remain colonized. Almost all (100%) of

the broiler chickens brought to the slaughter houses were reported to be colonized with *Campylobacter* (Jacobs-Reitsma et al., 2008), and the contaminated chicken meat acts as a probable risk of human campylobacteriosis. Detection of *C. jejuni*, *C. coli* and *E. coli* on the carcasses is mainly due to contamination from the gastrointestinal contents of slaughtered healthy animals during processing as well as at retail (Nonga et al., 2010). *Campylobacter* has reported to develop resistance to a number of antibiotics including ciprofloxacin and other fluoroquinolones, macrolides and lincosamides, chloramphenicols, aminoglycosides, tetracyclines, ampicillins and other β -lactams, cotrimoxazole, and tylosin (Moore et al., 2006; Padungton and Kaneene, 2003).

Escherichia coli is part of the normal enteric microbial flora in human, poultry and other animals and the pathogenic *E. coli* causes disease in both (Amin et al., 2012). Pathogenic *E. coli* cause a number of diseases in both poultry such as colibacillosis, aerosacculitis, polyserositis, septicemia and other mainly extraintestinal diseases. In humans diseases caused include hemorrhagic colitis, haemolytic uremic syndrome, acute and chronic endemic and epidemic diarrhoea. They are transmitted from person to person via direct contact with animal carriers, faeces, contaminated soil and water or via ingestion undercooked meat and other animal products as well as contaminated vegetables and fruits. As a result of contamination from faeces, it is often found in soil, water and food. Commensal *E. coli* flora can be regarded as a rich source of emergence and spreading of antibiotic resistance (Da Costa et al., 2013). The chicken may also be infected with antibiotic resistance *E. coli* or may develop resistant to one or more antibiotics in the gut.

There is an alarming increase of antibiotic resistant bacteria in particular on MDR among human and animal population globally (Carlet et al., 2012; Marshal and Levy 2011).

However, limited data exists on the occurrence of multidrug resistant (MDR) *E. coli* and *Campylobacter* in poultry and poultry meat in Malaysia.

The hypothesis of this study was: There is a high occurrence of MDR *E. coli* and *Campylobacter* in broiler chicken (in farms) prior to harvesting and on chicken meat (retailed in markets).

Therefore, the objectives of this study were:

1. to determine the occurrence of *E. coli* and *Campylobacter* in chicken in the farm prior to harvesting and chicken meat retailed in the markets.
2. to determine the multidrug resistance (MDR) and antibiotic resistant profiles among the isolates.
3. to determine the associated risk factors in the occurrence of multidrug resistant (MDR) *Campylobacter* in chickens.

REFERENCES

- Aarestrup FM. (2004) Monitoring of antimicrobial resistance among food animals: principles and limitations. *Journal of Veterinary Medicine B*; 51: 380–388.
- Aarestrup, F. M., & Wegener, H. C. (1999). The effects of antibiotic usage in food animals on the development of antimicrobial resistance of importance for humans in *Campylobacter* and *Escherichia coli*. *Microbes and Infection*, 1(8), 639-644.
- Abulreesh, H. H., Paget, T.A., & Goulder, R. (2005). Recovery of thermophilic *Campylobacter* from pond water and sediment and the problem of interference by background bacteria in enrichment culture. *Water Research*, 39(13), 2877-82.
- Achen, M., Morishita, T. Y., & Ley, E. C. (1998). Shedding and colonization of *Campylobacter jejuni* in broilers from day-of-hatch to slaughter age. *Avian Diseases*, 732-737.
- Acke, E., McGill, K., Golden, O., Jones, B. R., Fanning, S., & Whyte, P. (2009). A comparison of different culture methods for the recovery of *Campylobacter* species from pets. *Zoonoses and Public Health*, 56(9-10), 490-495.
- ACMSF, (2004). Second Report on *Campylobacter*. London: Advisory Committee on the Microbiological Safety of Food.
- Adak, G. K., Long, S. M., & O'Brien, S. J. (2002). Trends in indigenous foodborne disease and deaths, England and Wales: 1992 to 2000. *Gut*, 51(6), 832-841.
- Adesiji, Y. O., Alli, O. T., Adekanle, M. A., & Jolayemi, J. B. (2011). Prevalence of *Arcobacter*, *Escherichia coli*, *Staphylococcus aureus* and *Salmonella* species in retail raw chicken, pork, beef and goat meat in Osogbo, Nigeria. *Sierra Leone Journal of Biomedical Research*, 3(1), 8-12.
- Adetunji, V. O. and Odetokun, I. A. (2011). Bacterial hazards and critical control points in goat processing at a typical tropical abattoir in Ibadan, Nigeria. *International Journal of Animal and Veterinary Advances* 3, 349-354
- Adhikari, B., Connolly, J., Madie, P., & Davies, P. (2004). Prevalence and clonal diversity of *Campylobacter jejuni* from dairy farms and urban sources. *New Zealand Veterinary Journal*, 52(6), 378–383.
- Adkin, A., E. Hartnett, L. Jordan, D. Newell, and H. Davison. (2006). Use of a systematic review to assist the development of *Campylobacter* control strategies in broilers. *Journal of Applied Microbiology*. 100:306–315
- Adzitey, F., Huda, N., & Rahmat Ali, G. R. (2012). Prevalence and antibiotic resistance of *Campylobacter*, *Salmonella*, and *L. monocytogenes* in ducks: A review. *Foodborne Pathogens and Disease*, 9(6), 498-505.

- Adzitey, F., N. Huda and G. Rusul, (2010b). Prevalence of *Escherichia coli* in duck intestines, faeces, soil and wash water samples in Penang, Malaysia. Proceedings of the 4th International Conference Postgraduate Education: Globalisation and Liberalisation of Postgraduate Education, Nov. 26-28, Cititel Mid Valley, Kuala Lumpur, pp: 165-167
- Ahmed, M. O., Clegg, P. D., Williams, N. J., Baptiste, K. E., & Bennett, M. (2010). Antimicrobial resistance in equine faecal *Escherichia coli* isolates from North West England. *Annals of Clinical Microbiology and Antimicrobials*, 9(1), 12.
- Akinbowale, O. L., Peng, H., & Barton, M. D. (2006). Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *Journal of Applied Microbiology*, 100(5), 1103-1113.
- Akwar, T. H., Poppe, C., Wilson, J., Reid-Smith, R. J., Dyck, M., Waddington, J & McEwen, S. A. (2007). Risk factors for Antimicrobial resistance among fecal *Escherichia coli* from residents on forty-three swine farms. *Microbial Drug Resistance*, 13(1), 69-76.
- Alhaj, N., Mariana, N. S., Raha, A. R., & Ishak, Z. (2007). Prevalence of antibiotic resistance among *Escherichia coli* from different sources in Malaysia. *International Journal of Poultry Science*, 6(4), 293-297.
- Allen, V. M., Bull, S. A., Corry, J. E. L., Domingue, G., Jørgensen, F., Frost, J. A., & Humphrey, T. J. (2007). *Campylobacter* spp. contamination of chicken carcasses during processing in relation to flock colonisation. *International Journal of Food Microbiology*, 113(1), 54-61.
- Allerberger, F., Al-Jazrawi, N., Kreidl, P., Dierich, M. P., Feierl, G., Hein, I., & Wagner, M. (2003). Barbecued chicken causing a multi-state outbreak of *Campylobacter jejuni* enteritis. *Infection*, 31(1), 19-23.
- Allos, B. M. (2001). *Campylobacter jejuni* Infections: update on emerging issues and trends. *Clinical infectious diseases : An official publication of the Infectious Diseases Society of America*, 32(8), 1201–6.
- Allos, B. M., & Blaser, M. J. (1995). *Campylobacter jejuni* and the expanding spectrum of related infections. *Clinical Infectious Diseases*, 1092-1099.
- Allwood, P. B., Jenkins, T., Paulus, C., Johnson, L., & Hedberg, C. W. (2004). Hand washing compliance among retail food establishment workers in Minnesota. *Journal of Food Protection*, 67(12), 2825-2828.
- Amin, A., Ayaz Khan, M., Haroon, U., Azam, S. M. F., Andaleeb, S., Ghumro, P. B., (2012). Determination of Antibiotic Resistance Prototype and in vitro R Plasmid Conjugation of *Escherichia coli* Isolates from Broiler Feces., *Thai Journal Veterinary Medicine*, (42) 87-93.

- Anderson, R. C., Carr, M. A., Miller, R. K., King, D. A., Carstens, G. E., Genovese, K. J & Nisbet, D. J. (2005). Effects of experimental chlorate preparations as feed and water supplements on *Escherichia coli* colonization and contamination of beef cattle and carcasses. *Food Microbiology*, 22(5), 439-447.
- Andersson, D. I., & Hughes, D. (2010). Antibiotic resistance and its cost: is it possible to reverse resistance. *Nature Reviews Microbiology*, 8(4), 260-271.
- Angulo, F. J., Nargund, V. N., & Chiller, T. C. (2004). Evidence of an association between use of anti-microbial agents in food animals and anti-microbial resistance among bacteria isolated from humans and the human health consequences of such resistance. *Journal of Veterinary Medicine, Series B*, 51(8-9), 374-379.
- Anonymous (2008). Bovine Genital *Campylobacteriosis*. OIE Terrestrial Manual 2008 Chapter 2.4.5. pp.661-670. OIE
- Anonymous (2010): Analysis of the baseline survey on prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on broiler carcasses, in the EU 2008-Part A: *Campylobacter* and *Salmonella* prevalence estimates. *EFSA Journal* 8, 1503.
- Anonymous (2012a): The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2010. *EFSA Journal* 10, 2597.
- Apun, K., Chang, P. P., Sim, E. U. H., & Micky, V. (2006). Clonal diversity of *Escherichia coli* isolates from marketed beef in East Malaysia. *World Journal of Microbiology and Biotechnology*, 22(7), 661-667.
- Apun, K., K.L. Kho, Y.L. Chong, F.H. Hashimatul and M.T. Abdullah et al. (2010). Detection of *Escherichia coli* O157:H7 in wildlife from distributed habitats in Sarawak, Malaysia. *Research Journal of Microbiology* 6:132-139
- Arsenault, J., Letellier, A., Quessy, S., Normand, V., & Boulianne, M. (2007). Prevalence and risk factors for *Salmonella* spp and *Campylobacter* spp. caecal colonization in broiler chicken and turkey flocks slaughtered in Quebec, Canada. *Preventive Veterinary Medicine*, 81(4), 250-264.
- Arslan, H., Azap, Ö. K., Ergönül, Ö., & Timurkaynak, F. (2005). Risk factors for ciprofloxacin resistance among *Escherichia coli* strains isolated from community-acquired urinary tract infections in Turkey. *Journal of Antimicrobial Chemotherapy*, 56(5), 914-918.
- Arumugaswamy, R. K., Rusul, G., Hamid, S. A., & Cheah, C. T. (1995). Prevalence of *Salmonella* in raw and cooked foods in Malaysia. *Food Microbiology*, 12, 3-8.

- Atanassova, V., & Ring, C. (1999). Prevalence of *Campylobacter* spp. in poultry and poultry meat in Germany. *International Journal of Food Microbiology*, 51(2), 187-190.
- Aubry-Damon, H., Grenet, K., Sall-Ndiaye, P., Che, D., Cordeiro, E., Bougnoux, M. E., & Andremont, A. (2004). Antimicrobial resistance in commensal flora of pig farmers. *Emerging Infectious Diseases*, 10(5), 873-879.
- Aydin, F., Atabay, H. I., & Akan, M. (2001). The isolation and characterization of *Campylobacter jejuni* subsp. *jejuni* from domestic geese (Anser anser). *Journal of Applied Microbiology*, 90(4), 637-642.
- Bae, W., Kaya, K. N., Hancock, D. D., Call, D. R., Park, Y. H., & Besser, T. E. (2008). Prevalence and antimicrobial resistance of thermophilic *Campylobacter* spp. from cattle farms in Washington State. *Applied and environmental microbiology*, 71(1), 169-174.
- Barnes, H. J., J. Vaillancourt, and W. B. Gross. (2003). Colibacillosis in Diseases of Poultry Eleventh Edition. Eds. Y.M. Saif, H.J. Barnes, J.R. Glisson, A.M. Fadly, L.R. McDougald, and D.E. Swayne. pgs. 631-656.
- Barnes, J.H., Vaillancourt, J. and Gross, W.B. (2008). Colibacillosis. In: Saif, Y.M., Barnes, H.J., Glisson, J.R., Fadly, A.M. and McDougald, L.R. (Ed.). Diseases of poultry (12th ed.) *Iowa State University Press*, IA. pp: 691-738.
- Berge ACB, Atwill ER, Sischo W.M. (2003). Assessing antibiotic resistance in fecal *Escherichia coli* in young calves using cluster analysis techniques. *Preventive Veterinary Medicine*; 61: 91-102.
- Bergsma, N. J., Fischer, A. R., Van Asselt, E. D., Zwietering, M. H., & De Jong, A. E. (2007). Consumer food preparation and its implication for survival of *Campylobacter jejuni* on chicken. *British Food Journal*, 109(7), 548-561.
- Berndtson, E., Danielsson-Tham, M. L., & Engvall, A. (1996). *Campylobacter* incidence on a chicken farm and the spread of *Campylobacter* during the slaughter process. *International Journal of Food Microbiology*, 32(1), 35-47.
- Berndtson, E., U. Emanuelson, A. Engvall, and M. L. Danielsson-Tham. (1996). A 1-year epidemiological study of *Campylobacters* in 18 Swedish chicken farms. *Preventive Veterinary Medicine*. 26:167-185.
- Berrang, M. E., Dickens, J. A., & Musgrove, M. T. (2000). Effects of hot water application after defeathering on the levels of *Campylobacter*, coliform bacteria, and *Escherichia coli* on broiler carcasses. *Poultry Science*, 79(11), 1689-1693.
- Berrang, M. E., Northcutt, J. K., & Cason, J. A. (2004). Recovery of *Campylobacter* from broiler feces during extended storage of transport cages. *Poultry Science*, 83(7), 1213-1217.

- Bester, L. A., and Essack, S. Y. (2008). Prevalence of antibiotic resistance *Campylobacter* isolates from commercial poultry suppliers in Kwa Zulu-Natal South Africa. *Journal of Antimicrobial Chemotherapy* 62:1298-1300.
- Black, R. E., Levine, M. M., Clements, M. L., Hughes, T. P., & Blaser, M. J. (1988). Experimental *Campylobacter jejuni* infection in humans. *Journal of Infectious Diseases*, 157(3), 472-479.
- Blaser, M. J. (1997). Epidemiologic and clinical features of *Campylobacter jejuni* infections. *Journal of Infectious Diseases*, 176 (Supplement 2), S103-S105.
- Blaser, M. J., Hardesty, H. L., Powers, B., & Wang, W. L. (1980). Survival of *Campylobacter fetus* subsp. *jejuni* in biological milieus. *Journal of Clinical Microbiology*, 11(4), 309–313.
- Blaser, M. J., Taylor, D. N., & Feldman, R. A. (1983). Epidemiology of *Campylobacter jejuni* infections. *Epidemiologic Reviews*, 5(1), 157-176.
- Boes, J., Nersting, L., Nielsen, E. M., Kranker, S., Enøe, C., Wachmann, H. C., & Baggesen, D. L. (2005). Prevalence and diversity of *Campylobacter jejuni* in pig herds on farms with and without cattle or poultry. *Journal of Food Protection*, 68(4), 722-727.
- Bolton, F. J., Coates, D., Hutchinson, D. N., & Godfree, A. F. (1987). A study of thermophilic *Campylobacters* in a river system. *Journal of Applied Bacteriology*, 62(2), 167-176.
- Bouki, C. (2010). Environmental transmission routes of *Campylobacter*: filling the gap of unexplained human campylobacteriosis. *MSc Thesis at Universiteit Utrecht*.
- Brieseman, M. A. (1990). A further study of the epidemiology of *Campylobacter jejuni* infections. *The New Zealand medical journal*, 103(889), 207-209.
- Brouwer, R., Mertens, M. J. A., Siem, T. H., & Katchaki, J. (1979). An explosive outbreak of *Campylobacter* enteritis in soldiers. *Antonie van Leeuwenhoek*, 45(3), 517-519.
- Bruinsma N, Hutchinson JM, van den Bogaard AE, Giamarellou H, Degener J, Stobberingh EE (2005). Influence of population density on antibiotic resistance. *Journal of Antimicrobial Chemotherapy*; 51:385–90.
- Bruinsma, N.; Filius, PM.; van den Bogaard, AE.; Nys, S.; Degener, J.; Endtz, HP.; Stobberingh, EE. (2003). Hospitalization, a risk factor for antibiotic-resistant *Escherichia coli* in the community? *Journal of Antimicrobial Chemotherapy*, 51(4):1029-1032.
- Buhr, R. J., N. A. Cox, N. J. Stern, M. T. Musgrove, J. L. Wilson, and K. L. Hiatt. (2002). Recovery of *Campylobacter* from segments of the reproductive tract of broiler breeder hens. *Avian Diseases*. 46:919–924.

- Bull S.A. et al., (2006). Sources of *Campylobacter* spp. colonizing housed broiler flocks during rearing. *Applied Environmental and Microbiology Journal.*, 72(1), 645-652.
- Bunner, CA.; Norby, B.; Bartlett, PC; Erskine, RJ.; Downes, FP.; Kaneene, JB. (2007). Prevalence and pattern of antimicrobial susceptibility in *Escherichia coli* isolated from pigs reared under antimicrobial-free and conventional production methods. *Journal of the American Veterinary Medical Association* 231(2):275-283.
- Butzler, J. P. (2004). *Campylobacter*, from obscurity to celebrity. *Clinical microbiology and infection*, 10(10), 868-876.
- Butzler, J. P., M. De Boeck, H. Goosens. (1983). A new selective medium for isolation of *Campylobacter jejuni* from human faeces. *European Journal of Clinicals*. 2: 389-394.
- Byappanahalli, M., Fowler, M., Shively, D., & Whitman, R. (2003). Ubiquity and persistence of *Escherichia coli* in a midwestern coastal stream. *Applied and Environmental Microbiology*, 69(8), 4549-4555.
- Bywater, R., Deluyker, H., Deroover, E., De Jong, A., Marion, H., McConville, M. & Walters, J. (2004). A European survey of antimicrobial susceptibility among zoonotic and commensal bacteria isolated from food-producing animals. *Journal of Antimicrobial Chemotherapy*, 54(4), 744-754.
- Caballero, A.T., Carrera, J.A.V., Lengomin, M.E.F. (1998). Evaluacion de la vigilancia microbiologica de alimentos que se venden en las calles. *Revista Cubana de Alimentacion y de Nutricion* 12, 7 – 10
- Camarda, A., D. G. Newell, R. Nasti, and G. Di Modugnoa. (2000). Genotyping *Campylobacter jejuni* strains isolated from the gut and oviduct of laying hens. *Avian Diseases*. 44:907–912.
- Carlet, J., Jarlier, V., Harbarth, S., Voss, A., Goossens, H., & Pittet, D. (2012). Ready for a world without antibiotics? The penières antibiotic resistance call to action. *Antimicrobial resistance and infection control*, 1(1), 1-13.
- Cawthraw, S. A., L. Lind, B. Kaijser, and D. G. Newell. (2000). Antibodies, directed towards *Campylobacter jejuni* antigens, in sera from poultry abattoir workers. *Clinical Experimental Immunology*. 122:55–60.
- Cawthraw, S. A., Wassenaar, T. M., Ayling, R., & Newell, D. G. (1996). Increased colonization potential of *Campylobacter jejuni* strain 81116 after passage through chickens and its implication on the rate of transmission within flocks. *Epidemiology and Infection*, 117(01), 213-215.
- Centers for Disease Control and Prevention. (CDC) (2010). *Campylobacter*. Available from: <http://www.cdc.gov/nczved/divisions/dfbmd/diseases/campylobacter/>

- Centre for Disease Control and Prevention (CDC) (The US Centers for Disease Control and Prevention) (2005). *Campylobacter* infection. Available from www.cdc.gov/ncidod/dbmd/diseaseinfo/campylobacter_g.htm.
- Chaban, B., Ngeleka, M., & Hill, J. E. (2010). Detection and quantification of 14 *Campylobacter* species in pet dogs reveals an increase in species richness in feces of diarrheic animals. *BMC Microbiology*, *10*(1), 73.
- Chai, L. C., Robin, T., Ragavan, U. M., Gunsalam, J. W., Bakar, F. A., Ghazali, F. M., & Kumar, M. P. (2007). Thermophilic *Campylobacter* spp. in salad vegetables in Malaysia. *International Journal of Food Microbiology*, *117*(1), 106–111.
- Châtre, P., Haenni, M., Meunier, D., Botrel, M. A., Calavas, D., & Madec, J. Y. (2010). Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from cattle between 2002 and 2006 in France. *Journal of Food Protection*, *73*(5), 825-831.
- Chauvin, C., Clement, C., Bruneau, M. and Pommeret, D. (2007). Time-patterns of antibiotic exposure in poultry production—a Markov chains exploratory study of nature and consequences. *Preventive Veterinary Medicine*, *80*: 230±240.
- Choo, L. C., Saleha, A. A., Wai, S. S., & Fauziah, N. (2011). Isolation of *Campylobacter* and *Salmonella* from houseflies (*Musca domestica*) in a University campus and a poultry farm in Selangor, Malaysia. *Tropical Biomedicine*, *28*(1), 16-20.
- Chopra, I., & Roberts, M. (2001). Tetracycline antibiotics: Mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiology and Molecular Biology Reviews*, *65*(2), 232-260.
- Clinical and Laboratory Standards Institute. (CLSI) (2010). Performance standards for antimicrobial susceptibility testing: *Twentieth information Supplement M100-S20* Wayne, PA. USA.
- Cloak, O.M., Duffy, G., Sheridan, J.J., Blair, I.S. and Mc Dowell, D.A. (2001). A survey on the incidence of *Campylobacter* spp. and the development of a survey adhesion polymerase chain reaction (SA-PCR) assay for the detection of *Campylobacter jejuni* in retail meat products. *Food Microbiology* *18*: 287-298.
- Cogan, T. A., Slader, J., Bloomfield, S. F., & Humphrey, T. J. (2002). Achieving hygiene in the domestic kitchen: the effectiveness of commonly used cleaning procedures. *Journal of Applied Microbiology*, *92*(5), 885-892.
- Cohen N, Ennaji H, Bouchrif B, Hassar M, Karib H (2007) Comparative study of microbiological quality of raw poultry meat at various seasons and for different slaughtering processes in Casablanca (Morocco). *Journal of Applied Poultry Research* *16*:502–508

- Coker, A. O., Isokpehi, R. D., Thomas, B. N., Amisu, K. O., & Obi, C. L. (2002). Human campylobacteriosis in developing countries. *Emerging Infectious Diseases*, 8(3), 237-244.
- Cools, I., D'Haese, E., Uyttendaele, M., Storms, E., Nelis, H. J., & Debevere, J. (2005). Solid phase cytometry as a tool to detect viable but non-culturable cells of *Campylobacter jejuni*. *Journal of Microbiological Methods*, 63, 107–114.
- Corry, J. E. L., & Atabay, H. I. (2001). Poultry as a source of *Campylobacter* and related organisms. *Journal of Applied Microbiology*, 90(S6), 96S-114S.
- Costa, D., Poeta, P., Sáenz, Y., Coelho, A. C., Matos, M., Vinué, L & Torres, C. (2008). Prevalence of antimicrobial resistance and resistance genes in faecal *Escherichia coli* isolates recovered from healthy pets. *Veterinary Microbiology*, 127(1), 97-105.
- Cox, N., N. Stern, J. Wilson, M. Musgrove, R. Buhr, and R. Hiatt. (2001). Isolation of *Campylobacter* from semen samples of commercial 50 week old parent roosters. *International Journal Medical Microbiology*. 291:39.
- Da Costa, P. M., Loureiro, L., and Matos, A. J. (2013). Transfer of multidrugresistant bacteria between intermingled ecological niches: the interface between humans, animals and the environment. *International Journal of Environmental Research and Public Health* 10, 278–294.
- Da Costa, P. M., Oliveira, M., Bica, A., Vaz-Pires, P., & Bernardo, F. (2007). Antimicrobial resistance in *Enterococcus* spp and *Escherichia coli* isolated from poultry feed and feed ingredients. *Veterinary Microbiology*, 120(1), 122-131.
- Da Costa, P.M., Belo, A., Goncalves, J., Bernardo, F. (2009) Field trial evaluating changes in prevalence and patterns of antimicrobial resistance among *Escherichia coli* and *Enterococcus* spp isolated from growing broilers medicated with enrofloxacin, apramycin, and amoxicillin. *Veterinary Microbiol.* 18: 284-292.
- Danis, K., Di Renzi, M., O'Neill, W., Smyth, B., McKeown, P., Foley, B., Devine, M.(2009). Risk factors for sporadic *Campylobacter* infection: an all-Ireland case control study. *European Surveillance*, 14(7).
- Davis, M., & Morishita, T. Y. (2005). Relative ammonia concentrations, dust concentrations, and presence of *Salmonella* species and *Escherichia coli* inside and outside commercial layer facilities. *Avian Diseases*, 49(1), 30-35.
- De Boer, E., Hahne, M., (1990). Cross-contamination with *Campylobacter jejuni* and *Salmonella* spp. from raw chicken products during food preparation. *Journal of Food Protection* 53, 1067 – 1068

- de Wit, M. A., M. P. Koopmans, L. M. Kortbeek, N. J. van Leeuwen, A. I. Bartelds, and Y. T. Duynhoven (2001). Gastroenteritis in sentinel general practices in the Netherlands. *Clinical and Infectious Diseases* 33: 280-288.
- Debruyne, L., Broman, T., Bergström, S., Olsen, B., On, S. L. W., & Vandamme, P. (2010b). *Campylobacter volucris* sp. nov, isolated from black-headed gulls (*Larus ridibundus*). *International Journal of Systematic and Evolutionary Microbiology*, 60(8), 1870–5.
- Debruyne, L., D. Gevers, and P. Vandamme (2008). Taxonomy of the Family Campylobacteraceae, p. 3-25. In I. Nachamkin, C. M. Szymanski, and M. J. Blaser (ed), *Campylobacter*, 3rd ed. *American Society for Microbiology Press, Washington, DC*.
- Dekeyser, P., Gossuin-Detrain, M., Butzler, J. P., & Sternon, J. (1972). Acute enteritis due to related vibrio: first positive stool cultures. *Journal of Infectious Diseases*, 125(4), 390-392.
- Denis, M., Refrégier-petton, J., Laisney, M. J., Ermel, G. and Salvat, G. (2001). *Campylobacter* contamination in French chicken production from farm to consumers. Use of a PCR assay for detection and identification of *Campylobacter jejuni* and *C. coli*. *Journal of Applied Microbiology* 91: 255-267
- Denis, M., Tanguy, M., Chidaine, B., Laisney, M. J., Mégraud, F., & Pravallo, P. (2011). Description and sources of contamination by *Campylobacter* spp. Of river water destined for human consumption in Brittany, France. *Pathologie Biologie*, 59, 256-263.
- Dhama, K., Rajagunalan, S., Chakraborty, S., Verma, A.K., Kumar, A., Tiwari, R. and Kapoor, S. (2013a). Food-borne pathogens of animal origin – diagnosis, prevention and control and their zoonotic significance: A review. *Pakistan Journal of Biological Sciences*, 16(20): 1076-1085.
- Dhillon, A. S., H. L. Shivaprasad, D. Schaberg, F. Wier, S. Weber, and D. Bandli. (2006). *Campylobacter jejuni* infection in broiler chickens. *Avian Diseases* 50:55–58.
- Diarrassouba, F.; Diarra, MS.; Bach, S.; Delaquis, P.; Pritchard, J.; Topp, E.; Skura, BJ. (2007). Antibiotic resistance and virulence genes in commensal *Escherichia coli* and *Salmonella* isolates from commercial broiler chicken farms. *Journal for Food Protection* 70(6):1316- 1327.
- Diker, K. S., Esendal, O. M., & Akan, M. (2000). Epidemiology of ovine *Campylobacter* infection determined by numerical analysis of electrophoretic protein profiles. *Journal of Veterinary Medicine, Series B*, 47(10), 739-744.
- Dominguez, C., Gomez, I. and Zumalacarregui, J. (2002). Prevalence of *Salmonella* and *Campylobacter* in retail chicken meat in Spain. *International Journal of Food Microbiology* 72: 165-168.

- Donnelly, J. P. (1999). Commentary on the MAFF technical report: a review of antimicrobial resistance in the food chain. *International Journal of Antimicrobial Agents*, 12, 63-65
- Duffy, G., Cagney, C., and Lynch, O. (2007). Recovery and identification of emerging campylibacteraceae from food. In Research Report No 85 edited by Downey, G. *Ashtown Food Research Centre*: Dublin
- Ellis-Iversen, J., Cook, A. J. C., Smith, R. P., Pritchard, G. C., & Nielsen, M. (2009a). Temporal patterns and risk factors for *Escherichia coli* O157 and *Campylobacter* spp, in young cattle. *Journal of Food Protection*, 72(3), 490–6.
- Ellis-Iversen, J., Pritchard, G. C., Wooldridge, M., & Nielsen, M. (2009). Risk factors for *Campylobacter jejuni* and *Campylobacter coli* in young cattle on English and Welsh farms. *Preventive Veterinary Medicine*, 88(1), 42-48.
- Endtz, H. P., Ruijs, G. J., van Klingeren, B., Jansen, W. H., van der Reyden, T., & Mouton, R. P. (1991). Quinolone resistance in *Campylobacter* isolated from man and poultry following the introduction of fluoroquinolones in veterinary medicine. *Journal of Antimicrobial Chemotherapy*, 27(2), 199-208.
- Endtz, H. P., Vliegthart, J. S., Vandamme, P., Weverink, H. W., van den Braak, N. P., Verbrugh, H. A., & van Belkum, A. (1997). Genotypic diversity of *Campylobacter lari* isolated from mussels and oysters in The Netherlands. *International Journal of Food Microbiology*, 34(1), 79-88.
- Englen, M. D., Hill, A. E., Dargatz, D. A., Ladely, S. R., & Fedorka-Cray, P. J. (2007). Prevalence and antimicrobial resistance of *Campylobacter* in US dairy cattle. *Journal of Applied Microbiology*, 102(6), 1570–7.
- Engvall, A., Bergqvist, A., Sandstedt, K., Danielsson-Tham, M.L. (1986). Colonisation of broilers with *Campylobacter* in conventional broiler-chicken flocks. *Acta Veterinaria Scandinavica* 27, 540-547.
- European Food Safety Authority (EFSA) (2008a). Foodborne antimicrobial resistance as a biological hazard. Scientific opinion of the panel on biological hazards. *The European Food Safety Authority Journal*, 765, 1-87.
- European Food Safety Authority (EFSA) (2008b). Working Group on Developing Harmonised Schemes for Monitoring Antimicrobial Resistance in Zoonotic Agents. Harmonised monitoring of antimicrobial resistance in *Salmonella* and *Campylobacter* isolates from food animals in the European Union. *Clinical Microbiology and Infection* 14: 522–533.
- European Food Safety Authority (EFSA) (2009). Joint opinion on antimicrobial resistance (AMR) focused on zoonotic infections. *European Food Safety Authority Journal* 7: 1372.

- European Food Safety Authority (EFSA) (2010). Community summary report on trends and sources of zoonoses and zoonotic agents and food-borne outbreaks in the European Union in 2008. *European Food Safety Authority Journal* 8(1) 111–136.
- European Food Safety Authority (EFSA) (2010b). Analysis of the baseline survey on the prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on broiler carcasses, in the EU, 2008, Part B: *Campylobacter*. *European Food Safety Authority Journal* 8:1522.
- European Food Safety Authority (EFSA) (2012). EU summary report on Zoonoses, zoonotic agents and food-borne outbreaks 2010. *European Food Safety Authority Journal* 10:2957.
- European Food Safety Authority (EFSA) (2013). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2011. *European Food Safety Authority Journal*, 11(4), 1-250
- Evans, J.A., Russell, S.L., James C., & Corry, J.E.L. (2004). Microbial contamination of food refrigeration equipment. *Journal of Food Engineering*, 62, 225-232.
- Everest, P. and Ketley, J. M. (2002). *Campylobacter*. Molecular Medical Microbiology edited by Sussman, M. pp. 1311-1330. Academic Press: UK
- Feng, P. C., Jinneman, K., Scheutz, F., & Monday, S. R. (2011). Specificity of PCR and serological assays in the detection of *Escherichia coli* Shiga toxin subtypes. *Applied and Environmental Microbiology*, 77(18), 6699-6702.
- Fernandez, H., Salazar, R., & Landskron, E. (1993). Occurrence of thermotolerant species of *Campylobacter* in three groups of hens maintained under different environmental conditions. *Revista de Microbiologia*, 24(4), 265-8.
- Fofana, A.; Bada Alambedji, R.; Seydi, M.; Akakpo, AJ. (2006). Antibiotic resistance of *Escherichia coli* strains isolated from raw chicken meat in Senegal. *Dakar Medicine*, 51(3): 145-150.
- Food and Agricultural Organization (FAO) (2007a). Core production data. <http://faostat.fao.org/site/340/default.aspx> Accessed Jan. 25, 2007.
- Food and Agricultural Organization (FAO) (2007b). ProdSTAT: Livestock (primary and processed). <http://faostat.fao.org/site/569/DesktopDefault.aspx?PageID=569> Chicken Meat Accessed Jan. 25, 2007
- Food and Drug Administration (FDA) (2001). Risk assessment on the human health impact of fluoroquinolone resistant *Campylobacter* associated with the consumption of chicken. Rockville, MD: US FDA.

- Fosse, J., Seegers, H., & Magras, C. (2007). Foodborne zoonoses due to meat: a quantitative approach for a comparative risk assessment applied to pig slaughtering in Europe. *Veterinary Research*, 39(1), 1-1.
- Franco, D. A. (1988). *Campylobacter* species: Considerations for controlling a food-borne pathogen. *Journal of Food Protection*, 51, 145-153
- Fratamico, P. M., & Smith, J. L. (2006). *Escherichia coli* infections. *Foodborne Infections and Intoxications*, 205-208.
- Frederick, A., & Huda, N. (2011). *Campylobacter* in Poultry: Incidences and Possible Control Measures.
- Frediani-Wolf, V., & Stephan, R. (2003). Resistance patterns of *Campylobacter* spp. strains isolated from poultry carcasses in a big Swiss poultry slaughterhouse. *International Journal of Food Microbiology*, 89(2), 233-240.
- Friedman, C. R., Hoekstra, R. M., Samuel, M., Marcus, R., Bender, J., Shiferaw, B., & Tauxe, R. V. (2004). Risk factors for sporadic *Campylobacter* infection in the United States: a case-control study in FoodNet sites. *Clinical Infectious Diseases*, 38(Supplement 3), S285-S296.
- Friedman, C. R., J. Neimann, H. C. Wegner, and R. V. Tauxe. (2004). Epidemiology of *Campylobacter jejuni* infections in the United States and other industrialized nations, p. 121-138. In I. Nachamkin and M. J. Blaser (eds.), *Campylobacter*, 2nd ed. ASM Press, Washington, D.C.
- Friesema, I. H. M., De Boer, R. F., Duizer, E., Kortbeek, L. M., Notermans, D. W., Smeulders, A., ... & Van Duynhoven, Y. T. H. P. (2012). Aetiology of acute gastroenteritis in adults requiring hospitalization in The Netherlands. *Epidemiology and Infection*, 140(10), 1780-1786.
- Frost, J. A., Kelleher, A., & Rowe, B. (1996). Increasing ciprofloxacin resistance in *Salmonellas* in England and Wales 1991–1994. *Journal of Antimicrobial Chemotherapy*, 37(1), 85-91.
- Fullerton, K. E., Ingram, L. A., Jones, T. F., Anderson, B. J., McCarthy, P. V., Hurd, S & Angulo, F. J. (2007). Sporadic *Campylobacter* infection in infants: a population-based surveillance case-control study. *The Pediatric Infectious Disease Journal*, 26(1), 19-24.
- Fuster-Valls, N., Hernandez-Herrero, M., Marin-de-Mateo, M., & Rodriguez- Jerez, J.J. (2008). Effect of different environmental conditions on the bacteria survival on stainless steel surfaces. *Food Control*, 19, 308-314.
- Gallay, A., Prouzet-Mauleon, V., Kempf, I., Lehours, P., Labadi, L., Camou, C., Denis, M., Valk, H., Desenclos, J.C. and Megraud, F. (2007). *Campylobacter* antimicrobial drug resistance among humans, broiler, chicken and pigs, France. *Emerging Infectious Diseases* 13:259-266.

- Garin, B., Gouali, M., Wouafo, M., Perchec, A. M., Thu, P. M., Ravaonindrina, N., & Pouillot, R. (2012). Prevalence, quantification and antimicrobial resistance of *Campylobacter* spp. on chicken neck-skins at points of slaughter in 5 major cities located on 4 continents. *International Journal of Food Microbiology*, 157(1), 102-107.
- Gaunt, P. N., & Piddock, L. J. (1996). Ciprofloxacin resistant *Campylobacter* spp. in humans: an epidemiological and laboratory study. *Journal of Antimicrobial Chemotherapy*, 37(4), 747-757.
- Ge, B., White, D. G., McDermott, P. F., Girard, W., Zhao, S., Hubert, S., & Meng, J. (2003). Antimicrobial-resistant *Campylobacter* species from retail raw meats. *Applied and Environmental Microbiology*, 69(5), 3005-3007.
- Genigeorgis, C., Hassuneh, M., & Collins, P. (1986). *Campylobacter jejuni* infection on poultry farms and its effect on poultry meat contamination during slaughtering. *Journal of Food Protection*, 49(11), 895-903.
- Ghafir, Y., China, B., Dierick, K., De Zutter, L., & Daube, G. (2007). A seven-year survey of *Campylobacter* contamination in meat at different production stages in Belgium. *International Journal of Food Microbiology*, 116(1), 111-120.
- Gharst, G., Hanson, D., & Kathariou, S. (2006). Effect of direct culture versus selective enrichment on the isolation of thermophilic *Campylobacter* from feces of mature cattle at harvest. *Journal of Food Protection*, 69(5), 1024-1027.
- Ghidan, A., Dobay, O., Kaszanyitzky, É. J., Samu, P., Amyes, S. G., Nagy, K., & Rozgonyi, F. (2008). Vancomycin resistant enterococci (VRE) still persist in slaughtered poultry in Hungary 8 years after the ban on avoparcin. *Acta Microbiologica et Immunologica Hungarica*, 55(4), 409-417.
- Gibbens, J. C., Pascoe, S. J. S., Evans, S. J., Davies, R. H., & Sayers, A. R. (2001). A trial of biosecurity as a means to control *Campylobacter* infection of broiler chickens. *Preventive Veterinary Medicine*, 48(2), 85-99.
- Gibreel, A., & Taylor, D. E. (2006). Macrolide resistance in *Campylobacter jejuni* and *Campylobacter coli*. *Journal of Antimicrobial Chemotherapy*, 58(2), 243-255.
- Giessen, A. V. D., B. Bloemberg, et al. (1996). Epidemiological study on risk factors and risk-reducing measures for *Campylobacter* infections in Dutch broiler flocks. *Epidemiology and Infection*. 117:245–250.
- Gilpin, B. J., Scholes, P., Robson, B., & Savill, M. G. (2008). The transmission of thermotolerant *Campylobacter* spp. to people living or working on dairy farms in New Zealand. *Zoonoses and Public Health*, 55(7), 352-360.

- Gorman, R. and Adley, C. C. (2004). An evaluation of five preservation techniques and conventional freezing temperatures of -20°C and -85°C for long-term preservation of *Campylobacter jejuni*. *Letters of Applied Microbiology* 38:306-310.
- Gormley, F.J., MacRae, M., Forbes, K.J., Ogden, I.D., Dallas, J.F. and Strachan, N.J.C. (2008). Has retail chicken played a role in the decline of human campylobacteriosis? *Applied and Environmental Microbiology* 74: 383-390.
- Gregory, E., Barnhart, H., Dreesen, D. W., Stern, N. J., & Corn, J. L. (1997). Epidemiological study of *Campylobacter* spp. in broilers: source, time of colonization, and prevalence. *Avian Diseases*, 890-898.
- Griffiths, P. L., & Park, R. W. (1990). *Campylobacter* associated with human diarrhoeal disease. *The Journal of Applied Bacteriology*, 69(3), 281–301.
- Gudmundsdottir, S., Hardardottir, H., & Gunnarsson, E. (2003). Subtyping of *Salmonella enterica* serovar typhimurium outbreak strains isolated from humans and animals in Iceland. *Journal of Clinical Microbiology*, 41(10), 4833-4835.
- Guévremont, E., Nadeau, É., Sirois, M., & Quessy, S. (2006). Antimicrobial susceptibilities of thermophilic *Campylobacter* from humans, swine, and chicken broilers. *Canadian Journal of Veterinary Research*, 70(2), 81.
- Guo, W., Ling, C., Cheng, F., Guo, W.Z., Ling, C.S. and Cheng, F.H. (1998). Preliminary investigation on enterohaemorrhagic *Escherichia coli* O157 from domestic animals and fowl in Fujian province. *Chinese Journal of Zoonoses*, 14: 3–6.
- Hakkinen, M., Heiska, H., & Hänninen, M. L. (2007). Prevalence of *Campylobacter* spp. in cattle in Finland and antimicrobial susceptibilities of bovine *Campylobacter jejuni* strains. *Applied and Environmental Microbiology*, 73(10), 3232-3238.
- Hald, B., A. Wedderkopp, and M. Madsen. (2000). Thermophilic *Campylobacter* spp. in Danish broiler production: a cross sectional survey and a retrospective analysis of risk factors for occurrence in broiler flocks. *Avian Pathology* 29:123–131.
- Hald, B., Bang, D. D., Pedersen, K., Dybdahl, J., Madsen, M., & Study, T. (2004). Flies and *Campylobacter* Broiler Flocks. *Emerging Infectious Diseases*, 10(8), 1490–1492.
- Hald, B., Wedderkopp, A., & Madsen, M. (2000). Thermophilic *Campylobacter* spp. in Danish broiler production: a cross-sectional survey and a retrospective analysis of risk factors for occurrence in broiler flocks. *Avian Pathology*, 29(2), 123-131.

- Hald, T., Vose, D., Wegener, H. C., & Koupeev, T. (2004). A Bayesian Approach to Quantify the Contribution of Animal Food Sources to Human Salmonellosis. *Risk Analysis*, 24(1), 255-269.
- Hanninen, M. L., Haajanen, H., Pummi, T., Wermundsen, K., Katila, M. L., Sarkkinen, H & Rautelin, H. (2003). Detection and typing of *Campylobacter jejuni* and *Campylobacter coli* and analysis of indicator organisms in three waterborne outbreaks in Finland. *Applied and Environmental Microbiology*, 69(3), 1391-1396.
- Hanninen, M., Niskanen, M., & Korhonen, L. (1998). Water as a Reservoir for *Campylobacter jejuni* Infection in Cows Studied by Serotyping and Pulsed-field Gel Electrophoresis. *Journal of Veterinary Medicine*, 45(2), 37-42.
- Hartnett, E., Fazil, A., Paoli, G., Nauta, M., Christensen, B. B., Rosenquist, H., et al. (2009). Série évaluation des risques microbiologiques 11. Rapport. In Evaluation des risques liés à *Campylobacter* spp. dans les poulets de chair. FAO/OMS.
- Harvey, R.B., Young, C.R., Ziprin, R.L., Hume, M.E., Genovese, K.J., Anderson, R.C., Droleskey, R.E., Stanker, L.H., Nisbet, D.J., (1999). Prevalence of *Campylobacter* spp. isolated from the intestinal tract of pigs raised in an integrated swine production system. *Journal of American Veterinary Medical Association* 215, 1601-1604.
- HCV New Drug Research (2011) *Escherichia coli*, detected in 94% of retail chicken meat samples in the Netherlands.
- Herman, L., Heyndrickx, M., Grijspeerdt, K., Vandekerchove, D., Rollier, I., & De Zutter, L. (2003). Routes for *Campylobacter* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiology and Infection*, 131(03), 1169-1180.
- Heuvelink, A. E., Zwartkruis-Nahuis, J.T., van den Biggelaar, F.L., van Leeuwen, W.J. and de Boer, E. (1999). Isolation and characterization of verocytotoxin-producing *Escherichia coli* O157 from slaughter pigs and poultry. *International Journal of Food Microbiology*, 52: 67-75.
- Hiett, K. L., Cox, N. A., Buhr, R. J., & Stern, N. J. (2002). Genotype analyses of *Campylobacter* isolated from distinct segments of the reproductive tracts of broiler breeder hens. *Current Microbiology*, 45(6), 0400-0404.
- Hong, J., Kim, J. M., Jung, W. K., Kim, S. H., Bae, W., Koo, H. C., & Park, Y. H. (2007). Prevalence and antibiotic resistance of *Campylobacter* spp. isolated from chicken meat, pork, and beef in Korea, from 2001 to 2006. *Journal of Food Protection*, 70(4), 860-866.
- Huang, J. L., Xu, H. Y., Bao, G. Y., Zhou, X. H., Ji, D. J., Zhang, G & Jiao, X. A. (2009). Epidemiological surveillance of *Campylobacter jejuni* in chicken,

- dairy cattle and diarrhoea patients. *Epidemiology and Infection*, 137(08), 1111-1120.
- Humphrey, T. J., & Beckett, P. (1987). *Campylobacter jejuni* in dairy cows and raw milk. *Epidemiology and infection*, 98(03), 263-269.
- Humphrey, T. J., A. Henley, and D. G. Lanning. (1993). The colonization of broiler chickens with *Campylobacter jejuni*: some Epidemiological investigations. *Epidemiology and Infection* 110:601–607.
- Humphrey, T., O’Brien, S., & Madsen, M. (2007). *Campylobacter* as zoonotic pathogens: a food production perspective. *International Journal of Food Microbiology*, 117(3), 237–57.
- Inglis, G. D., Kalischuk, L. D., & Busz, H. W. (2004). Chronic shedding of *Campylobacter* species in beef cattle. *Journal of Applied Microbiology*, 97(2), 410-420.
- Inglis, G. D., Morck, D. W., McAllister, T. A., Entz, T., Olson, M. E., Yanke, L. J., & Read, R. R. (2006). Temporal prevalence of antimicrobial resistance in *Campylobacter* spp. from beef cattle in Alberta feedlots. *Applied and Environmental Microbiology*, 72(6), 4088-4095.
- Institute of Food Technology (IFT) (2006). Antimicrobial resistance: implications for the food system. Institute of Food Technologists, Chicago, IL. <http://www.ift.org/knowledge-center/read-iftpublications/sciencereports/expert-reports/antimicrobial-resistance.aspx>.
- Jackson, T.C., Marshall, D.L., Acuff, G.R., Dickson, J.S., (2001). Meat, poultry, and seafood. In: Doyle, M.P., Beuchat, L.R., Montville, T.J. (Eds.), *Food Microbiology: Fundamentals and Frontiers*, 2nd Edition. ASM Press, Washington, DC, pp. 91–109 .
- Jacobs-Reitsma, W. F., U. Lyhs, and J. Wagenaar. (2008). *Campylobacter* in the food supply, p. 627–644. In I. Nachamkin, C. M. Szymanski, and M. J. Blaser (ed.), *Campylobacter*, 3rd ed. ASM Press, Washington, DC.
- Jacobs-Reitsma, W. F., Van de Giessen, A. W., Bolder, N. M., & Mulder, R. W. A. W. (1995). Epidemiology of *Campylobacter* spp. at two Dutch broiler farms. *Epidemiology and Infection*, 114(03), 413-421.
- Jacobs-Reitsma, W., C. Becht, T. De Vries, J. Van der Plas, B. Duim, and J. Wagenaar. (2001). No evidence for vertical transmission of *Campylobacter* in a study on Dutch breeder and broiler farms. *International Journal of Medical Microbiology* 291:39.
- Jacobs-Reitsma, W.F. (2000): *Campylobacter* in the Food Supply. In: *Campylobacter*. 2nd ed. Nachamkin, I. and M.J. Blaser (Eds). Washington, DC. ASM Press, pp. 467-481

- Jacobs-Reitsma, W.F., (1997). Aspects of Epidemiology of *Campylobacter* in poultry. *Veterinary Quarterly* 19, 113 – 117. I.F.S.T., 1999. Organic food. *International Food Safety News* 8, 2 – 6.
- Jayarao, B. M., & Henning, D. R. (2001). Prevalence of foodborne pathogens in bulk tank milk. *Journal of Dairy Science*, 84(10), 2157-2162.
- Jeffrey, J.S., Singer, R.S., O'Connor, R. and Atwill, E.R. (2004). Prevalence of pathogenic *Escherichia coli* in the broiler house environment. *Avian Diseases*, 48: 189-195.
- Jensen, A. N., Dalsgaard, A., Baggesen, D. L., & Nielsen, E. M. (2006). The occurrence and characterization of *Campylobacter jejuni* and *C. coli* in organic pigs and their outdoor environment. *Veterinary Microbiology*, 116(1), 96-105.
- Jimenez, S. M., Tiburzi, M. C., Salsi, M. S., Pirovani, M. E. and Moguevsky, M. A. (2003). The role of visible faecal material as a vehicle for generic *Escherichia coli*, coliform, and other *enterobacteria* contaminating poultry carcasses during slaughtering. *Journal of Applied Microbiology* 95, 451-456.
- Jones K., (2001). *Campylobacters* in water, sewage and the environment. *Journal of Applied Microbiology* 90: 68S-79S.
- Jordan D, et al. (2009). Antimicrobial use in the Australian pig industry: results of a national survey. *Australian Veterinary Journal* 87: 222–229
- Jørgensen, F., Bailey, R., Williams, S., Henderson, P., Wareing, D. R. A., Bolton, F. J., & Humphrey, T. J. (2002). Prevalence and numbers of *Salmonella* and *Campylobacter* spp. on raw, whole chickens in relation to sampling methods. *International Journal of Food Microbiology*, 76(1), 151-164.
- Kabir, S.M.L. (2010). Avian colibacillosis and salmonellosis: A closer look at epidemiology, pathogenesis, diagnosis, control and public health concerns. *International Journal of Environmental Research and Public Health*, 7: 89-114.
- Kaper, J. B., Nataro, J. P., & Mobley, H. L. (2004). Pathogenic *Escherichia coli*. *Nature Reviews Microbiology*, 2(2), 123-140.
- Kapperud, G., E. Skjerve, L. Vik, K. Hauge, A. Lysaker, I. Aalmen, S. Ostroff, and M. Potter. (1993). Epidemiological investigation of risk factors for campylobacter colonisation in Norwegian broiler flocks. *Epidemiology and Infection* 111:245–255.
- Karagiannis, I., Sideroglou, T., Gkolfinopoulou, K., Tsouri, A., Lampousaki, D., Velonakis, E. N. & Bonovas, S. (2010). A waterborne *Campylobacter jejuni* outbreak on a Greek island. *Epidemiology and Infection*, 138(12), 1726-1734.
- Karmali, M. A., Simor, A. E., Roscoe, M., Fleming, P. C., Smith, S. S., & Lane, J. (1986). Evaluation of a blood-free, charcoal-based, selective medium for the

isolation of *Campylobacter* organisms from feces. *Journal of Clinical Microbiology*, 23(3), 456-459.

- Kaszanyitzky, E. J., Tenk, M., Ghidan, A., Fehervari, G. Y., & Papp, M. (2007). Antimicrobial susceptibility of *Enterococci* strains isolated from slaughter animals on the data of Hungarian resistance monitoring system from 2001 to 2004. *International Journal of Food Microbiology*, 115(1), 119-123.
- Kazwala, R.R., Collins, J.D., Hannan, J., Crinion, R.A. and O'Mahony, H. (1990) Factors responsible for the introduction and spread of *Campylobacter jejuni* infection in commercial poultry production. *Veterinary Record* 126, 305-306.
- Kemp, R., Leatherbarrow, A. J. H., Williams, N. J., Hart, C. A., Clough, H. E., Turner, J. & French, N. P. (2005). Prevalence and genetic diversity of *Campylobacter* spp. in environmental water samples from a 100-square-kilometer predominantly dairy farming area. *Applied and Environmental Microbiology*, 71(4), 1876-1882.
- Ketley, J. M. (1997). Pathogenesis of enteric infection by *Campylobacter*. *Microbiology*, 143(1), 5-21.
- Kikuvi GM, Schwarz S, Ombui JN, Mitema ES, Kehrenberg C. (2007). Streptomycin and chloramphenicol resistance genes in *Escherichia coli* isolates from cattle, pigs, and chicken in Kenya. *Microbiology and Drug Resistant* 13:62– 68.
- King, E. O. (1957). Human infections with *Vibrio fetus* and a closely related *Vibrio*. *Infectious Diseases* 101, 119-128.
- Kist, M. (1986). [Who discovered *Campylobacter jejuni/coli*? A review of hitherto disregarded literature]. Zentralblatt für Bakteriologie, Mikrobiologie, und Hygiene. Series A, Medical microbiology, infectious diseases. *Virology and Parasitology*, 261(2), 177-186.
- Kittl, S., Kuhnert, P., Hächler, H., & Korczak, B. M. (2011). Comparison of genotypes and antibiotic resistance of *Campylobacter jejuni* isolated from humans and slaughtered chickens in Switzerland. *Journal of Applied Microbiology*, 110(2), 513-520.
- Kumar, A., Agarwal, R. K., Bhilegaonkar, K. N., Shome, B. R., & Bachhil, V. N. (2001). Occurrence of *Campylobacter jejuni* in vegetables. *International Journal of Food Microbiology*, 67(1), 153-155.
- Kurinčič, M., Berce, I., Zorman, T., & Smole Možina, S. (2005). The prevalence of multiple antibiotic resistance in *Campylobacter* spp. from retail poultry meat. *Food Technology and Biotechnology*, 43(2), 157-163.
- Kusumaningrum, H. D., van Asselt, E. D., Beumer, R. R., & Zwietering, M. H. (2004). A quantitative analysis of cross-contamination of *Salmonella* and

Campylobacter spp. via domestic kitchen surfaces. *Journal of Food Protection*, 67(9), 1892–1903.

- Kusumaningrum, H.D., Van Putten, M.M., Rombouts, F.M. and Beumer, R.R. (2002) Effects of antibacterial dishwashing liquid on foodborne pathogens and competitive microorganisms in kitchen sponges. *Journal of Food Protection* 65, 61–65.
- Kuusi, M., Nuorti, J. P., Hänninen, M. L., Koskela, M., Jussila, V., Kela, E. & Ruutu, P. (2005). A large outbreak of campylobacteriosis associated with a municipal water supply in Finland. *Epidemiology and Infection*, 133(04), 593-601.
- Lastovica, A. J., & Allos, B. M. (2008). Clinical significance of *Campylobacter* and related species other than *Campylobacter jejuni* and *Campylobacter coli*. In eds Nachamkin, I., Szymanski, C. M., & Blaser, M. J. *Campylobacter*, (Ed. 3), Washington, DC: ASM Press, 123-149.
- Laxminarayan, R., Duse, A., Wattal, C., Zaidi, A. K., Wertheim, H. F., Sumpradit, N & Cars, O. (2013). Antibiotic resistance—the need for global solutions. *The Lancet infectious diseases*, 13(12), 1057-1098.
- Le Roux, E., and Lastovica, A. J. (1998). The Cape Town Protocol: How to isolate the most *Campylobacters* for your dollar, pound, frank, yen, etc. In: *Campylobacter, Helicobacter and Related Organisms*. Ed. Lastovica, A. J., Newell, D. G., and Lastovica, E. E. pp 30-33. Institute of Child Helath, Univ. of Cape Town, Rondebosch: Cape Town, South Africa
- Lee, A., O'Rourke, J. L., Barrington, P. J., & Trust, T. J. (1986). Mucus colonization as a determinant of pathogenicity in intestinal infection by *Campylobacter jejuni*: a mouse cecal model. *Infection and Immunity*, 51(2), 536–46.
- Lee, M. D., & Newell, D. G. (2006). *Campylobacter* in poultry: filling an Ecological niche. *Avian Diseases*, 50(1), 1-9.
- Lévesque, S., Frost, E., Arbeit, R. D., & Michaud, S. (2008). Multilocus sequence typing of *Campylobacter jejuni* isolates from humans, chickens, raw milk, and environmental water in Quebec, Canada. *Journal of Clinical Microbiology*, 46(10), 3404-3411
- Levine, M. M. (1987). Escherichia coli that cause diarrhea: enterotoxigenic, nteropathogenic, enteroinvasive, enterohemorrhagic, and enteroadherent. *Journal of infectious Diseases*, 155(3), 377-389.
- Levy, S. B., and B. Marshall. (2004). Antibacterial resistance worldwide: causes, challenges and responses. *Nature Medicine*. 10(12 Suppl.): S122&129.
- Li, J., Pai, H., Kim, Y. K., Kim, N. H., Eun, B. W., Kang, H. J & Ahn, H. S. (2007). Control of extended-spectrum β -lactamase-producing *Escherichia coli* and

Klebsiella pneumoniae in a children's hospital by changing antimicrobial agent usage policy. *Journal of Antimicrobial Chemotherapy*, 60(3), 629-637.

- Li, L., Jiang, Z. G., Xia, L. N., Shen, J. Z., Dai, L., Wang, Y. & Wu, C. M. (2010). Characterization of antimicrobial resistance and molecular determinants of beta-lactamase in *Escherichia coli* isolated from chickens in China during 1970–2007. *Veterinary Microbiology*, 144(3), 505-510
- Lindblom, G. B., E. Sjorgren, and B. Kaijser. (1986). Natural campylobacter colonization in chickens raised under different environmental conditions. *Journal of Hygiene*. (London) 96:385–391.
- Litrup, E., M. Torpdahl, and E. M. Nielsen. (2007): "Multilocus sequence typing performed on *Campylobacter coli* isolates from humans, broilers, pigs and cattle originating in Denmark." *Journal of Applied Microbiology* 103.1.210-218.
- Luangtongkum, T., Jeon, B., Han, J., Plummer, P., Logue, C. M., & Zhang, Q. (2009). Antibiotic resistance in *Campylobacter*: emergence, transmission and persistence. *Future Microbiology*, 4(2), 189–200.
- Luangtongkum, T., Morishita, T. Y., Ison, A. J., Huang, S., McDermott, P. F., & Zhang, Q. (2006). Effect of conventional and organic production practices on the prevalence and antimicrobial resistance of *Campylobacter* spp. in poultry. *Applied and Environmental Microbiology*, 72(5), 3600-3607.
- Lues, J. F. R., & Van Tonder, I. (2007). The occurrence of indicator bacteria on hands and aprons of food handlers in delicatessen sections of a retail group. *Food Control*, 18, 326-332.
- Lutful Kabir, S. M., Jamrul Islam, Hasanuzzaman Suman M., Shahidur Rahman Khan, M and Shinji Yamasaki. (2014). Isolation, Identification and Antimicrobial Susceptibility Profiles of *Campylobacter* species with assessment of their risk factors in broiler flocks of Bangladesh Agricultural University Poultry Farm. *Journal of Basic Applied Sciences and Research*, 4(7)160-168
- Lynch, O. A, Cagney, C., McDowell, D. A., & Duffy, G. (2011). Occurrence of fastidious *Campylobacter* spp. in fresh meat and poultry using an adapted cultural protocol. *International Journal of Food Microbiology*, 150(2-3), 171–7.
- Mabote, K. I., Mbewe, M., & Ateba, C. N. (2011). Prevalence of *Campylobacter* Contamination in Fresh Chicken Meat and Milk Obtained from Markets in the Northwest Province, South Africa. *Journal of Human Ecology*, 36(1), 23–28.
- Mackiw E, Korsak D, Rzewuska K, Tomczuk K, Rozynek E (2012): Antibiotic resistance in *Campylobacter jejuni* and *Campylobacter coli* isolated from food in Poland. *Food Control* 23, 297–301.

- Madden, R. H., Murray, K. A., & Gilmour, A. (2007). Carriage of four bacterial pathogens by beef cattle in Northern Ireland at time of slaughter. *Letters in Applied Microbiology*, 44(2), 115-119.
- Madigan, M.T., Martinko, J.M., Parker, J., (2003). Brock Biology of Microorganisms, 10th ed. Pearson Education, London.
- Magiorakos, A. P., Srinivasan, A., Carey, R. B., Carmeli, Y., Falagas, M. E., Giske, C. G & Monnet, D. L. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clinical Microbiology and Infection*, 18(3), 268-281.
- Man S. M. (2011). Reviews: The Clinical importance of emerging *Campylobacter* species. *National Reviews of Gastroenterology and Hepatology*. 8:669-685.
- Mansouri-najand, L., Saleha, A. A., & Wai, S. S. (2012). Prevalence of Multidrug resistance *Campylobacter jejuni* and *Campylobacter coli* in chickens slaughtered in selected markets, Malaysia. *Tropical Biomedicine*, 29(2), 231-238.
- Marija Kurinčič, Berce, I., Zorman, T., & Smole Možina, S. (2005). The prevalence of multiple antibiotic resistance in *Campylobacter* spp. from retail poultry meat. *Food Technology and Biotechnology*, 43(2), 157-163.
- Marshall, B. M., & Levy, S. B. (2011). Food animals and antimicrobials: impacts on human health. *Clinical microbiology reviews*, 24(4), 718-733
- Marshall, B. M., & Levy, S. B. (2011). Food animals and antimicrobials: impacts on human health. *Clinical Microbiology Reviews*, 24(4), 718-733.
- McEwen, S. A. (2006). Antibiotic use in animal agriculture: What have we learned and where are we going?. *Animal biotechnology*, 17(2), 239-250.
- Mead, P. S., L. Slutsker, V. Dietz, L. F. McCaig, J. S. Bresee, C. Shapiro, P. M. Griffin, R. V. Tauxe. (1999). Food-related illness and death in the United States. *Emerging Infectious Diseases*. 5: 607-625.
- Meeyam, T., Padungtod, P. and Kaneene, J. B. (2004). Molecular characterization of *Campylobacter* isolated from chickens and humans in northern Thailand. *Southeast Asia Journal of Tropical Medical Public Health* 35:670-676.
- Megraud, F., & Bultel, C. (2004). Appréciation des risques alimentaires liés aux *Campylobacters*. Rapport de. In Application au couple poulet/*Campylobacter jejuni*. AFSSA, p.96
- Meldrum, R. J., & Wilson, I. G. (2007). *Salmonella* and *Campylobacter* in United Kingdom retail raw chicken in 2005. *Journal of Food Protection*, 70(8), 1937-1939.

- Meng, X., Yang, J., Cao, Y., Li, L., Jiang, X., Xu, X & Zhang, Y. (2007). Increasing fatty acid production in *E. coli* by simulating the lipid accumulation of oleaginous microorganisms. *Journal of Industrial Microbiology & Biotechnology*, 38(8), 919-925.
- Mensah, P., Yeboah-Manu, D., Owusu-Darko, K., Ablordey, A., (2002). Street foods in Accra, Ghana: how safe are they? *Bulletin of the World Health Organization* 80, 1 – 14.
- Miwa, N., Takegahara, Y., Terai, K., Kato, H., & Takeuchi, T. (2003). *Campylobacter jejuni* contamination on broiler carcasses of *C. jejuni* negative flocks during processing in a Japanese slaughterhouse. *International Journal of Food Microbiology*, 84(1), 105-109.
- Mølbak, K. (2004). Spread of resistant bacteria and resistance genes from animals to humans—the public health consequences. *Journal of Veterinary Medicine, Series B*, 51(89), 364-369.
- Moldlab (2003): Mold analytical services & independent laboratory. Sewage \ Black-water: Contamination and *Escherichia coli*. (http://www.moldlab.com/sewage_screen.htm, accessed on 16 August, 2007).
- Moniri, R., & Dastehgoli, K. (2005). Fluoroquinolone-resistant *Escherichia coli* isolated from healthy broilers with previous exposure to fluoroquinolones: Is there a link?. *Microbial Ecology in Health and Disease*, 17(2), 69-74.
- Moore, J. E., Barton, M. D., Blair, I. S., Corcoran, D., Dooley, J. S., Fanning, S., Kempft, I., Lastovica, A. J., Loweri, C. J., Matsuda, M., McDowell, D. A., McMahon, A., Millar, B. C., Rao, J. R., Rooney, P. J., Seal, B. S., Snelling, W. J. and Tolba, O. (2006). The Epidemiology of antibiotic resistance in *Campylobacter*. *Microbes and Infection* 8:1955–1966
- Moore, J. E., Caldwell, P. S., Millar, B. C., & Murphy, P. G. (2001a). Occurrence of *Campylobacter* spp. in water in Northern Ireland: implications for Public Health. *The Ulster Medical Journal*, 70(2), 102–7
- Moore, J. E., Corcoran, D., Dooley, J. S., Fanning, S., Lucey, B., Matsuda, M & Whyte, P. (2005). *Campylobacter*. *Veterinary Research*, 36(3), 351-382.
- Morabito, S., Dell’Omo, G., Agrimi, U., Schmidt, H., Karch, H., Cheasty, T., & Caprioli, A. (2001). Detection and characterization of Shiga toxin-producing *Escherichia coli* in feral pigeons. *Veterinary Microbiology*, 82(3), 275-283.
- Musgrove, M. T., Jones, D. R., Northcutt, J. K., Cox, N. A., Harrison, M. A., Fedorka-Cray, P. J., & Ladely, S. R. (2006). Antimicrobial resistance in *Salmonella* and *Escherichia coli* isolated from commercial shell eggs. *Poultry science*, 85(9), 1665-1669.

- Nachamkin, I., and M. Blaser. (2000). Diagnosis and antimicrobial susceptibility of *Campylobacter* species, p. 45-66. I. Nachamkin, J. Engberg, and F. M. Aarestrup (ed.), *Campylobacter*, 2nd ed. American Society for Microbiology Press, Washington, D.C.
- Nandi, S., Maurer, J. J., Hofacre, C., & Summers, A. O. (2004). Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistance integrons in poultry litter. *Proceedings of the National Academy of Sciences of the United States of America*, 101(18), 7118-7122.
- Nataro, J. P., & Kaper, J. B. (1998). Diarrheogenic *Escherichia coli*. *Clinical Microbiology Reviews*, 11(1), 142-201.
- National Public Health Services (NPHS). The National Public Health Service for Wales (2006). Infection & Communicable Disease Services; Information & Surveillance Data. *Campylobacter*. Available from <http://www.wales.nhs.uk/sites/home.-cfm?OrgID=368> [Last updated: 06/06/2006].
- Nauta, M. J., Fischer, A. R., Van Asselt, E. D., De Jong, A. E., Frewer, L. J., & De Jonge, R. (2008). Food safety in the domestic environment: the effect of consumer risk information on human disease risks. *Risk Analysis*, 28(1), 179-192.
- Nauta, M., Hill, A., Rosenquist, H., Brynestad, S., Fetsch, A., van der Logt, P., ... & Havelaar, A. (2009). A comparison of risk assessments on *Campylobacter* in broiler meat. *International Journal of Food Microbiology*, 129(2), 107-123.
- Nazmul, M. H. M. (2008). Molecular characterization of verotoxin gene in enteropathogenic *Escherichia coli* isolated from Miri Hospital, Sarawak, Malaysia. *Biomedical Research*, 19(1), 9-12.
- Nesbakken, T., Eckner, K., & Røtterud, O.-J. (2008). The effect of blast chilling on occurrence of human pathogenic *Yersinia enterocolitica* compared to *Campylobacter* spp. and numbers of hygienic indicators on pig carcasses. *International Journal of Food Microbiology*, 123(1-2), 130-3.
- Newell D.G. & Fearnley C., (2003). Sources of *Campylobacter* colonization in broiler chickens. *Applied Environmental Microbiology* 69 , 4343-4351.
- Newell, D. G. (2001). Animal models of *Campylobacter jejuni* colonization and disease and the lessons to be learned from similar *Helicobacter pylori* models. *Journal of Applied Microbiology*, 90(S6), 57S-67S.
- Newell, D. G., Elvers, K. T., Dopfer, D., Hansson, I., Jones, P., James, S., & Allen, V. M. (2011). Biosecurity-based interventions and strategies to reduce *Campylobacter* spp. on poultry farms. *Applied and Environmental Microbiology*, 77(24), 8605-8614.

- Newell, D. G., Frost, J. A., Duim, B., Wagenaar, J. A., Madden, R. H., Plas, J. V. D., & On, S. L. W. (2000). New developments in the subtyping of *Campylobacter* species.
- Nonga, H. E., Sells, P., & Karimuribo, E. D. (2010). Occurrences of thermophilic *Campylobacter* in cattle slaughtered at Morogoro municipal abattoir, Tanzania. *Tropical Animal Health and Production*, 42(1), 73–8.
- Norstrom, M., Hofshagen, M., Stavnes, T., Schau, T., Lassen, J. and Kruse, H. (2006). Antimicrobial resistance in *Campylobacter jejuni* from humans and broilers in Norway. *Epidemiology and Infection* 134:127-130.
- Nzouankeu, A., Ngandjio, A., Ejenguele, G., Njine, T., & Wouafo, M. N. (2010). Multiple contaminations of chickens with *Campylobacter*, *Escherichia coli* and *Salmonella* in Yaounde (Cameroon). *Journal of Infection in Developing Countries*, 4(9), 583-686
- O'Brien, T. F. (2002). Emergence, spread, and environmental effect of antimicrobial resistance: how use of an antimicrobial anywhere can increase resistance to any antimicrobial anywhere else. *Clinical Infectious Diseases*, 34(Supplement 3), S78-S84.
- O'Reilly, C. E., A. B. Bowea, N. E. Perez, J. P. Sarisky, C. A. Shepherd, M. D. Miller, B. C. Hubbard, M. Herring, S. D. Buchanan, C. C. Fitzgerald, V. Hill, M. J. Arrowood, L. X. Xiao, R. M. Hoekstra, E. D. Miatz, M. F. Lynch, and Outbreak working group. (2007). A Waterborne Outbreak of gastroenteritis with Multiple Etiologies among Resort Island Visitors and Residents: Ohio. *CID*. 44: 506-512.
- O'Brien, F. G., Price, C., Grubb, W. B., & Gustafson, J. E. (2002). Genetic characterization of the fusidic acid and cadmium resistance determinants of *Staphylococcus aureus* plasmid pUB101. *Journal of Antimicrobial Chemotherapy*, 50(3), 313-321.
- Ogden, I. D., MacRae, M., Johnston, M., Strachan, N. J. C., Cody, A. J., Dingle, K. E., & Newell, D. G. (2007). Use of multilocus sequence typing to investigate the association between the presence of *Campylobacter* spp. in broiler drinking water and *Campylobacter* colonization in broilers. *Applied and Environmental Microbiology*, 73(16), 5125–9.
- Olah, P.A., Sherwood, J.S., Elijah, L.M., Dockter, M.R., Doetkott, C., Miller, Z. and Logue, C.M. (2004) Comparison of antimicrobial resistance in *Salmonella* and *Campylobacter* isolated from turkeys in the Midwest USA. *Food Microbiology* 21, 779–789.
- Oliver, S. P., Murinda, S. E., & Jayarao, B. M. (2011). Impact of antibiotic use in adult dairy cows on antimicrobial resistance of veterinary and human pathogens: a comprehensive review. *Foodborne Pathogens and Disease*, 8(3), 337-355.

- On, S.L.W. (2001): Taxonomy of *Campylobacter*, *Arcobacter*, *Helicobacter* and related bacteria: current status, future prospects and immediate concerns. *Journal of Applied Microbiology* 90(S): 1S-5S
- Ono, K., and Yamamoto, K. (1999). Contamination of meat with *Campylobacter jejuni* in Saitama, Japan. *International Journal of Food Microbiology* 47:211-219.
- Oporto, B., Juste, R. A., & Hurtado, A. (2009). Phenotypic and genotypic antimicrobial resistance profiles of *Campylobacter jejuni* isolated from cattle, sheep, and free-range poultry faeces. *International Journal of Microbiology*, 2009 (2010)
- Oyarzabal, O. A., Conner, D. E., & Hoerr, F. J. (1995). Incidence of *Campylobacters* in the intestine of avian species in Alabama. *Avian Diseases*, 147-151.
- Ozaki, H., & Murase, T. (2009). Multiple routes of entry for *Escherichia coli* causing colibacillosis in commercial layer chickens. *The Journal of Veterinary Medical Science/the Japanese Society of Veterinary Science*, 71(12), 1685-1689.
- Padungton, P. and Kaneene, J. B. (2003). *Campylobacter* spp. in human, chickens, pigs and their antimicrobial resistance. *Journal of Veterinary Medical Science* 65:161-170.
- Pallecchi, L.; Lucchetti, C.; Bartoloni, A.; Bartalesi, F.; Mantella, A.; Gamboa, H.; Carattoli, A.; Paradisi, F.; Rossolini, GM. (2007). Population structure and resistance genes in Antibiotic-resistant bacteria from a remote community with minimal antibiotic exposure. *Antimicrobial Agents and Chemotherapy* 51(4):1179-1184.
- Pandey, M. C., Bawa, A. S., & Guerrero-Legarreta, I. (2010). Basic Operations and Conditions. *Handbook of Poultry Science and Technology, Secondary Processing*, 2, 403.
- Park, S. F. (2002). The physiology of *Campylobacter* species and its relevance to their role as foodborne pathogens. *International Journal of Food Microbiology*, 74(3), 177-188.
- Pattison M. (2001). Practical intervention strategies for *Campylobacter*. *Journal of Applied Microbiology* 90:121-125.
- Payne, R. E., Lee, M. D., Dreesen, D. W., & Barnhart, H. M. (1999). Molecular epidemiology of *Campylobacter jejuni* in broiler flocks using randomly amplified polymorphic DNA-PCR and 23S rRNA-PCR and role of litter in its transmission. *Applied and Environmental Microbiology*, 65(1), 260-263.
- Pearson, A. D., M. H. Greenwood, R. K. Feltham, T. D. Healing, J. Donaldson, D. M. Jones, and R. R. Colwell. (1996). Microbial ecology of *Campylobacter jejuni* in a United Kingdom chicken supply chain: intermittent common

source, vertical transmission, and amplification by flock propagation. *Applied Environmental Microbiology*. 62:4614–4620.

Penner, J. L. (1988). The genus *Campylobacter*: a decade of progress. *Clinical Microbiology Review*. 1: 157-172.

Peterson, M. C. (2003). *Campylobacter jejuni* enteritis associated with consumption of raw milk. *Journal of Environmental Health*, 65(9), 20-21.

Pezzotti, G., Serafin, A., Luzzi, I., Mioni, R., Milan, M., & Perin, R. (2003). Occurrence and resistance to antibiotics of *Campylobacter jejuni* and *Campylobacter coli* in animals and meat in northeastern Italy. *International Journal of Food Microbiology*, 82(3), 281-287.

Phillips, I., Casewell, M., Cox, T., De Groot, B., Friis, C., Jones, R., & Waddell, J. (2004). Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *The Journal of Antimicrobial Chemotherapy*, 53(1), 28–52.

Piddock, L. J. V., Griggs, D., Johnson, M. M., Ricci, V., Elviss, N. C., Williams, L. K., & Owen, R. J. (2008). Persistence of *Campylobacter* species, strain types, antibiotic resistance and mechanisms of tetracycline resistance in poultry flocks treated with chlortetracycline. *Journal of Antimicrobial Chemotherapy*, 62(2), 303-315.

Porter, I. A., & Reid, T. (1980). A milk-borne outbreak of *Campylobacter* infection. *Journal of Hygiene*, 84(03), 415-419.

Posch, J., Feierl, G., Wuest, G., Sixl, W., Schmidt, S., Haas, D. U & Marth, E. (2006). Transmission of *Campylobacter* spp. in a poultry slaughterhouse and genetic characterisation of the isolates by pulsed-field gel electrophoresis. *British Poultry Science*, 47(3), 286-293.

Pradhan, A. K., Van Kessel, J. S., Karns, J. S., Wolfgang, D. R., Hovingh, E., Nelen, K. A. & Schukken, Y. H. (2009). Dynamics of endemic infectious diseases of animal and human importance on three dairy herds in the northeastern United States. *Journal of Dairy Science*, 92(4), 1811-1825

Price, L. B., Johnson, E., Vailes, R., & Silbergeld, E. (2005). Fluoroquinolone-resistant *Campylobacter* isolates from conventional and antibiotic-free chicken products. *Environmental Health Perspectives*, 557-560.

Price, L. B., Lackey, L. G., Vailes, R., & Silbergeld, E. (2007). The persistence of fluoroquinolone-resistant *Campylobacter* in poultry production. *Environmental Health Perspectives*, 1035-1039.

Radu, S., Ling, O. W., Rusul, G., Karim, M. I. A., & Nishibuchi, M. (2001). Detection of *Escherichia coli* O157: H7 by multiplex PCR and their characterization by plasmid profiling, antimicrobial resistance, RAPD and PFGE analyses. *Journal of Microbiological Methods*, 46(2), 131-139.

- Ragimbeau, C., Schneider, F., Losch, S., Even, J. and Mossong, J. (2008) Multilocus sequence typing, pulsed-field gel electrophoresis, and fla short variable region typing of clonal complexes of *Campylobacter jejuni* strains of human, bovine, and poultry origins in Luxembourg. *Applied Environmental Microbiology* 74, 7715–7722.
- Rahman, M., Rahman, B. M., Rahman, B. (2008). Antibigram and Plasmid Profile Analysis of Isolated *Escherichia coli* from Broiler and Layer. *Research Journal of Microbiology*, 3, 82-90.
- Ram, S., Vajpayee, P., & Shanker, R. (2008). Rapid culture-independent quantitative detection of enterotoxigenic *Escherichia coli* in surface waters by real-time PCR with molecular beacon. *Environmental Science & Technology*, 42(12), 4577-4582.
- Ramabu, S. S., Boxall, N. S., Madie, P., & Fenwick, S. G. (2004). Some potential sources for transmission of *Campylobacter jejuni* to broiler chickens. *Letters in Applied Microbiology*, 39(3), 252-256.
- Rasschaert, G., Houf, K., Van Hende, J., & De Zutter, L. (2006). *Campylobacter* contamination during poultry slaughter in Belgium. *Journal of Food Protection* 69(1), 27-33.
- Refregier-Petton, J., N. Rose, M. Denis, and G. Salvat. (2001). Risk factors for *Campylobacter* spp. contamination in French broiler chicken flocks at the end of the rearing period. *Preventive Veterinary Medicine* 50:89–100.
- Reinhard, R., M.T. James, F.J. George, C.E. Robert, W.F. Robert, D.A. Alpha and F.Custy (1996): Analysis of *Campylobacter jejuni*, *Campylobacter coli*, *Salmonella*, *Klebsiella pneumoniae*, and *Escherichia coli* O157:H7 in Fresh Hand-Picked Blue Crab (*Callinectes sapidus*) Meat. *Journal of Food Protection*. 59(8): 803-807
- Rejab, S. B. M., Zessin, K H., Fries, R., & Patchanee, P. (2012). *Campylobacter* in chicken carcasses and slaughterhouses in Malaysia. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 43(1), 96–104.
- Rhoades, J.R., Duffy, G., Koutsoumanis, K., (2009). Prevalence and concentration of vero cytotoxigenic *Escherichia coli*, *Salmonella enterica* and *Listeria monocytogenes* in the beef production chain: a review. *Food Microbiology* 26, 357–376.
- Rodloff, A. C., Leclercq, R., Debbia, E. A., Cantón, R., Oppenheim, B. A., & Dowzicky, M. J. (2008). Comparative analysis of antimicrobial susceptibility among organisms from France, Germany, Italy, Spain and the UK as part of the tigecycline evaluation and surveillance trial. *Clinical Microbiology and Infection*, 14(4), 307-314.

- Rodrigo, S., Adesiyun, A., Asgarali, Z., & Swanston, W. (2007). Antimicrobial resistance of *Campylobacter* spp. isolated from broilers in small poultry processing operations in Trinidad. *Food Control*, 18(4), 321-325.
- Roels, T. H., Wickus, B., Bostrom, H. H., Kazmierczak, J. J., Nicholson, M. A., Kurzynski, T. A., & Davis, J. P. (1998). A foodborne outbreak of *Campylobacter jejuni* (O [ratio] 33) infection associated with tuna salad: a rare strain in an unusual vehicle. *Epidemiology and Infection*, 121(02), 281-287.
- Rollins, D. M., & Colwell, R. R. (1986). Viable but nonculturable (VBNC) stage of *Campylobacter jejuni* and its role in survival in the natural aquatic environment. *Applied and Environmental Microbiology*, 52(3), 531-8.
- Ronner, A. C., Engvall, E. O., Anderson, L. and Kaijser, B. (2004). Species identification by genotyping and determination of antibiotic resistance in *Campylobacter jejuni* and *Campylobacter coli* from humans and chickens in Sweden. *International Journal of Food Microbiology* 96:173-179.
- Rosenquist, H., Nielsen, N. L., Sommer, H. M., Nørrung, B., & Christensen, B. B. (2003). Quantitative risk assessment of human campylobacteriosis associated with thermophilic *Campylobacter* species in chickens. *International Journal of Food Microbiology*, 83(1), 87-103.
- Rozynek, E., Dzierżanowska-Fangrat, K., Korsak, D., Konieczny, P., Wardak, S., Szych, J & Dzierżanowska, D. (2008). Comparison of antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from humans and chicken carcasses in Poland. *Journal of Food Protection*, 71(3), 602-607.
- Russell S.M. (2009) Salmonella intervention strategies and testing methods. March 2009 issue of the University's poultry tips. <http://www.thepoultrysite.com/articles/1357/salmonella-intervention-strategies-and-testing-methods>
- Russell, R. G., O'Donnoghue, M., Blake, D. C., Zulty, J., & DeTolla, L. J. (1993). Early colonic damage and invasion of *Campylobacter jejuni* in experimentally challenged infant *Macaca mulatta*. *Journal of Infectious Diseases*, 168(1), 210-215.
- Russo, T. A., & Johnson, J. R. (2000). Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *Journal of Infectious Diseases*, 181(5), 1753-1754.
- Sa'enz, Y., Zarazaga, M., Lantero, M., Gastan˜ares, M.J., Baquero, F., Torres, C., (2000). Antimicrobial resistance in *Campylobacter* strains isolated from animals, foods, and humans in Spain in 1997 – 1998. *Antimicrobial Agents and Chemotherapy*. 44, 267 – 271.

- Sahilah, A. M. (1998). Plasmid profiling and antibiotic resistance of *Escherichia coli* and *E. coli O157* strains, Food Science and Biotechnology Department, *University Putra Malaysia*, Serdang Selangor.
- Sahin, O., Morishita, T. Y. and Zhang, Q. (2002). *Campylobacter* colonization in poultry: source of infection and modes of transmission. *Animal Health Research Reviews* 3:95-105.
- Saito, S., Yatsuyanagi, J., Harata, S., Ito, Y., Shinagawa, K., Suzuki, N., Amano, K. and Enomoto, K. (2005). *Campylobacter jejuni* isolated from retail poultry, bovine feces and bile, and human diarrheal samples in Japan: comparison of serotypes & genotypes. *FEMS Immunology and Medical Microbiology* 45: 311-319.
- Saleha, A. A. (2002). Isolation and characterization of *Campylobacter jejuni* from broiler chickens in Malaysia. *International Journal of Poultry Science* 1:94-97.
- Saleha, A. A. (2004). Epidemiological study on the colonization of chickens with *Campylobacter* in broiler farms in Malaysia: possible risk and management factors. *International Journal of Poultry Science*. 3: 129-134.
- Saleha, A. A., Christopher, J. R. G., Aini, I. and Ganapathy, K. (2001). Occurrence of *Campylobacter* in free-flying wild birds. Proceedings of 13th VAM Congress, *Kuala Lumpur*, Pp. 70-71.
- Salihu, M. D., Abdulkadir, J. U., Oboegbulem, S. I., Egwu, G. O., Magaji, A. A., Lawal, M., & Hassan, Y. (2009). Isolation and prevalence of *Campylobacter* species in cattle from Sokoto state, Nigeria. *Veterinaria Italiana*, 45(4), 501–5.
- Sallam, Kh.I. (2007). Prevalence of *Campylobacter* in chicken and chicken by-products retailed in Sapporo area, Hokkaido, Japan. *Food Control* 18: 1113-1120.
- Sanad, Y. M., Kassem, I. I., Abley, M., Gebreyes, W., LeJeune, J. T., & Rajashekara, G. (2011). Genotypic and phenotypic properties of cattle-associated *Campylobacter* and their implications to public health in the USA. *PloS one*, 6(10), 25778.
- Sato, K., Bartlett, P. C., Kaneene, J. B., & Downes, F. P. (2004). Comparison of prevalence and antimicrobial susceptibilities of *Campylobacter* spp. isolates from organic and conventional dairy herds in Wisconsin. *Applied and environmental microbiology*, 70(3), 1442-1447.
- Savill, M., Hudson, A., Devane, M., Garrett, N., Gilpin, B. and Ball, A. (2003) Elucidation of potential transmission routes of *Campylobacter* in New Zealand. *Water Science Technology* 47, 33–38.

- Scherer, K., Bartelt, E., Sommerfeld, C., & Hildebrandt, G. (2006b). Quantification of *Campylobacter* on the surface and in the muscle of chicken legs at retail. *Journal of Food Protection*, 69(4), 757-761.
- Schwarz, S., Chaslus-Dancla, E. (2001) Use of Antimicrobials in veterinary medicine and mechanisms of resistance. *Veterinary Research* 32: 201-225.
- Sengeløv, G., Agersø, Y., Halling-Sørensen, B., Baloda, S. B., Andersen, J. S., & Jensen, L. B. (2003). Bacterial antibiotic resistance levels in Danish farmland as a result of treatment with pig manure slurry. *Environment International* 28(7), 587-595.
- Shane, S.M. (1997): Campylobacteriosis. In: Diseases of Poultry. 10th ed., Calnek, B.W. (Ed.), Iowa State University Press, Ames. pp. 235-245
- Shar, A. H., Kazi, Y. F., Kanhar, N. A., Soomro, I. H., Zia, S. M. and Ghumro, P. B. (2010). Drinking water quality in Rohri City, Sindh, Pakistan. *African Journal of Biotechnology* 9 (42): 7102-7107.
- Shreeve, J. E., Toszeghy, M., Pattison, M., & Newell, D. G. (2000). Sequential spread of *Campylobacter* infection in a multipen broiler house. *Avian Diseases*, 983-988.
- Silva, J., Leite, D., Fernandes, M., Mena, C., Gibbs, P. A., & Teixeira, P. (2011). *Campylobacter* spp. as a foodborne pathogen: a review. *Frontiers in microbiology*, 2.
- Silva, J., Leite, D., Fernandes, M., Mena, C., Gibbs, P. A., & Teixeira, P. (2011). *Campylobacter* spp. as a foodborne pathogen: a review. *Frontiers in Microbiology*, 2.
- Singer, R. S., J. S. Jeffrey, T. E. Carpenter, C. L. Cooke, E. R. Atwill, W. O. Johnson, and D. C. Hirsh. (2000). Persistence of cellulitis-associated *Escherichia coli* DNA fingerprints in successive broiler chicken flocks. *Veterinary Microbiology* 75:59-71.
- Skirrow, M. B. (2006). John McFadyean and the centenary of the first isolation of *Campylobacter* species. *Clinical infectious diseases*, 43(9), 1213-1217.
- Skirrow, M. B., & Blaser, M. J. (2000). Clinical aspects of *Campylobacter* infection. *Campylobacter*, 2nd ed. ASM Press, Washington, DC, 69-88.
- Skov, M. N., Spencer, A. G., Hald, B., Petersen, L., Nauerby, B., Carstensen, B., & Madsen, M. (2004). The role of litter beetles as potential reservoir for *Salmonella enterica* and thermophilic *Campylobacter* spp. between broiler flocks. *Avian Diseases*, 48(1), 9-18.

- Smet, A., Martel, A., Persoons, D., Dewulf, J., Heyndrickx, M., Catry, B., Herman, L., Haesebrouck, F. et al. (2008) Diversity of extended-spectrum *b*-lactamases and class C *b*-lactamases among cloacal *Escherichia coli* in Belgian broiler farms. *Antimicrobial Agents Chemotherapy* 52, 1238–1243.
- Smith, J.L., Drum, D.J.V., Dai, Y., Kim, J.M., Sanchez, S., Maurer, J.J., et al. (2007) Impact of antimicrobial usage on antimicrobial resistance in commensal *Escherichia coli* strains colonizing broiler chickens. *Applied Environmental Microbiology* 73:1404-1414.
- Sneed, J., Strohbehn, C., Gilmore, S. A., & Mendonca, A. (2004). Microbiological evaluation of foodservice contact surfaces in Iowa assisted-living facilities. *Journal of the American Dietetic Association*, 104(11), 1722-1724
- Snelling, W. J., Matsuda, M., Moore, J. E., & Dooley, J. S. G. (2005). *Campylobacter jejuni*. *Letters in Applied Microbiology*, 41(4), 297-302.
- Son, R., Mutalib, S.A., Rusul, G., Ahmad, Z., Morigaki, T., Asai, N., Kim, Y.B., Okuda, J., Nishibuchi, M., (1998). Detection of *Escherichia coli* 0157:H7 in the beef marketed in Malaysia. *Applied Environmental Microbiology* 64, 1153–1156.
- Songer, J. G., & Post, K. W. (2005). The genera *Campylobacter*, *Helicobacter* and *Arcobacter*. *Veterinary Microbiology*, 223-231.
- Soonthornchaikul, N., Garelick, H., Jones, H., Jacobs, J., Ball, D., & Choudhury, M. (2006). Resistance to three antimicrobial agents of *Campylobacter* isolated from organically-and intensively-reared chickens purchased from retail outlets. *International Journal of Antimicrobial Agents*, 27(2), 125-130.
- Sovillo, F.J, L.E. LIEB and S.H. Waterman (1991): Incidence of campylobacteriosis among patients with AIDS in Los Angeles County. *Journal of Acquired Immune Deficiency Syndrome* 4(6): 598-602
- Speer, B. S., Shoemaker, N. B., & Salyers, A. A. (1992). Bacterial resistance to tetracycline: mechanisms, transfer, and clinical significance. *Clinical Microbiology Reviews*, 5(4), 387-399.
- Stanley K. & Jones K., (2003). Cattle and sheep as reservoirs of *Campylobacter*. *Journal of Applied Microbiology* 94: 104S-113S.
- Stanley, K. N., Wallace, J. S., Currie, J. E., Diggle, P. J., & Jones, K. (1998a). The seasonal variation of thermophilic *Campylobacter* in beef cattle, dairy cattle and calves. *Journal of Applied Microbiology*, 85(3), 472–80.
- Steele, M. L., McNab, W. B., Poppe, C., Griffiths, M. W., Chen, S., Degrandis, S. A., & Odumeru, J. A. (1997). Survey of Ontario bulk tank raw milk for food-borne pathogens. *Journal of Food Protection*, 60(11), 1341-1346.

- Stern, N. J., & Line, J. E. (1992). Comparison of three methods for recovery of *Campylobacter* spp. from broiler carcasses. *Journal of Food Protection*, 55(9), 663-666.
- Stern, N. J., Fedorka-Cray, P., Bailey, J. S., Cox, N. A., Craven, S. E., Hiatt, K. L. & Mead, G. C. (2001). Distribution of *Campylobacter* spp. in selected US poultry production and processing operations. *Journal of Food Protection*, 64(11), 1705-1710.
- Sukhumungoon, P., Mittraparp-Arthorn, P., Pomwised, R., Chareunjiratragul, W & Vuddhakul, V. (2011). High concentration of Shiga toxin 1-producing *Escherichia coli* isolated from Southern Thailand. *International Food Research Journal* 18: 683-688.
- Takkinen, J., Ammon, A., Robstad, O. & Breuer, T. (2003). European survey on *Campylobacter* surveillance and diagnosis 2001. *European Surveillance* 8: 207-213.
- Tambur, Z., Stojanov, I., Konstantinovic, S., Jovanovic, D., Cenic-Milosevic, D. and Opacic, D. (2010). Multidrug resistance of *Campylobacter jejuni* and *Campylobacter coli* to tested antibiotics in strains originating from humans, poultry and swine. *Zbornik Matice Srpske Zaprirodne Nauke* 118: 27-35.
- Tan, S. L., Lee, H. Y., Abu Bakar, F., Abdul Karim, M. S., Rukayadi, Y. and Mahyudin, N. A., (2013). Microbiological quality on food handlers' hands at primary schools in Hulu Langat District, Malaysia. *International Food Research Journal* 20 (5): 2973- 2977.
- Tang, J. Y. H., Mohamad Ghazali, F., Saleha, A. A., Nishibuchi, M. and Son, R. (2009). Comparison of thermophilic *Campylobacter* spp. occurrence in two types of retail chicken samples. *International Food Research Journal* 16:277-288.
- Tang, J. Y. H., Saleha, A. A., Jalila, A., Mohamad Ghazali, F., Tuan Chilek, T. Z., Ahmad, N., Sandra, A., Nishibuchi, M. and Son, R. (2010). Thermophilic *Campylobacter* spp. occurrence on chickens at farm, slaughter house and retail. *International Journal of Poultry Science* 9:134-138.
- Taremi, M., Mehdi Soltan Dallal, M., Gachkar, L., MoezArdalan, S., Zolfagharian, K., & Reza Zali, M. (2006). Prevalence and antimicrobial resistance of *Campylobacter* isolated from retail raw chicken and beef meat, Tehran, Iran. *International Journal of Food Microbiology*, 108(3), 401-403.
- Tauxe, R. (1992). Epidemiology of *Campylobacter jejuni* infections in the United States and other industrialized nations, p. 9-19. In I. Nachamkin, M. Blaser, and L. Tompkins (ed.), *Campylobacter jejuni: current status and future trends*. American Society for Microbiology, Washington, D.C.

- Teguh S.S., (2014). Characterisation of emerging *Campylobacter* species from broiler chicken and chicken meat in Selangor and their antibiotic resistance. MVSc Thesis: *Faculty of Veterinary Medicine, Universiti Putra Malaysia* (Malaysia).
- Tenaillon, O., Skurnik, D., Picard, B., & Denamur, E. (2010). The population genetics of commensal *Escherichia coli*. *Nature Reviews Microbiology*, 8(3), 207-217.
- Thorsteinsdottir, T. R., Kristinsson, K. G., Fridriksdottir, V., & Gunnarsson, E. (2008). Antimicrobial resistance of *Campylobacter* spp. isolated from broiler flocks in Iceland 2001-2005. *Microbial Drug Resistance*, 14(1), 49-53.
- Trieber, C.A., Taylor, D.E., (2000). Mechanisms of Antibiotic resistance in *Campylobacter*. In: Nachamkin, I., Blaser, M.J. (Eds.), *Campylobacter. American Society for Microbiology, Washington, DC*, pp. 441–454.
- Twaites, R. T. and Frost, J. A. (1997). Drug resistance in *Campylobacter jejuni*, *C. coli* and *C. lari* isolated from humans in North West England and Wales. *Journal of Clinical Pathology* 52:812-814.
- Uaboi-Egbenni, P. O., Bessong, P. O., Samie, A., & Obi, C. L. (2012). Potentially pathogenic *Campylobacter* species among farm animals in rural areas of Limpopo province, South Africa: A case study of chickens and cattles. *African Journal of Microbiology Research*, 6(12), 2835–2843.
- Unicomb, L. E., Dalton, C. B., Gilbert, G. L., Becker, N. G., & Patel, M. S. (2008). Age-specific risk factors for sporadic *Campylobacter* infection in regional Australia. *Foodborne Pathogens and Disease*, 5(1), 79-85.
- Ursinitsch, B., Pless, P., & Kofer, J. (2002). Prevalence and Resistance of *Campylobacter* spp in Styrian Poultry meat. *Food Safety Assurance and Veterinary Public Health. ed. Smulders FJ, Collins JD*, 2, 380-382.
- Usha, M. R., Fauziah, M., Tunung, R., Chai, L. C., Cheah, Y. K., Farinazleen, M. G., et al. (2010). Occurrence and antibiotic resistance of *Campylobacter jejuni* and *Campylobacter coli* in retail broiler chicken. *International Food Research Journal*, 17, 247-255.
- Van de Giessen, A. W., Tilburg, J. J. H. C., Ritmeester, W. S., & Van Der Plas, J. (1998). Reduction of *Campylobacter* infections in broiler flocks by application of hygiene measures. *Epidemiology and Infection*, 121(01), 57-66.
- Van den Bogaard, A. E., London, N., Driessen, C., & Stobberingh, E. E. (2001). Antibiotic resistance of faecal *Escherichia coli* in poultry, poultry farmers and poultry slaughterers. *Journal of Antimicrobial Chemotherapy*, 47(6), 763-771.

- Van Looveren, M., Daube, G., De Zutter, L., Dumont, J. M., Lammens, C., Wijdooghe, M., & Goossens, H. (2001). Antimicrobial susceptibilities of *Campylobacter* strains isolated from food animals in Belgium. *Journal of Antimicrobial Chemotherapy*, 48(2), 235-240.
- Van Vliet, A. H., & Ketley, J. M. (2001). Pathogenesis of enteric *Campylobacter* infection. *Symposium Series (Society for Applied Microbiology)*, 90, 45S–56S.
- Vandamme, P. (2000). “Taxonomy of the family *Campylobacteraceae*,” in *Campylobacter*, I.Namchamkin & M.J. Blaser (Washington, DC: ASM), 3-27.
- Varga, J., Mezes, B., Fodor, L., (1990). Serogroups of *Campylobacter jejuni* from man and animals. *Journal of Veterinary Medicine Series B* 37 (6), 407-411
- Velayudhan, J., & Kelly, D. J. (2002). Analysis of gluconeogenic and anaplerotic enzymes in *Campylobacter jejuni*: an essential role for phosphoenolpyruvate carboxykinase. *Microbiology (Reading, England)*, 148(Pt 3), 685–94.
- Veron, M. and R. Chatelaine (1973): Taxonomic study of the genus *Campylobacter* Sebald and Veron and designation of the neotype strain for the type species, *Campylobacter fetus* (Smith and Taylor) Sebald and Veron. *International Journal of Systemic Bacteriology*. 23(2): 122-134
- Waino, M, Bang, D. D., Lund, M., Nordentoft, S., Andersen, J. S., Pedersen, K. and Madsen, M. (2003). Identification of *Campylobacter* isolated from Danish broilers by phenotypic tests and species-specific PCR assays. *Journal of Applied Microbiology* 95:649-655.
- Wang, G., Clark, C. G., & Rodgers, F. G. (2002). Detection in *Escherichia coli* of the genes encoding the major virulence factors, the genes defining the O157: H7 serotype, and components of the type 2 Shiga toxin family by multiplex PCR. *Journal of Clinical Microbiology*, 40(10), 3613-3619.
- Weese, J. S. (2011). Bacterial enteritis in dogs and cats: diagnosis, therapy, and zoonotic potential. *The Veterinary Clinics of North America. Small Animal Practice*, 41(2), 287.
- Weese, J. S., & van Duijkeren, E. (2010). Methicillin-resistant *Staphylococcus aureus* and *Staphylococcus pseudintermedius* in veterinary medicine. *Veterinary Microbiology*, 140(3), 418-429.
- Wesley, I. V., Wells, S. J., Harmon, K. M., Green, A., Schroeder-Tucker, L., Glover, M., & Siddique, I. (2000). Fecal Shedding of *Campylobacter* and *Arcobacter* spp. in Dairy Cattle. *Applied and Environmental Microbiology*, 66(5), 1994-2000.

- Whyte, P., McGill, K., Cowley, D., Madden, R.H., Moran, L., Scates, P., Carrol, C., O'Leary, A., Fanning, S., Collins, J. D., McNamara, E., Moore, J. E. and Cormican, M. (2004). Occurrence of *Campylobacter* in retail foods in Ireland. *International Journal of Food Microbiology* 95: 111-118.
- Wieczorek, K., Szewczyk, R., & Osek, J. (2012). Prevalence, antimicrobial resistance, and molecular characterization of *Campylobacter jejuni* and *C. coli* isolated from retail raw meat in Poland. *Veterinary Medicine-Czech*, 57, 293-9.
- Wieland, B., Regula, G., Danuser, J., Wittwer, M., Burnens, A. P., Wassenaar, T. M., & Stärk, K. D. C. (2005). *Campylobacter* spp. in dogs and cats in Switzerland: risk factor analysis and molecular characterization with AFLP. *Journal of Veterinary Medicine, Series B*, 52(4), 183-189.
- Wilson, I. G. (2002). *Salmonella* and *Campylobacter* contamination of raw retail chickens from different producers: a six year survey. *Epidemiology and Infection*, 129(03), 635-645.
- Winfield, M. D., & Groisman, E. A. (2003). Role of nonhost environments in the lifestyles of *Salmonella* and *Escherichia coli*. *Applied and Environmental Microbiology*, 69(7), 3687-3694
- Wong, T. L., Hollis, L., Cornelius, A., Nicol, C., Cook, R. and Hudson, J. A. (2007). Prevalence, numbers, and subtypes of *Campylobacter jejuni* and *Campylobacter coli* in uncooked retail meat samples. *Journal of Food Protection* 70:566-573.
- World Health Organization (WHO) (2000). Food-borne disease: a focus for health education. 1st edition. *World Health Organization*, Geneva
- World Health Organization (WHO) (2011). *Campylobacter*. Fact sheet N°255. Available from: <http://www.who.int/mediacentre/factsheets/fs255/en/>
- World Health Organization (WHO) (2013). The global view of campylobacteriosis- Report of expert consultation. *World Health Organization*. Geneva
- World Health Organization. (WHO) (2001). The increasing incidence of human campylobacteriosis report and proceedings of a WHO consultation of experts, *Copenhagen, Denmark*, 21-25 November 2000.
- Yamazaki-Matsune, W., Taguchi, M., Seto, K., Kawahara, R., Kawatsu, K., Kumeda, Y., Kitazato, M., Nukina, M., Misawa, N. and Tsukamoto, T. (2007). Development of a multiplex PCR assay for identification of *Campylobacter coli*, *Campylobacter fetus*, *Campylobacter hyointestinalis* subsp. *Hyointestinalis*, *Campylobacter jejuni*, *Campylobacter lari* and *Campylobacter upsaliensis*. *Journal of Medical Microbiology* 56:1467-1473

- Yan, S. S., Pendrak, M. L., Foley, S. L., & Powers, J. H. (2005). *Campylobacter* infection and Guillain-Barré syndrome: public health concerns from a microbial food safety perspective. *Clinical and Applied Immunology Reviews*, 5(5), 285-305
- Yang, C., Jiang, Y., Huang, K., Zhu, C., & Yin, Y. (2003). Application of real-time PCR for quantitative detection of *Campylobacter jejuni* in poultry, milk and environmental water. *FEMS Immunology & Medical Microbiology*, 38(3), 265-271.
- Yoke Kqueen, C., Learn, Han, L., Noorzaleha, A. S., Son, R., Sabrina, S., Jiun Horng, S., & Chaiv, Hoon, K. (2008). Characterization of multiple antimicrobial resistant *Salmonella enterica Subsp. enterica* isolated from indigenous vegetables and poultry in Malaysia. *Letters in Applied Microbiology*, 46(3), 318-324.
- Young, C.R., Harvey, R., Anderson, R., Nisbet, D., Stanker, L.H., (2000). Enteric colonisation following natural exposure to *Campylobacter* in pigs. *Research Veterinary Sciences* 68: 75– 78.
- Young, K. T., Davis, L. M., & DiRita, V. J. (2007). *Campylobacter jejuni*: Molecular biology and pathogenesis. *Nature Reviews Microbiology*, 5(9), 665-679.
- Yuan, L., Liu, J. H., Hu, G. Z., Pan, Y. S., Liu, Z. M., Mo, J., & Wei, Y. J. (2009). Molecular characterization of extended-spectrum β -lactamase-producing *Escherichia coli* isolates from chickens in Henan Province, China. *Journal of Medical Microbiology*, 58(11), 1449-1453.
- Zaliha, I. and A.B. Rusli, (2004). A study on hygienic standard of food premises and microbiological quality of food in Kota Bharu. Working Paper. *Universiti Sains Malaysia*.
- Zhang, Q. and Plummer, P. J. (2008). Mechanisms of antibiotic resistance in *Campylobacter*. . In *Campylobacter*, 3rd ed. Edited by Nachakim, I., Szymanski, C. M., and Blaser, M. J. pp. 263-272. *ASM Press*: Washington.
- Zhang, R.; Eggleston, K.; Rotimi, V.; Zeckhauser, R.J. (2006). Antibiotic resistance as a global threat: Evidence from China, Kuwait and the United States. *Global Health*, 2:6.
- Zhao, C., Ge, B., De Villena, J., Sudler, R., Yeh, E., Zhao, S & Meng, J. (2001). Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the Greater Washington, DC, area. *Applied and Environmental Microbiology*, 67(12), 5431-5436.
- Zhao, S., Maurer, J. J., Hubert, S., De Villena, J. F., McDermott, P. F., Meng, J & White, D. G. (2005). Antimicrobial susceptibility and molecular characterization of avian pathogenic *Escherichia coli* isolates. *Veterinary Microbiology*, 107(3), 215-224.

Zoran, T., Biljana, M. S., Radoje, D., & Zoran, K. (2010). Susceptibility of *Campylobacter jejuni* and *Campylobacter coli* isolated from animals and humans to tetracycline. *African Journal of Microbiology Research*, 4(12), 1246-1250.





© COP YRIGHT UPM