



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF CONJUGATED LINOLEIC ACID ON ADIPOGENIC GENES
REGULATION IN CHICKENS***

SURIYA KUMARI A/P RAMIAH

FPV 2015 11



**EFFECTS OF CONJUGATED LINOLEIC ACID ON ADIPOGENIC GENES
REGULATION IN CHICKENS**

By

SURIYA KUMARI A/P RAMIAH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2015

© COPYRIGHT UPM



COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

TO MY LATE FATHER, MR.RAMIAH

I Love You Forever



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Degree of Doctor of Philosophy

EFFECTS OF CONJUGATED LINOLEIC ACID ON ADIPOGENIC GENES REGULATION IN CHICKENS

By

SURIYA KUMARI A/P RAMIAH

December 2015

Chairman : Associate Professor Goh Yong Meng, PhD
Faculty : Veterinary Medicine

Modern commercial chickens exhibit excessive fat accumulation in the abdominal area. The major goals of the poultry industry are to increase the carcass yield and to reduce carcass fatness. Excessive fat deposition in chicken is detrimental for human consumption. Increasing the leanness of meat will improve the meat quality, and thus makes animals more valuable. Conjugated linoleic acid (CLA) consists of a complex mixture of geometrical (cis and trans) and positional isomers which consist of two major isomers; cis-9, trans-11 CLA and trans-10, cis-12 CLA. Animal studies have reported that dietary intake of CLA changes animal body composition by preventing obesity development. This study was performed with the hypothesis that CLA modified adipose tissue through alteration of the adipocyte genes. Therefore the general objective of current study was to evaluate the effects of CLA on lipid metabolism, cellular morphology and transcription of regulatory genes that were involved in adipogenesis of chickens. Cis-9, trans-11 CLA and trans-10, cis-12 CLA isomers were evaluated individually for their effects on morphological changes, and adipogenic genes expressions on primary adipose tissue isolated from chicken. Adipose tissue isolated from specific pathogen-free (SPF) chicken was cultured in induction media containing Dulbecco's Modified Eagle Medium (DMEM: HEM's (50:50); 1.5% bovine serum albumin; 100mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid); 2mg/mL collagenase type I, and 1% penicillin/streptomycin. The media were incorporated with two concentrations of both CLA isomers at 1.51% and 2.56% with a CONTROL group (without CLA isomers). After day 7, adipose tissues differentiation was monitored morphologically using ImageJ software, and adipogenic genes expressions were analyzed by real time polymerase chain reaction (PCR). It was observed that the efficacy of cis-9, trans-11 CLA isomer was more pronounced than trans-10, cis-12 CLA isomer in the in vitro study. Adipose tissue morphology data presented in this work revealed that the domination of cis-9, trans-11 isomer CLA effect was observed at higher concentration in the abdominal fat of broiler chickens. This was associated with a lower transcriptional level of peroxisome proliferator-activated receptor gamma (PPAR γ), and adipocyte protein 2 (aP2), together with lesser abdominal adipocyte volume and smaller amount of fat. The acyl-Coenzyme A binding domain containing 5 (ACBD 5), and lipoprotein lipase (LPL) were down-regulated by cis-9, trans-11 CLA isomer. The primary adipose tissue treated with trans-10, cis-12 concentrations had no changes on the adipose cellularity and adipogenic genes. The

influences of CLA supplementation on growth performance, fatty acid composition, lipid peroxidation, meat colour and plasma lipids in broiler chicken were investigated in this study. The CLA used in this study was of commercial feed grade (Lutrell® BSAF, SE, Ludwigshafen, Germany). A total of 180 broiler chickens were allocated to 3 dietary treatments (0, 2.5 and 5% CLA), and given a standard broiler starter diet from 1 to 21 days, and finisher diet from 22 to 42 days. Body weight of chickens and feed intake were recorded weekly. After slaughter, the breast meat was aged at 4°C for 0, 3 and 6 days. The fatty acid composition was measured in the breast meat. The dietary CLA supplementation significantly ($P < 0.05$) increased the content of CLA in chicken meat. The predominant CLA in meat from birds with supplemented diets was trans-10, cis-9. The proportion of monounsaturated (MUFA) fatty acid in meat decreased significantly ($P < 0.05$) with increasing CLA supplementation. Dietary CLA also increased the thiobarbituric acid reactive substances (TBARS) values in breast meat. Conjugated linoleic acid feeding also resulted in the reduction of plasma total cholesterol, low-density protein, and the ratio of high-density protein, particularly among the 5% CLA fed-chickens. The effects of CLA on adipocytes morphology, fatty acid profile and PPARs associated genes were performed in CLA-fed chickens. The adipocyte morphology was analyzed using ImageJ software. It was also observed that the content of cis-9, trans-11 CLA in abdominal fat tissue was higher than that of trans-10, cis-12 CLA. The CLA feeding increased total saturated fatty acid and decreased the MUFA in concentration in broilers compared to the control group. Conjugated linoleic acid fed-chickens have lesser mean abdominal adipocyte volume and a smaller amount of fat because of reduced capacities to store fats. The influence on body composition appears to be dependent on PPAR α (alpha). Current study demonstrated that CLA down-regulated aP2 transcription, which was parallel to PPAR γ transcription in adipose tissue of broiler chickens. Adipocyte protein 2 (aP2) was regulated by PPAR γ as PPAR γ is regarded as „master regulator“ of adipocyte differentiation. The upregulation of LPL suggested that this gene might play an independent role in regulation of adipogenesis in chicken model. The upregulation of ACBD 5 seems to be novel; therefore, more studies need to be done on this gene. However, LPL gene was not altered which indicates that the expression of LPL occurs spontaneously at confluence and is independent. Therefore, the current results supported the hypotheses that CLA modified adipose tissue through alteration of the adipocyte genes. It is evident that the CLA down-regulated PPAR γ and aP2, which subsequently resulted in decreased adipocyte size, number and area of abdominal fat cells. The comparatively lower CLA content in poultry species could be mitigated through the use of feed and biotechnology approach to enhance CLA content. In conclusion, the present results proved that CLA feeding is a practical strategy to reduce fat deposition in broiler chicken, while increasing its appeal to the general populace consuming chicken meat.

Abstrak thesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

KESAN ASID LINOLEIK TERKONJUGAT KE ATAS PENGAWALATURAN GEN ADIPOS PADA AYAM

Oleh

SURIYA KUMARI A/P RAMIAH

Disember 2015

Pengerusi : Profesor Madya Goh Yong Meng, PhD
Fakulti : Perubatan Veterinar

Ayam komersial moden mempunyai lemak berlebihan di bahagian abdomen. Matlamat utama industri ternakan ayam adalah untuk meningkatkan hasil karkas and mengurangkan lemak karkas bagi meningkatkan margin keuntungan penternak. Selain itu, lemak yang berlebihan pada ayam akan memudaratkan kesihatan manusia. Pengurangan lemak daging bukan sahaja meningkatkan kualiti daging, malah ia juga akan menjadikan hasil daging ternakan lebih bernilai. Asid linoleik terkonjugat (CLA) terdiri daripada campuran kompleks isomer, cis dan trans. Terdapat dua isomer utama CLA yakni, cis-9, trans-11 CLA dan trans-10, cis-12 CLA. Kajian haiwan telah menunjukkan bahawa pengambilan CLA mengubah komposisi badan haiwan, dan berkesan untuk mencegah obesiti. Kajian ini telah dilaksanakan dengan hipotesis bahawa induksi gen adipos akibat pendedahan CLA akan memberi kesan kepada pengubahsuaian tisu adipos Oleh itu objektif kajian ini adalah untuk mengkaji kesan CLA pada metabolisme lemak, morfologi selular dan transkripsi gen kawal selia yang terlibat dalam pengawalaturan tisu adipos ayam. Cis-9, trans-11 CLA dan trans-10, cis-12 CLA isomers telah dinilai secara individu untuk menentukan kesan perubahan morfologi dan ekspresi adipos gen, pada kultur primer dari tisu adipos ayam. Tisu adipos yang diasingkan dari sampel ayam dikultur dalam media induksi yang mengandungi Dulbecco's Modified Eagle Medium (DMEM:HEM's (50:50); 1.5% albumin serum lembu; 100mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid); 2mg/mL collagenease jenis I, dan 1% penicillin/streptomycin. Media tersebut dicampur dengan dua konsentrasi CLA isomer pada 1.51% dan 2.56% dengan kumpulan kawalan (CONTROL) (tanpa CLA isomer). Selepas hari ke-7, pembezaan morfologi tisu adipos dipantau dengan menggunakan perisian ImageJ, dan ekspresi gen adipos dianalisis dengan teknik PCR. Isomer CLA cis-9, trans-11 menunjukkan kesan *in vitro* yang lebih ketara berbanding isomer trans -10, cis 12 CLA. Data morfologi tisu adipos kajian ini menunjukkan bahawa konsentrasi cis-9, trans-11 isomer CLA yang tinggi dikaitkan dengan tahap transkripsi reseptor peridi peroksisom gamma (PPAR γ), dan protein adiposit (aP2) yang rendah, serta jumlah adipos tissue yang minium dan bersaiz kecil. Aktiviti gen ACBD 5 (Acyl-Coenzyme A binding domain containing 5) dan lipoprotein lipase (LPL) nyata susut dengan kehadiran cis-9, trans-11 isomer CLA. Kultur primer tisu adipos ayam yang didedahkan kepada isomer trans-10, cis-12 CLA tidak menunjukkan perubahan dalam pertumbuhan adipos and gen adipos. Pengaruh suplemen CLA ke atas prestasi pertumbuhan, komposisi asid lemak, peroksidasi lipid, warna daging dan lipid plasma turut dikaji. Asid linoleik terkonjugat yang digunakan

dalam kajian ini adalah gred makanan komersial (Lutrell® BSAF, SE, Ludwigshafen, Germany). Sebanyak 180 ekor ayam pedaging diberi makan diet yang mengandungi 3 kumpulan rawatan, terdiri daripada 0, 2.5 and 5% CLA). Ayam tersebut diberikan diet ayam permulaan dari hari 1 hingga 21, dan diet penghabisan dari hari 22 hingga 42. Berat badan ayam dan pengambilan makanan direkodkan setiap minggu. Selepas disembelih, daging dada ayam disimpan pada suhu 4° C untuk 0, 3, dan 6 hari. Komposisi asid lemak daging juga dianalisis. Berat badan dan tahap pengambilan makanan bagi ayam pedaging berkurangan dengan peningkatan tahap CLA. Supplementasi CLA dalam pemakanan meningkatkan kandungan CLA daging ayam dengan signifikan ($P < 0.05$). Isomer trans-10, cis-9 CLA merupakan isomer yang utama dalam daging dada ayam. Kadar asid lemak monotaktepung (MUFA) dalam daging berkurangan dengan penambahan CLA dengan signifikan ($P < 0.05$). Rawatan dengan CLA juga meningkatkan nilai TBARS reaktif dalam daging. Di samping itu, rawatan CLA juga membawa kepada pengurangan jumlah kolesterol plasma, dan lipoprotein berketumpatan rendah (LDL), serta nisbah lipoprotein berketumpatan tinggi (HDL) kepada LDL, terutamanya pada kumpulan ayam yang menerima rawatan 5% CLA. Kesan CLA ke atas morfologi adipos, profil asid lemak dan reseptor peridi peroksisom (PPARs) juga dicerap pada ayam yang menerima rawatan CLA. Morfologi tisu adipos dianalisa menggunakan perisian ImageJ. Kandungan isomer cis-9, trans-11 CLA didapati lebih tinggi daripada isomer trans-10, cis-12 CLA di dalam lemak abdomen ayam. Supplementasi CLA meningkatkan jumlah kandungan asid lemak tepu, serta mengurangkan tahap MUFA berbanding diet kawalan. Ayam yang diberi CLA mempunyai jumlah sel adipos yang minimum dan saiz lemak yang kecil, disebabkan oleh keupayaannya yang lebih rendah dalam penyimpanan lemak. Komposisi badan ayam pedaging didapati tidak mempunyai kaitan signifikan dengan PPAR α (alpha). Kajian ini menunjukkan bahawa CLA mengurangkan aktiviti transkripsi aP2 selari dengan aktiviti transkripsi PPAR γ dalam tisu adipos ayam pedaging. Gen protein adiposit (aP2) dikawal oleh PPAR γ dan dianggap sebagai pengawalatur utama bagi pertumbuhan adipos. Peningkatan aktiviti LPL menunjukkan bahawa gen ini tidak memberi sebarang kesan ke atas tumbesaran adipos dalam model ayam. Peningkatan aktiviti transkripsi ACBD 5 merupakan penemuan baharu dalam kajian ini. Oleh itu, lebih banyak kajian lanjutan perlu dilakukan ke atas gen ini. Walau bagaimanapun, tahap ekspresi gen LPL adalah tidak berubah. Keputusan ini menyokong hipotesis bahawa CLA mengakibatkan pengubahsuaian tisu adipos melalui kesannya ke atas gen adipos. Adalah didapati juga bahawa CLA mengurangkan aktiviti PPAR γ and aP2. Ini seterusnya mengakibatkan pengurangan pada saiz adipos, bilangan dan liputan sel lemak abdomen. Kesimpulannya, kandungan CLA dalam ayam boleh diperkayakan melalui penggunaan suplemen CLA, serta kaedah bioteknologi. Keputusan kajian ini membuktikan bahawa penggunaan CLA dalam makanan ialah satu strategi praktikal untuk mengurangkan pengumpulan lemak pada ayam pedaging, di samping memberi manfaat kesihatan kepada pengguna.

ACKNOWLEDGEMENTS

The successful completion of this thesis was made possible through the invaluable contribution of a number of people. To say “thank you” to all of you is not even enough to express my thankfulness. I would like to express my gratitude to my supervisor, Assoc. Prof. Dr. Goh Yong Meng for his expertise, understanding and patience. I appreciate his vast knowledge and skill in many areas and his assistance in writing publications. I doubt that I will ever be able to convey my appreciation fully, but I owe him my eternal gratitude.

To Dr. Mahdi Ebrahimi who provided me with direction, technical support and became more of a mentor and friend, than a co-supervisor. I would like to thank the other members of my committee, Dr.Tan Seau Wei, Dr.Yeap Swee Keong and Dr. Hafandi Ahmad for the assistance they provided at all levels of the research project. Finally, I would like to thank my director Prof. Zulkifli Idrus for encouraging me to further my studies.

To my father Mr. Ramiah and my mother Mrs. Kamalambal, who have been the light of my life and have given me the extra strength and motivation to get things done. This thesis is dedicated to my parents. A special thanks to my dearest sisters Sharmila, Indra and Rajeswary for support unconditional love, advices and constant encouragements throughout the course of my studies.

To my ever supportive soul mate, my dearest husband and my best friend, Pubaalan, thank you for helping me with my thesis, no matter how technical this is for you. Thank you for the undying love and support you have provided me and for believing in me that I can finish my manuscript on time. I also dedicate this PhD thesis to my three children, Niklesh, Shuveythaa and Vinoshan Mudhaliar who are the pride and joy of my life. I love you more than anything and I appreciate all your patience and support during mommy’s PhD studies.

I recognize that this research would not have been possible without the financial assistance of Malaysian Government Fundamental Research Grant Scheme (FRGS) under grant number 5450632. Last but not least, I am also very grateful for the logistical and personnel support from the Faculty of Veterinary Medicine, and Institute of Tropical Agriculture, Universiti Putra Malaysia.

I certify that a Thesis Examination Committee has met on 16 December 2015 to conduct the final examination of Suriya Kumari a/p Ramiah on her thesis entitled "Effects of Conjugated Linoleic Acid on Adipogenic Genes Regulation in Chickens" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Mohamed Ariff bin Omar, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Norhani binti Abdullah, PhD

Professor
Institute of Tropical Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Jalila binti Abu, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Andrew J. Sinclair, PhD

Professor
Deakin University
Australia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 16 February 2016

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Goh Yong Meng, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Hafandi Bin Ahmad, PhD

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Tan Sheau Wei, PhD

Research Officer
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Yeap Swee Keong, PhD

Research Fellow
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Mahdi Ebrahimi, PhD

Post Doctorate
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree At Any Other Institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric Number: Suriya Kumari a/p Ramiah GS 35018

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Goh Yong Meng, PhD

Signature: _____
Name of
Member of
Supervisory
Committee: Hafandi Bin Ahmad, PhD

Signature: _____
Name of
Member of
Supervisor
Committee: Tan Sheau Wei, PhD

Signature: _____
Name of
Member of
Supervisor
Committee: Yeap Swee Keong, PhD

Signature: _____
Name of
Member of
Supervisor
Committee: Mahdi Ebrahimi, PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Introduction	4
2.2 Metabolism of CLAs	6
2.3 The impact of CLAs on human health	7
2.4 Overview of global poultry production	7
2.5 Fat deposition in poultry	8
2.6 Approach to provide a nutritionally enhanced poultry meat source to consumers	8
2.7 CLAs altered adipose cellularity	10
2.8 Alteration of CLAs on fatty acid profiles in poultry products	10
2.9 Effects of CLAs in avian species	11
2.9.1 Laying hens	11
2.9.2 Ducks	12
2.9.3 Geese	13
2.9.4 Quails	13
2.10 CLAs as antioxidant	13
2.11 CLAs as anticarcinogenic agent	14
2.12 Transcriptional regulation of genes by CLAs	16
2.13 Summary and Conclusion	16
3 THE EFFECTS OF CONJUGATED LINOLEIC ACID ISOMERS ON THE MORPHOLOGICAL CHANGES IN ADIPOSE TISSUE AND ADIPOGENIC GENES EXPRESSION ON PRIMARY ADIPOSE TISSUE	18
3.1 Introduction	18
3.2 Material and methods	19
3.2.1 Experimental birds	19
3.2.2 Preparation of CLAs	19
3.2.3 Animal welfare	19
3.2.4 Isolation of adipose tissue and primary culture of chicken adipocytes	20
3.2.5 Trizol RNA extraction	20
3.2.6 Lipid staining	20

	3.2.7	Real-time polymerase chain reaction (PCR)	21
	3.2.8	Statistical analysis	21
3.3	Results		22
	3.3.1	Morphological changes of adipose tissue by cis-9, trans-11 and trans-10, cis-12 CLA isomers	22
	3.3.2	Transcriptional expression level of aP2, LPL and ACBD5 from adipose tissue isolated from broiler chicken	24
3.4	Discussion		27
	3.4.1	Morphological changes of adipose tissue by CLA isomers	27
	3.4.2	Effects of transcriptional expression of aP2, LPL and ACBD 5 by CLA isomers	28
3.5	Conclusion		29
4	GROWTH PERFORMANCE, FATTY ACID PROFILES OF BREAST MUSCLE, MEAT COLOUR, LIPID PEROXIDATION AND PLASMA LIPID PROFILE IN BROILER CHICKEN FED WITH CONJUGATED LINOLEIC ACID		30
	4.1	Introduction	30
	4.2	Material and methods	31
	4.2.1	Animal welfare	31
	4.2.2	Experimental birds and diets	31
	4.2.3	Fatty acid extraction	34
	4.2.4	Fatty acid analysis	34
	4.2.5	Meat colour	34
	4.2.6	Lipid peroxidation	34
	4.2.7	Plasma lipids profile	35
	4.2.8	Statistical analysis	35
	4.3	Results	35
	4.3.1	Growth performances	35
	4.3.2	Fatty acid composition of breast meat	36
	4.3.3	Meat colour	38
	4.3.4	Lipid peroxidation	38
	4.3.5	Plasma lipid profile	39
	4.4	Discussion	40
	4.4.1	Growth performances	40
	4.4.2	Fatty acid composition of breast meat	41
	4.4.3	Meat colour	42
	4.4.4	Lipid peroxidation	42
	4.4.5	Plasma lipids profile	43
	4.5	Conclusion	43

5	THE EFFECTS OF CONJUGATED LINOLEIC ACID ON THE MORPHOLOGY OF ADIPOSE TISSUE AND THE EXPRESSION OF PPARS AND ASSOCIATED GENES IN BROILER CHICKENS	44
5.1	Introduction	44
5.2	Material and methods	45
5.2.1	Animal welfare	45
5.2.2	Experimental birds and diets	45
5.2.3	Total lipid extraction	45
5.2.4	Fatty acid analysis	46
5.2.5	RNA extraction	46
5.2.6	Real-time polymerase chain reaction	46
5.2.7	Isolation and culture of adipocyte cells	48
5.2.8	Determination of abdominal fat cellularity	48
5.2.9	Statistical analysis	48
5.3	Results	48
5.3.1	Fatty acid composition of the abdominal fat	48
5.3.2	Abdominal fat expression of PPARs, aP2, ACBD 5 and LPL transcriptional levels	50
5.3.3	Abdominal fat cellularity	54
5.4	Discussion	57
5.4.1	Fatty acid composition of the abdominal fat	57
5.4.2	Abdominal fat expression of PPARs, aP2, ACBD 5 and LPL transcriptional levels	58
5.4.3	Abdominal fat cellularity	59
5.5	Conclusion	61
6	GENERAL DISCUSSION	62
7	CONCLUSIONS	66
7.1	Limitations of the study	67
7.2	Recommendations and future research	67
	REFERENCES	68
	BIODATA OF STUDENT	91
	LIST OF PUBLICATIONS	92

LIST OF TABLES

Table		Page
2.1	CLAs of various foods	5
2.2	Mean CLAs content in various poultry products	9
2.3	Mean CLAs content in various poultry species	10
2.4	Effects of CLAs on egg production of laying hens	12
4.1	Ingredients and chemical compositions of experimental diets	32
4.2	Fatty acid composition of experimental diets	33
4.3	Body weight, feed intake and gain/feed from broiler chickens fed with CONTROL, LCLA and HCLA	36
4.4	Fatty acid composition of breast meat (mg/100g meat) of broiler chicken fed with different inclusion levels of CLA in the diet.	37
4.5	Color values of chicken breast meat from different treatments	38
4.6	The effects of CLA on cholesterol, triglycerides, and lipoproteins (mmol/L) indices in broiler chickens.	40
5.1	Sequences of forward and reverse primers for real time PCR	47
5.2	Fatty acid profile of the adipose tissue of broiler chicken (percentage of total identified fatty acids) across treatment groups	49

LIST OF FIGURES

Figure		Page
2.1	Structure of linoleic acid and CLA isomers. 1. Linoleic acid (typical n-6 PUFA); 2. cis-9, trans-11 CLA and 3. trans-10, cis-12 CLA.	4
2.2	CLAs production by <i>Lactobacillus acidophilus</i>	6
2.3	CLAs biosynthesis in the ruminant's body	7
2.4	Regulating CLA-induced adipocyte apoptosis via intrinsic and extrinsic pathways	15
3.1	Effect of conjugated linoleic acid (CLA) isomers of two concentrations on adipocytes area size in chicken adipocytes compared to CONTROL group	22
3.2	Effects of conjugated linoleic acid (CLA) isomers at two concentrations on the adipocyte diameter of chicken adipocytes compared to CONTROL	23
3.3	Effect of conjugated linoleic acid (CLA) isomers of two concentrations on the number of chicken adipocytes compared to CONTROL	24
3.4	Comparison of relative aP2 transcriptional level in the adipocytes of broiler chickens cultured in two concentrations of conjugated linoleic acid isomers	25
3.5	Comparison of LPL relative transcriptional level in the adipocyte tissue of broiler chickens cultured in two concentrations of conjugated linoleic acid isomers	26
3.6	Comparison of ACBD 5 relative transcriptional level in the adipocyte tissue of broiler chickens cultured in two concentrations of conjugated linoleic acid isomers	27
4.1	TBARS values (mg/kg) of chicken breast meat from different treatment after 0, 3 and 6 days storage	39
5.1	Comparison of PPAR α relative transcriptional level in the abdominal fat of chickens fed diets with CONTROL, LCLA, and HCLA	50
5.2	Comparison of PPAR γ relative transcriptional level in the abdominal fat of chickens fed diets with CONTROL, LCLA, and HCLA	51

5.3	Comparison of aP2 relative transcriptional level in the abdominal fat of chickens fed diets with CONTROL, LCLA, and HCLA	52
5.4	Comparison of ACBD 5 relative transcriptional level in the abdominal fat of chickens fed diets with CONTROL, LCLA, and HCLA	53
5.5	Comparison of LPL relative transcriptional level in the abdominal fat of chickens fed diets with CONTROL, LCLA, and HCLA	54
5.6	Effect of dietary conjugated linoleic acid (CLA) on adipocyte area in the abdominal fat in chickens fed with CONTROL, LCLA, and HCLA	55
5.7	Effects of dietary conjugated linoleic acid (CLA) on adipocyte diameter in chickens fed with CONTROL, LCLA and HCLA	56
5.8	Effects of dietary conjugated linoleic acid (CLA) on adipocyte cell number in chickens fed with CONTROL, LCLA and HCLA	57

LIST OF ABBREVIATIONS

°C	degrees centigrade
ACBD 5	acyl-Coenzyme A binding domain containing 5
acetyl-CoA	acetyl-coenzyme
ADG	average daily gain
ANOVA	analysis of variance
aP2	adipocyte protein 2
BSA	bovine serum albumin
BW	body weight
cal	calorie
CCAAT	enhancer binding protein alpha C/EBP- α
CD-36/FAT	fatty acid translocase
DISC	death-inducing signaling complex
DM	dry matter
DMEM	dulbecco's modified eagle medium
FAME	fatty acids methyl esters
FAR	fatty acid ratio
FAS	fatty acid synthase
FBS	foetal bovine serum
FFA	free fatty acid
g	gram
GLUT4	glucose transporter 4
GPDH	glycerol-3-phosphate dehydrogenase
HDL	high-density lipoprotein
kcal	kilo calories
Kg	kilogram
L	liter
LA	linoleic acid
LDL	low-density lipoprotein
LPL	lipoprotein lipase
MDA	malondialdehyde
ME	metabolizable energy

mg	milligram
min	minute
mL	milliliter
mL/min	milliliters per minute
mmol/L	millimoles per liter
MUFA	monounsaturated fatty acid
n-6: n-3 ratio	total n-6 PUFA to total n-3 PUFA ratio
PBS	phosphate buffer saline
PCR	polymerase chain reaction
PPAR	peroxisome proliferator-activator receptors
PUFA	polyunsaturated fatty acids
PUFA: SFA ratio	total PUFA to total SFA ratio
SCD	stearoyl-CoA desaturase
SEM	standard error of mean
SFA	saturated fatty acid
SPF	specific pathogen-free
TAG	triglycerides
TBARS	thiobarbituric acid reactive substances
TC	total cholesterol
UFA	unsaturated fatty acids
UFS: SFA ratio	total UFA to total SFA ratio
VLDL	very low-density lipoprotein

CHAPTER 1

GENERAL INTRODUCTION

According to Food and Agriculture Organization (FAO) data, the human population benefits significantly from poultry meat and eggs (Marangoni et al., 2015). From a nutritional point of view, poultry meat contains major source of protein (18-25%) and significant amounts of monounsaturated fatty acids (only a third of total fat is made up of saturated fatty acids (Stangierski and Lesnierowski, 2015). Malaysia is the third largest poultry producer in Asia Pacific (MIDA, 2014). This industry has been self-sufficient since 1984. In Malaysia, poultry meat output are estimated to be in the region of RM7, 603.77 million in year 2014 (FLFAM, 2015). The broilers are the most efficient major food animal surpassed both beef and pork meat.

The high growth rate of the modern broiler chicken has resulted in a tremendous increase in nutrient requirement for muscle protein synthesis. Excess fat is a major economic loss to the broiler industry as it not only reduces the carcass yield but also rejection from of the meat by consumers (Sahraei, 2012). The accumulation of fat in modern broiler chicken strains is about 13–14.5% fat of their body weight (Tumova and Teimouri, 2010). This situation most commonly occurs with broiler chickens that consume feed *ad libitum*.

In poultry, the major site of "*de novo*" fat synthesis is the liver (Cornejo et al., 2007). The fat will be transported via low-density lipoproteins or chylomicrons for storage in adipose tissues as triglycerides (Hermier, 1997). In general, the abdominal fat deposition reduced via decreasing the size and/or number of abdominal adipose cells by inhibiting the absorption of dietary fat and fatty acid synthesis, and/or promoting fatty acid β -oxidation (Fouad and El-Senousey, 2014). The abdominal fat pad can be used as a parameter to estimate the total body fat content as it is linked directly to total body fat content in avian species (Fouad and El-Senousey, 2014). The broiler industry needs new method to reduce the fatness in broilers, which would economically benefit producers, and lead to improved health and welfare of chickens.

It has been well documented that nutrition has a significant effect on fat deposition in poultry (Fouad and El-Senousey, 2014). The deposition of fat and fatty acids in animal tissues has been ascribed to a complex regulation network of lipogenic genes (Costa et al., 2013). It gave a negative impact for consumers who are conscious of dietary fat intake. The increased amount of adipose tissue mass may be due to increase preadipocytes, as well as through adipogenesis and/or increased deposition of cytoplasmic triglyceride per cell (Soukas et al., 2001). Adipogenesis is regulated by adipogenesis-related candidate genes such as peroxisome proliferator-activator receptors (PPARs), C/EBP- α (CCAAT/enhancer binding protein alpha), aP2 (adipocyte protein 2), and lipoprotein lipase (LPL) (Yaqiong et al., 2014). Therefore, it is

important to know the regulation of adipose tissue deposition and metabolism in chicken.

Peroxisome proliferator activated receptors (PPAR) is a ligand-dependent transcription factor and a member of the nuclear receptor super family (Wang *et al.*, 2008b). Conjugated linoleic acid (CLA) serves as ligand activator for PPAR isoforms able to regulate cellular metabolism, differentiation and in part of carbohydrate homeostasis (Peters *et al.*, 2001). Royan *et al.* (2011a); Wang *et al.* (2008b); Sato *et al.* (2009) suggested that, PPAR γ (gamma), is strongly associated with abdominal fat deposition. In avian species, however, its mechanism has yet to be extensively investigated, as the PPAR may mediate their effects through a different signaling path. In view of the far-reaching effects of conjugated linoleic acid, this study will attempt to elucidate the PPAR expression and its influence on lipid mobilization, adipogenesis and fat deposition in chickens.

In recent years, focus by researchers on health-promoting properties in poultry meat by fatty acid has been increasing (Yari *et al.*, 2015). Modification of these attributes in meat with bioactive substances such as CLA, vitamins and antioxidants, and a balanced n-6 to n-3 PUFA is a simpler and much more commonly applied method than the genetic approach (Stangierski and Lesnierowski, 2015). Polyunsaturated fatty acids (PUFA), particularly the conjugated linoleic acid is increasingly recognized for its beneficial health impact (Park *et al.*, 1997; West 1998; Pariza *et al.*, 2001). Conjugated linoleic acid refers to a group of geometric and positional dienoic isomers of linoleic acid [C18:2(n-6)]. The cis-9,trans-11 CLA isomer is the principal dietary form of conjugated linoleic acid, comprises approximately 90% found primarily in beef and dairy products and the 10,12 isomer comprises the remaining 10% (Carvalho *et al.*, 2010). Collectively, animal studies demonstrated dietary intake of a crude mixture of conjugated linoleic acid isomers prevents the development of obesity (Ostrowska *et al.*, 1999). It is possible to theorize that conjugated linoleic acid could be used as a metabolic modifier in the poultry industry as there are many studies by Zhang *et al.* (2009a); Zhou (2008); Ramiah *et al.* (2014a), evidenced that conjugated linoleic acid is a new tool that can be used to reduce fatness in modern broiler chickens.

Most animals or human subjects have consumed a mixture of the two main CLA isomers: cis-9, trans-11 CLA and trans-10, cis-12 CLA (Larsen *et al.*, 2003). Feitoza *et al.* (2009) reported that trans-10, cis-12 CLA is the main isomer responsible for alterations in lipid profile and body composition. Nonetheless, 90% CLA in dairy products comprise of cis-9, trans-11 CLA (Feitoza *et al.*, 2009). The body fat-lowering effect of CLA has also been reported inconsistent humans and less significant in animals (Riserus *et al.*, 2001; Blankson *et al.*, 2000). Physiological alterations induced by conjugated linoleic acid in relation to the gene expression depend on the animal species (Azain *et al.*, 2000). Conjugated linoleic acid intake has been associated with numerous health benefits and inclusion of conjugated linoleic acid into animal feed improves the meat quality and provides nutritional values for human consumption. Therefore, the general objective of this study is to investigate effects of conjugated linoleic acid on the fat deposition in broiler chickens with the following hypothesis and objectives:

Hypothesis Statements

1. The effects of conjugated linoleic acid isomers on the morphological changes in adipose tissue and adipogenic gene expression on primary adipose tissue mainly related to trans-10, cis-12 CLA isomer.
2. Incorporation of conjugated linoleic acid will have an effect on the growth performance, fatty acid profile of meat and meat quality of broiler chicken.
3. Incorporation of conjugated linoleic acid will have an effect on adipose tissue cellularity and the regulation of adipogenic gene transcription.

Based on these hypotheses, the **SPECIFIC OBJECTIVES** of the present study were:

1. To assess the effects of conjugated linoleic acid isomers on the morphological changes in adipose tissue, and adipogenic genes expression on primary adipose tissue.
2. To investigate the effects of conjugated linoleic acid on the growth performance, fatty acid composition of breast muscle, lipid peroxidation, meat colour, and plasma lipids in broiler chicken.
3. To determine the possible effects of conjugated linoleic acid on the morphology of adipose tissue, the expression of PPARs and associated genes in broiler chickens.

REFERENCES

- Adlof, R., Duval, S., & Emken, E. (2000). Biosynthesis of conjugated linoleic acid in humans. *Lipids*, 35, 131-135.
- Ahn, D., Sell, J., Jo, C., Chamruspollert, M., & Jeffrey, M. (1999). Effect of dietary conjugated linoleic acid on the quality characteristics of chicken eggs during refrigerated storage. *Poultry Science*, 78, 922-928.
- Aletor, V., Eder, K., Becker, K., Paulicks, B., Roth, F., & Roth-Maier, D. (2003). The effects of conjugated linoleic acids or an alpha-glucosidase inhibitor on tissue lipid concentrations and fatty acid composition of broiler chicks fed a low-protein diet. *Poultry Science*, 82(5), 796-804.
- Aydin, R. (2005). The effect of conjugated linoleic acid on the fatty acid composition of different tissues and yolk lipids in pigeons. *South Africa Journal of Animal Science*, 35, 253-259.
- Aydin, R., & Cook, M. (2004). The effect of dietary conjugated linoleic acid on egg yolk fatty acids and hatchability in Japanese quail. *Poultry Science*, 83, 2016-2022.
- Aydin, R., Pariza, M.W., & Cook, M.E. (2001). Olive oil prevents the adverse effects of dietary conjugated linoleic acid on chick hatchability and egg quality. *Journal of Nutrition*, 131, 800-806.
- Azain, M.J., Hausman, D.B., Sisk, M.B., Flatt, W.P., & Jewell, D.E. (2000). Dietary conjugated linoleic acid reduces rat adipose tissue cell size rather than cell number. *Journal of Nutrition*, 130, 1548-1554.
- Badinga, L., Selberg, K., Dinges, A., Corner, C., & Miles, R. (2003). Dietary conjugated linoleic acid alters hepatic lipid content and fatty acid composition in broiler chickens. *Poultry Science*, 82(1), 111-116.
- Banni, S., Angioni, E., Contini, M., Carta, G., Casu, V., & Iengo, G. (1998). Conjugated linoleic acid and oxidative stress. *Journal of the American Oil Chemists' Society*, 75(2), 261-267.
- Banni, S., Petroni, A., Blasevich, M., Carta, G., Cordeddu, L., & Murru, E. (2004). Conjugated linoleic acids (CLA) as precursors of a distinct family of PUFA. *Lipids*, 39(11), 1143-1146.
- Barnes, K., Winslow, N., Shelton, A., Hlusko, K., & Azain, M. (2011). Effect of dietary conjugated linoleic acid on marbling and intramuscular adipocytes in pork. *Journal of Animal Science*, 90, 1142-1149.
- Barroeta, A.C. (2006). Nutritive value of poultry meat: Relationship between vitamin E and PUFA. *Worlds Poultry Science Journal*, 63, 277-284.

- Bauman, D., & Griinari, J.M. (1999). Biosynthesis of CLA and its incorporation into meat and milk of ruminants. *Journal of Animal Science*, 77 (Supplement 1): 1.
- Belury, M.A. (2002). Inhibition of carcinogenesis by conjugated linoleic acid: Potential mechanisms of action. *Journal of Nutrition*, 10, 2995-2998.
- Belury, M.A., & Kempa-Steczko, A. (1997). Conjugated linoleic acid modulates hepatic lipid composition in mice. *Lipids*, 32, 197-204.
- Benjamin, S., & Spener, F. (2009). Conjugated linoleic acids as functional food: an insight into their health benefits. *Nutrition and Metabolism*, 6(1), 36.
- Berg, J.M., Tymoczko, J.L., & Stryer, L. (2002). The utilization of fatty acids as fuel requires three stages of processing. *Biochemistry*. 5th edition. New York: Freeman, W.H. <http://www.ncbi.nlm.nih.gov/books/NBK22581/>. Retrieved 23 February 2015.
- Bessa, R., Santos-Silva, J., Ribeiro, J., & Portugal, A. (2000). Reticulo-rumen biohydrogenation and the enrichment of ruminant edible products with linoleic acid conjugated isomers. *Livestock Production Science*, 63(3), 201-211.
- Blankson, H., Stakkestad, J.A., Fagertun, H., Thom, E., Wadstein, J., & Gudmundsen, O. (2000). Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *Journal of Nutrition*, 130, 2943-2948.
- Blankson, H., Stakkestad, J.A., Fagertun, H., Thom, E., Wadstein, J., & Gudmundsen, O. (2000). Conjugated linoleic acid reduces body fat mass in overweight and obese humans. *Journal of Nutrition*, 130, 2943-2948.
- Bolukbasi, S.C. (2006). Effect of dietary conjugated linoleic acid (CLA) on broiler performance, serum lipoprotein content, muscle fatty acid composition and meat quality during refrigerated storage. *British Poultry Science*, 47, 470-476.
- Botelho, A., Santos-Zago, L., & Oliveira, A. (2009). Effect of conjugated linoleic acid supplementation on lipoprotein lipase activity in 3T3-L1 adipocyte culture. *Revista De Nutrição*, 22(5), 767-771.
- Bou, R., Codony, R., Tres, A., Decker, E., & Guardiola, F. (2009). Dietary strategies to improve nutritional value, oxidative stability, and sensory properties of poultry products. *Critical Reviews in Food Science and Nutrition*, 49(9), 800-822.
- Bouthegourd, J.C., Even, P.C., Gripois, D., Tiffon, B., Blouquit, M.F., Roseau, S., Lutton, C., Tome, D., & Martin, J.C. (2002). A CLA mixture prevents body triglyceride accumulation without affecting energy expenditure in Syrian hamsters. *Journal of Nutrition*, 132, 2682-2689.
- Brandebourg, T.D. & Hu, C.Y. (2005). Isomer-specific regulation of differentiating pig preadipocytes by conjugated linoleic acids. *Journal of Animal Science*. 83, 2096-2105.

- Brasaemle, D.L. (2010). Lipolysis control: The plot thickens. *Cell Metabolism*, 11, 173-174.
- Brodie, A.E., Manning, V.A., Ferguson, K.R., Jewell, D.E., & Hu, Y. (1999). Conjugated linoleic acid inhibits differentiation of pre- and post-confluent 3T3-L1 preadipocytes but inhibits cell proliferation only in pre-confluent cells. *Journal of Nutrition*, 129, 602-606.
- Brown, J.M., & McIntosh, M.K. (2003). Conjugated linoleic acid in humans: Regulation of adiposity and insulin sensitivity. *Journal Nutrition*, 133, 3041-3046.
- Brown, J.M., Halvorsen, Y.D., Lea-Currie, Y.R., Geigerman, C., & McIntosh, M. (2001) Trans-10, cis-12, but not cis-9, trans-11, conjugated linoleic acid attenuates lipogenesis in primary cultures of stromal vascular cells from human adipose tissue. *Journal Nutrition*, 131, 2316-2321.
- Brown, J.M., Sandberg-Boyse, M., Skov, S., Morrison, R., Storkson, J., Lea-Currie, R., Pariza, M., Mandrup, S., & McIntosh, M. (2003). Isomer-specific regulation of metabolism and PPAR- γ by conjugated linoleic acid (CLA) in human preadipocytes. *Journal Lipid Research*, 44, 1287-1300.
- Buccioni, A., Antongiovanni, M., Mele, M., Gualtieri, M., Minieri, S., & Rapaccini, S. (2009). Effect of oleic and conjugated linoleic acid in the diet of broiler chickens on the live growth performances, carcass traits and meat fatty acid profile. *Italian Journal of Animal Science*, 8, 603-614.
- Butterwith, S.C. (1997). Regulators of adipocyte precursor cells. *Poultry Science*, 76, 118-123.
- Camarena, M.S.Y., Vanden Heuvel, J.P., Blanchard, S.G., Leesnitzer L.A., & Belury, M.A. (1999). Conjugated linoleic acid is a potent naturally occurring ligand and activator of PPAR alpha. *Journal Lipid Research*, 40, 1426-1433.
- Cartwright, A.L. (1991). Adipose cellularity in *callus domesticus*: Investigations to control body composition in growing chickens. Symposium: Avian lipid metabolism and biochemistry. *Journal of Nutrition*, 1486-1497.
- Carvalho, E.B.T., Melo, I.L.P. , & Mancini-Filho, J. (2010). Chemical and physiological aspects of isomers of conjugated fatty acids. *Ciência e Tecnologia de Alimentos*, 30, b295-307.
- Chae, S.H., Keeton, J.T., & Smith, S.B. (2004). Conjugated linoleic acid reduces lipid oxidation in aerobically stored, cooked ground beef patties. *Journal Food Science*, 69, S306-S309.
- Chajes, V., Lavillonniere, F., Maillard, V., Giraudeau, B., Jourdan, M.L., Sebedio, J.L. & Bougnoux, P. (2003). Conjugated linoleic acid content in breast adipose tissue of breast cancer patients and the risk of metastasis. *Nutrition and Cancer*, 45, 17-23.

- Chamruspollert, M., & Sell, J.L. (1999). Transfer of dietary conjugated linoleic acid to egg yolks of chickens. *Poultry Science*, 78, 1138-1150.
- Chapman, C.L., Morgan, M., & Murphy, M.C. (2000). Maternal and early dietary fatty acid intake: changes in lipid metabolism and liver enzymes in adult rats. *Journal of Nutrition*, 130, 146-151.
- Chen, Z.Y., Chan, P.T., Kwan, K.Y., & Zhang, A. (1997). Reassessment of the antioxidant activity of conjugated linoleic acids. *Journal of American Chemical Society*, 74, 749-753.
- Cheng, Z., Oguejiofor, C.F., Swangchan-Uthai, T., Carr, S. & Wathes, D.C. (2015). Relationships between circulating urea concentrations and endometrial function in postpartum dairy cows. *Animals*, 5, 748-773.
- Cherian, G., Holsonbake, T.B., Goeger, M.P., & Bildfell, R. (2002). Dietary CLA alters yolk and tissue FA composition and hepatic histopathology of laying hens. *Lipids*, 37, 751-757.
- Chin, S.F., Eiu, W., Torkson, J.M., Ha, Y.E., & Pariza, M.W. (1992). Dietary sources of conjugated dienoic isomers of linoleic acid. A newly recognized class of anticarcinogens. *Journal of Food Composition Analysis*, 5, 185-197.
- Chin, S.F., Storkson, J.M., Albright, K.J., Cook, M.E., & Pariza, M.W. (1994). Conjugated linoleic acid is a growth factor for rats as shown by enhanced weight gain and improved feed efficiency. *Journal of Nutrition*, 124, 2344-2349.
- Chin, S.F., Storkson, J.M., & Pariza, M.W. (1993). Conjugated dienoic derivatives of linoleic acid: A new class of anticarcinogens. Food flavor and safety. *Journal of American Chemical Society*, 262-271.
- Choa, S., Ryua, C., Yang, J., Mbiriri, D.T., Choi, C.W., Chae, J.I., Kim, Y.H., Shim, K.S., Kim, K.J., & Choi, N.J. (2013). Effect of conjugated linoleic acid feeding on the growth performance and meat fatty acid profiles in broiler: meta-analysis. *Asian Australasian Journal of Animal Science*, 26, 995-1002.
- Choi, Y., Kim, Y., Han, Y., Park, Y., Pariza, M., & Ntambi, J. (2000). The trans-10, cis-12 isomer of conjugated linoleic acid downregulates stearoyl-coadesaturase 1 gene expression in 3T3-L1 adipocytes. *Journal Nutrition*, 130, 1920-1924.
- Choi, Y.H. (2009). Conjugated linoleic acid as a key regulator of performance, lipid metabolism, development, stress and immune functions and gene expression in chickens. *Asian Australasian Journal of Animal Science*. 22, 448-458.
- Chung, S.H., Kim, I.H., Park, H.G., Kang, H.S., Yoon, C.S., & Jeong, H.Y. (2008). Synthesis of conjugated linoleic acid by human-derived Bifidobacterium breve LMC 017: utilization as a functional starter culture for milk fermentation. *Journal Agriculture and Food Chemistry*, 14, 3311-3316.

- Cianzio, D.S., Topel, D.G., Whitehurst, G.B., Beitz, D.C., & Self, H.L. (1985). Adipose tissue growth and cellularity: Changes in bovine adipocyte size and number. *Journal of Animal Science*, 60, 970–976.
- Clandinin, M.T., Cheema, S., Field, C.J., Garg, M.L., Venkatraman, J., & Clandinin, T.R. (1991). Dietary fat: exogenous determination of membrane structure and cell function. *Federation of American Societies for Experimental Biology Journal*.1991, 13, 2761-2769.
- Close, R.N., Schoeller, D.A., Watras, A.C., & Nora, E.H. (2007). Conjugated linoleic acid supplementation alters the 6-mo change in fat oxidation during sleep. *The American Journal of Clinical Nutrition*, 86:797-804.
- Corino, C., Lo Fiego, D.P., Macchioni, P., Pastorelli, G., Di Giancamillo, A., Domeneghini, C., & Rossi, R. (2007). Influence of dietary conjugated linoleic acids and vitamin E on meat quality, and adipose tissue in rabbits. *Meat Science*, 76, 19-28.
- Corino, C., Magni, S., Pastorelli, R., Rossi, R., & Mourot, J. (2003). Effect of conjugated linoleic acid on meat quality, lipid metabolism, and sensory characteristics of dry-cured hams from heavy pigs. *Journal of Animal Science*, 81, 2219-2229.
- Corino, C., Mourot, J., Magni, S., Pastorelli, G., & Rosi F. (2002). Influence of dietary conjugated linoleic acid on growth, meat quality, lipogenesis, plasma leptin and physiological variables of lipid metabolism in rabbits. *Journal of Animal Science*, 80, 1020-1028.
- Cornejo, S., Gadelha, A.C., Pokniak, J., & Villouta, G. (2007). Qualitative feed restriction on productive performance and lipid metabolism in broiler chickens. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 59, 1554-1562 .
- Cortinas, L., Barroeta, A., Villaverde, C., Galobart, J., Guardiola, F., & Baucells, M.D. (2005) Influence of dietary polyunsaturated level on chicken meat quality: lipid oxidation. *Poultry Science*. 84, 48-55.
- Costa, A.S.H., Lopes, P.A., Estevão, M., Martins, S.V., Alves, S.P., Pinto, R.M.A., Pissarra, H., Correia, J.J., Pinho, M., Fontes, C.M.G.A., & Prates, J.A.M. (2012). Contrasting cellularity and fatty acid composition in fat depots from Alentejana and Barrosã bovine breeds fed high and low forage diets. *International Journal of Biological Science*, 8, 214-227.
- Costa, D.A., Pires, V., Fontes, C., & Mestre Prates, J. (2013). Expression of genes controlling fat deposition in two genetically diverse beef cattle breeds fed high or low silage diets. *BMC Veterinary Research*, 9, 118.
- Crespo, N., & Esteve-Garcia, E. (2002). Dietary polyunsaturated fatty acids decrease fat deposition in separable fat depots but not in the remainder carcass. *Poultry Science*, 81, 512-518.

- Crow, M.T., Mani, K., Nam, Y.J., & Kitsis, R.N. (2004). The mitochondrial death pathway and cardiac myocyte apoptosis. *Circulation Research*, 95, 957-970.
- Czauderna, M., Kowalczyk, J., Niedźwiedzka, K.M., Wąsowska, I., Pastuszewska, B., Bulska, E., & Ruszczyńska, A. (2004). Liver and body mass gain, content of conjugated linoleic acid (CLA) isomers and other fatty acids in the liver of rats fed CLA isomers and selenium. *Journal of Animal Feed Science*. 13, 353-369.
- D'Andre, H.C., Paul, W., Shen, X., Jia, X., Zhang, R., & Sun, L. (2013). Identification and characterization of genes that control fat deposition in chickens. *Journal of Animal Science and Biotechnology*, 1, 43.
- De Smet, S. (2012). Meat, poultry and fish composition: Strategies for optimizing human intake of essential nutrients. *In Animal Frontier*, 2, 10-16.
- de-Deckere, E.A.M., Van Amelsfoort, J.M.M., McNeill, G.P. & Jones, P. (1999). Effects of conjugated linoleic acid (CLA) isomers on lipid levels and peroxisome proliferation in the hamster. *British Journal of Nutrition*, 82, 309-17.
- Dirain, C.O., Toyomizu, M., Wing, T., Cooper, M., & Bottje, W.G. (2007). Gene expression in breast muscle and duodenum from low and high feed efficient broilers. *Poultry Science*, 86, 372-381.
- Du, M., & Ahn, D. (2002). Effect of dietary conjugated linoleic acid on the growth rate of live birds and on the abdominal fat content and quality of broiler meat. *Poultry Science*, 81, 428-433.
- Du, M., & Ahn, D. (2003). Dietary CLA affects lipid metabolism in broiler chicks. *Lipids*, 38, 505-11.
- Du, M., Ahn, D., & Sell, J. (2000). Effects of dietary conjugated linoleic acid and linolenic acid ratio on polyunsaturated fatty acid status in laying hens. *Poultry Science*, 79, 1749-1757.
- Du, M., Ahn, D., Nam, K.C., & Sell, J.L. (2000). Influence of dietary conjugated linoleic acid on volatile profiles, color and lipid oxidation of irradiated raw chicken meat. *Meat Science*. 56, 387-395.
- Dugan, M. E. R., Aalhus, J. L., Lien, K. A., Schaefer, A. L., & Kramer, J. K. G. (2001). Effects of feeding different levels of conjugated linoleic acid and total oil to pigs on live animal performance and carcass composition. *Canadian Journal of Animal Science*, 81, 505-510.
- Dugan, M.E.R., Aalhus, J.L., Jeremiah, L.E., Kramer, J.K.G., & Schaefer, A.L. (1999). The effects of feeding conjugated linoleic acid on subsequent pork quality. *Canada Journal of Animal Science*, 79, 45-5.
- Dugan, M.E.R., Aalhus, J.L., Schaefer, A.L., & Kramer, J.K.G. (1997). The effect of conjugated linoleic acid on fat to lean repartitioning and feed conversion in pigs. *Canada Journal of Animal Science*. 77, 723-725.

- Ebrahimi, M., Rajion, M.A., Goh, Y.M., & Sazili, A.Q. (2012). Impact of different inclusion levels of oil palm (*Elaeisguineensis* Jacq.) fronds on fatty acid profiles of goat muscles. *Journal of Animal Physiology and Animal Nutrition*, 96, 962-969.
- Ebrahimi, M., Rajion, M.A., Goh, Y.M., Sazili, A.Q., & Schonewille, J.T. (2013). Effect of linseed oil dietary supplementation on fatty acid composition and gene expression in adipose tissue of growing goats. *BioMed Research International*. DOI:10.1155/2013/194625 (2013).
- Eggert, J.M., Belury, M.A., Kempa-Steczko, A., Mills, S.E., & Schinckel, A.P. (2001). Effects of conjugated linoleic acid on the belly firmness and fatty acid composition of genetically lean pigs. *Journal of Animal Science*, 79, 2866–2872.
- Elmore, S. Apoptosis: A review of programmed cell death. (2007). *Toxicology and Pathology*, 35, 495–516.
- Evans, M., Geigerman, C., Cook, J., Curtis, L., Kuebler, B., & McIntosh, M. (2000) Conjugated linoleic acid suppresses triglyceride accumulation and induces apoptosis in 3T3-L1 preadipocytes. *Lipids*, 35, 899-910.
- Fernandez, M.L., & West, K.L. (2005). Mechanisms by which dietary fatty acids modulate plasma lipids. *Journal of Nutrition*, 135, 2075-2078.
- Ferrini, G., Baucells, M.D., Esteve-García, E., & Barroeta, A.C. (2008). Dietary polyunsaturated fat reduces skin fat as well as abdominal fat in broiler chickens. *Poultry Science*, 87, 528-535.
- Fesler, J.A., & Peterson, D.G. (2013). Conjugated linoleic acids alter body composition differently according to physiological age in Moulard ducks. *Poultry Science*, 10, 2697-270.
- FLFAM, Federation of Livestock Farmers' Association of Malaysia (2014). Broiler Production. <http://www.flfam.org.my/index.php/industry-info/the-poultry-industry/broiler-production>. Retrieved on 12 January 2015.
- Folch, J., Lees, M., & Sloane Stanely, G.H. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *Journal of Biological Chemistry*, 226, 497-509.
- Food Outlook. (2014). Biannual Report On Global Food Markets. <http://www.fao.org/docrep/018/al999e/al999e.pdf>. Retrieved 11 August 2014.
- Fouad, A. M., & El-Senousey, H. K. (2014). Nutritional factors affecting abdominal fat deposition in poultry: A Review. *Asian-Australasian Journal of Animal Sciences*, 27, 1057–1068.
- Francis, G.A., Fayard, E., Picard, F., & Auwerx, J. (2003). Nuclear receptors and the control of metabolism, *Annual Review of Physiology*, 65, 261-311.

- Fritsche, J., & Steinhardt, H. (1998). Amounts of conjugated linoleic acid (CLA) in German foods and evaluation of daily intake. *Zeitschrift für Lebensmittel-Untersuchung und -Forschung, A, European Journal of Food Research and Technology*, 206, 77–82.
- Gavino, V.C., Gavino, G., Leblanc, M.J., & Tuchweber, B. (2000). An isomeric mixture of conjugated linoleic acids but not pure cis-9, trans-11-octadecadienoic acid affects body weight gain and plasma lipids in hamsters. *Journal of Nutrition*. 130, 27-29.
- Georgiadi, A., & Kersten, S. (2012). Mechanisms of Gene Regulation by Fatty Acids. *Advance Nutrition*, 2, 127-134.
- Granlund, L., Jan, I.P., & Hilde, I.N. (2005). Impaired lipid accumulation by trans10, cis12 CLA during adipocyte differentiation is dependent on timing and length of treatment, *Biochimica et Biophysica Acta*, 1687, 11-22.
- Granlund, L., Juvet, L.K., Pedersen, J.I., & Nebb, H.I. (2003). Trans-10, cis 12-conjugated linolenic acid prevents triacylglycerol accumulation in adipocytes by acting as a PPAR γ modulator. *Journal of Lipid Research*, 44, 1441-1452.
- Grashorn, M.A. (2005). Enrichment of eggs and poultry meat with biologically active substances by feed modifications and effects on the final quality of the product. *Polish Journal of Food and Nutrition Sciences*, 14, 15-20.
- Grau, A., Codony, R., Grimpa, S., Baucells, M.D., & Guardiola, F. (2001). Cholesterol oxidation in frozen dark chicken meat: Influence of dietary fat source, and α -tocopherol and ascorbic acid supplementation. *Meat Science*, 57, 197–208.
- Griffin, H. D. (1992). Manipulation of egg yolk cholesterol: a physiologist's view. *World's Poultry Science Journal*, 48, 101-112.
- Griffin, M.J., & Sul, H.S. (2004). Insulin regulation of fatty acid synthase gene transcription: Roles of USF and SREBP-1c. *Critical Reviews. International Union of Biochemistry and Molecular Biology Life*. 56, 595–600.
- Griinari, M., Corl, B.A., Lacy, S.H., Chouinard, P.Y., Nurmela, K.V.V., & Bauman, D.E. (2000). Conjugated linoleic acid is synthesized endogenously in lactating dairy cows by $\delta 9$ -desaturase. *Journal of Nutrition*. 130, 2285-2291.
- Guo, L., Sun, B., Shang, Z., Leng, L., Wang, Y., Wang, N., & Li, H. (2011) Comparison of adipose tissue cellularity in chicken lines divergently selected for fatness. *Poultry Science*, 90, 2024–2034.
- Ha, Y.L., Grimm, N.K., & Pariza, M.W. (1987). Anticarcinogens from fried ground beef: heat-altered derivatives of linoleic acid. *Carcinogenesis*, 8, 1881-1887.
- Ha, Y.L., Storkson, J., & Pariza, M.W. (1990). Inhibition of benzo(a) pyrene-induced mouse forestomach neoplasia by conjugated dienoic derivatives of linoleic acid. *Cancer Research*, 50, 1097–1101.

- Halle, I., Jahreis, G., Henning, M., Köhler, P., & Dänicke, S. (2012). Effects of dietary conjugated linoleic acid on the growth performance of chickens and ducks for fattening and fatty acid composition of breast meat. *Journal für Verbraucherschutz und Lebensmittelsicherheit*, 7, 3-9.
- Harvatine, K.J., Perfield, J.W.II., & Bauman, D.E. (2009) Expression of enzymes and key regulators of lipid synthesis are up-regulated in adipose tissue during CLA-induced milk fat depression in the dairy cow. *Journal of Nutrition*, 139, 849–854.
- Hermier, D. (1997). Lipoprotein metabolism and fattening in poultry. *Journal of Nutrition*, 127, 805–808.
- Herzallah, S. (2013). Enrichment of conjugated linoleic acid (CLA) in hen eggs and broiler chickens meat by lactic acid bacteria. *British Poultry Science*, 54, 747-752.
- Houseknecht, K.L., VandenHeuvel, J.P., Moya-Camarena, S.Y., Portocarrero, C.P., Peck, L.W., Nickel, K.P., & Belury M.A. (1998). Dietary conjugated linoleic acid normalizes impaired glucose tolerance in the Zucker diabetic fatty FA/FA rat. *Biochemical and Biophysical Research Communications*, 244, 678-682.
- Hubbard, N.E., Lim, D., Summers, L., & Erickson, K.L. (2000). Reduction of murine mammary tumor metastasis by conjugated linoleic acid. *Cancer Letters*.150, 93–100.
- Hur, S.J., Ye, B.W., Lee, J.L., Ha, Y.L., Park, G.B., & Joo, S.T. (2004). Effects of conjugated linoleic acid on color and lipid oxidation of beef patties during cold storage. *Meat Science*, 66, 771–775.
- Igarashi, M., & Miyazawa, T. (2001). The growth inhibitory effect of conjugated linoleic acid on a human hepatoma cell line, HepG2, is induced by a change in fatty acid metabolism, but not the facilitation of lipid peroxidation in the cells. *BBA Molecular and Cell Biology of Lipids*, 1530, 162–171.
- Ip, C., Chin, S., Scimeca, J., & Pariza, M. (1991). Mammary cancer prevention of conjugated dienoic derivative of linoleic acid. *Cancer Research*, 51, 6118-6124.
- Ip, C., Ip, M.M., Loftus, T., Shoemaker, S., & Shea-Eaton, W. (2000). Induction of apoptosis by conjugated linoleic acid in cultured mammary tumor cells and premalignant lesions of the rat mammary gland. *Cancer Epidemiology, Biomarkers & Prevention*. 9, 689-696.
- Jakobsen, M.U., O'Reilly, E.J., & Heitmann, B.L. (2009). Major types of dietary fat and risk of coronary heart disease: A pooled analysis of 11 cohort studies. *The American Journal of Clinical Nutrition*. 5, 425-432.
- Ji, B., Ernest, B., Gooding, J.R., Das, S., Saxton, A.M., Simon, J., Dupont, J., Métayer-Coustard, S., Campagna, S.R., & Voy, B.H. (2012). Transcriptomic and

metabolomic profiling of chicken adipose tissue in response to insulin neutralization and fasting. *BMC Genomics*, 13, 441.

- Jiang, W., Nie, S., Qu, Z., Bi, C., & Shan, A. (2014). The effects of conjugated linoleic acid on growth performance, carcass traits, meat quality, antioxidant capacity, and fatty acid composition of broilers fed corn dried distillers grains with solubles. *Poultry Science*, 93, 1202-1210.
- Jones, S., Ma, D.W., Robinson, F.E., Field, C.J., & Clandinin, M.T. (2000). Isomers of conjugated linoleic acid (CLA) are incorporated into egg yolk lipids by CLA-fed laying hens. *Journal of Nutrition*, 130, 2002-2005.
- Joo, S.T., Lee, J.L., Ha, Y.L., & Park, G.B. (2002). Effects of dietary conjugated linoleic acid on fatty acid composition, lipid oxidation, color, and water-holding capacity of pork loin. *Journal of Animal Science*, 80, 108-112.
- Jouki, M., & Khazaei, N. (2012). Color and oxidation changes in camel meat during storage. Research Article. *International Journal of Pharma and Bio Sciences*, 3.
- Kang, K., Liu, W., Albright, K.J., Park, Y., & Pariza, M.W. (2003). Trans-10,cis-12 CLA inhibits differentiation of 3T3-L1 adipocytes and decreases PPAR gamma expression. *Biochemical and Biophysical Research Communications*, 303, 795-799.
- Kang, K.J., & Choi, S.S. (2008). Effects of conjugated linoleic acid supplementation on fatty acid composition in the plasma, liver, and epididymal fat pads of male-Sprague Dawley rats. *Journal of Medicinal Food*, 11, 435-442.
- Kawahara, S., Takenoyama, S., Takuma, K., Muguruma, M., & Yamauchi, K. (2009). Effects of dietary supplementation with conjugated linoleic acid on fatty acid composition and lipid oxidation in chicken breast meat. *Journal of Animal Science*, 80, 468-474.
- Kelley, N.S., Hubbard, N.E., & Erickson, K.L. (2007). Conjugated linoleic acid isomers and cancer. *Journal of Nutrition*, 137, 12, 2599-2607.
- Kennedy, A., Martineza, K., Schmidt, S., Mandrup, S., LaPointa, K., & McIntosh, M. (2010). Antiobesity mechanisms of action of conjugated linoleic acid. *Journal of Nutritional Biochemistry*, 21, 171-179.
- Kennedy, S.R. (2007). Bioactive fatty acids as dietary supplements for farmed fish: effects on growth performance, lipid metabolism, gene expression and immune parameters. Ph.D. Thesis, University of Stirling, Scotland. <https://dspace.stir.ac.uk/bitstream/1893/389/1/kennedy-farmed-fish-thesis.pdf>. Retrieved 15 October 2014.
- Kepler, C.R., Hiron, K.P., McNeill, J.J., & Tove, S.B. (1966). Intermediates and products of the biohydrogenation of linoleic acid by *Butyrivibrio fibrisolvens*. *Journal of Biological Chemistry*, 241, 1350-1354.

- Khanal, R.C., & Thiman, T.R. (2004). Biosynthesis of conjugated linoleic acid (CLA): A review. *Pakistan Journal of Nutrition*, 3, 72-81.
- Kim, E., Holthuizen, P., Park, H., Ha, Y., Jung, K., & Park, J. (2002). Trans-10, cis-12-conjugated linoleic acid inhibits Caco-2 colon cancer cell growth. *The American Journal of Physiology-Gastrointestinal and Liver Physiology*, 283, G357-G367.
- Kim, J.H., Hwangbo, J., Choi, N.J., Park, H.G., Yoon, D.H., Park, E.W., Lee, S.H., Park, B.K., & Kim, Y.J. (2007). Effect of dietary supplementation with conjugated linoleic acid, with oleic, linoleic, or linolenic acid, on egg quality characteristics and fat accumulation in the egg yolk. *Poultry Science*, 86, 1180-1186.
- Kim, Y.H., Nam, K.C. & Ahn, D. (2002). Volatile profiles, lipid oxidation and sensory characteristics of irradiated meat from different animal species. *British Journal of Nutrition*. 61, 257-265.
- Kimoto, N., Hirose, M., Futakuchi, M., Iwata, T., Kasai, M., & Shirai, T. (2001). Site-dependent modulating effects of conjugated fatty acids from safflower oil in a rat two-stage carcinogenesis model in female Sprague-Dawley rats. *Cancer Letter*, 168, 15–21.
- Konig, B., Kluge, K., Haase, K., Brandsh, C., Stangle, G.I., & Eder, K. (2007). Effects of clofibrate treatment in laying hens, *Poultry Science*, 86, 1187-1195.
- Kramer, J.K., Fellner, V., Dugan, M.E., Sauer, F.D., Mossoba, M.M., & Yurawecz, M.P. (1997). Evaluating acid and base catalysts in the methylation of milk and rumen fatty acids with special emphasis on conjugated dienes and total trans fatty acids. *Lipids*, 11, 1219-28.
- Kramer, J.K.G., Cruz-Hernandez, C., Deng, Z.Y., Zhou, J.Q., Jahreis, G., & Dugan, M.E.R. (2004). Analysis of conjugated linoleic acid and trans 18:1 isomers in synthetic and animal products. *The American Journal of Clinical Nutrition*. 79, 1137S-1145S.
- Laliotis, G.P., Bizelis, I., & Rogdakis, E. (2010). Comparative approach of the de novo fatty acid synthesis (lipogenesis) between ruminant and non ruminant mammalian species: from biochemical level to the main regulatory lipogenic genes. *Current Genomics*. 11, 168-83.
- Lam, C.K., Chen, J., Cao, Y., Yang, L., Wong, Y.M., Yeung, S.Y.V., Yao, X., Huang, Y., & Chen, Z.Y. (2008) Conjugated and non-conjugated octadecaenoic acids affect differently intestinal acyl coenzyme A: Cholesterol acyltransferase activity. *Atherosclerosis*, 198, 85–93.
- Larkina, T. A., Sazanova, A. L., Fomichev, K. A., Barkova, O. Y., Malewski, T., Jaszczak, K., & Sazanov, A. A. (2011). HMG1A and PPARG are differently expressed in the liver of fat and lean broilers. *Journal of Applied Genetics*, 52, 225-228

- Larsen, T.M., Toubro, S., & Astrup, A. (2003). Efficacy and safety of dietary supplements containing CLA for the treatment of obesity: evidence from animal and human studies. *Journal of Lipid Research*, 44, 2234-41.
- Lee, K.N, Pariza, M.W. & Ntambi, J.M. (1998). Conjugated linoleic acid decreases hepatic stearoyl-CoA desaturase mRNA expression. *Biochemical and Biophysical Research Communications*, 248, 817-821.
- Lee, K.N., Kitchevsky, D., & Pariza, M.W. (1994). Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis*, 108, 19-25.
- Leung, Y.H., & Liu, R.H. (2000). Trans-10,cis-12-conjugated linoleic acid isomer exhibits stronger oxyradical scavenging capacity than cis-9,trans-11-conjugated linoleic acid isomer. *Journal of Agriculture and Food Chemistry*, 48, 5469-5475.
- Li, Y., & Watkins, B.A. (1998). Conjugated linoleic acids alter bone fatty acid composition and reduce ex vivo prostaglandin E2 biosynthesis in rats fed n-6 or n-3 fatty acids. *Lipids*. 33, 417-425.
- Li, Y., Seifert, M.F., & Ne, D.M. (1999). Dietary conjugated linoleic acids alter serum IGF-1 and IGF binding protein concentrations and reduce bone formation in rats fed (n-6) or (n-3) fatty acids. *Journal of Bone and Mineral Research*, 14, 1153-1162.
- Linehan, C., Gupta, S., Samali, A., & Connor, L.O. (2012). Bisphenol A-mediated suppression of LPL gene expression inhibits triglyceride accumulation during adipogenic Differentiation of human adult stem cells. *PLoS One*, 2012.
- Marangoni, F., Corsello, G., Cricelli, C., Ferrara, N., Ghiselli, A., Lucchin, L., & Poli, A. (2015). Role of poultry meat in a balanced diet aimed at maintaining health and wellbeing: an Italian consensus document. *Food & Nutrition Research*, 59.
- Martin, D., Muriel, E., Gonzalez, E., Viguera, J., & Ruiz, J. (2008). Effect of dietary conjugated linoleic acid and monounsaturated fatty acids on productive, carcass and meat quality traits of pigs. *Livestock Science*, 117, 155-164.
- Martin, J.C., Grégoire S Siess, M.H., Genty, M., Chardigny, J.M., Berdeaux, O., Juanéda, P., & Sébédio, J.L. (2000). Effects of conjugated linoleic acid isomers on lipid-mobilizing enzymes in male rats. *Lipids*, 35, 91 - 98.
- Matsubara, Y., Sato, K., Ishii, H., & Akiba, Y. (2005). Changes in mRNA expression of regulatory factors involved in adipocyte differentiation during fatty acid induced adipogenesis in chicken. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 141, 108-115.
- McLeod, R.S., LeBlanc, A.M., Langille, M.A., Mitchell, P.L., & Currie, D.L. (2004). Conjugated linoleic acids, atherosclerosis, and hepatic very-low-density lipoprotein metabolism. *The American Journal of Clinical Nutrition*, 79, 1169-1174.

- McNeel, R.L., & Mersmann, H.J. (2003). Effects of isomers of conjugated linoleic acid on porcine adipocyte growth and differentiation. *The Journal of Nutritional Biochemistry*, 14, 266-274.
- Mehr, M.A., Hassanabadi, A., Mirghelenj, S.A., & Kermanshasi, H. (2014). Effect of in ovo injection of conjugated linoleic acid on immune status and blood biochemical factors of broilers chickens. *Spanish Journal Of Agricultural Research*, 12, 455-461.
- Mercier, Y., Gatellier, P., Viau, M., Remignon, H., Renerre, M. (1998). Effect of dietary fat and vitamin E on colour stability and on lipid and protein oxidation in turkey meat during storage. *Meat Science*, 48, 301–318.
- MIDA, Malaysian Investment Development Authority. (2014). Food Technology and Sustainable Resources. <http://www.mida.gov.my/home/food-technology-and-sustainable-resources/posts/>. Retrieve on 5 July 2015.
- Miller, C.C., Park, Y., Pariza, M.W., & Cook, M.E. (1994). Feeding conjugated linoleic acid to animals partially overcomes catabolic responses due to endotoxin injection. *Biochemical and Biophysical Research Communications*. 198, 1107-1112.
- Min, B., & Ahn, D. (2005). Mechanism of lipid peroxidation in meat and meat products- A review. *The Food Science and Biotechnology*, 14, 152-163.
- Muller, L.D., & Delahoy, J.E. (2004). CLA implications for animal production and human health. <http://extension.psu.edu/animals/dairy/nutrition/nutrition-and-feeding/nutrition-and-health/conjugated-linoleic-acid-cla-implications-for-animal-production-and-human-health>. 2004, Retrieved 10 July 2014.
- National Research Council (NRC). Nutrient requirements of poultry. 9th Revision Edition. 1994, NAS-NRC, Washington, DC.
- Nazare, J.A., de la Perriere, A.B., Bonnet, F., Desage, M., Peyrat, J., Maitrepierre, C., Louche-Pelissier, C., Bruzeau, J., Goudable, J., & Lassel, T. (2007). Daily intake of conjugated linoleic acid-enriched yoghurts: effects on energy metabolism and adipose tissue gene expression in healthy subjects. *British Journal of Nutrition*, 97, 273-280.
- Nicolosi, R.J., Rogers, E.J., Kritchevsky, D., Scimeca, J.A., & Huth, P.J. (1997). Dietary conjugated linoleic acid reduces plasma lipoproteins and early atherosclerosis in hypercholesterolemic hamsters. *Artery*, 22, 266-277.
- Ogawa, J., Matsumura, K., Kishino, S., Omura, Y., & Shimizu, S. (2001). Conjugated linoleic acid accumulation via 10-hydroxy-12-octadecaenoic acid during microaerobic transformation of linoleic acid by *Lactobacillus acidophilus*. *Applied and Environmental Microbiology*, 67, 1246-1252.
- Okruszek, A. (2012). Fatty acid composition of muscle and adipose tissue of indigenous Polish geese breeds. *Archiv fur Tierzucht*, 3, 294-302.

- Ostrowska, E., Muralitharan, M., Cross, R.F., Bauman, D.E., & Dunshea, F.R. (1999). Dietary conjugated linoleic acids increase lean tissue and decrease fat deposition in growing pigs. *Journal of Nutrition*, 129, 2037-2042.
- Palombo, J., Ganguly, A., Bistrain, B., & Menard, M. (2002). The antiproliferative effects of biologically active isomers of conjugated linoleic acid on human colorectal and prostatic cancer cells. *Cancer Letter*, 177, 163-172.
- Pandit, A., Anand, S., Kalscheur, K., & Hassan A. (2012). Production of conjugated linoleic acid by lactic acid bacteria in milk without any additional substrate. *International Journal of Dairy Technology*, 65, 603-608.
- Pariza, M.W., & Hargraves, W.A. (1985). A beef derived mutagenesis modulator inhibits initiation on mouse epidermal tumors by 7, 12-dimethylbenz(a)anthracene. *Carcinogenesis*, 6, 591-594.
- Pariza, M.W., Park, Y., & Cook, M.E. (2000). Mechanisms of action of conjugated linoleic acid: Evidence and speculation (Review). *Proceedings of the Society for Experimental Biology and Medicine*, 223, 8-13.
- Pariza, M.W., Park, Y., & Cook, M.E. (2001). The biologically active isomers of conjugated linoleic acid. *Progress in Lipid Research*, 40, 283-289.
- Park, H.S., Ryu, J.H., Ha, Y.L., & Park, H.Y. (2001). Dietary conjugated linoleic acid (CLA) induces apoptosis of colonic mucosa in 1,2-dimethylhydrazine-treated rats: a possible mechanism of the anticarcinogenic effect by CLA. *British Journal Nutrition*, 86, 549-55.
- Park, Y., Albright, K., Liu, W., Storkson, J., Cook, M., & Pariza, M. (1997). Effect of conjugated linoleic acid on body composition in mice. *Lipids*, 32, 853-858.
- Park, Y., Storkson, J.M., Albright, K.J., Liu, W., & Pariza, M.W. (1999). Evidence that the trans-10, cis-12 isomer of conjugated linoleic acid induces body composition changes in mice. *Lipids*, 34, 235-241.
- Pereira, S.L., Leonard, A.E., & Mukerji, P. (2003) Recent advances in the study of fatty acid desaturases from animals and lower eucaryotes. *Prostaglandins, leukotrienes, and essential fatty acids*, 68, 97-106.
- Peters, J.M., Park, Y., Gonzalez, F.J., & Pariza, M.W. (2001). Influence of conjugated linoleic acid on body composition and target gene expression in peroxisome proliferator-activated receptor α -null mice. *Biochimica et Biophysica Acta*, 1533, 233-242.
- Petracci, M., & Cavani, C. (2012). Meat growth and poultry meat quality issues. *Nutrients*, 4, 1-12.
- Petrik, M.B.H., McEntee, M.F., Johnson, B.T., Obukowicz, M.G., & Whelan, J. (2000). Highly unsaturated (n-3) fatty acids, but not α -linolenic, conjugated linoleic or γ -linolenic acids, reduce tumorigenesis in apcmin/+ mice. *Journal of Nutrition*, 130, 2434-2443.

- Poulos, S.P., Sisk, M., Hausman, D.B., Azain, M.J., & Hausman, G.J. (2001). Pre- and post-natal dietary conjugated linoleic acid alters adipose tissue development, body weight gain and body composition in Sprague-Dawley rats. *Journal of Nutrition*, 2001, 131, 2722-2731.
- Qi, R., Yang, F., Huang, J., Peng, H., Liu, Y., & Liu, Z. (2014). Supplementation with conjugated linoleic acid decreases pig back fat deposition by inducing adipocyte apoptosis. *BMC Veterinary Research*, 10, 141.
- Rabobank, (2014). Global poultry prices taking off in Q2. <http://www.worldpoultry.net/Broilers/Markets--Trade/2014/6/Rabobank-Global-poultry-prices-taking-off-in-Q2-1549335W/>. Retrieved on 11 August 2014.
- Raes, K., Huyghebaert, G., De Smet, S., Nollet, L., Arnouts, S., & Demeyer, D. (2002). The deposition of conjugated linoleic acids in eggs of laying hens fed diets varying in fat level and fatty acid profile. *Journal of Nutrition*, 132, 182-189.
- Rajion, M.A., McLean, J.G., & Cahill, R.N. (1985). Essential fatty acids in the fetal and newborn lamb. *Australian Journal of Biological Sciences*, 38, 33-40.
- Ramiah, S.K., Meng, G.Y., & Ebrahimi, M. (2014a). Dietary conjugated linoleic acid alters oxidative stability and alleviates plasma cholesterol content in meat of broiler chickens. *Scientific World Journal*, 2014, Article ID 949324:10 pages.
- Ramiah, S.K., Meng, G.Y., Sheau, T.W., Keong, Y.S., & Ebrahimi, M. (2014b). Dietary conjugated linoleic acid supplementation leads to downregulation of PPAR transcription in broiler chickens and reduction of adipocyte cellularity. *PPAR Research*, 2014, Article ID 137652:10 pages.
- Ramsay, T. G., & Rosebrough, R. W. (2003). Hormonal regulation of postnatal chicken preadipocyte differentiation in vitro. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 136, 245-253.
- Resnyk, C.W., Carré, W., Wang, X., Porter, T.E., Simon, J., Le Bihan-Duval, E., Duclos, M.J., Aggrey, S.E., & Cogburn, L.A. (2013). Transcriptional analysis of abdominal fat in genetically fat and lean chickens reveals adipokines, lipogenic genes and a link between hemostasis and leanness. *BioMed Central Genomics*, 14, 557.
- Richards, M.P., Poch, S.M., Coon, C.N., Rosebrough, R.W., Ashwell, C.M., & McMurtry, J.P. (2003). Feed restriction significantly alters lipogenic gene expression in broiler breeder chickens. *Journal of Nutrition*, 133, 707-715.
- Riserus, U., Arner, P., Brismar, K., & Vessby, B. (2002). Treatment with dietary trans-10cis-12 conjugated linoleic acid causes isomer-specific insulin resistance in obese men with the metabolic syndrome. *Diabetes Care*, 2002, 25, 1516-1521.
- Riserus, U., Berglund, L., & Vessby, B. (2001). Conjugated linoleic acid (CLA) reduced abdominal adipose tissue in obese middle-aged men with signs of the

- metabolic syndrome: a randomised controlled trial. *International Journal of Obesity Related Metabolic Disorders*, 25, 1129–1135.
- Risérus, U., Vessby, B., Årnlöv, J., & Basu, S. (2004). Effects of *cis*-9, *trans*-11 conjugated linoleic acid supplementation on insulin sensitivity, lipid peroxidation, and proinflammatory markers in obese men. *The American Journal of Clinical Nutrition*, 80, 2279-283.
- Rodbell, M. (1964). Metabolism of isolated fat cells. I. Effects of hormones on glucose metabolism and lipolysis. *The Journal of Biological Chemistry*, 239, 375-380.
- Rosen, E.D., Walkey, C.J., Puigserver, P., & Spiegelman, B.M. (2000). Transcriptional regulation of adipogenesis. *Genes Development*, 14, 1293-1307.
- Royan, M., Meng, G.Y., Othman, F., Sazili, A.Q., & Navidshad B. (2011b) Effects of dietary conjugated linoleic acid (CLA), n-3 and n-6 fatty acids on performance and carcass traits of broiler chickens. *African Journal of Biotechnology*, 10, 17379-17384.
- Royan, M., Meng, G.Y., Othman, F., Sazili, A.Q., & Navidshad, B. (2011a). Effects of conjugated linoleic Acid, fish Oil and soybean Oil on PPARs (α & γ) Mrna expression in broiler chickens and their relation to body fat deposits. *International Journal of Molecular Science*, 12, 8581-8595.
- Rule, D.C., Broughton, K.S., Shellito, S.M., & Maiorano, G. (2002). Comparison of musclefatty acid profiles and cholesterol concentrations of bison, beef cattle, elk, and chicken. *Journal of Animal Science*, 80, 1202-1211.
- Saadoun, A. (2014). New approaches on the importance of fatty acids in poultry meat on human health. *Cuban Journal of Agricultural Science*, 48, 59-61.
- Sadur, C.N., Yost, T.J., & Eckel, R.H. (1984). Fat feeding decreases insulin responsiveness of adipose tissue lipoprotein lipase. *Metabolism*, 33, 1043-1047.
- Sahin, H., Uyanik, F., & Inanc, N. (2008). Effects of conjugated linoleic acid on body composition and selected biochemical parameters in obese women. *Pakistan Journal of Nutrition*, 7, 546-549.
- Sahraei, M. (2012). Feed Restriction In Broiler Chickens Production. *Biotechnology in Animal Husbandry*, 28, 333-352.
- Santora, J.E., Palmquist, D.L., & Roehriq, K.L. (2000). Trans-vaccenic acid is desaturated to conjugated linoleic acid in mice. *Journal of Nutrition*, 130, 208-215.
- Sanz, M., Flores, A., Perez de Ayala, P., & Lo'pez-Bote, C.J. (1999). Higher lipid accumulation in broilers fed saturated fats than those fed unsaturated fats. *British Poultry Science*, 40, 95-101.

- Sato, K., & Akiba, Y. (2002). Lipoprotein lipase mRNA expression in abdominal adipose tissue is little modified by age and nutritional state in broiler chickens. *Poultry Science*, 2002, 81, 846-852.
- Sato, K., Abe, H., Kono, T., Yamazaki, M., Nakashima, K., Kamada, T., & Akiba, Y. (2009). Changes in peroxisome proliferator-activated receptor gamma gene expression of chicken abdominal adipose tissue with different age, sex and genotype. *Animal Science Journal*, 80, 322–327.
- Sato, K., Matsushita, K., Matsubara, Y., Kamada, T., & Akiba, Y. (2008). Adipose tissue fat accumulation is reduced by a single intraperitoneal injection of peroxisome proliferator-activated receptor gamma agonist when given to newly hatched chicks. *Poultry Science*, 87, 2281-2286.
- Sato, K., Yonemura, T., Ishii, H., Toyomizu, M., Kamada, T., & Akiba, Y. (2009). Role of peroxisome proliferator-activated receptor beta/delta in chicken adipogenesis. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 154, 370-375.
- Sato, K.Y., Akiba, Y., Chida, Y., & Takahashi, K. (1999). Lipoprotein hydrolysis and fat accumulation in chicken adipose tissues are reduced by chronic administration of lipoprotein lipase monoclonal antibodies. *Poultry Science*, 78, 1286-1291.
- Satory, D.L., & Smith, S.B., (1999). Conjugated linoleic acid inhibits proliferation but stimulates lipid filling of murine 3T3-L1 preadipocytes. *Journal of Nutrition*, 129, 92-97.
- Schmid, A., Collomb, M., Sieber, R., & Bee, G. (2006). Conjugated linoleic acid in meat and meat products: A review. *Meat Science*, 73, 29-41.
- Shang, X.G., Wang, F.L., Li, D.F., Yin, J.D., & Li, J.Y. (2004). Effects of dietary conjugated linoleic acid on the productivity of laying hens and egg quality during refrigerated storage. *Poultry Science*, 83, 1688-1695.
- Shin, D., Kakani, G., Karimi, A., Cho, Y.M., Kim, S.W., Ko, Y.G., Shim, K.S., & Park, J.H. (2011a). Influence of dietary conjugated linoleic acid and its combination with flaxseed oil or fish oil on saturated fatty acid and n-3 to n-6 fatty acid ratio in broiler chicken meat. *Asian Australasian Journal of Animal Science*, 24, 1249–1255.
- Shin, D., Narciso-Gaytán, C., Park, J.H., Smith, S.B., Sánchez-Plata, M.X., & Ruiz-Feria, C.A. (2011b). Dietary combination effects of conjugated linoleic acid and flaxseed or fish oil on the concentration of linoleic and arachidonic acid in poultry meat. *Poultry Science*, 90, 1340–1347.
- Shinn, S., Liyanage, R., Lay, J., & Proctor, A. (2014). Improved fatty acid analysis of conjugated linoleic acid-rich egg yolk triacylglycerols and phospholipid species. *Journal of Agriculture and Food Chemistry*, 62, 6608-6615.

- Simon, O., Maenner, K., Schaefer, K., Sagredos, A., & Eder, K. (2000). Effects of conjugated linoleic acids on protein to fat proportions, fatty acids, and plasma lipids in broilers. *European Journal of Lipid Science and Technology*, 102, 402-10.
- Sirri, F., Tallarico, N., Meluzzi, A., & Franchini, A. (2003). Fatty acid composition and productive traits of broiler fed diets containing conjugated linoleic acid. *Poultry Science*, 82, 1356-1361.
- Sisk, M.B., Hausman, D.B., Martin, R.J., & Azain, M.J. (2001). Dietary conjugated linoleic acid reduces adiposity in lean but not obese Zucker rats. *Journal of Nutrition*, 13, 1668-1674.
- Skibsted, L.H. (1996). Chemical changes in meat and meat products during storage, transportation and retail display—theoretical consideration. S.A Taylor, A Raimundo, M Severini, F.J.M Smulders (Eds.), *Meat quality and meat packaging*, ECCEAMST, Utrecht, The Netherlands, 169–181.
- Smith, S.B., Hively, T.S., Cortese, G.M., Han, J.J., Chung, K.Y., & Casteñada, P. (2002). Conjugated linoleic acid depresses the D9 desaturase index and stearoyl coenzyme A desaturase enzyme activity in porcine subcutaneous adipose tissue. *Journal of Animal Science*, 80, 2110-2115.
- Sneddon, A.A., Tsofliou, F., Fyfe, C.L., Matheson, I., Jackson, D.M., Horgan, G., Winzell, M.S., Wahle, K.W., Ahren, B., & Williams, L.M. (2008). Effect of a conjugated linoleic acid and omega-3 fatty acid mixture on body composition and adiponectin. *Obesity*. 16, 1019-1024.
- Snel, M., Jonker, J.T., & Schoones, J. (2012). Ectopic fat and insulin resistance: pathophysiology and effect of diet and lifestyle interventions. *International Journal of Endocrinology*, 2014, Article ID 983814.
- Soukas, A., Socci, N.D., Saatkamp, B.D., Novelli, S., & Friedman, J.M. (2001). Distinct transcriptional profiles of adipogenesis in vivo and in vitro. *Journal of Biological Chemistry*, 276, 34167-34174.
- Stangierski, J., & Lesnierowski, G. (2015). Nutritional and health-promoting aspects of poultry meat and its processed products. *World's Poultry Science Journal*, 71, 71-82.
- Stangl, G.I. (2000). High dietary levels of a conjugated acid mixture alter hepatic glycerophospholipid class profile and cholesterol-carrying serum lipoproteins of rats. *Journal of Nutritional Biochemistry*, 11, 184–191.
- Stewart, A.B., Kelley, G., & Kimathi, B. (2014). Expression of potential regulatory genes in abdominal adipose tissue of broiler chickens during early development. *Genetic Research International*, 2014: Article ID 318304.
- Suksombat, W., Boonmee, T., & Lounglawan, P. (2007). Effects of various levels of conjugated linoleic acid supplementation on fatty acid content and carcass composition of broilers. *Poultry Science*, 86, 318-324.

- Szymczyk, B., & Pisulewski, P.M. (2003). Effects of dietary conjugated linoleic acid on fatty acid composition and cholesterol content of hen egg yolks. *British Journal of Nutrition*, 90, 93-99.
- Szymczyk, B., Pisulewski, P., Szczurek, W., & Hanczakowski, P. (2001). Effects of conjugated linoleic acid on growth performance, feed conversion efficiency, and subsequent carcass quality in broiler chickens. *British Journal of Nutrition*, 85, 465-473.
- Takada, I., & Kobayashi, M. (2013). Structural features and transcriptional activity of chicken PPARs (α , β , and γ). *PPAR Research*, 2013, Article ID 186312.
- Takahashi, Y., Kushiro, M., Shinohara, K., & Ide, T. (2002). Dietary conjugated linoleic acid reduces body fat mass and affects gene expression of proteins regulating energy metabolism in mice. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 133, 395-404.
- Taylor, C. G., & Zahradka, P. (2004). Dietary conjugated linoleic acid and insulin sensitivity and resistance in rodent models. *The American Journal of Clinical Nutrition*, 79, 1164S-1168S.
- Tekeleselassie, A.W., Goh, Y.M., Rajion, M.A., Motshakeri, M., & Ebrahimi, M. (2013). A high-fat diet enriched with low omega-6 to omega-3 fatty acid ratio reduced fat cellularity and plasma leptin concentration in Sprague-Dawley rats. *The Scientific World Journal*, 2013, 2013, 7 pages.
- Terpstra, A. H. (2004). Effect of conjugated linoleic acid on body composition and plasma lipids in humans: an overview of the literature. *The American Journal of Clinical Nutrition*, 79, 352-361.
- Terpstra, A.H., Beynen, A.C., Everts, H., Kocsis, S., Katan, M.B., & Zock, P.L. (2002). The decrease in body fat in mice fed conjugated linoleic acid is due to increases in energy expenditure and energy loss in the excreta. *Journal of Nutrition*, 132, 940-945.
- Thiel-Cooper, R.L., Parrish, J.R.F.C., Sparks, J.C., Wiegand, B.R., & Ewan, R.C. (2001). Conjugated linoleic acid changes swine performance and carcass composition. *Journal of Animal Science*, 79, 1821-1828.
- Tontonoz, P., Hu, E., & Graves, R.A. (1994). mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. *Genes Development*, 8, 1224-1234.
- Tsuboyama K.N., Takahashi, M., Tanemura, K., Kim, H.J., Tange, T., Okuyama, H., Kasai, M., Ikemoto, S., & Ezaki, O. (2000). Conjugated linoleic acid supplementation reduces adipose tissue by apoptosis and develops lipodystrophy in mice. *Diabetes*, 49, 1534 -1542.
- Tumova, E., & Teimouri A. (2010). Fat deposition in the broiler chicken: A review. *South African Breweries*, 41, 121-128.

- Turpeinen, A.M., Mutanen, M., Aro, A., Salminen, I., Basu, S., Palmquist, D.L., & Griinari, J.M. (2001). Bioconversion of vaccenic acid to conjugated linoleic acid in humans. *The American Journal of Clinical Nutrition*, 76, 504-510.
- Van Den Berg, J.J.M., Cook, N.E., & Tribble, D.L. (1995). Reinvestigation of the antioxidant properties of conjugated linoleic acid. *Lipids*, 30, 599-605.
- Vandesompele J., De, P.K., Pattyn, F., Poppe, B., Van, R.N., De, P.A., & Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, 3, RESEARCH0034.
- Vidal-Puig, A.J., Considine, R.V., & Jimenez-Liñan, M. (1997). Peroxisome proliferator-activated receptor gene expression in human tissues: effects of obesity, weight loss, and regulation by insulin and glucocorticoids. *Journal Clinical Investigation*, 99, 2416-2422 (1997).
- Visioli, F., Giordano, E., Nicod, N.M., & Dávalos, A. (2012). Molecular targets of omega 3 and conjugated linoleic fatty acids – “Micromanaging” cellular response. *Frontiers in Physiology*, 3, 42.
- Wang, H.B., Li, H., Wang, Q.G., Zhang, X.Y., Wang, S.Z., Wang, Y.X., & Wang, X.P. (2007). Profiling of chicken adipose tissue gene expression by genome array. *BMC Genomics*, 27, 193-207 (2007).
- Wang, L.S., Huang, Y.W., Liu, S.L., Yan, P., & Lin, Y.C. (2008a). Conjugated linoleic acid induces apoptosis through estrogen receptor alpha in human breast tissue. *BMC Cancer*, 8, 208.
- Wang, Y., & Jones, P. (2004). Dietary conjugated linoleic acid and body composition. *The American Journal of Clinical Nutrition*, 79, 1153S-1158S.
- Wang, Y., Mu, Y., Li, H., Ding, N., & Wang, Q. (2008b). Peroxisome proliferator-activated receptor- γ gene: a key regulator of adipocyte differentiation in chickens. *Poultry Science*, 87, 226-232.
- Warren, J.M., Simon, V.A., Bartolini, G., Erickson, K.L., Mackey, B.E., & Kelley, D.S. (2003). Trans -10,cis -12 CLA increases liver and decreases adipose tissue lipids in mice: Possible roles of specific lipid metabolism genes. *Lipids*, 38, 497-504.
- Watkins, B.A., Feng, S., Strom, A.K., DeVitt, A.A., Yu, L., & Li, Y. (2003). Conjugated linoleic acids alter the fatty acid composition and physical properties of egg yolk and albumen. *Journal of Agriculture and Food Chemistry*, 51, 6870-6876.
- Watts, J.L., & Browse, J. (2002). Genetic dissection of polyunsaturated fatty acid synthesis in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 5854-5859.

- Weinstock, P.H., Frank, S.L., Hudgins, L.C., Radner, H., Friedman, J.M., Zechner, R., & Breslow, J.L. (1997). Lipoprotein lipase controls fatty acid entry into adipose tissue, but fat mass is preserved by endogenous synthesis in mice deficient in adipose tissue lipoprotein lipase. *Proceedings of the National Academy of Sciences of the United States of America*, 94, 10261-10266.
- West, D., Delany, J., Camet, P., Blohm, F., Truett, A., & Scimeca, J. (1998). Effects of conjugated linoleic acid on body fat and energy metabolism in the mouse. *The American Journal of Physiology*, 275, 667-672.
- Whigham, L.D., Watras, A.C., & Schoeller, D.A. (2007). Efficacy of conjugated linoleic acid for reducing fat mass: A meta-analysis in humans. *The American Journal of Clinical Nutrition*, 85, 1203-1211.
- Wiegand, B.R., Sparks, J.C., Parrish, J.F.C., Zimmerman, D.R. (2002). Duration of feeding conjugated linoleic acid influences growth performance, carcass traits, and meat quality of finishing barrows. *Journal of Animal Science*, 80, 637-643.
- Wilson, T.A., Nicolosi, R.J., Chrysam, M., & Kritchevsky, D. (2000). Conjugated linoleic acid reduces early atherosclerosis greater than linoleic acid in hypercholesterolemic hamsters. *Nutrition Research*, 20, 1795-805.
- Wong, J., Kim, Y., Park, Y., Lee, S.H., Baek, S.J., & Par, Y. (2015) Isomer specificity of conjugated linoleic acid on suppression of osteosarcomas. *Journal of Nature and Science*, 1, e67.
- Xu, Z.R., Wang, M.Q., Mao, H.X., Zhan, X.A., & Hu, C.H. (2003). Effects of L-carnitine on growth performance, carcass composition, and metabolism of lipids in male broilers. *Poultry Science*, 82, 408-413.
- Yamamoto, H., Kurebayashi, S., Hirose, T., Kouhara, H., & Kasayama, S. (2002) Reduced IRS-2 and GLUT4 expression in PPAR γ 2- induced adipocytes derived from C/EBP β and C/EBP δ - deficient mouse embryonic fibroblasts. *Journal of Cell Science*. 115, 3601-3607.
- Yan, H.J., Lee, E.J., Nam, K.C., Min, B.R., & Ahn, D. (2006). Dietary functional ingredients: Performance of animals and quality and storage stability of irradiated raw turkey breast. *Poultry Science*, 85, 1829-1837.
- Yang, L., Huang, Y., & Chen, Z.Y. (2000). Oxidative stability of conjugated linoleic acid isomers. *Journal of Agriculture and Food Chemistry*. 48, 3072-3076.
- Yang, Z.H., Miyahara, H., Iwasaki, Y., Takeo, J., & Katayama, M. (2013). Dietary supplementation with long-chain monounsaturated fatty acids attenuates obesity-related metabolic dysfunction and increases expression of PPAR gamma in adipose tissue in type 2 diabetic KK-Ay mice. *Nutrition and Metabolism*, 10, 16.
- Yaqiong, Y., Shumao, L., Heping, M., Xiaohong, T., Yangdan, O., Jian, C., Yongjiang M., & Yugu, Li. (2014). Analysis of differentially expressed genes and signaling pathways related to intramuscular fat deposition in skeletal muscle

of sex-linked dwarf chickens, *BioMed Research International*, 2014, 2014, Article ID 724274, 7 pages, 2014.

- Yari, P., Goudarzi, A. M., Ahmadi, M., Masoumi, J., & Azad, S. K. (2015). Influences of Conjugated linoleic acid on poultry products. *International Journal of Life Sciences*, 9(4), 1-5.
- Yin, M.C., & Faustman, C. (1993). The influence of temperature, bpH and phospholipid composition upon the stability of myoglobin and phospholipids: A liposome model. *Journal of Agriculture and Food Chemistry*, 41,853–857.
- Zabala, A., Churruga, I., Fernández-Quintela, A., Rodríguez, V.M., Macarulla, M. T., Martínez, J. A., & Portillo, M. P. (2006). trans-10, cis-12 Conjugated linoleic acid inhibits lipoprotein lipase but increases the activity of lipogenic enzymes in adipose tissue from hamsters fed an atherogenic diet. *British Journal of Nutrition*, 95, 1112-1119.
- Zhang, A., & Chen, Z.Y. (1997). Oxidative stability of conjugated linoleic acids relative to other polyunsaturated fatty acids. *Journal of American Chemical Society*, 74, 1611-1613.
- Zhang, G.M., Wen, J., Chen, J.L., Zhao, G.P., Zheng, M.Q., & Li, W.J. (2007). Effect of conjugated linoleic acid on growth performances, carcass composition, plasma lipoprotein lipase activity and meat traits of chicken. *British Poultry Science*, 48, 217–223.
- Zhang, H., Guo, Y., & Yuan, J. (2005). Conjugated linoleic acid enhanced the immune function in broiler chicks. *British Journal of Nutrition*, 94, 746-752.
- Zhang, H.J., Guo, Y.M., Yang, Y., & Yuan, J.M. (2006). Dietary conjugated linoleic acid enhances spleen PPAR- γ mRNA expression in broiler chicks, *British Poultry Science*, 47, 726-733.
- Zhang, X.H., Wang, B., Long, F., Wang, L., & Yang, Z. (2009). The effects of dietary conjugated linoleic acid (CLA) on fatty acid composition and key enzymes of fatty acid oxidation in liver and muscle of geese. *Turkish Journal of Veterinary Animal Science*, 33, 215-222.
- Zhang, Y., Hongtrakul, K., Ma, Q.G., Liu, L.T., & Hu, X.X. (2009b). Effects of dietary alpha-lipoic acid on anti-oxidative ability and meat quality in Arbor Acres broilers. *Asian Australasian Journal of Animal Science*, 22, 1195–1201.
- Zhao, X., Mo, D., Li, A., Gong, W., Xiao, S., Zhang, Y., Qin, L., Niu, Y., Guo, Y., Liu, X., Cong, P., He, Z., Wang, C., Li, J., & Chen, Y. (2011). Comparative analyses by sequencing of transcriptomes during skeletal muscle development between pig breeds differing in muscle growth rate and fatness. *PLoS One*, 5, 1-18.

- Zhou, J. (2008). Effect of dietary conjugated linoleic acid (CLA) on abdominal fat deposition in yellow-feather broiler chickens and its possible mechanism. *Asian Australasian Journal Animal Science*, 21, 1760–1765.
- Zhou, X., Li, D., Yin, J., Ni, J., Dong, B., Zhang, J., & Du, M. (2007). CLA differently regulates adipogenesis in stromal vascular cells from porcine subcutaneous adipose and skeletal muscle. *Journal of Lipid Research*, 48, 1701-1709.
- Zlatanos, S.N., Laskaridis, K., & Sagredos, A. (2008). Conjugated linoleic acid content of human plasma. *Lipids*, 7, 34.



© COPYRIGHT UPM

