



UNIVERSITI PUTRA MALAYSIA

***ESTABLISHMENT OF JAVA MEDAKA (*Oryzias javanicus* Bleeker)
AS TEST ORGANISM FOR ECOTOXICOLOGICAL STUDIES
IN TROPICAL REGION***

SHAHRIZAD BINTI YUSOF

FS 2014 40



**ESTABLISHMENT OF JAVA MEDAKA (*Oryzias javanicus* Bleeker)
AS TEST ORGANISM FOR ECOTOXICOLOGICAL STUDIES
IN TROPICAL REGION**

By

SHAHORIZAD BINTI YUSOF

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfillment of the Requirements for the Degree of Doctor of
Philosophy**

June 2014

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**ESTABLISHMENT OF JAVA MEDAKA (*Oryzias javanicus* Bleeker)
AS TEST ORGANISM FOR ECOTOXICOLOGICAL STUDIES
IN THE TROPICAL REGION**

By

SHAHORIZAD BINTI YUSOF

June 2014

Chair : Ahmad Bin Ismail, PhD
Faculty : Science

The medaka fish is a group of small fish distributed in large areas in Asia. This study attempts to establish Java medaka (*Oryzias javanicus*) as a test organism for ecotoxicological studies in order to gain knowledge of the changes in biological processes of other organisms when exposed to pollutants. The ecology of Java medaka was investigated to understand their responses to environmental pollutants. The findings covered the distribution and localities of the fish, habitat preference or niche they occupied, relationship with other medaka species, and their spawning ground. Thirty eight localities in the coastal areas in Peninsular Malaysia were surveyed from 2008 to 2013. All localities surveyed in the west and south coast are inhabited by Java medaka in abundance but they do not occur in the east coast. In several localities, Java medaka co-exists with Indian medaka (*Oryzias dancena*), a phenomenon that has never been reported before. In other localities they inhabit different niche with Indian medaka, preferring areas with higher salinity while Indian medaka choose lower salinity areas. Java medaka can tolerate a wide salinity range (0.2 – 30.0 ppt), making them suitable to represent freshwater and marine environment. To support research activities using Java medaka, the laboratory culture was established. It is carried out in ambient temperature, thus, it can represent almost similar condition with the fish habitat. Their suitability for laboratory cultivation, ease of maintenance, regular spawning, short embryonic period and generation time make Java medaka a convenient test organism. The laboratory culture established in this study further ensures that the test fish is of high quality, free from parasites and diseases. The sensitivity of different life stages of Java medaka to environmental pollutants was tested. All life stages of the fish has been utilised and they have shown particular sensitivity. The embryos were sensitive to low concentrations (0.01 – 0.05 ppm) of heavy metals (Cd, Hg, Pb, Cu and Zn) in terms of developmental impairments. The most significant impairment was embryonic death which was metal dependent where the rank of metal severity is as follows: Cd (47.0 - 100%) > Hg (22.0 - 64.0%) > Pb (10.0 - 55.0%) > Zn (6.0 - 40.0%) > Cu (4.0

- 38.0%). In exposure to glyphosate-based herbicide, embryonic death was found to be directly proportional to the exposure concentrations. Other developmental endpoints in embryos exposed to the pollutants tested include embryotoxicity and teratogenicity. The developmental stages of Java medaka constitute simple and effective models for evaluating pollutant impact. The induction of micronucleus formation and nuclear abnormalities when the adult fish were exposed to very low levels of Cd and Hg (1.0 and 10.0 ppb) and the induction of only nuclear abnormalities in exposure to higher concentrations of Cu and Zn (10.0 and 100.0 ppb) showed the difference in their genotoxic responses towards different metals. Utilizing Java medaka in testing will closely reflect the impacts of toxic pollutants on the environment particularly for the tropical region. As the fish is new to many local researchers, the findings of this study provide the much needed information. Future researches can utilise the fish in various fields of scientific investigation.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
Sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENUBUHAN JAVA MEDAKA (*Oryzias javanicus* Bleeker)
SEBAGAI ORGANISMA UJIAN DALAM BIDANG EKOTOKSIKOLOGI**

Oleh

SHAHORIZAD BINTI YUSOF

Jun 2014

Pengerusi : Ahmad Bin Ismail, PhD
Fakulti : Sains

Ikan medaka terdiri daripada sekumpulan ikan kecil yang tersebar di kawasan yang luas di Asia. Kajian ini bertujuan untuk menubuhkan 'Java medaka' (*Oryzias javanicus*) sebagai organisma ujian untuk bidang ekotoksikologi dalam usaha untuk mendapatkan maklumat tentang perubahan yang berlaku dalam proses biologi dalam organisma lain apabila terdedah kepada bahan pencemar. Ekologi 'Java medaka' telah dikaji untuk memahami gerak balas mereka terhadap bahan pencemar alam sekitar. Hasil kajian meliputi taburan dan lokaliti, pemilihan habitat dan nic, hubungan dengan spesies medaka lain serta tempat pembiakannya. Tiga puluh lapan lokaliti di kawasan pesisiran Semenanjung Malaysia telah dikaji dari tahun 2008 hingga 2013. Kesemua lokaliti yang dikaji di pantai barat dan selatan mempunyai taburan "Java medaka" yang tinggi tetapi ikan ini tidak terdapat di pantai timur. Dalam beberapa lokaliti, "Java medaka" wujud bersama "Indian medaka" (*Oryzias dancena*), suatu fenomena yang belum pernah dilaporkan sebelum ini. Dalam lokaliti lain mereka hidup dalam nic yang berbeza di mana "Java medaka" memilih kawasan yang mempunyai saliniti tinggi dan "Indian medaka" memilih kawasan bersaliniti rendah. "Java medaka" boleh hidup dalam julat saliniti yang besar (0.2 – 30.0 ppt), yang menyebabkan mereka sesuai untuk mewakili persekitaran air tawar dan marin. Untuk menyokong aktiviti-aktiviti penyelidikan menggunakan 'Java medaka', sistem pengkulturan ikan ini di dalam makmal telah ditubuhkan. Ia dijalankan dalam suhu ambien dan dengan itu dapat mewakili keadaan yang hampir sama seperti habitat ikan tersebut. Kesesuaian untuk dibiakkan di makmal, kemudahan penyenggaraan, kekerapan bertelur, tempoh perkembangan embrio dan generasi yang pendek menjadikan "Java medaka" organisma kajian yang sesuai. Kejayaan pengkulturan ikan ini di dalam makmal memastikan penghasilan ikan yang berkualiti tinggi, bebas dari parasit dan penyakit. Kepekaan dan respon oleh pelbagai peringkat hidup 'Java medaka' kepada beberapa bahan pencemar telah diuji. Semua peringkat hidup ikan ini telah digunakan dan mereka telah menunjukkan kepekaan tertentu. Embrio ikan ini bergerak balas kepada kepekatan logam

berat (Cd, Hg, Pb, Cu and Zn) yang rendah (0.01 - 0,05 ppm) dari segi gangguan perkembangan. Gangguan perkembangan yang paling signifikan ialah kematian embrio yang bergantung kepada jenis logam berat di mana turutan keseriusan adalah seperti berikut: Cd (47.0 - 100%) > Hg (22.0 - 64.0%) > Pb (10.0 - 55.0%) > Zn (6.0 - 40.0%) > Cu (4.0 - 38.0%). Dalam pendedahan kepada herbisid berasaskan glifosat, kematian embrio adalah berkadar terus dengan kepekatan herbisid. Gangguan perkembangan lain akibat pendedahan kepada bahan pencemar yang diuji termasuk keracunan dan kesan teratogenik. Peringkat awal perkembangan 'Java medaka' merupakan model ringkas dan berkesan untuk menilai kesan pencemaran. Aruhan pembentukan mikronukleus dan ketidaknormalan nukleus apabila ikan dewasa didedahkan kepada kepekatan Cd dan Hg yang sangat rendah (1.0 and 10.0 ppb) dan aruhan penghasilan nukleus tidak normal dalam pendedahan kepada kepekatan Cu dan Zn yang lebih tinggi (10.0 and 100.0 ppb) menunjukkan perbezaan dalam gerak balas genotoksik mereka terhadap logam berat yang berbeza. Penggunaan "Java medaka" dalam ujian akan menggambarkan dengan lebih dekat tentang impak bahan pencemar toksik terhadap persekitaran terutama di kawasan tropika. Oleh kerana ikan ini masih belum terkenal di kalangan penyelidik tempatan, dapatan dari kajian ini dapat memberikan maklumat yang sangat diperlukan. Ikan ini boleh digunakan dalam pelbagai bidang kajian saintifik.

ACKNOWLEDGEMENTS

I would like to express the deepest appreciation to my committee chair Professor Dr. Ahmad Ismail, who has shown the attitude and the substance of a genius: he continually and persuasively conveyed a spirit of adventure in regard to research and scholarship, and an excitement in regard to teaching. Without his supervision and constant help this dissertation would not have been possible.

A special gratitude I give to my husband, whose contribution in stimulating suggestions and encouragement, helped me to coordinate my project especially in writing this report.

I would like to thank the Department of Biology, Universiti Putra Malaysia for providing me with a good environment and facilities to complete this project. Special thanks also goes to Universiti Putra Malaysia for sponsoring my trip to Japan for an attachment program related to my PhD research.

I am indebted to Dr. Koji Inoue, Prof. Dr. Jiro Koyama and Dr. Masato Kinoshita for their guidance and encouragement during my visit to their medaka laboratories in Japan.

I am also grateful for the assistance given by my research group mates, particularly during sampling sessions. Sampling trips would not be enjoyable without all of you.

Finally, an honourable mention goes to my loved ones, who have supported me throughout the entire process, both by keeping me harmonious and helping me putting pieces together. I will be grateful forever for your understanding and kindness.

I certify that a Thesis Examination Committee has met on 6 June 2014 to conduct the final examination of Shahrizad binti Yusof on her thesis entitled "Establishment of Java Medaka (*Oryzias javanicus* Bleeker) as Test Organism for Ecotoxicological Studies in Tropical Region" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Faridah binti Qamaruz Zaman, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Che Roos bin Saad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Aziz bin Arshad, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Jiro Koyama, PhD

Professor
Kagoshima University
Japan
(External Examiner)



NORITAH OMAR, PhD

Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 July 2014

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Ahmad Bin Ismail, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Hishamuddin Bin Omar, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

Siti Khadijah Binti Daud, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Shahrizad Binti Yusof (GS18374)

Declaration by Members of Supervisory Committee

This is to confirm that:


- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature:
Name of
Chairman of
Supervisory
Committee:



Ahmad Bin Ismail, PhD

Signature:
Name of
Member of
Supervisory
Committee:



Hishamuddin Bin Omar, PhD

Signature:
Name of
Member of
Supervisory
Committee:



Siti Khadijah Binti Daud, PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvi
 CHAPTER	
 1 INTRODUCTION	 1
 2 LITERATURE REVIEW	 5
2.1 Aquatic Pollution	5
2.2 Bioindicator, Biomonitor and Test Organism for Aquatic Pollution	8
2.3 Fish as Sentinel Species for Ecotoxicology	13
2.4 The Medaka Fish and Scientific Researches	15
2.5 The Potential of Java Medaka as Test Organism	16
2.5.1 Taxonomy of Java medaka	16
2.5.2 Important characteristics of Java medaka	17
2.5.3 Importance of laboratory culture of test Organism	18
 3 ECOLOGY OF JAVA MEDAKA	 19
3.1 Introduction	19
3.2 Materials and Methods	21
3.2.1 Distribution of Java medaka in Peninsular Malaysia	21
3.2.2 Ecological parameters of microhabitat for Java medaka	22
3.2.3 Tidal migration	22
3.2.4 Spawning ground	24
3.2.5 Co-existence of Java medaka and Indian medaka	24
3.3 Results and Discussion	26
3.3.1 Distribution of Java medaka in Peninsular Malaysia	26
3.3.2 Ecological parameters of microhabitat of Java medaka	29

3.3.3	Tidal migration	33
3.3.4	Spawning ground	36
3.3.5	Co-existence of Java medaka and Indian medaka	38
3.4	Conclusion	40
4	LABORATORY CULTURE OF JAVA MEDAKA	42
4.1	Introduction	42
4.2	Materials and Methods	42
4.2.1	Sampling of Java medaka	42
4.2.2	Acclimatisation of Java medaka	44
4.2.3	Re-circulating system	45
4.2.4	Culture and feeding	47
4.2.5	Methods for collecting Java medaka eggs	51
4.2.6	Surface sterilization of Java medaka eggs	52
4.2.7	Maintenance of Java medaka culture	52
4.3	Results and Discussion	53
4.3.1	Sampling of Java medaka	53
4.3.2	Acclimatisation of Java medaka	55
4.3.3	Re-circulating system	56
4.3.4	Culture and feeding	57
4.3.5	Methods for collecting Java medaka eggs	66
4.3.6	Surface sterilization of Java medaka eggs	67
4.4	Conclusion	68
5	JAVA MEDAKA AS TEST ORGANISM	69
5.1	Introduction	69
5.2	Materials and Methods	70
5.2.1	Responses of early life stages of Java medaka to pollutants	70
5.2.2	Responses of juvenile Java medaka to glyphosate-based herbicide.	72
5.2.3	Responses of adult Java medaka to heavy metals	72
5.3	Results and Discussion	73
5.3.1	Responses of early life stages of Java medaka to pollutants	73
5.3.2	Responses of juveniles Java medaka to glyphosate based herbicide	92
5.3.3	Responses of adult Java medaka to heavy metals	96
5.4	Conclusion	101
6	GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	102
	BIBLIOGRAPHY	107
	BIODATA OF STUDENT	133
	LIST OF PUBLICATIONS	134

LIST OF TABLES

Table	Page
2.1 Common organisms used in ecotoxicological studies	9
2.2 Important, early studies of medaka since 1846	16
3.1 Comparison of several meristic characters of Java medaka	26
3.2 Occurrence of Java medaka in Peninsular Malaysia at different sampling occasions	27
3.3 Localities of Java medaka and their physicochemical parameters in the west coast of Peninsular Malaysia (mean \pm SE)	30
3.4 Physico-chemical parameters in sites visited in the east coast of Peninsular Malaysia (mean \pm SE)	31
3.5 Physico-chemical parameters within the occurrence of Java medaka during low tide and high tide at Sungai Raya tributaries (mean \pm SD)	33
3.6 Mean of female with eggs and discharged eggs collected from surface sediments	37
3.7 Comparison of several meristic characteristics of Indian medaka	38
3.8 Physico-chemical parameters of locations with co-existence of Java and Indian medaka in Peninsular Malaysia (mean \pm SE)	39
4.1 Culture conditions for Java medaka	47
4.2 Consequences of using different transport containers for Java medaka	54
4.3 Mortality rate of Java medaka during acclimatisation to freshwater and saltwater and during first month of culture period (mean \pm SD; N = 90 per treatment)	55
4.4 Spawning frequency (%) and percentage of fertilised eggs from different parental sex ratio	58
4.5 Mortality, spawning and fertilized eggs of Java medaka cultured in two different salinities (N = 30 per treatment)	60
4.6 Performance of Java medaka eggs incubated in different salinity levels (N = 100 per treatment)	61
4.7 Spawning activity of Java medaka under different photoperiod	63
4.8 Hatching rate of Java medaka eggs from different methods of egg collection (N = 100 per treatment)	66
4.9 Hatching percentage of eggs from different method of surface sterilisation	67
5.1 Percentage of developmental consequences of the exposure of Java medaka eggs to different concentrations of heavy metals (N = 100 per treatment)	75

5.2	Frequency of developmental impairments in pre-hatch embryos of Java medaka exposed to different heavy metals (N = 100 per treatment)	82
5.3	Frequency of morphological impairments in hatched populations of Java medaka after pre-exposure to different concentrations of heavy metals (N = 100 per treatment)	83
5.4	Percentages of developmental impairments in Java medaka embryo exposed to different concentrations of glyphosate (N = 90 per treatment)	90
5.5	Percentage of developmental impairments in post-hatch Java medaka juveniles exposed to different concentrations of glyphosate (N differs in each treatment)	91
5.6	The percentages of survival of juvenile Java medaka in 8 days exposure to glyphosate (N = 30 per treatment)	95
5.7	Micronucleus and nuclear abnormalities (Mean \pm SD) in the red blood cells of Java medaka after exposure to heavy metals	97

LIST OF FIGURES

Figure		Page
2.1	Male (a) and female (b) Java medaka (note the yellow submarginal band of the caudal fin in male)	17
3.1	Aerial view of Java medaka showing silvery eyes (SE) and two silvery stripes (SS)	21
3.2	Study site showing Sungai Raya and its tributaries	23
3.3	The pair of shiny blue eyes of Indian medaka observed in their natural habitat	25
3.4	Localities of Java medaka in Peninsular Malaysia	27
3.5	Abundance of the juvenile and adult Java medaka in relation to salinity changes during rising tide	35
3.6	Abundance of the juvenile and adult Java medaka at different salinity during receding tide	36
3.7	Percentages of different types of medaka fish sampled in different brackish water pools in Kuala Linggi	37
3.8	Abundance of Java and Indian medaka in relation to tidal changes in Sungai Raya tributaries	40
4.1	Sampling activity during low tide (a) and high tide (b)	43
4.2	The re-circulating system set-up	45
4.3	The re-circulating system for Java medaka culture in the Department of Biology, UPM	46
4.4	The <i>Artemia</i> culture system and harvesting procedure	50
4.5	The filaments present in the outer layer of the embryo	52
4.6	Fertilised egg of Java medaka with double-layer coating (a) and unfertilised egg with single-layer coating (b)	59
4.7	Shrunken Java medaka embryo (arrow shows the wide gap of the perivitelline space)	60
4.8	Growth rate of Java medaka fed with different diet	65
4.9	Spawning performance of Java medaka fed with different diet	65
5.1	Examples of delayed development (A) and accelerated development (B) after 11 days of exposure to 1.0 ppm Hg	80
5.2	Abnormally curved tail exposed in embryos exposed to (A) 0.9 mg/l Pb at day 9 and 9B) 0.8 mg/l Pb at day 8.	84
5.3	Some examples of permanently bent tail in juveniles after exposure of the eggs to Pb and Hg	84
5.4	Absence of right pectoral fin, spinal curvature and flexure of the posterior tail region after exposure of the eggs to Pb	85
5.5	Absence of both pectoral fins in juvenile from exposure of eggs to 0.05 Pb	85
5.6	Survival rate of early life stages of Java medaka after exposure to different concentrations of glyphosate	88
5.7	Percentage of hatchability of Java medaka embryos in	88

	different concentrations of glyphosate	
5.8	Heart rate (mean beats/minute) of early life stages of Java medaka in the course of 16-day exposure to different concentrations of glyphosate	89
5.9	Normal embryos (a) and developmental abnormalities in pre-hatch Java medaka embryo exposed to glyphosate: shrunken yolk (b), abnormal body curvature (c) and lateral body curvature (d)	90
5.10	Developmental defects in post-hatch juvenile of Java medaka exposed to glyphosate; abdominal enlargement (a) and cell disruption (b)	92
5.11	Frequency (%) of smooth and shaking movement exhibited by Java medaka juveniles in different concentrations of glyphosate after 48-hour exposure	93
5.12	Frequency (%) of smooth and shaking movement exhibited by Java medaka juveniles in different concentrations of glyphosate after 96-hour exposure	94
5.13	Photomicrographs of erythrocyte cells with normal nucleus (A); with micronuclei (B) and with nuclear abnormalities: blebbed nuclei (C); lobed nuclei (D); notched nuclei (E) and binuclei (F) (Jiraungkoorskul et al., 2007).	96
5.14a	Normal cells with normal nucleus (A) and cells with micronucleus (MN).	99
5.14b	Cells with notched nucleus (NN).	99
5.14c	Cells with lobed nucleus (LN).	99
5.14d	Cells with blebbed nucleus (BN).	100
5.11e	Cells with vacuolated nucleus (VN).	100
5.11f	Cells with ringed nucleus (RN). The contrast of the figure on the right is adjusted to emphasise the gap between the nucleus and the cytoplasm.	100

LIST OF ABBREVIATIONS

Ca^{2+}	Calcium ion
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	Calcium chloride dehydrate
Cd	Cadmium
cm	Centimetre
CO_2	Carbon dioxide
Cu	Copper
DNA	Deoxyribonucleic acid
DO	Dissolved oxygen
E	East
EPA	Eicosapentaenoic acid
F^-	Fluoride ion
Hg	Mercury
hr	Hour
KCl	Potassium chloride
km	Kilometre
L	Litre
LC50	Lethal concentration 50
Lmin^{-1}	Litre per minute
m	Meter
m^2	Meter square
Mg^{2+}	Magnesium ion
mgL^{-1}	Milligram per litre
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	Magnesium sulphate heptahydrate

mm	Millimetre
N	North
N	Number of sample
N	Sample size
NaCl	Sodium chloride
°C	Degree Celsius
POEA	Polyethoxylene amine
ppb	Parts per billion
ppm	Parts per million
ppt	Parts per thousand
SD	Standard Deviation
SE	Standard Error
Temp.	Temperature
UPM	Universiti Putra Malaysia
UV	Ultraviolet
Zn	Zinc
ZnCl ₂	Zinc chloride
µg	Microgram

CHAPTER 1

INTRODUCTION

The release of many hazardous chemicals to the environment is currently identified as a global threat to human and other organisms' health as well as the environment itself. Yet, the world is continuously producing and releasing vast numbers and amounts of chemical compounds to the environment from time to time. The impacts of these chemicals on living organisms and the environment are not well studied much. These chemicals enter the environment, disperse, persist or bio-magnify through the food chain. They may enter the environment intentionally through measured release, for example the application of pesticides in agriculture activities or through unregulated release of industrial by products, household wastes and pharmaceuticals from hospitals. The increasing numbers of chemicals present in the environment due to human activities creates a need for researches on the impacts of these chemicals. In order to protect the environmental health, the possible biological consequences arise from the exposure and the accumulation of the pollutants needs to be strategically assessed. Many monitoring activities and reliable researches were established to evaluate the levels of these chemicals and their biological impacts.

Several organisms such as microorganisms, crustaceans, gastropods, worms and fish have been used either as indicators for these substances or as bio-monitoring agents or as test organisms. These organisms which act as biological indicators have been chosen by many scientists to measure environmental pollution instead of physical or chemical indicators. Physical and chemical indicators can neither show the cumulative effects of different pollutants in the ecosystem nor the duration of the existence of the pollutants which can be evaluated by biological indicators or bio-indicators (Karr, 1981). The application of biological methods can be integrated into other studies and does not cost much and the results can be obtained in a short time. Less equipment is required while a large area can be intensively surveyed and a large amount of information can be obtained for later assessment (Chapman, 1996). The effects of pollutants on the environment are often understood better when biotic factors are correlated with physical and chemical data. The organisms living in aquatic ecosystem reflect the history of the water body whereas chemical analysis is only valid for recent condition. Thus, biological assessment has become more popular and important globally with many methods have been developed.

As an indicator species, a bio-monitoring agent or a test organism should fulfil certain criteria. In order to obtain fast and comparable results, a responsive, readily available and representative organism is needed as test organism. The medaka fish is one of the most important model organisms among teleosts.

The medaka fish belongs to the genus *Oryzias* which is classified under family Adrianichthyidae. This family consists of a diverse group of small fish

that occupy the freshwater, brackish water and saltwater in large areas in Asia. The most established among the medaka fish is the Japanese medaka (*Oryzias latipes*) which has been widely used in experimental vertebrate biology for over a century. It is distributed in the freshwater of Japan, Korea and China (Naruse et al., 1993; Naruse, 1996). It is also a well established experimental organisms in fields such as embryology (Iwamatsu et al., 2005), physiology (Inoue and Takei, 2002) and molecular studies (Yamasaki et al., 2006; Imanishi et al., 2007). One of its relatives, the Java medaka (*O. javanicus*) is distributed in Peninsular Malaysia, Singapore, Indonesia, Thailand and Western Borneo (Iwamatsu et al., 1982; Roberts 1998; Magtoon and Termvidchakorn, 2009). Although Java medaka has only been used in scientific investigation recently, it is gaining reputation as an indicator species among the scientific community.

Java medaka also holds an important position in the estuarine food chain. Occupying the lower trophic levels, they are consumed by larger fish such as mudskipper and other important economic species. Their ecological role contributes to their importance in the aquatic ecosystem where they form a critical connection in the food web to larger fish, seabirds and marine mammals, thus creating a crucial ecosystem value. In addition, the coastal habitat in which Java medaka can be found includes coastline and mangrove areas that are of major economic importance and should be closely monitored and protected. Nevertheless, little is known on the ecology of Java medaka. In establishing an organism as test organism, information on the biology and ecology of the organism is very important in order to understand their responses to any tests or treatments. Sufficient knowledge on the biology and ecology of the organism would make the data from any tests easier to interpret. Thus, the first objective of this research is to investigate some ecological aspects of Java medaka.

To obtain consistent results from the experiments using the fish, a laboratory cultured Java medaka is required. Fish freshly sampled from the environment may not give uniform and comparable results due to their differences in age, size and the salinity of water where they are taken from. Furthermore, testing sensitivity may also differ between sampling locations due to prolonged conditioning to specific environment in the wild. This would make it very difficult for the local scientists in particular to make thorough assessment and to generalise the response shown by the fish in establishing a universal test organism. Therefore, the use of laboratory culture is very important as it could lessen the uncertainties brought about by these differences. Hence, the second objective of this study was to develop inexpensive, easy to maintain and sustainable culture and breeding technique for Java medaka in both saltwater and freshwater in the laboratory.

So far the utilisation of medaka as laboratory animal was carried out in places away from the region. Countries such as Japan, Korea, China, Malaysia and Taiwan. As the use of local or native species is one of the criteria for test organism it is very appropriate to establish our very own species as test organism for the tropical region. Moreover there is no perfect test organism been established yet to represent this type of ecosystem. As most of the

information and knowledge on both fish and its culturing system were developed and adapted to the different environments and seasons available in the said countries, establishing our very own culturing system is necessary. Developing the first medaka laboratory culture system in Malaysia, particularly in the Department of Biology, Universiti Putra Malaysia required systematic planning. Well-known medaka laboratories in Japan were visited to gain information on the laboratory culture for medaka fish. One of the key features that determine the success of the culture system lies in the manipulation of the physico-chemical factor which provides suitable condition for the fish to live healthily and breed at optimum rate. This is made possible through the information gathered while studying the ecology of Java medaka (the first objective of this study) which served as crucial knowledge for the culture establishment.

Java medaka is a small tropical fish closely related to the well-known Japanese medaka. This species has many characteristics similar to the well-established laboratory fish, Japanese medaka. They are hardy, have short life span and life cycle and grow very fast. They can be easily identified in their natural habitat and have a wide geographical range and availability. The high tolerance towards different salinity levels further distinguished the species from other bio-monitoring agents. This seawater and freshwater-adaptable species is suitable to be used to assess the health of aquatic systems in freshwater, estuarine, and marine environments. Nevertheless, there is no detailed information on laboratory culture methods and maintenance for Java medaka. While its foreign counterpart, the Japanese medaka has been extensively researched and utilised in laboratory for toxicology studies, Java medaka has the potential to be developed into a suitable test organism for the tropical region. Therefore, the information on the fish ecology in the various localities found in the wild would greatly aid in the development of the culture method and system for the fish in the laboratory.

There are growing numbers of studies that highlighted Java medaka responses to the various pollutants found in the natural environment (Imai et al., 2005; 2007; Koyama et al., 2006; 2008; Añasco et al., 2008; Cheikyula et al., 2008; Woo et al., 2006; 2009a; 2009b; 2014; Won et al., 2011; 2013; Khodadoust et al., 2013). A test organism for eco-toxicological studies should show excellent responsiveness to chemical contaminant exposure. Responses may either be acute or chronic and should be prominent and consistent throughout the different life stages of the fish population for particular doses or level of pollutants exposed. Therefore, the third objective of this study is to assess the fish capabilities to respond to hazardous chemicals in the environment in order to support the establishment of Java medaka as test organism.

In summary, in the establishment of Java medaka as a test organism for ecotoxicological studies, three objectives were specified:

- i. To investigate several ecological aspects of Java medaka (*Oryzias javanicus* Bleeker).
- ii. To develop a laboratory culture protocol for Java medaka.
- iii. To assess the sensitivity of Java medaka to several environmental pollutants.

It is hoped that this study does not only provide fundamental information on the biology and ecology of the Java medaka in its native ranges, but it is also the first to highlight the establishment of laboratory culture system for Java medaka in its native country. This study utilised the knowledge on the biology and ecology of the fish to establish the culture system. Products from the laboratory culture were then utilised in exposure tests using several pollutants common to the local environment. Different life stages of the fish, from embryonic to adult were employed in the tests. It is a great hope that the findings will pave the way towards establishing our own medaka as sentinel test fish for tropical countries around the world.

BIBLIOGRAPHY

- Abbas Alkarkhi, F.M., Ismail, N., Mat Easa, A. 2008. Assessment of arsenic and heavy metal contents in cockles (*Anadara granosa*) using multivariate statistical techniques. *Journal of Hazardous Materials* 150(3): 783-789.
- Aguigwo, J.N. 2002. The toxic effect of Cymbush pesticide on growth and survival of African catfish, *Clarias gariepinus* (Burchell). *Journal of Aquatic Science* 17(2): 81-84.
- Ahmed, M.K., Parvin, E., Arif, A., Akter, M.S., Khan, M.S., Islam, M.M. 2010. Measurement of genotoxic potential of cadmium in different tissues of fresh water climbing perch *Anabas testudineus* (Bloch), using the comet assay. *Environmental Toxicology and Pharmacology* 30(1): 80-84.
- Åkerman, G., Balk, L. 1995. A reliable and improved methodology to expose fish in the early embryonic stage. *Marine Environmental Research* 39: 155-158.
- Akhir, M.F., Sinha, P.C., Hussain, M.L. 2011. Seasonal variation of South China Sea physical characteristics off the east coast of Peninsular Malaysia from 2002 – 2010 datasets. *International Journal of Environmental Sciences* 2(2): 569-575.
- Añasco, N.C., Koyama, J., Imai, S., Nakamura, N. 2008. Toxicity of residual chlorines from hypochlorite-treated seawater to marine amphipod *Hyale barbicornis* and estuarine fish *Oryzias javanicus*. *Water Air and Soil Pollution* 195:129–136.
- Anonymous, 2010. *World Water Day*. Retrieved on 25th November 2010 from www.worldwaterday2010info.
- Atta, K.I., Abdel-Karim, A.E., Elsheikh, E.H. 2012. Ultrastructural study of the effect of heavy metals on the regenerating tail fin of the teleost fish, *Oreochromis niloticus*. *The Journal of Basic and Applied Zoology* 65(4): 232-239.
- Avault, J.W. 1996. *Fundamentals of Aquaculture*. Baton Rouge, LA: AVA Publishing Company.
- Ayllon, F., Garcia-Vazquez, E. 2000. Induction of micronuclei and other nuclear abnormalities in European minnow *Phoxinus phoxinus* and mollie *Poecilia latipinna*: an assessment of the fish micronucleus test. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 467(2): 177-186.

- Ayoola, S.O. 2008. Histopathological effects of glyphosate on juvenile African catfish (*Clarias gariepinus*). *American-Eurasian Journal of Agriculture and Environmental Science* 4(3): 362-367.
- Azrina, M.Z., Yap, C.K., Ismail, A.R., Ismail, A., Tan, S.G. 2006. Anthropogenic impacts on the distribution and biodiversity of benthic macro-invertebrates and water quality of the Langat River, Peninsular Malaysia. *Ecotoxicology and Environmental Safety* 64: 337-347.
- Bagdonas, E., Vosyliene, M.Z. 2006. A study of toxicity and genotoxicity of copper, zinc and their mixture to rainbow trout (*Oncorhynchus mykiss*). *Biologija* 1: 8-13.
- Banu, B.S., Devi, K.D., Mahboob, M., Jamil, K. 2001. *In vivo* genotoxic effect of zinc sulfate in mouse peripheral blood leukocytes using Comet assay. *Drug and Chemical Toxicology* 24(1): 63-73.
- Baras, E., Jobling, M. 2002. Dynamics of intracohort cannibalism in cultured fish. *Aquaculture Research* 33(7): 461-479.
- Barjhoux, I., Baudrimont, M., Morin, B., Landi, L., Gonzalez, P., Cachot, J. 2012. Effects of copper and cadmium spiked-sediments on embryonic development of Japanese medaka (*Oryzias latipes*). *Ecotoxicology and Environmental Safety* 79: 272-282.
- Barnhoorn, I.E.J., Bornman, M.S., Pieterse, G.M., van Vuren, J.H.J. 2004. Histological evidence of intersex in feral sharptooth catfish (*Clarias gariepinus*) from an estrogen-polluted water source in Gauteng, South Africa. *Environmental Toxicology* 19(6): 603-608.
- Belanger, S.E., Rawlings, J.M., Carr, G.J. 2013. The use of fish embryo toxicity tests for the prediction of acute fish toxicity to chemicals. *Environmental Toxicology and Chemistry* 32(8): 1768-1783.
- Berglund, M., Åkesson, A., Bjellerup, P., Vahter, M. 2000. Metal-bone interactions. *Toxicology Letters* 112-113: 219-225.
- Birge, W.J., Black, J.A. 1979. Effects of Copper on Embryonic and Juvenile Stages of Aquatic Animals In *Copper in the Environment*. Part II. Health Effects, ed. J.O. Nriagu, pp 373-399. New York: John Wiley and Sons.
- Birge, W.J., Black, J.A., Westerman, A.G., Ramey, B.A., 1983. Fish and amphibian embryos: a model system for evaluating teratogenicity. *Fundamental and Applied Toxicology* 3: 237-242.
- Blaber, S.J.M. 1997. *Fish and Fisheries of Tropical Estuaries*. London, United Kingdom: Chapman and Hall.

- Blaber, S.J.M., Blaber, T.G., 1980. Factors affecting the distribution of juvenile estuarine and inshore fish. *Journal of Fish Biology* 17: 143-162.
- Blomhoff, R., Blomhoff, H.K. 2006. Overview of retinoid metabolism and function. *Developmental Neurobiology* 66(7): 606-630.
- Boeuf, G., Payan, P. 2001. How should salinity influence fish growth? *Comprehensive Biochemistry and Physiology C* 130: 411-423.
- Bols, N.C., Brubacher, J.L., Ganassin, R.C., Lee, L.E.J. 2001. Ecotoxicology and innate immunity in fish. *Developmental and Comparative Immunology* 25: 853-873.
- Bortagaray, V., Aramburu, R.C., Barios, L., Ojeda, P., del Puerto, G., Rodríguez-Ithurralde, D. 2010. Embryotoxicity and teratogenesis in zebrafish embryos exposed *in vitro* to glyphosate-type herbicides. *Journal of Developmental Toxicity* 1 (1): 2-6.
- Bozek, M. A., and F. J. Rahel. 1991. Comparison of streamside visual counts to electrofishing estimates of Colorado River cutthroat trout fry and adults. *North American Journal of Fisheries Management* 11:38–42.
- Brand, M., Granato, M., Nüsslein-Volhard, C. (2002). Keeping and raising zebrafish In *Zebrafish: A Practical Approach*, ed. C. Nüsslein-Volhard and R. Dahm R, pp 7-37. Oxford, UK: Oxford University Press.
- Bresch, H., Markert, M., Munk, R., Spieser, O.H. 1986. Long-term fish toxicity test using the zebrafish: Effect of group formation and temporary separation by sex on spawning. *Bulletin of Environmental Contamination and Toxicology* 37(4): 606-614.
- Buikema, A.L., Niederlehner, Jr, B.R., Cairns Jr.J. 1982. Biological monitoring part IV - Toxicity testing. *Water Research* 16(3): 239-262.
- Burlibaşa, L., Gavrilă, L. 2011. Amphibians as model organisms for study environmental genotoxicity. *Applied Ecology and Environmental Research* 9(1): 1-15.
- Cairns Jr. J., Niederlehner, B. R. 1993. Ecological function and resilience: neglected criteria for environmental impact assessment and ecological risk analysis. *Environment Professional* 15: 116-124.
- Campana, M.A., Panzeri, A.M., Moreno, V.J., Dulout, F. 1995. Genotoxic evaluation of the pyrethroid lambda-cyhalothrin using the micronucleus test in erythrocytes of the fish *Cheirodon interruptus*. *Mutation Research* 438: 155-161.

- Campana, O., Sarasquete, C., Blasco, J. 2003. Effect of lead on ALA-D activity, metallothionein levels, and lipid peroxidation in blood, kidney, and liver of the toadfish *Halobatrachus didactylus*. *Ecotoxicology and Environmental Safety* 55(1): 116-125.
- Cardona, L. 2000. Effects of salinity on the habitat selection and growth performance of Mediterranean flathead grey mullet, *Mugil cephalus* (Osteichthyes, Mugilidae). *Estuarine, Coastal and Shelf Science* 50: 727-737.
- Carlson, G., Silverstein, J. 1997. Effect of ozonation on sorption of natural organic matter by biofilm. *Water Research* 31(10): 2467-2478.
- Carmichael, G.J., Tomasso, J.R. and Schwedler, T.E. 2002. Fish transportation. In *Fish Hatchery Management, second edition*, ed. G.A. Wedemeyer, pp. 641-660. Bethesda: American Fisheries Society,
- Çavaş, T., Ergene-Gözükara. 2005. Induction of micronuclei and nuclear abnormalities in *Oreochromis niloticus* following exposure to petroleum refinery and chromium processing plant effluents. *Aquatic Toxicology* 74(3): 264-271.
- Çavaş, T., Garanko, N.N., Arkhipchuk, V.V. 2005. Induction of micronuclei and binuclei in blood, gill and liver cells of fishes subchronically exposed to cadmium chloride and copper sulphate. *Food and Chemical Toxicology* 43: 569-574.
- Chan, K.M. 1995. Metallothionein: Potential biomarker monitoring heavy metal pollution in fish around Hong Kong. *Marine Pollution Bulletin* 31(4-12): 411-415.
- Chanda, M., Paul, M., Maity, J., Dash, G., Gupta, S.S. 2011. The use of antibiotics and disinfectants in ornamental fish farms of West Bengal, India. *Journal of Natural Science, Biology and Medicine* 2(2): 139-140.
- Chapman, D. 1996. *Water Quality Assessments: A Guide to Use of Biota, Sediments and Water in Environmental Monitoring. Second Edition*. Cambridge: University Press.
- Charoy, C., Janssen, C.R. 1999. The swimming behaviour of *Brachionus calyciflorus* (rotifer) under toxic stress: II. Comparative sensitivity of various behavioural criteria. *Chemosphere* 38 (14): 3247-3280.

- Cheikyula, J.O., Koyama, J., Uno, S. 2008. Comparative study of bioconcentration and EROD activity induction in the Japanese flounder, red sea bream, and Java medaka exposed to polycyclic aromatic hydrocarbons. *Environmental Toxicology* 23 (3): 354-362.
- Chen, J-C., Liu, P-C., Lin, Y-T. 1991. Reduction of ammonia and nitrite in shrimp larviculture in a recirculation system. *Asian Fisheries Science* 4: 211-218.
- Chen, T-H., Lina, C-C., Meng, P.J. 2014. Zinc oxide nanoparticles alter hatching and larval locomotor activity in zebrafish (*Danio rerio*). *Journal of Hazardous Materials* 277: 134-140.
- Chovanec, A., Hofer, R., Schiemer, F. 2003. Fish as bioindicators. In *Bioindicators and Biomonitoring*, ed. B.A. Markert, A.M. Breure, H.G. Zechmeister, pp 639-676. Oxford: Elsevier.
- Clark, M.S., Elgar, G. 2000. *Fugu rubripes*: a fish model genome. In *Comparative genomics*, ed. M.S. Clark, pp. 71-96. Boston: Kluwer Academic Publishers.
- Cohen, T., Que Hee, S.S., Ambrose, R.F. 2001. Trace metals in fish and invertebrates of three California coastal wetlands. *Marine Pollution Bulletin* 42(3): 224-232.
- Craik, J.C.A., Harvey, S.M. 1988. A preliminary account of metal levels in eggs of farmed and wild Atlantic salmon and their relation to egg viability. *Aquaculture* 73: 309-321.
- Cui, Y., Freedman, J.H. 2009. Cadmium induces retinoic acid signaling by regulating retinoic acid metabolic gene expression. *The Journal of Biological Chemistry* 284 (37): 24925-24932.
- Dave, G., Xiu, R. 1991. Toxicity of mercury, copper, nickel, lead, and cobalt to embryos and larvae of zebrafish, *Brachydanio rerio*. *Archives of Environmental Contamination and Toxicology* 121: 126-134.
- DeTolla, L.J., Srinivas, S., Whitaker, B.R., Andrews, C., Hecker, B., Kane, A.S., Reimschuessel, R. 1995. Guidelines for the care and use of fish in research. *Institute for Laboratory Animal Research Journal* 37(4): 159-173.
- Du, Z-Y., Liu, Y-J., Tian, L-X., He, J-G., Cao, J-M., Liang, G-Y. 2006. The influence of feeding rate on growth, feed efficiency and body composition of juvenile grass carp (*Ctenopharyngodon idella*). *Aquaculture International* 14(3): 247-257.
- Dufour, E.K., Kumaravel, T., Nohynek, G.J., Kirkland, D., Toutain, H. 2006. Clastogenicity, photo-clastogenicity or pseudo-photo-

clastogenicity: Genotoxic effects of zinc oxide in the dark, in pre-irradiated or simultaneously irradiated Chinese hamster ovary cells. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 607(2): 215–224

Dumpert, K., Zietz, E. 1984. Platanna (*Xenopus levis*): A test organism for determining the embryotoxic effects of environmental chemicals. *Ecotoxicology and Environmental Safety* 8: 55-74.

Dutta, H.M., Arends, D.A. 2003. Effects of endosulfan on brain acetylcholinesterase activity in juvenile bluegill sunfish. *Environmental Research* 91(3):157–162

Eaton, J.G., Mc Kim, J.M., Holcombe, J.W., 1978. Metal toxicity to embryos and larvae of seven freshwater fish species - cadmium. *Bulletin of Environmental Contamination and Toxicology* 19: 95-103.

Edwards, J.W., Edyvane, K.S., Boxall, V.A., Hamann, M., Soole, K.L. 2001. Metal levels in seston and marine fish flesh near industrial and metropolitan centres in South Australia. *Marine Pollution Bulletin* 42 (5): 389-396.

Ellenberger, S.A., Baumann, P.C., May, T.W. 1994. Evaluation of effects caused by high copper concentrations in Torch Lake, Michigan, on reproduction of yellow perch. *Journal of Great Lakes Research* 20(3): 531-536.

Ellsaesser, C.F., Clem, L.W. 1986. Haematological and immunological changes in channel catfish stressed by handling and transport. *Journal of Fish Biology* 28: 511-521.

Emlen S.T, Oring L.W. 1977. Ecology, sexual selection, and the evolution of mating systems. *Science* 197: 215–223.

EFSA Scientific Panel on Animal Health and Welfare, 2005. Opinion on the 'Aspects of the biology and welfare of animals used for experimental and other scientific purposes'. *The EFSA Journal* 292: 1-46.

Espina, N.G., Weiss, P., 1995. DNA-repair in fish from polluted estuaries. *Marine Environmental Research* 39(1-4): 309-312.

FAO (Food and Agriculture Organization), 2003. *Review of World Water Resources by Country*. FAO: Rome, Italy.

FAO (Food and Agriculture Organization), 2012. Water Report. Retrieved on 25th November 2010 from http://www.fao.org/nr/water/aquastat/countries_regions/MYS/index.stm

- Fichez, R., Adjerdoud, M., Bozec, Y-M., Breau, L., Chancerelle, Y., Chevillon, C., Douillet, P., Fernandez, J-M., Frouin, P., Kulbicki, M., Moreton, B., Ouillon, S., Payri, C., Perez, T., Sasal, P., Thébault, J. 2005. A review of selected indicators of particle, nutrient and metal inputs in coral reef lagoon system. *Aquatic Living Resource* 18: 125-147.
- Fisher K.C. 1963. The formation and properties of the external membrane of the trout egg. *Royal Society of Canada* 1: 323–332.
- Fränzle, O. 2003. Bioindicators and environmental stress assessment. In *Bioindicators and Biomonitoring*, ed. B.A. Markert, A.M. Breure, H.G. Zechmeister, pp 41-84. Oxford: Elsevier.
- Frenzilli, G., Falleni, A., Scarcelli, V., Del Barga, I., Pellegrini, S., Savarino, G., Mariotti, V., Benedetti, M., Fattorini, D., Regoli, F., Nigro, M. 2008. Cellular responses in the cyprinid *Leuciscus cephalus* from a contaminated freshwater ecosystem. *Aquatic Toxicology* 89(3): 188-196.
- Gering, E., Atkinson, C.T. 2004. A rapid method for counting nucleated erythrocytes on stained blood smears by digital image analysis. *Journal of Parasitology* 90(4): 879-881.
- Gianneccchini, L.G., Massago, H., Fenandes, J.B.K. 2012. Effects of photoperiod on reproduction of Siamese fighting fish *Betta splendens*. *Revista Brasileira de Zootecnia* 41(4): 821-826.
- Gibson R.N., Yoshiyama R.M. 1999. Intertidal fish communities. In *Intertidal Fish: Life in two worlds*, ed. .M.H. Horn, K.L.M. Martin, M.A. Chotkowski, pp. 97–125. San Diego, CA: Academic Press.
- Giesy, J.P., Dobson, S., Solomon, K., 2000. Ecotoxicological risk assessment for Roundup® herbicide. *Reviews of Environmental Contamination and Toxicology* 167: 35 -120.
- Gluszczak, L., Miron, D.S., Moraes, B.S., Simões, R.R., Schetinger, M.R.C., Morsch, V.M., Loro, V.L. 2007. Acute effects of glyphosate herbicide on metabolic and enzymatic parameters of silver catfish (*Rhamdia quelen*). *Comparative Biochemistry and Physiology Part C* 146: 519-524.
- Gomma, M.N.E., Abou-Arab, A.A.K., Badawy, A., Khayria, N. 1995. Distribution patterns of some heavy metals in Egyptian fish organs. *Food Chemistry* 53: 385-389.
- Grant, J.W.A., Gaboury, C.L., Levitt, H.L. 2000. Competitor-to-resource ratio, a general formulation of operational sex ratio, as a predictor of competitive aggression in Japanese medaka (Pisces: Oryziidae). *Behavioral Ecology* 11: 670–675.

- Grechay, P.A., Targett, T.E. 1996. Effects of turbidity, light level and prey concentration on feeding of juvenile weakfish *Cynoscion regalis*. *Marine Ecology Progress Series* 131: 11-16.
- Guecheva, T., Henriques, J.A.P., Erdtmann, B. 2001. Genotoxic effects of copper sulphate in freshwater planarian *in vivo*, studied with the single-cell gel test (comet assay). *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 497(1-2): 19-27.
- Gui, Y.X., Fan, X.N., Wang, H.M., Wang, G., Chen, S.D. 2012. Glyphosate induced cell death through apoptotic and autophagic mechanisms. *Neurotoxicology and Teratology* 34(3): 344-349.
- Guillard, J. 1998. Daily migration cycles of fish populations in a tropical estuary (Sine-Saloum, Senegal) using a horizontal-directed split-beam transducer and multibeam sonar. *Fisheries Research* 35: 23-31.
- Hagenmaier, H.E. 1974. The hatching process in fish embryos - IV. The enzymological properties of a highly purified enzyme (chorionase) from the hatching fluid of the rainbow trout, *Salmo gairdneri* Rich. *Comprehensive Biochemistry and Physiology* 49(B): 313-324.
- Hajeb, P., Jinap, S., Ismail, A., Fatimah, A.B., Jamilah, B., Abdul Rahim, M. 2009. Assessment of mercury level in commonly consumed marine fishes in Malaysia. *Food Control* 20(1): 79-84.
- Håkanson, L. 1999. *Water pollution - methods and criteria to rank, model and remediate chemical threats to aquatic ecosystems*. Leiden: Backhuys Publishers.
- Hall, J.A., Frid, C.L.J., Gill, M.E. 1997. The response of estuarine fish and benthos to an increasing discharge of sewage effluent. *Marine Pollution Bulletin* 34(7): 527-535.
- Hammer, M. J. 1986. *Water and wastewater technology, Second edition*. New York. USA: John Wiley Inc.
- Hanley, J.S., Shashar, N., Smolowitz, R., Bullis, R.A., Mebane, W.N., Gabr, H.R., Hanlon, R.T. 1998. Modified laboratory culture techniques for the European cuttlefish *Sepia officinalis*. *Biology Bulletin* 195: 223-225.
- Hawkins, W.E., Walker, W.W., Fournie, J.W., Manning, C.S., Krol R.M. 2003. Use of Japanese medaka (*Oryzias latipes*) and guppy (*Poecilia reticulata*) in carcinogenesis testing under national toxicology program protocols. *Toxicologic Pathology* 31(1): 88-91.

- Hazel C.R, Meith S.J. (1970) Bioassay of king salmon eggs and sac fry in copper solution. *California Fish Game* 56: 121-124.
- Heisler, N. (1993). Acid-base regulation in response to changes of the environment: characteristics and capacity. In *Fish Ecophysiology*, ed. J.C. Rankin, F.B. Jensen, pp. 207-230. New York: Chapman and Hall.
- Heslinga, A. 1976. Effect of copper on the coral reef echinoid *Echinometra mathaei*. *Marine Biology* 3(2): 155-160.
- Heddle, A., Cimino, M.C., Hayashi, M., Romagna, F., Shelby, M.D., Tucker, J.D., Vanparys, Ph., MacGregor, J.T. 1991. Micronuclei as an index of cytogenetic damage: Past, present, and future. *Environmental and Molecular Mutagenesis* 18(4): 277-291.
- Hii, Y. S., Law, A.t., Shazili, N.A.M., Abdul Rashid, M.K., Mohd Lokman, H. 2006. The Straits of Malacca: Hydrological Parameters, Biochemical Oxygen Demand and Total Suspended Solid. *Journal of Sustainability Science and Management* 1(1): 1-14.
- Hinton, D.E., Kullman, S.W., Hardman, R.C., Volz, D.C., Chen, P.J., Carney, M., Bencic, D.C. 2005. Resolving Mechanisms of toxicity while pursuing ecotoxicological relevance? *Marine Pollution Bulletin* 51: 635-648.
- Holm, G., Lundström, J., Andersson, T., Norrgren, L. 1994. Influences of halogenated organic substances on ovarian development and hepatic EROD activity in the three-spined stickleback, *Gasterosteus aculeatus* and rainbow trout, *Oncorhynchus mykiss*. *Aquatic Toxicology* 29 (3-4): 241-256.
- Hodson, P.V., Blunt, B.R., Spry, D.J. 1978. Chronic toxicity of water-borne and dietary lead to rainbow trout (*Salmo gairdneri*) in Lake Ontario water. *Water Research* 12: 869-878.
- Huang, W., Cao, L., Shan, X., Lin, L., Dou, S. 2011. Toxicity testing of waterborne mercury with red sea bream (*Pagrus major*) embryos and larvae. *Bulletin of Environmental Contamination and Toxicology* 86(4): 398-405.
- Huffman D.E., Gennaccaro, A., Rose J.B., Dussert, B.W. 2002. Low- and medium-pressure UV inactivation of microsporidia *Encephalitozoon intestinalis*. *Water Research* 36: 3161-3164.
- Ijiri, K. 1995. Medaka fish had the honor to perform the first successful vertebrate mating in space. *The Fish Biology Journal MEDAKA* 7: 1-10.

- Ikram, M.M., Ismail, A., Yap, C.K., Nor Azwady, A.A. 2010. Levels of heavy metals (Zn, Cu, Cd and Pb) in mudskippers (*Periophthalmodon schlosseri*) and sediments collected from intertidal areas at Morib and Remis, *Toxicology and Environmental Chemistry* 92: 1471-1486.
- Imai, S., Koyama, J., Fujii, K. 2005. Effects of 17 β -estradiol on the reproduction of Java medaka (*Oryzias javanicus*), a new test fish species. *Marine Pollution Bulletin* 51(8-12): 708-714.
- Imai, S., Koyama, J., Fujii, K. 2007. Effects of estrone on full life cycle of Java medaka (*Oryzias javanicus*), a new marine test fish. *Environmental Toxicology and Chemistry* 26(4): 726-731.
- Imanishi, Y., Hisatomi, O., Yamamoto, S., Satoh, T., Kotaka, S., Kobayashi, Y., Tokunaga, F. 2007. A third photoreceptor-specific GRK found in the retina of *Oryzias latipes* (Japanese killifish). *Zoological Science* 24(1): 87-93.
- Inoue, K., Takei, Y. 2002. Diverse adaptability in *Oryzias* species to high environmental salinity. *Zoological Science* 19(7): 727-734.
- Inoue, K., Takei, Y. 2003. Asian medaka fishes offer new models for studying mechanisms of seawater adaptation. *Comparative Biochemistry and Physiology Part B* 136: 635-645.
- Irving, E.C., Baird, D.J., Culp, J.M. 2009. Cadmium toxicity and uptake by mats of the freshwater diatom: *Navicula pelliculosa* (Bréb) Hilse. *Archives of Environmental Contamination and Toxicology* 57: 524-530.
- Ishikawa, Y. 2000. Medaka fish as a model system for vertebrate developmental genetics. *BioEssays* 22: 487-495.
- Ismail, A., Badri, M.A., Ramlan, M.N. 1991. Heavy metal contamination in fiddler crabs (*Uca annulipes*) and hermit crabs (*Clibanarius* sp.) in a coastal area of northern Peninsular Malaysia. *Environmental Technology* 12: 923-926.
- Ismail, A., Jusoh, N.R., Ghani, I.A. 1995. Trace metal concentrations in marine prawns off the Malaysian coast. *Marine Pollution Bulletin* 31(1-3): 108-110.
- Ismail, A., Ramli, R. 1997. Trace metals in sediments and molluscs from an estuary receiving pig farms effluent. *Environmental Technology* 18(5): 509-515.
- Ismail, A., Safahieh, A. 2005. Copper and Zinc in intertidal surface sediment and *Telescopium telescopium* from Lukut River, Malaysia. *Coastal Marine Science* 29(2): 111-115.

- Iwamatsu, T. 1998. *The Integrated Book for the Biology of the Medaka*. Okayama, Japan: Daigaku Kyouiku Publishing Co.
- Iwamatsu, T., 1978. Studies on oocyte maturation of the medaka, *Oryzias latipes*. VI. Relationship between the circadian cycle of oocyte maturation and activity of the pituitary gland. *Journal of Experimental Zoology* 206: 355–364.
- Iwamatsu, T., Imaki, A., Kawamoto, A., Inden, A. 1982. On *Oryzias javanicus* collected at Jakarta, Singapore and West Kalimantan. *Annotationes Zoologici Japonenses* 55: 190-198.
- Iwamatsu, T., Kobayashi, H., Sato, M. 2005. *In vivo* fertilization and development of medaka eggs initiated by artificial insemination. *Zoological Science* 22(1): 119-123.
- Iwamatsu, T., Hirata, K. 1984. Normal course of development of the Java medaka, *Oryzias javanicus*. *Bulletin Aichi University Education (Natural Science)* 33: 87-109.
- Iyagba, M.A., Adoki, A., Sokari, T.G. 2008. Testing biological methods to treat rubber effluent. *African Journal of Agriculture Research* 3: 448-454.
- Jafari, M., Kamarudin, M.S., Saad, C.R., Arshad, A., Oryan, S., Gailani, M.H.T. 2011. Effects of different diet on growth, survival and body composition of *Rutilus frisii kutum* larvae. *Journal of Fisheries and Aquatic Science* 6 (6): 662-668.
- Jennett J.C., Effier, S.W., Wixson, B.G. 1980. Mobilization and toxicological aspects of sedimentary contaminants. In. *Contaminants and sediments Volume I: Fate and transport case studies, modeling, toxicity*, ed. R.A. Baker, pp. 429–444. Ann Arbor, USA: Ann Arbor Science.
- Jeon, S.Y., Woo, S., Won, H., Kim, B., Yum, S. 2010. Differential gene expression profiling in B [k] F-exposed marine medaka, *Oryzias javanicus* by subtractive hybridization. *Toxicology and Environmental Health Science* 2(4): 231-237.
- Jezierska, B., Lugowska, K., Witeska, M. 2009. The effects of heavy metals on embryonic development of fish (a review). *Fish Physiology and Biochemistry* 35: 625-640.
- Jezierska, B., Lugowska, K., Witeska, M., Sarnowski, P. 2000. Malformations of newly hatched common carp larvae. *Electronic Journal of Polish Agricultural University* 3: 2.
- Jezierska, B., Witeska, M. 2001. *Metal Toxicity to Fish*. Poslaskiej: Wydawnictwo Akademii.

- Jindal, R., Sharma, C. 2011. Biomonitoring of pollution in river Sutlej. *International Journal of Environmental Science* 2 (2): 863-872
- Jiraungkoorskul, W., Sahaphong, S., Kosai, P., Kim, M. 2007. Micronucleus test: the effect of ascorbic acid on cadmium exposure in fish (*Punctius altus*). *Research Journal of Environmental Toxicology* 1: 27-36.
- Johnson, E., Carew, K.A., Sloman. 2007. The effects of copper on the morphological and functional development of zebrafish embryos. *Aquatic Toxicology* 84(4): 431-438.
- Jothy, A.A., Rauck, G., Latiff, M.S.S.A., Ong, K.S., Leong, P.C., Carvalho, J.L. 1975. *Demersal fish resources in Malaysian waters. Second trawl survey of the coastal waters off the east coast of Peninsular Malaysia*. Fisheries Bulletin No. 4: Kuala Lumpur.
- Kahlown, M. A., Majeed, A. 2003. *Water-resources situation in Pakistan: challenges and future strategies. Water resources in the south: present scenario and future prospects Islamabad, Pakistan*. Commission on Science and Technology for Sustainable Development in the South (COMSATS): Islamabad.
- Kang, C.K., Tsai, S.C., Lee, T.H., Hwang, P.P. 2008. Differential expression of branchial Na⁺/K⁺-ATPase of two medaka species, *Oryzias latipes* and *Oryzias dancena*, with different salinity tolerances acclimated to fresh water, brackish water and seawater. *Comprehensive Biochemistry and Physiology A* 151(4): 566-75.
- Kakuno, A., Fujii, K., Koyama, J. 2001. Histological studies on the gonadal development in Java medaka (*Oryzias javanicus*). *Bulletin of Fish Research Agency* 1: 35-37.
- Kamarudin, M.S., Otoi, S., Saad, C.R. 2011. Changes in growth, survival and digestive enzyme activities of Asian redtail catfish, *Mystus nemurus*, larvae fed on different diets. *African Journal of Biotechnology* 10 (21): 4484-4493.
- Kapur, K., Yadav, N.A., 1982. The effects of certain heavy metal salts on the development of eggs in common carp, *Cyprinus carpio* var. Communis. *Acta Hydrochimica et Hydrobiologica* 10: 517-522.
- Karr, J.R. 1981. Assessment of biotic integrity using fish community. *Fisheries* 6(6): 21-27.
- Katselis, G., Koukou, K., Dimitriou, E., Koutsikopoulos, C. 2007. Short-term seaward fish migration in the Messolonghi-Etoliko lagoons (Western Greek coast) in relation to climatic variables and lunar cycle. *Estuarine, Coastal and Shelf Science* 73(3-4): 571-582.

- Kaur, K., Bajwa, K. 1987. Effect of zinc and cadmium on early life stages of common carp, *Cyprinus carpio* (Linn). *Annals of Applied Biology* 3: 28-33.
- Kawasaki, T., Saito, K., Deguchi, T., Fujimori, K., Tadokoro, M., Yuba, S., Kawarabayashi, Y., 2008. Pharmacological characterisation of isoproterenol-treated medaka fish. *Pharmacological Research* 58 (5-6): 348-355.
- Khodadoust, D., Ismail, A. Zulkifli, S.Z., Tayefeh, F.H. 2013. Short time effect of cadmium on juveniles and adults of Java medaka (*Oryzias javanicus*) fish as a bioindicator for ecotoxicological studies. *Life Science Journal* 10(1): 1857-1861.
- Kinoshita, M., Murata, K., Naruse, K., Tanaka, M., 2010. *Medaka: Biology, Management and Experimental Protocols*. Ames: Wiley-Blackwell.
- Kinoshita, M. 2012. Transgenic Medaka, Small Fresh Water Teleost (*Oryzias latipes*), is Now Available for Environmental Science. In *Interdisciplinary Studies on Environmental Chemistry - Environmental Pollution and Ecotoxicology*, ed. M. Kawaguchi, K. Misaki, H. Sato, T. Yokokawa, T. Itai, T.M. Nguyen, J. Ono, S. Tanabe, S. pp. 49-54. Tokyo: Terrapub.
- Kiso, K., Mahyam, M. I., 2003. Distribution and feeding habits of juvenile and young John's snapper *Lutjanus johnii* in the Matang Mangrove estuary, west coast of Peninsula Malaysia. *Fisheries Science* 69: 563-568.
- Kolkovski, S., Curnow, J., King, J. 2004. Intensive rearing system for fish larvae research II: *Artemia* hatching and enriching system. *Aquaculture Engineering* 31: 309-317.
- Kong, R.Y.C., Giesy, J.P., Wu, R.S.S., Chen, E.X.H., Chiang, M.W.L., Lim, P.L., Yuen, B.B.H., Yip, B.W.P., Mok, H.O. L., Au, D.W.T. 2007. Development of marine fish model for studying in vivo molecular responses in ecotoxicology. *Aquatic Toxicology* 86(2): 131-141.
- Kowalska-Wochna, E., Moniuszko-Jakoniuk, J., Kulikowska, E., & Miniuk, K. 1988. The effect of orally applied aqueous solutions of lead and zinc on chromosome aberrations and induction of sister chromatid exchanges in the rat (*Rattus* sp.). *Genetica Polonica* 29(2): 181-89.
- Koyama, J., Imai, S., Fujii, K., Kawai, S., Yap, C.K., Ismai, A. 2006. Pollution by estrogens in river and estuarine waters around Kuala Lumpur, Malaysia, and their effects on the estuarine Java medaka, *Oryzias javanicus*. *Japanese Journal of Environmental Toxicology* 9(2): 141-147.

- Koyama, J., Kawamata, M., Imai, S., Fukunaga, M., Uno, S., Kakuno, A. 2008. Java medaka: a proposed new marine test fish for ecotoxicology. *Environmental Toxicology* 23(4): 487-491.
- Krumme, U. 2004. Patterns in tidal migration of fish in a Brazilian mangrove channel as revealed by a split-beam echosounder. *Fisheries Research* 70(1): 1-15.
- Krumme, U., Saint-Paul, U. 2003. Observations of fish migration in a macrotidal mangrove channel in Northern Brazil using a 200-kHz split-beam sonar, *Aquatic Living Resources* 16: 175-184.
- Küçükoğlu, M., Binokay, U.S., Boğa Pekmezekmek, A. 2013. The effects of zinc chloride during early embryonic development in zebrafish (*Brachydanio rerio*). *Turkish Journal of Biology* 37: 158-164.
- Kucuksezgin, F., Altay, O., Uluthurhan, E., Kontas, A. 2001. Trace metal and organochlorine levels in red mullet (*Mullus barbatus*) from the Eastern Aegean, Turkey. *Water Research* 35(9): 2327-2332.
- Kume, K., Kubo, Y. 2005. Bibliography on an endangered fish, medaka, *Oryzias latipes* (Temminck & Schlegel). *Bulletin of Faculty of Agriculture, Kinki University* 38: 157-228.
- Ku Kassim, K.Y., Ahmad, A.M.I. 2007. *Keadaan laut perairan Semenanjung Malaysia untuk panduan nelayan*. Kuala Lumpur: Jabatan Perikanan SEAFDEC/MFRDMD.
- Kvarnemo, C., Ahnesjö, I. 1996. The dynamics of operational sex ratios and competition for mates. *Trends in Ecology and Evolution* 11: 404-408.
- Kwak, I-S., Chon, T-S., Kang, H-M., Chung, N-I., Kim, J.S., Kon, S-C., Lee, S-K., Kim, Y-S. 2002. Pattern recognition of the movement tracks of medaka (*Oryzias latipes*) in response to sub-lethal treatments of an insecticide by using artificial neural networks. *Environmental Pollution* 120(3): 671-681.
- Lakra W.S., Nagpure, N.S., 2009. Genotoxicological studies in fish: a review. *Indian Journal of Animal Science* 79: 93-98.
- Lammer, E., Carr, G.J., Wendler, K., Rawlings, J.M., Belanger, S.E., Braunbeck, T.H., 2009. Is the fish embryo toxicity test (FET) with the zebrafish (*Danio rerio*) a potential alternative for the fish acute toxicity test? *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 149(2): 196-209.

- Langiano, V.C., Martinez, C.B.R. 2008. Toxicity and effects of a glyphosate-based herbicide on the Neotropical fish *Prochilodus lineatus*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 147(2): 222-231.
- Latiff, M.A., Bodaly, R.A., Johnston, T.A., Fudge, R.J.P. 2001. Effects of environmental and maternally derived methylmercury on the embryonic and larval stages of walleye (*Stizostedion vitreum*). *Environmental Pollution* 111: 139-148.
- Law, A.T., Singh, A. 1991. Heavy metal content and body weight of fish from Kelang Estuary, Malaysia. *Marine Pollution Bulletin* 13 : 217-218.
- Lee, A., Woo, S., Won, H., Lee, G., Lee, T.K., Yum, S. 2013. Changes in gene expression profile due to acute toxicity of toxaphene in the marine medaka. *Molecular and Cellular Technology* 9(2): 121-128.
- Lee, D., Kim, M.S., Nam, Y.K., Kim, D.S., Gong, S.P. 2013. Establishment and characterization of permanent cell lines from *Oryzias dancena* embryos. *Fisheries and Aquatic Sciences* 16(3): 177-185.
- Lee, T-M., Huang, Y-L., Chen, M-H. 2005. Copper induction of phosphorus deficiency in *Ulva fasciata* (Ulvales, Chlorophyta). *Phycologia* 44 (6): 620-628.
- Li, H.T., Facon, T. 2001. *From Vision to Action: A Synthesis of Experiences in Southeast Asia. The FAO-ESCAP Project on National Water Visions*. FAO Corporate Document Repository: Bangkok.
- Li, Z., Fu, H. 2009. Ecological studies of medaka in a remained habitat in Qinhuangdao, Hebei, China. *Frontiers of Agriculture in China* 3(2): 216-220.
- Lim L.C. 2002. Use of decapsulated *Artemia* cysts in ornamental fish culture. *Aquaculture Research* 33(8): 575-589.
- Lin, H.C., Hwang, P.P. 1998. Acute and chronic effects of gallium chloride (GaCl₃) on tilapia (*Oreochromis mossambicus*) larvae. *Bulletin of Environmental Contamination and Toxicology* 60: 931-935.
- Lopez-Barea, J. 1996. Biomarkers to detect environmental pollution. *Toxicology Letters* 88: 77-79.
- Love, B.J., Einheuser, M.D., Nejadhashemi, A.P., 2011. Effects on aquatic and human health due to large scale bioenergy crop expansion. *Science of the Total Environment* 409(17): 3215-3229.

- Magtoon, W., Termvidchakorn, A. 2009. A revised taxonomic account of rice fish *Oryzias* (Beloniformes; Adrianichthyidae), in Thailand, Indonesia and Japan. *The Natural History Journal of Chulalongkorn University* 9(1): 35-68.
- Marchetti, R., 1965. *Critical review of the effects of synthetic detergents on aquatic life: studies and reviews*. General Fisheries Commission for the Mediterranean: FAO.
- McCarthy, L.H., Stephens, G.R., Whittle, D.M., Peddle, J., Harbicht, S., LaFontaine, C., Gregor, D.J. 1997. Baseline studies in the Slave River, NWT, 1990-1994: Part II. Body burden contaminants in whole fish tissue and livers. *The Science of the Total Environment* 197: 55-86.
- McDonnel, G. E. 2007. *Antisepsis, Disinfection, and Sterilisation- types, action and resistance*. Washington: Blackwell Publishing.
- McKim, J.M., 1977. Evaluation of tests with early life stages of fish for predicting long term toxicity. *Journal of Fish Research Board Canada* 34: 1148-1154.
- McKim, J.M., Eaton, J.G., Holcombe G.W. 1978. Metal toxicity to embryos and larvae of eight species of freshwater fish--II: copper. *Bulletin of Environmental Contamination and Toxicology* 19: 608-616.
- McMichael, G.A., Fritts, A.L., Pearsons, T.N. 1998. Electrofishing injury to stream salmonids; injury assessment at the sample, reach and stream scales. *North American Journal of Fisheries Management* 18: 894-904.
- Mellen, J., MacPhee, M.S. 2001. Philosophy of environmental enrichment: past, present, and future. *Zoo Biology* 20: 211-26.
- Michibata, H., Nojima, Y., Kojima, M.K. 1987. Stage sensitivity of eggs of teleost *Oryzias latipes* to cadmium exposure. *Environmental Research* 42: 321-327.
- Milam, C.D., Farris, J.L., Wilhide, J.D. 2000. Evaluating mosquito control pesticides for effect on target and nontarget organisms. *Archive of Environmental Contamination and Toxicology* 39: 324-328.
- Miller, J.M., Crowder L.B., Moser, M.L. 1985. Migration and utilization of estuarine nurseries by juvenile fishes: an evolutionary perspective. *Contribution to Marine Science* 27: 338-352.
- Moore, M.N. 2002. A strategy for impact and risk assessment in integrated environmental management {Electronic version}. *Revue de Médecine Vétérinaire* 153: 507-512.

- Morgan, J.E., Morgan, A.J. 1988. Earthworms as biological monitors of cadmium, copper, lead and zinc in metalliferous soils. *Environmental Pollution* 54 (2): 123-138.
- Murad, M.W. Pereira, J.J. 2011. Malaysia: Environmental Health Issues. *Encyclopedia of Environmental Health* 577-594.
- Moreno, N.C., Sofia, S.H., Martinez, C.B.R. 2014. Genotoxic effects of the herbicide Roundup Transorb® and its active ingredient glyphosate on the fish *Prochilodus lineatus*. *Environmental Toxicology and Pharmacology* 37: 448-454.
- Naruse K., Shima, A., Matsuda, M., Sakaizumi, M., Iwamatsu, T., Soeroto, B., Uwa, H. 1993. Description and phylogeny of rice fish and their relatives belonging to the suborder Adrianichthyoidei in Sulawesi, Indonesia. *The Fish Biology Journal MEDAKA* 5: 11-15.
- Naruse, K. 1996. Classification and phylogeny of fishes of the genus *Oryzias* and its relatives. *Fish Biology Journal MEDAKA* 8: 1-9.
- Neškovic, N.K., Poleksic, V., Elezovic, I., Karan, V., Budimir, M. 1996. Biochemical and histopathological effects of glyphosate on carp, *Cyprinus carpio* L. *Bulletin of Environmental Contamination Toxicology* 56: 295-302
- Niemi, G.J., Hanowski, J.M., Lima, A.R., Nicholls, T., Weiland, N. 1997. A critical analysis on the use of indicator species in management. *Journal of Wildlife Management* 61: 1240-1252.
- Niihori, M., Mogami, Y., Naruse, N., Baba, S. A. 2004. Development and swimming behavior of medaka fry in a space flight aboard the space shuttle Columbia (STS-107). *Zoological Science* 21: 923-931.
- Okay, O.S., Tolun, L., Tüfekçi, V., Karacik, B., Kungolos, A., Samaras, P., Papadimitriou, C., Petala, M., Tsiroidis, V. 2005. Comparison of several toxicity tests applied to complex wastewaters and mussel biomarkers in receiving waters. *Journal of Environmental Science and Health, Part A: Toxic/Hazardous Substances and Environmental Engineering* 40: 1525-1541.
- Omitoyin, B.O., Ajani, E.K., Fajimi, O.A. 2006. Toxicity of Gramoxone (paraquat) to juvenile African catfish *Clarias gariepinus* (Burchell, 1882). *American-Eurasian Journal of Agriculture and Environmental Science* 1(1): 26-30.

- Omoniyi, I., Agbon, A.O., Sodunke, S.A. 2002. Effect of lethal and sub-lethal concentrations of tobacco (*Nicotiana tobaccum*) leaf dust extract on weight and hematological changes in *Clarias gariepinus* (Burchell). *Journal of Applied Science and Environmental Management* 6(2): 37-41.
- Ono, T., Uematsu, T., 1957. Mating ethogram in *Oryzias latipes*. In: Nakayama, K., Oshima, Y., Yamaguchi, Y., Tsuruda, Y., Kang, I.J., Kobayashi, M., Imada, N., Honjo, T. 2004. Fertilization success and sexual behavior in male medaka, *Oryzias latipes*, exposed to tributyltin. *Chemosphere* 55: 1331–1337.
- Ostroumov, S.A. 2005. "On the multifunctional role of the biota in the self-purification of aquatic ecosystems". *Russian Journal of Ecology* 36: 452-459.
- Otozaia, S., Ishikawa-Fujiwarab, T., Odac, S., Kameib, Y., Ryod, H., Satoe, A., Nomurad, T., Mitanic, H., Tsujimurae, T., Inoharaa, H., Todob, T. 2014. p53-Dependent suppression of genome instability in germ cells. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 760: 24–32.
- Overstreet, R.M., Barnes, S.S., Manning, C.S., Hawkins, W.E. 2000. Facilities and husbandary. In *The laboratory fish*, ed. G.K. Ostrander, pp. 41-63. London: Academy Press.
- Paganelli, A., Gnazzo, V., Acosta, H., Lopez, S.L., Carrasco, A.E. 2010. Glyphosate-based herbicides produce teratogenic effects on vertebrates by impairing retinoic acid signaling. *Chemical Research in Toxicology* 23(10): 1586-1595.
- Parenti, L. R. 2008. A phylogenetic analysis and taxonomic revision of ricefishes, *Oryzias* and relatives (Beloniformes, Adrianichthyidae). *Zoological Journal of the Linnean Society* 154: 494-610.
- Park, S-Y., Choi, J. 2009. Genotoxic effects of nonylphenol and Bisphenol A exposure in aquatic biomonitoring species: freshwater crustacean, *Daphnia magna*, and aquatic midge, *Chironomus riparius*. *Bulletin of Environmental Contamination and Toxicology* 83:463–468.
- Peixoto, F. 2005. Comparative effects of the Roundup and glyphosate on mitochondrial oxidative phosphorylation. *Chemosphere* 61(8): 1115-1122.
- Perry, D.M., Weis, J.S., Weis, P. 1988. Cytogenetics effects of methylmercury in the embryos of killifish *Fundulus heteroclitus*. *Environmental Contamination and Toxicology* 17: 569-574.

- Persoone, G., Calamari, D., Wells, P., 1990. Possibilities and limitations of predictions from short-term tests in the aquatic environment. In *Short-term Toxicity Tests for Non-genotoxic Effects*, ed. P. Bourdeau, E. Somers, G.M. Richardson, J.R. Hickman, pp. 301-312. New York: John Wiley and Sons Ltd.
- Porto, J.I.R., Araujo, C.S.O., Feldberg, E. 2005. Mutagenic effects of mercury pollution as revealed by micronucleus test on three Amazonian fish species. *Environmental Research* 97(3): 287-292.
- Pruski, A.M., Dixon, D.R. 2002. Effect of cadmium on nuclear integrity and DNA repair efficiency in the cells of *Mytilus edulis* L. *Aquatic Toxicology* 57: 127-137.
- Rad, F., Bozaoğlu, S., Gözükar, S.E., Karahan, A., Kurt, G. 2006. Effects of different long-day photoperiods on somatic growth and gonadal development in Nile tilapia (*Oreochromis niloticus* L.). *Aquaculture* 255: 292-300, 2006
- Rahman, M.Z., Hossain, Z. Mollah, M.F.A., Ahmed, G.U. 2002. Effect of diazinon 60 EC on *Anabas testudineus*, *Channa punctatus* and *Barbodes gonionotus* Naga. *The ICLARM Quarterly* 25(2): 8-12.
- Raskoff, K.A., Sommer, F.A., Hamner, W.M., Cross, K.M. 2003. Collection and culture techniques for gelatinous zooplankton. *Biological Bulletin* 204: 68-80.
- Rice, D.W., Harrison, Jr. I.F.L., Jearld, A. Jr. 1980. Effects of copper on early life history stages of northern anchovy, *Engraulis mordax*. *Fisheries Bulletin* 78(2): 675-683.
- Roberts, T.R. 1998. Systematic observations on tropical Asian medakas or ricefishes of the genus *Oryzias* with descriptions of four new species. *Ichthyological Research* 45: 213-224.
- Ross, S.A., McCaffery, P.J., Drager, U C., De Luca, L.M. 2000. Retinoids in embryonal development. *Physiological Reviews* 80(3): 1021–1054.
- Samson, J.C., Goodridge, R., Olobatuyi, F., Weis, J.S. 2001. Delayed effects of embryonic exposure of zebrafish (*Danio rerio*) to methylmercury (MeHg). *Aquatic Toxicology* 51: 369-376.
- Sanchez, M.V., Cahansky, A.V., Greco, L.S.L., Rodriguez, E.M., 2005. Toxicity of mercury during the embryonic development of *Chasmagnathus granulatus* (Brachura, Varunidae). *Environmental Research* 99: 72-78.

- Sarkar, A., Arora, M.P. 2001. Role of photoperiod on seasonal ovarian development of red headed bunting *Emberiza bruniceps*. *Journal of Experimental Zoology* 4(1): 9-20.
- Sazima, I. 1986. Similarities in feeding behaviour between some marine and freshwater fishes in two tropical communities. *Journal of Fish Biology* 29: 53-65.
- Scholtz, S., Fischer, S., Gündel, U., Küster, E., Luckenbach, T., Voelker, D. 2008. The zebrafish embryo model in environmental risk assessment - applications beyond acute toxicity testing. *Environmental Science Pollution Research* 15: 394-404.
- Scudder, B.C., Carter, J.L., Leland, H.V. 1988. Effects of copper on development of the fathead minnow, *Pimephales promelas* Rafinesque. *Aquatic Toxicology* 12: 107-124.
- Seoane, A.I., Güerci, A.M., Dulout, F.N. 2002. Malsegregation as a possible mechanism of aneuploidy induction by metal salts in MRC-5 human cells. *Environmental and Molecular Mutagenesis* 40(3): 200-206.
- Seredyńska-Sobecka, B., Tomaszewska, M., Janus, M., Morawski, A.W. 2006. Biological activation of carbon filters. *Water Research* 40(2): 355-363.
- Shazili, N.A.M., Pascoe, D., 1986. Variable sensitivity of rainbow trout (*Salmo gairdner*) eggs and alevins to heavy metals. *Bulletin of Environmental Contamination and Toxicology* 36: 468-474.
- Shima, A., Mitani, H. 2004. Medaka as a research organism: past, present and future. *Mechanisms of Development* 121(7-8): 599-604.
- Sierosławska, A., Rymuszka, A., Kalinowska, R., Skowroński, T., Bownik, A., Pawlik-Skowrońska, B. 2010. Toxicity of cyanobacterial bloom in the eutrophic dam reservoir (Southeast Poland). *Environmental Toxicology and Chemistry* 29 (3): 556–560,
- Sivalingam, P.M. 1978. Biodeposited trace metals and mineral content studies of some tropical marine algae. *Botanica Marina* 21(5): 327-330.
- Somasundaram, B., King, P.E., Shackley, S. 1984. The effects of zinc on postfertilization development in eggs of *Clupea harengus* L. *Aquatic Toxicology* 5(2): 167-178.
- Sorgeloos, P., Dhert, P., Candreva, P. 2001. Use of brine shrimp, *Artemia* sp., in marine fish larviculture. *Aquaculture* 200: 147-159.

- Spry, D., Wiener, G. 1991. Metal bioavailability and toxicity to fish in low-alkalinity lakes: a critical review. *Environmental Pollution* 71: 243-304.
- Stasiūnaitė, P. 1999. Long-term heavy metal mixture toxicity to embryos and alevins of rainbow trout (*Oncorhynchus mykiss*). *Acta Zoologica Lituanica Hydrobiologia* 9(2): 40–46.
- Straškraba, M. 1995. *Nutrient cycles of terrestrial and aquatic ecosystems*. Ullmann's Encyclopedia of Industrial Chemistry, Vol. B7. Weinheim: VCH-Verlagsgesellschaft.
- Stouthart, X.J.H.X., Haans, J.L.M., Lock, R.A.C., Wendelaar Bonga, S.E. 1996. Effects of water pH on copper toxicity to early life stages of the common carp (*Cyprinus carpio*). *Environmental Toxicology and Chemistry* 15(3): 376–383.
- Strmac, M., Oberemm, A., Braunbeck, T., 2002. Effects of sediment eluates and extracts from differently polluted small rivers on zebrafish embryos and larvae. *Journal of Fish Biology* 61: 24-38.
- Sudaryanto, A., Takahashi, S., Iwata, H., Tanabe, S., Ismail, A. 2004. Contamination of butyltin compounds in Malaysian marine environments. *Environmental Pollution* 130: 347-358.
- Szarek, J., Siwicki, A., Andrzejewska, A., Terech-Majewska, E., Banaszkiewicz, T. 2000. Effects of the herbicide Roundup™ on the ultrastructural pattern of hepatocytes in carp (*Cyprinus carpio*). *Marine Environmental Research* 50(1–5): 263-266.
- Szczerbik, P., Mikołajczyk, T., Sokołowska-Mikołajczyk, M., Socha, M. Chyb, J., Epler, P. 2008. The influence of cadmium on Prussian carp oocyte maturation, development of eggs and hatching. *Czech Journal of Animal Science* 53(1): 36–44.
- Takai, A., Kagawa, N., Fujikawa, K. 2004. Susceptibility of male and female medaka (*Oryzias latipes*) fish to spontaneous and X-ray induced micronucleus formation in gill cells. *Mutation Research* 558: 131-136.
- Tan, S.G., Yusoff, F.M. 2002. Biodiversity in the Straits: What are the opportunity? In *Tropical Marine Environment: Charting Strategies for the Millenium*, ed. F.M. Yusoff, M. Shariff, H.M. Ibrahim, S.G. Tan, S.Y. Tai , pp. 137-154. Serdang, Malaysia: Malacca Straits Research and Development Centre (MASDEC).
- Tang, C-H., Hsu, T-C., Tsai, C-W., Wang, W-H. 2009. Characterization of the planktonic shrimp, *Acetes intermedius*, as a potential biomonitor for butyltin. *Journal of Environmental Monitoring* 11: 92-99.

- Timms, A. M., Kleerekoper, H., Matis, J. 1972. Locomotor response of goldfish, channel catfish, and largemouth bass to a 'copper-polluted' mass of water in an open field. *Water Resources Research* 8(6): 1574–1580.
- Tong, S.L., Ho, C.Y., Pang, F.Y. 2007. Monitoring of Ba, Mn, Cu, and Ni during estuarine mixing. *Analytical Science* 13 (1997): 373-378.
- Törnqvist, R., Jarsjö, J., Karimov, B. 2011. Health risks from large-scale water pollution: Trends in Central Asia. *Environment International* 37(2): 435-442.
- Torres de Lemos, C., Rdel, M.P., Terra, N.R., Erdtmann, B. 2001. Evaluation of basal micronucleus frequency and hexavalent chromium effects in fish erythrocytes. *Environmental Toxicology and Chemistry* 20(6): 1320-1324.
- Tucker, J.W.Jr. 1998. *Marine Fish Culture*. Massachusetts, USA: Kluwer Academic Publishers.
- Udroiu, I. 2006. The micronucleus test in piscine erythrocytes. *Aquatic Toxicology* 79(2): 201-204.
- Ullah, R, Malik, R.N., Qadir, A. 2009. Assessment of groundwater contamination in an industrial city, Sialkot, Pakistan. *African Journal of Environmental Science and Technology* 3(12): 429-446.
- UNESCO (United Nations Educational, Scientific and Cultural Organization). 2003. *Water for People, Water for Life*. Paris: UNESCO.
- Van Der Oost, R., Beyer, J., Vermeulen, N.P.E. 2003. Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environmental Toxicology and Pharmacology* 13: 57-149.
- Van Wettère, A.J., Kullman, S.W., Hintonx, D.E., Law, J.M. 2013. Immunohistochemical characterization of the hepatic progenitor cell compartment in medaka (*Oryzias latipes*) following hepatic injury. *Journal of Comparative Pathology* 149: 434-445.
- Vanzella, T.P., Martinez, C.B.R., Cólus, I.M.S. 2007. Genotoxic and mutagenic effects of diesel oil water soluble fraction on a neotropical fish species. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 631(1): 36-43.
- Ventura, B.C., Angelis, F.D., Marin-Morales, M.A. 2008. Mutagenic and Genotoxic effects of the Atrazine herbicide in *Oreochromis niloticus* (Perciformes, Cichlidae) detected by the micronuclei test and the comet assay. *Pesticide Biochemistry and Physiology* 90: 42-51.

- Vera, M.S., Lagomarsino, I., Sylvester, M., Pérez, G.L., Rodríguez, P., Mugni, H., Pizarro, H., 2010. New evidences of Roundup® (glyphosate formulation) impact on the periphyton community and the water quality of freshwater ecosystems. *Ecotoxicology* 19(4): 710-721.
- Vetemaa, M. & Saat, T. 1996. Effects of salinity on the development of fresh-water and brackish-water ruffe *Gymnocephalus cernuus* (L.) embryos. [Electronic version]. *Annales Zoologici Fennici* 33: 47–51.
- Vetillard, A., Bailhache, T. 2004. Cadmium: An endocrine disrupter that affects gene expression in the liver and brain of juvenile rainbow trout. *Biology of Reproduction* 72: 119–126.
- Videla, H.A. 1996. *Manual of Biocorrosion*. Boca Raton: CRC Press. Inc.
- Vincent-Hubert, F., Arini, A., Gourlay-Francé, C. 2011. Early genotoxic effects in gill cells and hematocytes of *Dreissena polymorpha* exposed to cadmium, B[a]P and a combination of B[a]P and Cd. *Mutation Research* 723: 26-35.
- Vinodhini, R., Narayanan, M. 2008. Bioaccumulation of heavy metals in organs of fresh water fish *Cyprinus carpio* (common carp). *International Journal of Environmental Science and Technology* 5(2): 179 – 182.
- Wadekind, C., Von Siebenthal, B., Gingold, R., 2007. The weaker points of fish acute toxicity test and how tests on embryo can solve some issues. *Environmental Pollution* 148: 385-389.
- Walseng, B., Yan, N.D., Schartau, A.K. 2003. Littoral microcrustacean (Cladocera and Copepoda) indicators of acidification in Canadian Shield Lakes. *AMBIO: A Journal of the Human Environment* 32(3): 208-213.
- Wan Maznah, W.O., Mansor, M. 2002. Aquatic pollution assessment based on attached diatom communities in the Pinang River Basin, Malaysia. *Hydrobiologia* 487: 229–241.
- Ward, G.S., Cramm, G.C., Parrish, P.R., Petrocelli, S.R., 1982. Effect of ammonium jarosite on early life stages of a saltwater fish, *Cyprinodon variegates*. *Marine Pollution Bulletin* 13(6): 191-195.
- Warts, W.A. 1998. Vertebrate and fish evolution: why can some fish live in freshwater, some in salt water, and some in both? *World Aquaculture* 29(1): 65.

- Watanabe, T., Arakawa, T., Kitajima, C., Fukusho, K., Fujita, S. 1978. Nutritional quality of living feed from the viewpoint of essential fatty acids for fish. *Bulletin of Japanese Society of Science Fisheries* 44: 1223–1227.
- Watanabe, N., Kato, M., Suzuki, N., Inoue, C., Fedorova, S., Hashimoto, H., Maruyama, S., Matsuo, S., Wakamatsu, Y. 2009. Kidney regeneration through nephron neogenesis in medaka. *Development, Growth and Differentiation* 51: 135–143.
- WATO, 1997. The Welfare of Animals (Transport) Order 1997. Retrieved 17 January 2010 from <http://www.legislation.gov.uk/ukxi/1997/1480/contents/made>
- Weis, J.S., Weis, P., 1977. Effects of heavy metals on development of the killifish, *Fundulus heteroclitus*. *Journal of Fish Biology* 11 (1): 49–54.
- WHO (World Health Organization) 1976. *Mercury*. Environmental Health Criteria 1. Geneva: WHO. Retrieved on 2nd February 2014 at <http://www.who.int/>
- WHO (World Health Organization) and UNICEF (United Nations Children's Fund) 2000. Global water supply and sanitation assessment 2000 report. USA: World Health Organization and United Nations Children's Fund. Retrieved on 27th January 2010 from http://www.who.int/water_sanitation_health/monitoring/jmp2000.pdf
- WHO (World Health Organization) 2004. Water, Sanitation and Hygiene Links to Health (Facts And Figures). Retrieved on 2nd February 2014 from <http://www.who.int/en/>
- Wiener, J.G., Spry, D.J. 1996. Toxicological significance of mercury in freshwater fish. In. *Environmental Contaminants in Wildlife, Interpreting Tissue Concentrations*, ed. A.W. Redmond-Norwood, pp. 297-339. Boca Raton: CRC-Lewis Publishers.
- Williams, N.D., Holdway, D.A. 2000. The effects of pulse-exposed cadmium and zinc on embryo hatchability, larval development and survival of Australian crimson spotted rainbow fish (*Melanotaenia fluviatilis*). *Environmental Toxicology* 15(3): 165-173.
- Williams, T.D., Readman, G.G., Owen, S.F. 2009. Key issues concerning environmental enrichment for laboratory-held fish species. *Laboratory Animals* 43: 107-120.
- Wirjoatmodjo, S., Pitcher, T.J. 1984. Flounders follow the tides to feed: evidence from ultrasonic tracking in an estuary. *Estuarine, Coastal and Shelf Science* 19: 231–241.

- Wittbrodt, J., Shima, A., Scharl, M. 2002. Medaka: a model organism for the Far East. *Nature Reviews Genetics* 3: 53-64.
- Witeska, M., Samowski, P., Lugowska, K., Kowal, E. 2014. The effects of cadmium and copper on embryonic and larval development of ide *Leuciscus idus* Linn. *Fish Physiology and Biochemistry* 40(1): 151-63.
- Woltering, D.M. 1984. The growth response in fish chronic and early life stage toxicity tests: A critical review. *Aquatic Toxicology* 5(1): 1-21.
- Won, H., Woo, S., Lee, A., Yum, S. 2013. Gene expression profile changes induced by acute toxicity of benzo[a]pyrene in marine medaka. *Toxicology and Environmental Health Science* 5(3): 138-144.
- Won, H., Yum, H., Woo, S. 2011. Identification of differentially expressed genes in liver of marine medaka fish exposed to benzo[a]pyrene. *Toxicology and Environmental Health Science* 3(1):39-45.
- Woo, S., Denis, V., Yum, S. 2014. Transcriptional changes caused by bisphenol a in *Oryzias javanicus*, a fish species highly adaptable to environmental salinity. *Marine Drugs* 12(2): 983-998.
- Woo, S., Yum, S., Jung, J.H., Shim, W.J., Lee, C-H., Lee, T-K. 2006. Heavy metal-induced differential gene expression of metallothionein in Javanese medaka, *Oryzias javanicus*. *Marine Biotechnology* 8(6): 654-662.
- Woo, S., Yum, S., Kim, D-W, Park, H-S. 2009a. Transcripts level responses in marine medaka (*Oryzias javanicus*) exposed to organophosphorus pesticide. *Comparative Biochemistry and Physiology, Part C* 149: 427-432.
- Woo, S., Yum, S., Park, H-S., Lee T-K, Ryu, J-C. 2009b. Effects of heavy metals on antioxidants and stress-responsive gene expression in Javanese medaka (*Oryzias javanicus*). *Comparative Biochemistry and Physiology, Part C* 149: 289-299.
- Woodworth J., Pascoe, D., 1982. Cadmium toxicity to rainbow trout *Salmo gairdneri* Richardson: a study of eggs and alevins. *Journal of Fish Biology* 21: 47-57.
- WWF Malaysia, 2012. Retrieved on 13th April 2012 from http://www.wwf.org.my/about_wwf/what_we_do/freshwater_main/freshwater_sustainable_water_use/projects_sustainability_of_malaysia_s_water_resources_utility

- Yadav, K.K., Trivedi, S.P. 2009. Sublethal exposure of heavy metals induces micronuclei in fish, *Channa punctata*. *Chemosphere* 77: 1495-1500.
- Yamasaki, Y., Komoike, Y., Higashinakagawa, T. 2006. Adaptive changes in TEF-1 gene expression during cold acclimation in the medaka. *Zoological Science* 23(10): 903-908.
- Yassi, A.L., Kjellstrom, T. de Kok, T., Guidotti, T. 2001. *Basic Environmental Health*. New York: Oxford University Press.
- Yap, C.K., Ismail, A., Tan, S.G., Abdul Rahim, I. 2003. Can the shell of the green-lipped mussel *Perna viridis* from the west coast of Peninsular Malaysia be a potential biomonitoring material for Cd, Pb and Zn? *Estuarine, Coastal and Shelf Science* 57(4): 623-630.
- Yount, J.D., Niemi, G.J. 1990. Recovery of lotic communities and ecosystems from disturbances: theory and applications. *Environmental Management* 14: 132-137.
- Yu, R.M.K., Wong, M.M.L., Kong, R.Y.C., Wu, R.S.S., Cheng, S.H. 2006. Induction of hepatic choriogenin mRNA expression in male marine medaka: A highly sensitive biomarker for environmental estrogens. *Aquatic Toxicology* 77(4): 348-358.
- Zelikoff, J.T. 1998. Biomarkers of immunotoxicity in fish and other non-mammalian sentinel species: a predictive value for mammals? *Toxicology* 129(1): 63-71.
- Zulkifli, S.Z., Mohamat-Yusuff, F., Ismail, A. 2012. Bioaccumulation of selected heavy metals in soldier crabs, *Dotilla myctiroides* (Decapoda:Ocypodidae) from Bagan Lalang, Selangor, Malaysia. [Electronic version]. *Acta Biologica Malaysiana* 1: 94-100.