



**UNIVERSITI PUTRA MALAYSIA**

***CONFORMATIONAL DESIGN AND CHARACTERIZATION OF A  
TRUNCATED DIAMINE OXIDASE***

**NUR NADIA BINTI RAZALI**

**IB 2014 15**



**UPM**  
UNIVERSITI PUTRA MALAYSIA  
BERILMU BERBAKTI

**CONFORMATIONAL DESIGN AND CHARACTERIZATION OF A  
TRUNCATED DIAMINE OXIDASE**

By

**NUR NADIA BINTI RAZALI**

**Thesis Submitted of School of Graduated Studies, Universiti Putra Malaysia, in  
Fulfillment of the Requirements for the Degree of Master Science**

**December 2014**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

## CONFORMATIONAL DESIGN AND CHARACTERIZATION OF A TRUNCATED DIAMINE OXIDASE

By

**NUR NADIA BINTI RAZALI**

**December 2014**

**Chairman : Professor Abu Bakar Salleh, PhD**

**Faculty: Institute of Bioscience**

Diamine oxidase, DAO (EC 1.4.3.22) is an oxidoreductase enzyme that degrades primary amines to produce aldehyde,  $H_2O_2$  and  $NH_3$ . DAO, also known as histaminase, is a homodimer protein consisting of 60 to 105 kDa subunits. Being a large protein, DAO is more susceptible to destabilising agents such as heat and pH, similar to other large proteins. Due to the large structure of native diamine oxidase, researchers are being actively conducted to develop a more versatile, and stable protein because big molecules do not perform well in microsystem. It also gave more flexibility and low cost in production. Thus, the aim of this study was to develop a low cost miniature version of the native protein while mimicking its catalytic function via protein engineering and biomolecular approaches.

In the protein engineering approach, a rational design was applied to produce a mini protein that can offer more potential i.e. for use in the bioinformatics and biological discoveries. The gene encoding diamine oxidase was identified in the UNITPROT/KB database and it showed a 61% sequence similarity when blasted with NCBI database. The protein structure was built using a homology modeling in YASARA<sup>TM</sup>. All the constituent functional domains of the protein were precisely defined and shown as predicted. The miniaturization process of the protein was started with size reduction after its conserved domain has been predicted. Based on several observations, there were three domains in the predicted structure with each domain having its own major contribution to the protein structure. The domain 4 that contained the active site was selected for the splicing process in order to retain the enzyme's active site. The molecular dynamic simulation study was executed within 20 ns for each of the structures. This step is important to detect any abnormality in the *in silico* structure. Based on results from the RMSD, SASA and RMSF it has been determined that the mini

DAO has maintained its stability which is in alignment with the parental protein. The model protein was then submitted to a protein-ligand docking simulation to study the interaction between the enzyme and a substrate. The ligand interaction showed that histamine and spermidine were excellent substrates for the mini protein because they maintained the preferences of the native protein, which often chooses short-chain length substrates. Quality assessments of the native and mini protein were also conducted using several types of verification software.

Subsequently, a number of model structures were generated using a biomolecular method where the native and mini protein has been successfully cloned into vector pET102/TOPO during the first stage. The transformation of the recombinant proteins was also conducted as the sequencing result was identical to the expected reference sequence. Next, the recombinant cell was expressed into *E. coli* BL21 (DE3) and induced with 20% of IPTG, at 25°C for 24 h. A final mini protein activity of 62.283 U/mL was determined. Crude enzyme obtained from the intracellular expression was then purified. Purification strategy for obtaining fusion mini protein with his-tagged was established. Furthermore, fusion native purification was conducted to recover the fusion protein through affinity chromatography Ni-Sepharose. High purification yield of 88% was obtained for mini protein fusion tagged (His-tagged + mature mini protein). The molecular weight of the mini protein was 42.2 kDa by SDS-PAGE. As expected, the mini protein showed a high level of interaction with histamine. Surprisingly, the mini protein had also interacted with spermidine, a long side-chain ligand, which showed that its specificity towards different substrates had broadened up. By maintaining the native protein's pH profile, the mini protein was highly active at 40 °C and stable at pH 7, with a half-life of 80 min at 50 °C. Results from the kinetic studies showed that the  $K_m$  value for native diamine oxidase was 0.274 mM and 1.3 mM for the mini protein. Based on these results, it can be concluded that the mini protein was more active than the native protein but it has kept the same interaction with histamine. This characteristic has been identified by purely rational as well as experimental combinatorial design techniques. The results have been significant to postulate that the mini protein was more stable and versatile compared to the native protein due to its size, thermostability and broad substrate specificity. This mini protein is a suitable candidate for industrial biocatalyst and biological sensor applications.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia Sebagai memenuhi keperluan untuk Ijazah Master Sains

## **REKA-BENTUK KONFORMASIONAL DAN PENCIRIAN DIAMINA OKSIDA TERANGKAS**

Oleh

**NUR NADIA BINTI RAZALI**

**Disember 2014**

**Pengerusi: Profesor Dato' Abu Bakar Salleh, PhD**

**Fakulti: Institut Biosains**

Diamina oksida, DAO (EC 1.4.3.22) adalah kumpulan enzim oksidoreduktase yang menguraikan kumpulan utama amina untuk menghasilkan aldehyd,  $H_2O_2$  dan  $NH_3$ . DAO juga dikenali sebagai 'histaminase' yang merupakan satu protin bersubunit sama yang mempunyai saiz 60 hingga 105 kDa untuk setiap subunit. Oleh kerana ianya protein yang besar, DAO adalah terdedah kepada ejen-ejen yang menggugat kestabilan seperti haba dan pH, sama seperti protin besar yang lain. Oleh kerana diamina oksida asal mempunyai struktur yang besar, penyelidikan giat dilaksanakan untuk membangunkan protin yang lebih serba boleh, fleksibel dan stabil kerana pada kebiasaannya molekul-molekul yang besar tidak menunjukkan prestasi baik dalam sistem mikro. Oleh itu, kajian ini bertujuan untuk menghasilkan versi mini protin asal yang berkos rendah sambil berupaya meniru fungsi pemangkin melalui pendekatan kejuruteraan dan biomolekul protin.

Dalam pendekatan kejuruteraan protin, reka-bentuk rasional yang digunakan untuk menghasilkan protin mini boleh menawarkan banyak potensi seperti dalam kegunaan bioinformatik dan penemuan biologi. Gen pengekodan diamina oksida dikenal pasti di dalam pangkalan data UNITPROT/KB dan ia menunjukkan 61% persamaan turutan apabila disuaigandingkan terhadap pangkalan data NCBI. Struktur protin ini dibina menggunakan teknik homologi permodelan di YASARA<sup>TM</sup>. Semua juzuk fungsian domain protin yang tepat telah ditentukan dan ditunjukkan seperti apa yang diramalkan. Proses pengecilan protin bermula dengan pengurangan saiz protin selepas pemuliharaan domain dijalankan. Berdasarkan beberapa pemerhatian, terdapat tiga domain di dalam struktur setiap domain mempunyai sumbangan utama tersendiri di dalam struktur protein. Domain ke-4 yang mengandungi tapak aktif dipilih untuk melalui proses pengecilan bagi mengekalkan tapak aktif enzim yang sedia ada. Kajian simulasi dinamik

molekul ini telah disempurnakan dalam tempoh 20 ns bagi setiap struktur. Ini adalah penting untuk mengesan kecacatan struktur secara komputer. Berdasarkan keputusan dari RMSD, SASA dan RMSF, bagi menentukan bahawa mini protin telah mengekalkan kestabilan yang sejajar dengan protin asal. Model-model protin ini kemudiannya menjalani simulasi interaksi antara substrat dan ligan bagi mengenalpasti kecenderungan enzim tersebut dengan suatu substrat. Interaksi ligan ini mendapati histamina dan spermidina merupakan substrat yang terbaik untuk protin mini kerana ia mengekalkan ciri-ciri asal protin asli. Penilaian kualiti bagi setiap protin juga dijalankan dengan menggunakan beberapa jenis perisian pengesanan.

Seterusnya, sebagai peringkat permulaan, beberapa struktur model telah dihasilkan dengan menggunakan pendekatan biomolekul, di mana, protin asli dan mini protin telah berjaya diklonkan ke dalam vektor pET102/TOPO. Transformasi rekombinan protin ini juga telah menunjukkan 100% jujukan penjajaran terhadap jujukan rujukan. Kemudian, sel rekombinan protin mini tersebut telah berjaya diekspresikan di dalam *E.coli* BL21(DE3) dengan 20% IPTG pada suhu 25 °C selama 24 j. Sebanyak 62.283 U/mL telah dikenalpasti sebagai aktiviti protin mini. Enzim mentah yang telah diperolehi daripada ekspresi intrasel, kemudiannya dituliskan. Pemurnian strategi untuk mendapatkan protin mini yang bergabung dengan tag-his telah dijalankan melalui kromatografi Ni-Sepharose. Hasil penulenan yang tinggi sebanyak 88% telah diperolehi daripada protin mini gabungan (Tag-his + protin mini yang matang). Molekul protin mini ini bersaiz adalah 42.2 kDa seperti yang ditunjukkan pada SDS-PAGE. Seperti jangkaan, protin mini telah menunjukkan tahap interaksi dengan histamina yang tinggi. Menariknya, mini DAO juga telah menunjukkan interaksi yang bagus terhadap spermidina, sebuah ligan yang mempunyai rangkaian luar yang panjang. Ini menunjukkan, spesifisiti substrat telah diperluaskan. Dengan mengekalkan profil pH seperti protin asal, protin mini ini didapati sangat aktif pada 40 °C dan stabil pada pH 7, dengan nilai separuh hayat selama 80 min pada 50 °C. Manakala, hasil daripada kajian kinetik, nilai  $K_m$  protin asal adalah 0.274 mM dan 1.3 mM bagi protin mini. Ciri-ciri ini telah disokong melalui eksperimen kaedah rasional. Berdasarkan pemerhatian ini, maka disimpulkan bahawa protin mini ini lebih stabil dan serba boleh jika dibandingkan dengan protein asal berikutan saiz, tahap ketahanan suhu dan perluasan substrat. Ini menjadikan protin mini ini sesuai untuk aplikasi dalam pelbagai bidang industri serta aplikasi penderiaan biologi.

## ACKNOWLEDGEMENTS

In the name of ALLAH, all praise is to ALLAH the Almighty. Had it not been due to His will, the thesis will not be completed.

First and foremost, I would like to thank to my supervisor Prof. Dato' Abu Bakar Salleh for his guidance, understanding encouragement and advices throughout this study. He inspired me greatly and his motivation has also not been forgotten. Also I would like to express my grateful thanks to my co-supervisor Dr. Adam Leow Thean Chor for his help and valuable comment for during this study. I give a special gratitude to my research members group (EMTECH) especially Prof. Raja Noor Zaliha Raja Abdul Rahman, Prof. Mohd Basyarudin Abdul Rahman and Dr. Shukuri Bin Mohamad Ali, without whose knowledge and assistance this study would not be successful. Furthermore, I also want to acknowledge to my lab members especially Ang Swi See, Hafizah, Hisham, Hafidza, Zuhafah, Syazwani dan Zulhimi and other EMTECH member for their sincere help and kindness in helping to coordinate my project throughout these years.

Finally, my special thanks to my family especially my parents, Razali Othman and Zainab Hashim, who are always supportive no matter how hard the situation not only during the Master study but also throughout my life. I wish to extend my appreciation to all others, not individually named here, who have contributed directly or indirectly to this project.



I certify that a Thesis Examination Committee has met on 3 December 2014 to conduct the final examination of Nur Nadia binti Razali on her thesis entitled "Conformational Design and Characterization of a Truncated Diamine Oxidase" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

**Latifah binti Saiful Yazan, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Lai Oi Ming, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Goh Kian Mau, PhD**

Senior Lecturer  
Faculty of Biosciences and Medical Engineering (FBME)  
Universiti Teknologi Malaysia  
(External Examiner)



---

**ZULKARNAIN ZAINAL, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 26 February 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Abu Bakar Salleh, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)


**Adam Leow Thean Chor, PhD**

Senior Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: \_\_\_\_\_

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_

Name of Chairman of  
Supervisory Committee: \_\_\_\_\_

Signature: \_\_\_\_\_

Name of Member of  
Supervisory Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	<b>i</b>
<b>ABSTRAK</b>	<b>iii</b>
<b>ACKNOWLEDGEMENT</b>	<b>v</b>
<b>APPROVAL</b>	<b>vi</b>
<b>DECLARATION</b>	<b>viii</b>
<b>LIST OF TABLES</b>	<b>xiv</b>
<b>LIST OF FIGURES</b>	<b>xv</b>
<b>LIST OF APPENDICES</b>	<b>xviii</b>
<b>LIST OF ABBREVIATION</b>	<b>xix</b>
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	
2.1 Diamine oxidase, characterization and approaches	3
2.2 Miniaturization of protein	9
2.2.1 Current status of mini protein	10
2.2.2 Designing of miniature enzyme	12
2.2.3 Molecular dynamic simulation	13
<b>3 METHODOLOGY</b>	
3.1 Materials	15
3.2 Bacterial source	15
3.3 Structural study of diamine oxidase	
3.3.1 Homology modeling of diamine oxidase.	15
3.3.2 Prediction of diamine oxidase domains and active site	16
3.3.3 Quality assessment and visualization	16
3.3.4 Docking analysis of diamine oxidase	17
3.3.5 Molecular dynamic simulation	17
3.4 Cloning and protein expression for recombinant and mini protein	
3.4.1 Genomic extraction of <i>Arthrobacter globiformis</i>	18
3.4.2 Quantitation of DNA template	18
3.4.3 Isolation of gene for native diamine oxidase and mini protein	
3.4.3.1 Isolation of <i>A. globiformis</i> amine oxidase gene	19
3.4.3.2 Amplification of mini protein gene	20
3.4.4 Purification of PCR Product	20
3.4.5 Preparation of <i>Escherichia coli</i> Competent Cell	20

3.4.6	Heat-shock transformation of PCR Product of native diamine oxidase and mini protein into vector, pET102/D-TOPO	21
3.4.7	Extraction of recombinant plasmid	21
3.4.8	Sequencing the native diamine oxidase and mini protein genes	21
3.4.9	Protein expression of native diamine oxidase and mini protein in <i>E. coli</i> BL21 (DE3)	21
3.4.10	Analysis of native diamine and mini protein oxidase expression by SDS-PAGE	22
3.4.10.1	The effect of temperatures and inducer concentration on protein expression	23
3.4.11	Determination of protein concentration	23
3.4.12	Diamine oxidase assay	23
3.4.13	Protein purification recombinant mini diamine oxidase	
3.4.13.1	Purification of recombinant mini diamine oxidase	23
3.4.13.2	Native-PAGE Analysis	24
3.4.13.3	Western-Blot Analysis	24
3.4.14	Characterization of purified recombinant native diamine oxidase and mini protein	
3.4.12.1	Effect of diverse substrate towards native diamine oxidase and mini protein activities	25
3.4.12.2	Effect of temperature on native diamine oxidase and mini protein activities and stabilities	25
3.4.12.3	Effect of pH on mini diamine oxidase activities	25
3.4.13	$K_m$ and $V_{max}$ of recombinant mini protein	25

## 4 RESULTS AND DISCUSSION

4.1	Structural study of native diamine oxidase	
4.1.1	Amino acid sequence determination of diamine oxidase	26
4.1.2	Determination of three-dimensional of diamine oxidase	27
4.1.3	Miniaturization of diamine oxidase	
4.1.3.1	Prediction of domain conserved in diamine oxidase	30
4.1.4	Quality assessment and sequence evaluation	34
4.1.5	Docking of diamine oxidase and its derivatives	35
4.1.6	Molecular dynamic simulation	37
4.1.6.1	Root mean square deviation (RMSD) analysis	37

4.1.6.2	Root mean square fluctuation (RMSF) analysis	38
4.1.6.3	Solvent accessibility surface area (SASA) analysis	39
4.2	Cloning and expression of recombinant native diamine oxidase and mini protein	
4.2.1	Isolation genomic of <i>A. globiformis</i> DSM 20124	40
4.2.2	Isolation of native diamine oxidase genes	
4.2.2.1	Isolation of gene for native diamine oxidase	42
4.2.2.2	Amplification of mini protein	43
4.2.3	Cloning of diamine oxidase in <i>E.coli</i> competent cell TOP10	
4.2.3.1	Cloning and transformation of native diamine oxidase gene	43
4.2.3.2	Cloning, transformation and sequencing analysis of gene mini protein	46
4.2.4	Protein expression on native diamine oxidase and mini protein	48
4.2.4.1	The effect of temperature and inducer concentration on protein expression	48
4.2.5	Protein Purification and Characterization	
4.2.5.1	Purification of recombinant mini protein	51
4.2.5.2	Native polyacrylamide gel electrophoresis (Native-PAGE) of native diamine oxidase and mini protein	55
4.2.5.3	Western blot of native diamine oxidase and mini protein	55
4.2.5.4	Characterization of recombinant native diamine oxidase and mini protein	
4.2.5.4.1	Substrate specificity for native diamine oxidase and mini protein	56
4.2.5.4.2	Effect on temperature on purified native diamine oxidase and mini protein activity	57
4.2.5.4.3	Effect of pH on mini protein activity	59
4.2.5.5	$K_m$ and $V_{max}$ of mini protein	60
<b>5</b>	<b>SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	<b>63</b>

<b>REFERENCES</b>	65
<b>APPENDICES</b>	78
<b>BIODATA OF STUDENT</b>	84





## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1	List of studies on mini protein and peptide	12
2	Characteristic and properties of primers	19
3	Component of PCR reaction	19
4	Composition of 12% polyacrylamide gel	22
5	An overview of the graphical output of UnitProtKB/SWISS-PROT.	26
6	Similarity search for the amino acid of diamine oxidase sequence using NCBI-Blast	27
7	List of models designed by miniaturization process	31
8	Ramachandran values for the predicted diamine oxidase and its conserved domain in favored, allowed and outlier region using rampage ramachandan plot server.	35
9	The binding energy of the protein-ligand interaction with several types of substrates.	36
10	Purification table of intracellular expressed fusion mini protein	54
11	Summary of characterization of native and mini DAO	62

## LIST OF FIGURES

Figure		Page
1	Three-dimensional structure of diamine oxidase.	5
2	Conserved active site of CAOs.	6
3	Biogenic amines as substrate for copper amine oxidase.	8
4	Multiple sequence alignment of amino acids sequence.	28
5	3D structure of diamine oxidase after homology modeling.	29
6	Superposition of C $\alpha$ traces of predicted structure and template.	29
7	Three-dimensional structure of diamine oxidase predicted and its conserved domain after homology modeling.	30
8	Protein conserved domain of monomer native diamine oxidase from AGAO amine oxidase.	32
9	Three-dimensional structures after homology modeling.	32
10	Superposition of C $\alpha$ traces of mini protein and template by minimization approach.	33
11	The Ramanchan Plot for diamine oxidase and its domain.	35
12	An interaction between mini protein and a substrate.	36
13	Trajectories for RMSD for native DAO, D04 and mini protein for 20 ns.	38
14	Trajectories for RMSF for native DAO, D04 and mini protein for 20 ns.	39
15	Trajectories for SASA for native DAO, D04 and mini protein for 20 ns.	40
16	Pure culture of isolate <i>A. globiformis</i> on corynebacterium agar plate	41
17	Genomic DNA extraction of <i>A. globiformis</i> DSM 20124 on 0.8% (w/v) gel electrophoresis	41

18	PCR product for diamine oxidase gene on 1.0% (w/v) agarose gel.	42
19	Gel electrophoresis of PCR product of gene encoding for miniprotein.	43
20	pET102/D-TOPO plasmid and native DAO gene on 1.0% (w/v) agarose gel.	45
21	Amplification result for native DAO gene on 1.0% (w/v) agarose gel.	45
22	Gel electrophoresis of recombinant plasmid verify the positive transformant.	46
23	Sequence alignment of mini protein sequence after sequencing and designed targeted mini protein generated <i>in silico</i> in Biology Workbench 2.0 ( <a href="http://workbench.sdsc.edu/">http://workbench.sdsc.edu/</a> ).	47
24	SDS-PAGE for native diamine amine oxidase at 30° C , induction with 0.1 M IPTG incubated for 18 h.	48
25	Temperature optimization of mini protein with standard IPTG concentration.	49
26	Inducer optimization of mini protein at 25 °C using soluble fraction	50
27	Inducer optimization of mini protein at 25 °C using insoluble fraction	51
28	Purification profile of fusion recombinant mini protein by affinity chromatography.	53
29	SDS-PAGE (12%) of purified fusion mini protein through Nickel Sepharose High Performance affinity chromatography with the crude enzyme.	54
30	Verification of recombinant mini protein by native PAGE.	55
31	Verification of recombinant mini protein by immunoblotting method.	56
32	Substrates specificity on different carbon chain length of biogenic amines.	57
33	Effect of temperature on mini protein activity and parental DAO.	58
34	The effect of temperature on mini protein stability.	59

35	The effect of pH on mini protein.	60
36	Initial velocity versus substrate concentration plot for enzyme-catalyzed reaction.	61



## LIST OF APPENDICES

Appendix		Page
A	Materials and equipments used in the project	75
B	Vector map and Multiple cloning Site of pET102/TOPO	80
C	pET102/TOPO cloning region sequence	81
D	Calibration curve for determination of protein concentration by Bradford Assay	82
E	Standard curve for hydrogen peroxidase, H <sub>2</sub> O <sub>2</sub> release determination	83

## LIST OF ABBREVIATIONS

$\mu\text{L}$	Microliter
$\mu\text{m}$	Micrometer
AGAO	<i>Arthrobacter globiformis</i> amine oxidase
BA	Biogenic amine
BSAO	<i>Bovine serum</i> amine oxidase
CAO	Copper amine oxidase
Å	Amstrong
C	Celcius
cm	Centimeter
Co	Cobalt
Da	Dalton
DH <sub>2</sub> O	Distilled water
DNA	Deoxyribonucleic acid
ECAO	<i>Escherichia coli</i> amine oxidase
g	Gram
g/L	Gram per liter
h	Hour
hDAO	Human diamine oxidase
hVAP	Human vascular adhesion protein
HPAO	<i>Hansenula polymorpha</i> amine oxidase

IPTG	Isopropyl $\beta$ -D Thiogalactoside
kbp	kilo base pair
kDa	kilo Dalton
LB	Luria bertani
mg	miligram
ml	mililiter
MAO	Mono amine oxidase
M	Molar
MD	Molecular Dynamic
nm	nanometer
Ni	Nickel
PEM	Particle Mesh Ewald
RAO	Retina Amine Oxidase
RMSD	Root Mean Square Deviation
RMSF	Root Mean Square Fluctuation
OD	Optical Density
PCR	Polymerase Chain Reaction
SASA	Solvent Accessible Surface Area
SDS	Sodium Dodecyl Sulphate
SDS-PAGE	Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis
SOC	Super Optimum Broth
TEMED	N, N, N, N-Tetramethyllenediamine
TPQ	Topaquinone

U/mg	Unit per miligram
U/mL	Unit per mililiter
UV	Ultraviolet
v/v	Volume per volume
V	Volt
w/v	Weight per volume



© COPYRIGHT UPM



## CHAPTER 1

### INTRODUCTION

An enzyme's function is greatly influenced by its amino acid composition and sequence leading to protein conformation. Uniquely, an enzyme can catalyse a specific substrate due to the clefts on its surface, known as active sites that are complimentary in shape to that substrate. In addition, the enzyme's amino acid residues are responsible for forming the binding site that will lead to its specificity and selective properties, which will enable the enzyme to catalyze a reaction at a fantastic rate. Protein can also achieve their functionality by folding into compact and well-arranged structures. A protein can be found in the tertiary and quaternary structures, while, secondary structures only consist of  $\alpha$ -helices,  $\beta$ -sheets and turns. A simple combination of secondary structure elements was found in protein structure, called motif. These structures play a role in the different functions of proteins (Stiger et al., 1999).

Numerous studies have been reported on full length proteins with various functions inside living cells. A large protein normally consists of three-dimensional structure which is stable and well folded. A protein may also have several domains whereby a domain may exist in specific orientations to create different functions. With more than 35% of hydrophilic residues in a protein of more than 300 residues, the protein is likely to consist of more hydrophobic cores (George et al., 2005). This is an important characteristic for protein solubility. Generally, both natural and chemically synthesised large proteins are known to be active and functional. However, due to high demands of large protein in bulk or in relatively high concentration from the pharmaceutical and biosensor industry, there is an urgent requirement to fulfil this demand. Thus, to overcome this issue, a recombinant proteins have to be generated (Smyth et al., 2003). However, large-scale production of large protein can lead to high costs. In addition, producing recombinant proteins may sometimes lead to mismatches due to the extended length of the amino acid sequence in the original protein. Furthermore, the complex orientation that exists in the native protein may cause difficulty in recognising elements especially in the field of biosensors (Casset et al., 2003). Thus, developing a smaller protein that is related to a specific region of the native protein can be a good strategy for protein mimicking. Significant research efforts are devoted towards miniaturizing protein via truncations, corresponding to a particular specificity in purpose while simultaneously exploring their limited function.

Mini proteins, are low molecular weight proteins that active in specific biological processes. They have remarkable stability along with a high binding affinity and selectivity (Zoller et al., 2011). A better control on the mini protein could easily be activated with a lesser cross-reactivity in addition to being low cost and easier to generate. Small proteins also have fast turnaround time in production (Lipovsek et al., 2004).

Therefore, a small protein that mimics the active site of the enzyme can be developed without changing the specific characteristic of the native protein. In this research the targeted diamine oxidase (DAO) from *Athrobacter globiformis* had been miniaturized, for possible use as a biological receptor for specific biogenic amines in biocatalysis and biosensor technology.

Why diamine oxidase? In microorganism, diamine oxidases generally have a nutritional role in the utilization of primary amines as the sole source of nitrogen or carbon. They have been proposed to provide a signal in wound healing in plant (McGrath et al., 2009; Langley et al., 2006). While in mammals, they play an important role in regulating the intracellular level of amines such as neurotransmitter, detoxification, differentiation, development of cell and some CuAOs are known to be tissue-specific (Duff et al., 2003; Matsumura et al., 2004; Lunelli et al., 2005). In fact, DAO was the enzyme originally identified to clear extrinsic histamine from crumple lung and liver (Murray et al., 1999). Furthermore, in application wise, the enzyme is used pharmacologically in many types of disease such as diabetes and Alzheimer's disease (Masini and Raimondi, 2009). However, the use of DAO as biosensor is becoming important nowadays. It concluded that the interest in DAO is increasing in several fields due to its involvement in numerous physiological and metabolic pathway that related to the biological function (Lunelli et al., 2005). The versatility of DAO, provided a good reason why diamine oxidase was chosen to be the targeted protein. This knowledge is useful in understanding protein structure and function. It is hoped that this research may significantly contribute to the development of new biocatalysts in the future.

The aim of this research is to design a mini protein that functionally mimics the native enzyme. To achieve this objective, the following approaches have been undertaken.

1. To identify the selective active platform and scaffold of diamine oxidase.
2. To construct peptide structure *in silico*.
3. To experimentally validate the designed protein

## REFERENCES

- Aarsen, P. N. and Kemp, A. 1964. Rapid Spectrophotometric Micromethod for Determination of Histaminase Activity. *Nature* 204:1195.
- Ahmed, H. 2004. *Principles and Reactions of Protein Extraction, Purification, and Characterization*, pp. 102-103. Boca Raton: CRC Press.
- Airenne, T.T., Nymalm, Y., Kidron, H., Smith, D.J., Pihlavisto, M., Salmi, M., Jalkanen, S., Johnson, M.S. and Salminen, T.A. 2005. Crystal Structure of the Human Vascular Adhesion Protein-1: Unique Structural Features with Function Implications. *Protein Science* 14: 1964-1974.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman D. J. 1990. Basic Local Alignment Search Tool. *Journal Molecular Biology* 215: 403-410.
- Bagchi, P. and Hopper, W. 2011. Virtual Screening of Compounds from Valeriana Jatamansi with  $\alpha$ -Synuclei. *International Conference on Bioscience, Biochemistry and Bioinformatics* 5: 11-14.
- Barua, B., Lin, J.C., Williams, V.D., Kummeler, P., Neidigh, J.W. and Andersen, N.H. 2008. The Trp-cage: optimizing the stability of a globular miniprotein. *Protein Engineering, Design and Selection* 21(3):171- 185.
- Basavakumar, K., Karunasagar, I. and Karunasagar, I. 1992. Ability of fish Associated Bacteria to Produce Amines Suspected to be Involved in Scombroid Poisoning. *Indian Journal of Microbiology* 32: 75-79.
- Bharalee, R., Gupta, S., Bandyopadhyay, T., Gohain, B., Agarwala, N., Kalita M.C. and Das, S. 2012. Molecular Cloning, Expression and Computational Analysis of a Water Stress Inducible Copper-Containing Amine Oxidase Gene (CuAO) from Tea Plant [*Camellia sinensis* (L.) O. Kuntze]. *African Journal of Biotechnology* 11(89): 15547-15555.
- Belov, K., Harrison, G.A. and Cooper, D.W. 1998. Molecular Cloning of the cDNA Encoding the Constant Region of the Immunoglobulin a Heavy Chain (C Alpha) from a Marsupial: *Trichosurus Vulpecula* (Common Brushtail Possum). *Immunology Letter* 60 (2-3): 165-170.
- Bouvrette, P., Male, K.B., Luong, J. H.T. and Gibbs, B.F. 1997. Amperometric Biosensor for Diamine Using Diamine Oxidase Purified from Porcine Kidney. *Enzyme and Microbial Technology* 20 (1): 32-38.
- Bover-Cid, S. and Holzapfel, W. H. 1999. Improved Screening Procedure for Biogenic Amine Production by Lactic Acid Bacteria. *International Journal of Food Microbiology* 53: 33-41.

- Bradford, M. 1976. A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. *Analytical Biochemistry* 7(72): 248-254.
- Butt, A. M., Batool, M. and Tong, Y. 2011. Homology Modeling, Comparative Genomics and Functional Annotation of *Mycoplasma Genitalium* Hypothetical Protein MG\_237. *Bioinformatics* 7(6): 299-303.
- Carpene, C., Bour, S., Visentin, T., Pellati, F., Benvenuti, S., Iglesias-Osma, M.C., Garcia-Barrado, M. J. and Valet, P. 2005. Amine Oxidase Substrates for Impaired Glucose Tolerance Correction. *Journal Physiological Biochemistry* 61(2): 405-420.
- Casida, J. L. E. and Liu, K. 1974. *Arthrobacter globiformis* and Its Bacteriophage in Soil. *Applied Microbiology* 28(6): 951-959.
- Casset, F., Roux, F., Mouchet, P., Bes, C., Chardes, T., Granier, C., Mani, J., Pugnieri, M., Laune, D., Pau, B., Kaczorek, M., Lahana, R. and Rees, A. 2003. A Peptide Mimetic of an Anti-CD4 Monoclonal Antibody by Rational Design. *Biochemical and Biophysical Research Communications* 307: 198-205.
- Cedrone, F., Ménez, A. and Quéméneur, E. 2000. Tailoring New Enzyme Functions by Rational Redesign. *Current Opinion in Structural Biology* 10: 405-410.
- Chang, H. and Yang, A. 2014. Therapeutic peptides: *Method and Protocol, Methods in Molecular Biology*, Vol 1088, ed. A.E. Nixon, pp. 1-17.
- Chaplin, J.A., Budde, C.L. and Khmelnskiy, Y.L. 2001. Catalysis by Amine Oxidases in Nonaqueous Media. *Journal of Molecular Catalyst B* 13: 69.
- Chassande, O., Renard, S., Barbry, P. and Lazdunski, M. 1994. The Human Gene for Diamine Oxidase, an Amiloride Binding Protein. Molecular Cloning, Sequencing, and Characterization of the Promoter. *The Journal of Biological Chemistry* 269(20): 14484-14489.
- Chin, J.W., Gotzfeld, R.M. Fabian, M.A. and Schepartz, A. 2001. Methodology for Optimizing Functional Mixture Proteins Based on Avian Pancreatic Polypeptide using Phage- Display. *Bioorganic and Medical Chemistry Letters* 11:1501- 1505.
- Choi, Y. H., Matsuzaki, R., Fukui, T., Shimizu, E., Yorifuji, T., Sato, H., Ozaki, Y. and Tanizawa, K., 1995. Copper/topa quinine-containing histamine oxidase from *Arthrobacter globiformis*. Molecular Cloning and Sequencing, Overproduction of Precursor Enzyme, and Generation of Topo Quinone Cofactor. *Journal of Biological Chemistry* 270: 4712-4720.

- Chothial, C. and Lesk, A. M. 1986. The Relation Between the Divergence of Sequence and Structure in Proteins. *The EMBO Journal* 5(4): 823-826.
- Copeland, R.A. 2000. *Enzymes; A Practical Introduction to Structure, Mechanism and Data Analysis*. 2<sup>nd</sup> Edition. Wiley-VCH Inc.
- Cui, S. and Selwood, L. 2000. cDNA Cloning, Characterization, Expression and Recombinant Protein Production of Leukemia Inhibitory Factor (LIF) from the Marsupial, the Brushtail Possum (*Trichosurus Vulpecula*). *Gene* 243(1-2):167-178.
- Davis, G.D., Elisee, C., Newham, D.M. and Harrison, R.G. 1999. New Fusion Protein Systems Designed to Give Soluble Expression in *Escherichia coli*. *Biotechnology Bioengineering* 65(4): 382-388.
- Di Paolo, M.L., Lunelli, M., Fuxreiter, M., Rigo, A., Simon, I. and Scarpa, M. 2011. Active Site Residue Involvement in Monoamine or Diamine Oxidation Catalysed by Pea Seedling Amine Oxidase. *Federation of Biochemical Societies Journal* 278(8): 1232-1243.
- Dixit, S.B., Ponomarev, S. Y. and Beveridge, D. L. 2006. Root Mean Square Deviation Probability Analysis of Molecular Dynamics Trajectories on DNA. *Journal of Chemical Information and Modeling* 46: 1084-1093.
- Dong, Z., Wang, Y., Yin, Y. and Liu, J. 2011. Supramolecular Enzyme Mimics by Self-Assembly. *Current Opinion in Colloid & Interface Science* 16(6): 451-458.
- Duff, A.P., Cohen, A.E., Ellis, P.J., Kuchar, J.A., Langley, D.B., Shepard, E.M., Dooley, D.M., Freeman, H.C. and Guss, J.M. 2003. The Crystal Structure of *Pichia Pastoris* Lysyl Oxidase. *Biochemistry* 42(51): 15148-15157.
- Durham, E., Dorr, B., Woetzel, N., Staritzbichler, R. and Meiler, J. 2009. Solvent Accessible Surface Area Approximations for Rapid and Accurate Protein Structure Prediction. *Journal of Molecular Modeling* 15(9): 1093-1108.
- Elovaara, H., Kidron, H., Parkash, V., Nymalm, Y., Bligt, E., Ollikka, P., Smith, D. J., Pihlavisto, M., Salmi, M., Jalkanen, S. and Salminen, T.A. 2011. Identification of Two Imidazole Binding Sites and Key Residues for Substrate Specificity in Human Primary Amine Oxidase AOC3. *Biochemistry* 50(24): 5507-5520.

- Ernst, J.T., Becerril, J., Park, H. S., Yin, H. and Hamilton, A.D. 2003. Design and Application of an  $\alpha$ -Helix-Mimetic Scaffold Based on an Oligoamide-Foldamer Strategy: Antagonism of the Bak BH3/Bcl-xL Complex. *Angewandte Chemie* 5: 553-557.
- Eschbach, M., Möbitz, H., Rompf, A. and Jahn, D. 2003. Members of the Genus *Arthrobacter* Grow Anaerobically Using Nitrate Ammonification and Fermentative Processes: Anaerobic Adaptation of Aerobic Bacteria Abundant in Soil. *FEMS Microbiology Letter* 223(2): 227-230.
- Figuera-Losada, M. and LoGrasso, P.V. 2012. Enzyme Kinetics and Interaction Studies for Human JNK1 $\beta$ 1 and Substrates Activating Transcription Factor 2 (ATF2) and c-Jun N-terminal Kinase (c-Jun). *The Journal Of Biological Chemistry* 287(6): 13291-13302.
- Fusco, M.D., Federico, R., Boffi, A., Macone, A., Favero, G. and Mazzei, F. 2011. Characterization and Application of a Diamine Oxidase from *Lathyrus Sativus* as Component of an Electrochemical Biosensor for the Determination of Biogenic Amines in Wine and Beer. *Analytical and Bioanalytical Chemistry* 401: 707-716.
- Gallagher, S. R. 2001. One-Dimensional Electrophoresis Using Nondenaturing Conditions. *Current Protocols in Cell Biology* 10: 1-5
- Ganguly, B. and Prasad, S. 2012. Homology Modeling and Functional Annotation of Bubaline Pregnancy Associated Glycoprotein 2. *Journal of Animal Science and Biotechnology* 3(13): 1-9.
- Gao, N., Li, H., Li, Q., Liu, J. and Luo, G. 2011. Synthesis and Kinetic Evaluation of a Trifunctional Enzyme Mimic With a Dimanganese Active Centre. *Journal Inorganic Biochemistry* 105(2): 283-288.
- Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M.R., Appel, R.D. and Bairoch, A. 2005. Protein Identification and Analysis Tools on the ExPASy Server. *The Proteomics Protocols Handbook*, ed. J.M. Walker, pp. 571-607. New York: Humana Press.
- Gerlt, J.A. and Babbitt, P.C. 2009. Enzyme (re)design : Lesson From Natural Evolution and Computation. *Current Opinion in Chemical Biology* 13: 10-18.
- George, R.A., Lin, K. and Heringa, J. 2005. Scooby-domain: Prediction of Globular Domains in Protein Sequence. *Nucleic Acid Research* 33(2): 160-163.
- Goodfellow, J.M. 1995. *Computer Modeling in Molecular Biology*, ed. H. Kraus, pp. 1-103. Weinheim: VCH Verlagsgesellschaft Publishers.

- Greene, L.H., Lewis, T. E., Addou, S., Cuff, A., Dallman, T., Dibley, M., Redfern, O., Pearl, F., Nambudiry, R., Reid, A., Sillitoe, I., Yeats, C., Thornton, J.M. and Orengo, C.A. 2007. The CATH Domain Structure Database: New Protocols and Classification Levels Give a More Comprehensive Resource for Exploring Evolution. *Nucleic Acids Research* 35: 291-297.
- Haas, J., Roth, S., Arnold, K., Kiefer, F., Schmidt, T., Bordoli, L., and Schwede, T. 2013. The Protein Model Portal -A Comprehensive Resource for Protein Structure and Model Information. *Database: The Journal of Biological Databases and Curation* 2013: 1-8.
- Hames, B.D. 1990. *One-Dimensional Polyacrylamide Gel Electrophoresis*. In *Gel Electrophoresis of Proteins: A Practical Approach*, 2nd edition, ed. B.D. Hames and D. Rickwood, pp.383. New York: Oxford University Press.
- Hashimoto, T., Mitani, A. and Yamada, Y. 1990. Diamine Oxidase from Cultured Roots of *Hyoscyamus niger*: It's Function in Tropane Alkaloid Biosynthesis. *Plant Physiology* 93(1): 216-221.
- High, A., Prior, T., Bell, R.A. and Rangachari, P.K. 1999. Probing the "Active Site" of Diamine Oxidase: Structure-activity Relations for Histamine Potentiation by *O*-Alkylhydroxylamines on Colonic Epithelium. *Journal of Pharmacology and Experimental Therapeutic* 288: 490-501.
- Holtje, H.D., Sippl, W., Rognan, D. and Folkers, G. 2008. *Molecular Modeling: Basic Principles and Applications*, 3<sup>rd</sup> Edition, p. 126-145. Weinheim: WILEY-VCH Verlag GmbH & Co. KGaA.
- Hollingsworth, S.A. and Karplus, P.A. 2010. A Fresh Look at the Ramachandran Plot and the Occurrence of Standard Structures in Proteins. *Biomolecular Concepts* 1: 271-283.
- Holm, R.H., Kennepohl, P. and Solomon, E.I. 1996. Structural and Functional Aspects of Metal Sites in Biology. *Chemical Review* 96: 2239-2314.
- Jalilani, H.Z., Farajnia, S., Safdari, Y., Mohammadi, S.A., Barzegar, A. and Talebi, S. 2014. Optimized Condition for Enhanced Soluble-Expression of Recombinant Mutant *Anabaena Variabilis* Phenylalanine Ammonia Lyase. *Advanced Pharmaceutical Bulletin* 4(3): 261-266.
- James, J.J., Lakshmi, B.S., Seshasayee, A.S.N. and Gautam, P. 2007. Activation of *Candida rugosa* Lipase at Alkane-aqueous Interfaces: A Molecular Dynamics Study. *Federation of European Biochemical Societies Journal* 581: 4377-4383.
- Karovicova, J. and Kohajdova, Z. 2005. Biogenic Amines in Food. *Chemical Paper*. 59: 70-79.

- Karplus, M., and McCammon, J.A. 2002. Molecular Dynamic Simulations of Biomolecules. *Nature Structural Biology* 9(9): 646-788.
- Katakly, R. and Morgan, E. 2003. Potential of Enzyme Mimics in Biomimetic Sensors: a Modified-cyclodextrin as a Dehydrogenase Enzyme Mimic. *Biosensor Bioelectronics* 18(11): 1407-1417.
- Keerti, Gupta A., Kumar V., Dubey A., and Verma A. K. 2014. Kinetic Characterization and Effect of Immobilized Thermostable  $\beta$ -Glucosidase in Alginate Gel Beads on Sugarcane Juice. *Biochemistry* 2014: 1-8.
- Klinman, J.P. 2003. The Multi-Functional Topa-Quinone Copper Amine Oxidases. *Biochimica et Biophysica Acta* 1647: 131-137.
- Knapp, B., Frantal, S., Cibena, M., Schreiner, W. and Bauer, P. 2011. Is an Intuitive Convergence Definition of Molecular Dynamics Simulations Solely Based on the Root Mean Square Deviation Possible? *Journal Of Computational Biology* 18 (8): 995-1005.
- Kony, D. B., Hünenberger, P. H. and van Gunsteren, W. F. 2007. Molecular Dynamic Simulations of the Native and Partially Folded States of Ubiquitin: Influence of Methanol Cosolvent, pH, and Temperature on the Protein Structure And Dynamics. *Protein Science* 16(6): 1101-1118.
- Krajewska, B. 2004. Application of Chitin and Chitosan-Based Materials for Enzyme Immobilizations: a Review. *Enzyme and Microbial Technology* 35: 126-139.
- Krieger, E., Nabuurs, S.B. and Vriend, G. 2003. *Homology Modeling, in Structural Bioinformatics*, volume 44, eds P. E. Bourne and H. Weissig, Hoboken.,New Jersey: John Wiley & Sons, Inc.,
- Kritzer, J.A., Zutshi, R., Cheah, M., Ran, F.A., Webman, R., Wongjirad, T.M. and Schepartz, A. 2006. Miniature Protein Inhibitor of the p53-hDM2 Interaction. *Chembiochem* 7: 29-31.
- Kuntz-Jr, I.D. 1971. Hydration of macromolecules. III. Hydration of polypeptides *Journal of American Chemical Society* 93(2): 514-516.
- Kumar, V., Dooley, D.M., Freeman, H.C., Guss, J.M., Harvey, I., McGuirl, M.A., Wilce, M.C.J. and Zubak, V.M. 1996. Crystal Structure of a Eukaryotic (Pea Seedling) Copper-Containing Amine Oxidase at 2.2 Å Resolutions. *Structure* 4(8): 943-955.
- Laemmli, U.K. 1970. Cleavage of Structural Proteins During the Assembly of the Head of Bacteriophage T4. *Nature* 227(5259): 680-685.



- Langley, D.B., Duff, A.P., Freeman, H.C. and Guss, J.M. 2006. The Copper - containing Amine Oxidase from *Arthrobacter globiformis*: Refinement at 1.55 and 2.20 Å Resolution in Two Crystal Forms. *Acta Crystallographica Section F: Structure Biology Crystallization Communication* 62(11): 1052-1057.
- Langley, D.B., Brown, D.E., Cheruzel, L.E., Contakes, S.M., Duff, A.P., Hilmer, K. M., Dooley, D.M., Gray, H.B., Guss, J.M. and Freeman, H.C. 2008. Enantiomer-specific Binding of Ruthenium (II) Molecular Wires by the Amine Oxidase of *Arthrobacter globiformis*. *Journal of the American Chemical Society* 130(25): 8069-8078.
- Lapa-Guimarães, J. and Pickova, J. 2004. New Solvent Systems for Thin-layer Chromatographic Determination of Nine Biogenic Amines in Fish and Squid. *Journal Chromatography A* 1045(1-2): 223-232.
- Lele, B.S., Kulkarni, M.G. and Mashelkar, R.A. 1999. Productive and Nonproductive Substrate Binding in Enzyme Mimics. *Polymer* 40: 4063-4070.
- Letunic, I., Doerks, T. and Bork, P. 2012. SMART 7: Recent Updates to the Protein Domain Annotation Resource. *Nucleic Acids Research* 40: 302-305.
- Li, R., Dowd, V., Stewart, D.J., Burton, S.J. and Lowe, C.R. 1998. Design, Synthesis and Application of a Protein Mimetic. *Nature Biotechnology* 16: 190-195.
- Lipovsek, D. and Pluckthun, A. 2004. In-Vitro Protein Evolution by Ribosome Display and mRNA Display. *Journal of Immunological Methods* 290: 51-67.
- Liu, B., Gao, B., Xua, X., Hong, W., Sub, Y. and Yue, Q. 2011. Treatment of Pulp and Paper Wastewater by Using the Mimic Enzyme Prepared from Iron-Based Biomimetic Compound. *Desalination* 278(1-3): 36-41.
- Liu, H., Zhao, Y. and Li, Y. 2006. The synthesis and Characterization of a Helical Mixture Protein Mimicking the OGT Active Domain. *International Journal of Peptide Research and Therapeutic* 12: 237-241.
- Liu, S. 2006. *Method Development and Application in Protein Structure Prediction, Computational Molecular Docking and Network Biology*. pp. 50. PhD Thesis, State University of New York, Buffalo.
- Lorenzo, J.M., Martínez, S., Franco, I. and Carballo, J. 2008. Biogenic Amine Content During the Manufacture of Dry-Cured Lacon, a Spanish Traditional Meat Product: Effect of Some Additives. *Meat Science* 77: 287-293.

- Loukou, Z. and Zotou, A. 2003. Determination of Biogenic Amines as Dansyl Derivatives in Alcoholic Beverages by High-performance Liquid Chromatography with Fluorimetric Detection and Characterization of the Dansylated Amines by Liquid Chromatography-atmospheric Pressure Chemical Ionization Mass Spectrometry. *Journal of Chromatography A* 996(1-2): 103-113.
- Lovell, S.C., Davis, I.W., Arendall, W.B., de Bakker, P.I., Word, J.M., Prisant, M.G., Richardson, J.S. and Richardson, D.C. 2003. Structure validation by  $\alpha$ -Geometry:  $\phi$ ,  $\psi$  and  $C$ -beta Deviation. *Proteins: Structure Function Genetics* 50: 437-450.
- Lunelli, M., Di Paolo, M.L., Biadene, M., Calderone, V., Battistutta, R., Scarpa, M., Rigo, A. and Zanotti, G. 2005. Crystal Structure of Amine Oxidase from Bovine Serum. *Journal of Molecular Biology* 346: 991-1004.
- Macedo-Ribeiro, S., Almeida, C., Calisto, B.M., Friedrich, T., Mentele, R., Stürzebecher, J., Fuentes-Prior, P. and Pereira, P.J.B. 2008. Isolation, Cloning and Structural Characterisation of Boophilin, a Multifunctional Kunitz-Type Proteinase Inhibitor from the Cattle Tick. *PLoS ONE* 3(2): e1624.
- Mahmood, T. and Yang, P.C. 2012. Western blot: Technique, Theory, and Troubleshooting. *North American Journal of Medical Sciences* 4(9): 429-434.
- Maintz, L. and Novak, N. 2007. Histamine and histamine intolerance. *The American Journal of Clinical Nutrition* 85: 1185-1196.
- Malwane, S. and Deutch, C. E. 1999. Adaptive Characteristics of Salt-induced Myceloids of *Arthrobacter globiformis*. *Antonie van Leeuwenhoek* 75: 335-344.
- Malek, J.A. and Haft, D.H. 2001. Conserved Protein Domains are Maintained in an Average Ratio to Proteome Size. *Genome Biology* 2(5): 1-16.
- Martin, L., Barthe, P., Combes, O. Roumestand, C. and Vita, C. 2000. Engineering Novel Bioactive Mini Proteins on Natural Scaffolds. *Tetrahedron* 56: 9451-9460.
- Marx, U.C., Korsinczky, M.L., Schirra, H.J., Jones, A., Condie, B., Otvos, L. J. and Craik, D.J. 2003. Enzymatic Cyclization of a Potent Bowman-Birk Protease Inhibitor, Sunflower Trypsin Inhibitor 1, and Solution Structure of an Acyclic Precursor Peptide. *The Journal of Biological Chemistry* 278(24): 21782-21789.
- Marx, V. 2005. Watching Peptide Drugs Grow Up. *Chemical and Engineering News* 83(11): 17-24.

- Masini, E. and Raimondi, L. 2009. *Copper Amine Oxidases; Structures, Catalytic Mechanisms, and Role in Pathophysiology*, ed. G. Floris and B. Mondovi, pp. 239-240. Boca Raton, Florida: CRC Press.
- Mason, J.M. 2010. Design and Development of Peptides and Peptide Mimetic as Antagonists for Therapeutic Intervention. *Future Medicinal Chemistry* 2(12):1813-1822.
- Matsumura, K., Hisada, H., Obata, H., Hata, Y., Kawato, A., Abe, Y. and Akita, O. 2004. A Novel Amine Oxidase-Encoding Gene from *Aspergillus oryzae*. *Journal of Bioscience and Bioengineering* 98: 359-365.
- McGrath, A.P., Hilmer, K.M., Collyer, C.A., Dooley, D.M. and Guss, J.M. 2010. A New Crystal Form of Human Diamine Oxidase. *Acta Crystallographica Section F: Structure Biology Crystallization Communication* 66(2): 137-142.
- McGrath, A.P., Hilmer, K.M., Collyer, C.A., Shepard, E.M., Elmore, B.O., Brown, D.E., Dooley, D.M. and Guss, J.M. 2009. Structure and Inhibition of Human Diamine Oxidase. *Biochemistry* 48(41): 9810-9822.
- Meeuwissen, J. and Reek, J.N.H. 2010. Supramolecular Catalysis Beyond Enzyme Mimics. *Nature Chemistry* 2: 615-621.
- Merten, O. 2001. Recombinant Protein Production with Prokaryotic and Eukaryotic Cells. *A Comparative View on Host Physiology*, pp. 164-165. Dordrecht: Kluwer Academic Publisher.
- Morris, G.M., Goodsell, D.S., Halliday, R.S., Huey, R., Hart, W.E., Belew, R.K., Olson, A.J. 1998. Automated Docking using a Lamarckian Genetic Algorithm and an Empirical Binding Free Energy Function. *Journal of Computational Chemistry* 19: 1639-1662.
- Munjal, N. and Sawhney, S.K. 2002. Stability and Properties of Mushroom Tyrosinase Entrapped in Alginate, Polyacrylamide and Gelatin Gels. *Enzyme and Microbial Technology* 5(30): 613-619.
- Murray, J.M., Saysell, C.G., Wilmot, C.M., Tambyrajah, W.S., Jeager, J., Knowles, P.F., Phillips, S.E.V. and McPherson, M.J. 1999. The Active Site Base Controls Cofactor Reactivity in *Escherichia coli* Amine Oxidase: X-ray Crystallographic Studies with Mutational Variants. *Biochemistry* 38:8217-8227.
- Murzin, A.G., Brenner, S.E., Hubbard, T. and Chothia, C. 1995. SCOP: A Structural Classification of Proteins Database for the Investigation of Sequences and Structures. *Journal Molecular Biology* 247: 536-540.

- Naila, A., Flint, S., Fletcher, G., Bremer, P. and Meerdink, G. 2010. Control of Biogenic Amines in Food—Existing and Emerging Approaches. *Journal of Food Science* 75(7): 139-150.
- Pace, C.N. and Scholtz, J.M. 1998. A Helix Propensity Scale Based on Experimental Studies of Peptides and Proteins. *Biophysical Journal* 75: 422-427.
- Pearlman, D.A. 2005. Evaluating the Molecular Mechanics Poisson-boltzmann Surface Area Free Energy Method using a Congeneric Series of Ligands to p38 MAP Kinase. *Journal of Medicinal Chemistry* 48(24): 7796-7807.
- Pietrangeli, P., Federico, R., Mondovì, B. and Morpurgo, L. 2007. Substrate Specificity of Copper-Containing Plant Amine Oxidases. *Journal of Inorganic Biochemistry* 101(7) : 997-1004
- Rao, G.N., Rao, A.A., Rao, P.S. and Muppalaneni, N.B. 2012. A Tool for the Post Data Analysis of Screened Compounds Derived from Computer-aided Docking Scores. *Bioinformation* 9(4): 207-209.
- Rinaldi, A.C., Porcu, C.M., Oliva, S., Curreli, N., Rescigno, A., Sollai, F., Rinaldi, A., Finazzi-Agró, A. and Sanjust, E. 1998. Biosynthesis of the Topaquinone Cofactor in Copper Amine Oxidases- Evidence from Model Studies. *European Journal of Biochemistry* 251(1-2): 91-97.
- Roy, A., Kucukural, A. and Zhang, Y. 2010. I-TASSER: A Unified Platform for Automated Protein Structure and Function Prediction. *Nature Protocol* 5(4): 725-738.
- Saaïd, M., Saad, B., Hashim, N.H., Ali, A.S.M., Saleh, M.I. 2009. Determination of Biogenic Amines in Selected Malaysian Food. *Food Chemistry* 113(4): 1356-1362.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989. *Molecular Cloning: A Laboratory Manual*, second edition, pp.174. New York: Cold Spring Harbor Laboratory Press
- Seeliger, D., and de Groot, B. L. 2010. Ligand docking and binding site analysis With PyMOL and Autodock/Vina. *Journal of Computer-Aided Molecular Design* 24: 417-422.
- Sekiguchi, Y., Makita, H., Yamamura, A. and Matsumoto, K. 2004. A Thermostable Histamine Oxidase from *Arthrobacter crystallopoietes* KAIT-007. *Journal of Bioscience and Bioengineering* 97: 104-110.

- Schroder, G.F., Brunger, A.T. and Levitt, M. 2007. Combining Efficient Conformational Sampling with a Deformable Elastic Network Model Facilitates Structure Refinement at Low Resolution. *Structure* 15(12): 1630-1641.
- Sillitoe, I., Cuff, A.L., Dessailly, B.H., Dawson, N.L., Furnham, N., Lee, D., Lees, J.G., Lewis, T.E., Studer, R.A., Rentzsch, R., Yeats, C., Thornton, J.M. and Orengo, C.A. 2013. New Functional Families (FunFams) in CATH to Improve the Mapping of Conserved Functional Sites to 3D Structures. *Nucleic Acids Research* 41: 490-498.
- Singh, R., Chaturvedi, N. and Singh, V.K. 2012. In-silico Study of Herbal Compounds (Baicalin, Curcumin and Dronabinol) as Novel Mao Inhibitors for Parkinson's Disease Treatment. *International journal of Life science & Pharma Research* 2(3): 81-98.
- Sippl, J.M. 1993. Recognition of Errors in Three-Dimensional Structures of Proteins. *Protein: Structure, Function and Genetic*. 17: 355-362.
- Shalaby, A.R. 1996. Significance of Biogenic Amines to Food Safety and Human Health. *Food Research International* 29(7): 675-690.
- Sharma, J., Ramanathan, K. and Sethumadhavan, R. 2011. Identification of Potential Inhibitors against Acetylcholinesterase Associated with Alzheimer's Diseases: A Molecular Docking Approach. *Journal Computational Methodology Molecular Design* 1: 44-51.
- Shukla, S., Park, H.K., Kim, J.K. and Kim, M. 2010. Determination of Biogenic Amines in Korean Traditional Fermented Soybean Paste (Doenjang). *Food and Chemical Toxicology* 48(5): 1191-1195.
- Smialowski, P., Martin-Galiano, A.J., Mikolajka, A., Girschick, T., Holak, T.A. and Frishman, D. 2007. Protein Solubility: Sequence Based Prediction and Experimental Verification. *Bioinformatics*. 23(19): 2536-2542.
- Smith, M.A., Pirrat, P., Pearson, A.R., Kurtis, C.R.P., Trinh, C.H., Gaule, T.G., Knowles, P.F., Phillips, S.E.V. and McPherson, M.J. 2010. Exploring the Roles of the Metal Ions in *Escherichia coli* Copper Amine Oxidase. *Biochemistry* 49(6):1268-1280.
- Smyth, D.R., Mrozkiewicz, M.K., McGrath, W.J., Listwan, P. and Kobe, B. 2003. Crystal Structures of Fusion Proteins with Large-affinity Tags. *Protein Science* 12(7): 1313-1322.
- Souers, A.J. and Ellman, J.A. 2001. B-Turn Mimetic Library Synthesis: Scaffolds and Application. *Tetrahedron* 57: 7431-7448.

- Sternberg, M.J.E. 1996. Protein structure prediction- principles and approaches. *In Protein Structure Prediction. Practical Approach*, Ed. Sternberg, M.J.E., pp. 1-30. Oxford: Oxford University Press.
- Stigers, K.D., Soth, M.J. and Nowick, S.J. 1999. Designed Molecules that Fold to Mimic Protein Secondary Structures. *Current Opinion in Chemical Biology* 3: 714-723.
- Suresh, K. 2011. Comparative Modeling and Molecular Docking of Orphan Human CYP4V2 Protein with Fatty Acid Substrates: Insights into Substrate Specificity. *Bioinformation*. 7(7): 360-365.
- Tanizawa, K., Matsuzaki, R., Shimizu, E., Yorifuji, T., and Fukui, T. (1994). Cloning and Sequencing of Phenylethylamine Oxidase from *Arthrobacter globiformis* and Implication of Tyr-382 as the Precursor to its Covalently Bound Quinone Cofactor. *Biochemical and Biophysical Research Communications* 199(3): 1096-1102.
- Tolia, N.H. and Joshua-Tor, L. 2006. *Strategies for coexpression in Escherichia coli*. *Nature Methods* 3: 55-64.
- van Gunsteren, W.F., Billeter, S.R., Eising, A.A., Hünenberger, P.H., Krüger, P., Mark, A.E., Scott, W.R.P. and Tironi, I.G. 1996. *Biomolecular Simulation: The GROMOS96 Manual and User Guide*, pp. 1-1042. Zürich, Switzerland: Vdf Hochschulverlag AG an der ETH Zürich.
- Wallner, B. And Elofsson, A. 2005. All are not Equal: A Benchmark of Different Homology Modeling Programs. *Protein Science* 14: 1315-1327.
- Wang, F., Xiao, J., Pan, L., Yang, M., Zhang, G., Jin, S., Yu, J. 2008. A Systematic Survey of Mini-Proteins in Bacteria and Archaea. *PLoS One* 3(12): e4027.
- Wiederstein, M. and Sippl, M.J. 2007. ProSA-web: Interactive Web Service for the Recognition of Errors in Three-Dimensional Structures of Proteins. *Nucleic Acids Research* 35: 407-410.
- Wilkinson, D.L. and Harrison, R.G. 1991. Predicting the Solubility of Recombinant Proteins in *Escherichia Coli*. *Biotechnology (New York)* 5: 443-448.
- Wimalasekera, R., Villar, C., Begum, T. and Scherer, G.F. 2011. COPPER AMINE OXIDASE1 (CuAO1) of *Arabidopsis Thaliana* Contributes to Abscisic Acid- and Polyamine-induced Nitric Oxide Biosynthesis and Abscisic Acid Signal Transduction. *Molecular Plant* 4(4): 663-678.

- Wong, E.T.C., Na, D. and Gsponer, J. 2013. On the Importance of Polar Interactions for Complexes Containing Intrinsically Disordered Proteins. *PLOS Computational Biology* 9(8): 1-14.
- Yılmaz, M., Ozic, C. and Gok, I. 2012. *Principles of Nucleic Acid Separation by Agarose Gel Electrophoresis, Gel Electrophoresis - Principles and Basics*, ed. M. Sameh, pp 34- 42. Turkey: InTech.
- Yeung, N., Lin, Y., Gao, Y., Zhao, X., Russell, B.S., Lei, L., Miner, K.D., Robinson, H. and Lu Y. 2009. Rational Design of a Structural and Functional Nitric Oxide Reductase. *Nature* 462: 24-31.
- Zhang, L. and Skolnick, J. 1998. What Should the Z-Score of Native Protein Structures Be? *Protein Science* 7: 1201-1207.
- Zhou, A.Q., O'Hern, C.S. and Regan, L. 2011. Revisiting the Ramachandran Plot from a New Angle. *Protein Science* 20: 1166-1171.
- Zoller, F., Haberkorn, U. and Mier, W. 2011. Miniproteins as Phage Display-Scaffolds for Clinical Applications. *Molecules* 16: 2467-2485.