



UNIVERSITI PUTRA MALAYSIA

***DEVELOPMENT AND CHARACTERIZATION OF SWIFTLET NEST-BASED
FORMULATION IN NANO-COSMECEUTICALS***

SITI HUSNAA BINTI MOHD TAIB

IPPH 2015 5



**DEVELOPMENT AND CHARACTERIZATION OF SWIFTLET NEST-BASED
FORMULATION IN NANO-COSMECEUTICALS**

By

SITI HUSNAA BINTI MOHD TAIB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science
July 2015**

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

DEVELOPMENT AND CHARACTERIZATION OF SWIFTLET NEST-BASED FORMULATION IN NANO-COSMECEUTICALS

By

SITI HUSNAA BINTI MOHD TAIB

July 2015

Chair : Siti Salwa Abd Gani, PhD
Faculty : Halal Products Research Institute

Purified swiftlet nest (SN) was used as an active ingredient in the development of nano-cosmeceuticals using homogenizer. The sample preparation method comprises cleaning, drying with freeze dryer and grinding of SN. Then the fine powder of SN was added to the formulation. Prior to addition of SN into the formulation, the compound present in raw SN and purified SN were identified and compared by Fourier Transform Infrared (FTIR) spectroscopy analysis, nitrite test, protein analysis and moisture analysis.

One of the major nutrients in SN is protein. The protein content in purified SN (60.1%) higher than raw SN (57.5%) which is differs significantly at 95% confidence level. This was due to "Freeze-Drying" method that was carried out by low operating temperature. This process leads to minimal damage of the heat-sensitive materials. Hence, protein is able to be preserved and enhanced. The study showed raw SN has low concentration of nitrite (0.25 ppm) whereas purified SN devoid of nitrite. Making frequent water changes during the process of cleaning SN was able to reduce nitrite as nitrite is highly soluble in water. The FTIR results indicated that the main compounds present in the SN samples were carbohydrates and protein. Moisture content of purified SN should be less than 10%. For this study, moisture content of the purified SN (7.45%) was slightly higher from its raw SN (5.62%).

The formulations consisted of oil in water emulsions. In manufacturing cosmetics, formulation and optimization processes are two important issues. In this work, Response surface methodology (RSM) was utilised in order to investigate the influence of the nano-cosmeceuticals composition; purified SN (1–5% w/w), and Tween 80 (3–6% w/w) as well as the preparation method; time of homogenization (10–30 min), on the physicochemical properties of swiftlet nest-based nano-cosmeceuticals. The response variables were particle size and zeta potential which are very important characteristic in nano-cosmeceuticals. Formulation and optimization of three independent variables were carried out to obtain an optimum SN-based nano-cosmeceutical with the lowest particle size and high stability formulation.

After the optimization process, formulation coded as SN25 was selected as the best formulation with the particle size of 136.35 nm and zeta potential of -40.2 mV. This was obtained experimentally and was closer to the predicted value 136.22 nm and -40.07 mV, respectively. Formulation SN25 was determined to be 2.58% SN, 3.99% Tween 80, 90.03% deionized water and 3.4% other ingredients, with a time of homogenization of 17 min. The pH value of optimized formulation was 6.49. The formulation was stable after undergoing thaw cycles test, at room temperature and 45°C for three months.

The rheological property of optimized formulation was investigated using rheometer. The viscosity of the optimized formulation was found to decrease with the increase in the applied shear rate. Thus, these types of formulation could be categorized as pseudoplastic or shear thinning material. The optimized formulation was found to be non-irritating with a Human Irritancy Equivalent (HIE) score below 0.90. Potassium sorbate used as preservative in the formulations successfully prevented microbial growth. The inhibition zone activities of formulation showed inhibitions bacteria *Methicillin Resistant Staphylococcus aureus* (MRSA) and yeast *Candida albicans*.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Sarjana Sains

PEMBANGUNAN DAN PENCIRIAN FORMULASI BERASASKAN SARANG BURUNG WALIT DALAM NANO-KOSMESEUTIKAL

Oleh

SITI HUSNAA BINTI MOHD TAIB

July 2015

Pengerusi : Siti Salwa Abd Gani, PhD
Fakulti : Institut Penyelidikan Produk Halal

Sarang burung walit (SN) yang tulen (sudah dibersihkan) telah digunakan sebagai bahan aktif dalam pembangunan nano-kosmeseutikal menggunakan mesin pengsehati. Kaedah penyediaan sampel terdiri daripada pembersihan, pengeringan dengan pengering sejuk beku dan pengisaran SN. Kemudian serbuk halus SN telah ditambah kepada formulasi. Sebelum penambahan SN ke dalam formulasi, sebatian yang hadir dalam SN mentah dan SN yang tulen (dibersihkan) telah dikenal pasti dan dibandingkan dengan analisis spektroskopi Inframerah Transformasi Fourier (FTIR), ujian nitrit, analisis protein dan analisis kelembapan.

Salah satu komponen nutrien utama dalam SN adalah protein. Kandungan protein dalam SN tulen (60.1%) lebih tinggi daripada SN mentah (57.5%) yang berbeza dengan ketara pada 95% tahap keyakinan. Ini adalah kerana pengeringan sejuk beku yang telah dijalankan dengan suhu operasi yang rendah. Proses ini membawa kepada kerosakan yang minima bagi bahan-bahan sensitif haba. Oleh itu, protein mampu dikekalkan. Kajian ini menunjukkan SN mentah mengandungi kepekatan nitrit yang rendah (0.25 ppm) manakala SN yang tulen tidak mengandungi nitrit. Pertukaran air dengan kerap semasa proses pembersihan SN boleh membantu mengurangkan nitrit kerana nitrit adalah sangat larut dalam air. Keputusan FTIR menunjukkan bahawa sebatian utama yang hadir dalam sampel SN adalah karbohidrat dan protein. Kandungan lembapan SN yang tulen mestilah kurang daripada 10%. Untuk kajian ini, kandungan lembapan SN yang dibersihkan (7.45%) adalah lebih tinggi sedikit daripada SN mentah (5.62%).

Formulasi terdiri daripada emulsi minyak dalam air. Dalam pembuatan kosmetik, proses formulasi dan pengoptimuman adalah dua perkara penting. Dalam kajian ini, Kaedah Permukaan Respons (RSM) telah digunakan untuk mengkaji kesan komposisi nano-kosmeseutikal; SN ditulen (1-5% w/w), dan Tween 80 (3-6% w/w) serta kaedah penyediaan; masa homogenisasi (10-30 min), ke atas sifat-sifat fizikokimia nano-kosmeseutikal berasaskan sarang burung walit. Pembolehubah tindak balas ialah saiz zarah dan potensi zeta (kestabilan) yang merupakan ciri yang sangat penting dalam nano-kosmeseutikal. Formulasi dan pengoptimuman tiga pembolehubah bebas telah dijalankan untuk mendapatkan tahap optimum nano-kosmeseutikal berasaskan sarang

burung walit dengan saiz zarah yang paling rendah dan kestabilan formulasi yang tinggi.

Selepas proses pengoptimuman, formulasi dikodkan sebagai SN25 telah dipilih sebagai formulasi terbaik dengan saiz zarah sebanyak 136.35 nm dan potensi zeta sebanyak -40.2 mV. Ini diperolehi secara eksperimen dan adalah hampir kepada nilai 136.22 nm dan -40.07 mV, masing-masing. Formulasi SN25 telah ditentukan sebagai 2.58% SN, 3.99% Tween 80, 90.03% air ternyahion dan 3.4% bahan-bahan lain, dengan masa homogenisasi 17 min. Nilai pH formulasi yang optimum adalah 6.49. Formulasi adalah stabil selepas menjalani ujian kitaran mencair, pada suhu bilik dan 45°C selama tiga bulan.

Ciri reologi formulasi yang optimum telah dikaji menggunakan reometer. Kelikatan formulasi yang optimum didapati menurun dengan peningkatan kadar ricih yang digunakan. Oleh itu, jenis formulasi boleh dikategorikan sebagai pseudoplastik atau bahan penipisan ricih. Formulasi yang optimum didapati tidak merengsa dengan nilai Padanan Kerengsaan Manusia (HIE) kurang daripada 0.90. Kalium sorbat yang telah digunakan sebagai pengawet dalam formulasi berjaya menghalang pertumbuhan mikrob. Aktiviti zon perencatan formulasi jelas menghalang bacteria *Methicillin Resistant Staphylococcus aureus* (MRSA) dan yis *Candida albicans*.

ACKNOWLEDGEMENTS

First and foremost, praise and glory to Allah SWT the compassionate and merciful for giving me a change and strength to complete this project successfully. I would like to express my appreciation and thanks to my supervisor Dr. Siti Salwa Abd Gani, you have been a tremendous mentor for me. I would like to thank you for encouraging my research and for allowing me to grow as a research scientist. Your advices on my research and career have been priceless. I would especially like to thank my committee members, Professor Dr. Mahiran Basri, Professor Dr. Amin Ismail and Associate Professor Dr. Mohamad Zaki Ab Rahman for serving as my committee members even at hardship. I also want to thank you for letting my defense be an enjoyable moment, and for your brilliant comments and suggestions, thanks to you.

A special thanks to my family. Words cannot express how grateful I am to my mother, Salmah Rosli and father, Mohd Taib Kunting as well as to my lovely siblings and grandfather for all of the sacrifices that you've made on my behalf. Your prayer for me was what sustained me thus far. Special thanks goes to my friends, Suhaili, Farrah, Izzati, Kak Hasmah, Kak Nana, Kak Jaini, Syafinaz, Kak Ili, Alin, Nadirah and Hidayah who help and supported me in writing, gave suggestion and incited me to strive towards my goal.

Last but not least, I would like to acknowledge the financial support in this study from the Research university grant scheme (RUGS), Universiti Putra Malaysia (UPM) for project number 9361000, Graduate Research Fellowship (GRF) scholarship under UPM and MyBrain15 (MyMaster) under Ministry of Higher Education (MOHE).

To all thank you for being there for me. Only the AlMighty Allah SWT could pay back all your kindness to me.

I certify that a Thesis Examination Committee has met on 8 July 2015 to conduct the final examination of Siti Husnaa Mohd Taib on her thesis entitled "Development and Characterization of Swiftlet Nest-based Formulation in Nano-cosmeceuticals" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Shuhaimi Mustafa, PhD

Professor
Halal Products Research Institute
Universiti Putra Malaysia
(Chairman)

Mohd Zaizi Desa, PhD

Lecturer
Centre of Foundation Studies for Agricultural Science
Universiti Putra Malaysia
(Internal Examiner)

Roswanira Ab. Wahab, PhD

Lecturer
Faculty of Science
Universiti Teknologi Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 August 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Siti Salwa Abd Gani, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Amin Ismail, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Mahiran Basri, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Mohamad Zaki Ab Rahman, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Siti Husnaa Binti Mohd Taib (GS30424)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman of
Supervisory
Committee: Dr. Siti Salwa Abd Gani

Signature: _____
Name of Member of
Supervisory
Committee: Prof. Dr. Amin Ismail

Signature: _____
Name of Member of
Supervisory
Committee: Prof. Dr. Mahiran Basri

Signature: _____
Name of Member of
Supervisory
Committee: Assoc. Prof. Dr. Mohamad Zaki Ab Rahman

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xviii
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS	xx
CHAPTER	
1	
INTRODUCTION	1
Objectives	2
2	
LITERATURE REVIEW	3
2.1 Swiftlet Nest	3
2.1.1 Composition in Swiftlet Nest	3
2.1.2 Importance of Swiftlet Nest	5
2.1.3 Issues Regarding Nitrite Content in Swiftlet Nest	6
2.2 The Concept of Halal and Toyyiban in Cosmetics	7
2.3 Halal and Toyyiban Issues Related to the Consumption of Swiftlet Nest	8
2.4 The Halal Cosmetics Market and its Opportunities	9
2.5 Human Skin	10
2.5.1 Skin Structure and Function	10
2.6 Nano-cosmeceuticals	11
2.6.1 Cosmeceuticals	11
2.6.2 Emulsion	11
2.6.3 Nano-emulsion	13
2.6.4 Methods to Produce Nano-emulsion	13
2.6.5 Characterization of Nano-emulsion	15
2.6.6 Stability of Nano-emulsion	17
2.7 Response Surface Methodology (RSM)	19
3	
MATERIALS AND METHODS	21
3.1 Materials	21
3.2 Purification of swiftlet nest	21
3.3 Identification of swiftlet nest	21
3.3.1 Nitrite Test	21
3.3.2 Fourier Transform Infrared (FTIR) Spectroscopy Analysis	21
3.3.3 Protein Analysis	22
3.3.4 Moisture Analysis	22
3.4 Formulation of Nano-cosmeceuticals Containing Swiftlet Nest	23
3.5 Optimization of Swiftlet Nest-Based Nano-cosmeceuticals Using Response Surface Methodology (RSM)	23
3.5.1 Experimental Design	23
3.5.2 Statistical Analysis	24

3.6	Physico-chemical Analysis of Nano-cosmeceuticals	25
3.6.1	Particle Size	25
3.6.2	Zeta Potential	25
3.6.3	pH Measurement	25
3.6.4	Stability Study	25
3.6.5	Rheology Study	26
3.7	Safety Assessment	26
3.7.1	Antimicrobial/Antifungal Test	26
3.7.2	In-vitro Irritancy Test	27
4	RESULTS AND DISCUSSION	28
4.1	Identification of Swiftlet Nest	28
4.1.1	Nitrite Test	28
4.1.2	Fourier Transform Infrared (FTIR) Spectroscopy Analysis	29
4.1.3	Protein Content	30
4.1.4	Moisture Content	31
4.2	Selection of Ingredient in Nano-cosmeceuticals Formulation	31
4.3	Response Surface Methodology	32
4.3.1	Fitting the Response Surface Models (RSM)	32
4.3.2	Response Surface Analysis	34
4.3.3	Optimization of Swiftlet Nest-Based Nano- cosmeceuticals	37
4.4	Physico-chemical Analysis of Nano-cosmeceuticals	38
4.4.1	Particle Size	38
4.4.2	Zeta Potential	38
4.4.3	pH Measurement	38
4.4.4	Stability Study	38
4.4.5	Rheology Properties	39
4.5	Safety Test	40
4.5.1	Antimicrobial/Antifungal Test	40
4.5.2	Irritancy Test (In-Vitro Dermal Irritection Test)	42
5	CONCLUSIONS	44
	REFERENCES	45
	APPENDICES	56
	BIODATA OF STUDENT	67
	LIST OF PUBLICATIONS	68

LIST OF TABLES

Tables		Page
1	Composition of the SN	3
2	Stability characteristics based on zeta potential	16
3	The central composite design (CCD) matrix	24
4	Relationship of human equivalent (HIE) score to irritancy classification for the dermal irritation test method	27
5	Infrared data for raw SN and purified SN	30
6	The experimental data obtained for the two responses	32
7	Regression coefficients, adjusted and probability values for the particle size and zeta potential for nano-cosmeceuticals formulation	33
8	Analysis of variance (ANOVA) of regression coefficient of the fitted quadratic equation by using F -value and p -Value in the final model	34
9	The response values predicted from RSM for the optimized nano-cosmeceuticals	37
10	Inhibition zone of samples against MRSA and <i>Candida albicans</i> 9002	42
11	Irritancy test of formulation SN25	43

LIST OF FIGURES

Figure		Page
1	Illustration of Human Skin Layer	10
2	Schematic Illustration of a Dispersed System that Consists of Two Fully or Partially Immiscible Liquids.	12
3	Electric Double Layer Surrounding Nano-particle	16
4	Representative of Nitrite Content Results in (a) Raw SN and (b) Purified SN	28
5	Infrared Spectrum of (a) Raw SN and (b) Purified SN	29
6	Representative of Protein Content Results in Raw and Purified SN	31
7	Response Surface Plots Presenting the Interaction Effects of Time of Homogenization and Tween 80 Amount on Particle Size	35
8	Response Surface Plots Presenting the Interaction Effects SN Amount and Time of Homogenization on Particle Size	35
9	Response Surface Plots Presenting the Interaction Effects of Time of Homogenization and SN Amount on Zeta Potential	36
10	Response Surface Plots Presenting the Interaction Effects of SN Amount and Tween 80 Amount on Zeta Potential	37
11	Effect of Shear Rate on the Viscosity of the Optimized Formulation	39
12	Effect of Shear Rate of Optimized Formulation on the Shear Stress	40
13	Antimicrobial Test for Microbe (a) MRSA and (b) Candida albicans 9002	41

LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
CCD	Central Composite Design
CDC	Centers for Disease Control and Prevention, USA
cm	centimetre
FTIR	Fourier Transform Infrared
HIE	Human irritancy equivalent
ICD	irritant contact dermatitis
min	minute
mL	milliliter
MRSA	Methicillin Resistant Staphylococcus Aureus
nm	nano-meter
mV	millivolt
OECD	Organisation for Economic Cooperation and Development
o/w	Oil in water
Pa	Pascal
Pa.s	Pascal second
PIT	Phase Inversion Temperature
PPO	1,2-dipalmitoyl-3-oleoyl-rac-glycerol
PRE	formulation with addition of preservative
rpm	revolution per minute
RSM	Response Surface Methodology
SA	Sialic acid
SN	Swiftlet Nest
SN25	optimized formulation
Tween 80	Polyoxyethylene (20) sorbitan mono-oleate
w/o	Water in oil
w/w	weight/weight

CHAPTER 1

INTRODUCTION

The term cosmetic, which is used on a daily basis, only places emphasis on outward appearance. Currently, medical cosmetic products which are termed dermaceuticals or cosmeceuticals are in high demand. Furthermore, cosmeceuticals are committed to enhancing and maintaining the long-term function of the skin. Nano-cosmeceuticals are the strongest generation of cosmetic products which use nano-sized systems for the delivery of active ingredients to the skin cells for better penetration. They are subjected to particular manufacturing techniques and contain lower concentrations of surfactant as compared to classical skin care cosmetics (Harwansh *et al.*, 2011).

Recently, research about flora and fauna has been undertaken which might prove beneficial to the human body. Some of them are being used as active ingredients in new cosmetics. One of the fauna that was found to have a good value is swiftlet nest (SN). Several species of swiftlet produce edible nests that are consumed by humans worldwide, as a delicacy or as a medicinal food. There have been a number of studies conducted on the advantages of SN in cosmetics, food and medicine (Set, 2012; Norhayati *et al.*, 2010). The main component of the SN is glycoprotein. The existence of glycoprotein is capable of promoting cell division, and it has demonstrated the presence of an epidermal growth factor-like protein (Kong *et al.*, 1987; Ng *et al.*, 1986). SN has good effect on the skin and becomes attractions to ladies due to its properties of making the skin delicate and grows radiant. Besides, some people believe that a pregnant woman who consumes SN would have a baby with fine and smooth complexion. Although the importance of the SN has provided a high potential in new nano-cosmeceuticals area, research in this area is still limited as it is commonly consumed, not being applied to the skin. Hence, encourage us to develop new formulation of SN-based nano-cosmeceuticals with smaller molecules which are formulated to penetrate deep into the skin.

Formulation and optimization processes are two important issues in the manufacturing of cosmetics. The process of optimizing parameters not only increases the utility of the technologist, but also the quality of the product as well. Since, the lack of any model to optimize the use of the SN in nano-cosmeceuticals formulation which led this study to find an optimal formulation of SN-based nano-cosmeceuticals with nano particle size using response surface methodology (RSM). The RSM appears to be an effective and simple technique of analyzing, improving and optimizing of structures over practicable domain of parameter settings (Shokuhfar *et al.*, 2008). RSM is the most popular optimization method used in recent years (Chellamboli and Perumalsamy, 2014; Ryad *et al.*, 2010; Baş and Boyacı, 2007). Several studies have been investigated based on the application of the RSM in industrial processes. This method is a set of mathematical and statistical techniques in which a response of interest is influenced by several variables and the aim is to optimize this response by determining the correlation between the response and independent variables (Raissi and Farsani, 2009).

The benefits of SN in nano-cosmeceuticals were investigated. The use of local natural resources in nano-cosmeceuticals can be established. Furthermore, the usage of SN as nano-cosmeceuticals can be developed where it can open up new areas of nano-biotechnology exploitation, which leads to the necessity to extract and process this SN.

General objective of this study is to develop new formulation of nano-cosmeceuticals containing SN using RSM. The specific objectives are:

- 1 To evaluate the characteristics of SN by nitrite test, Fourier Transform-Infrared (FTIR), protein analysis and moisture analysis.
- 2 To develop new formulation and optimize the formulation of SN-based nano-cosmeceuticals using RSM.
- 3 To characterize the properties of formulation with respect to particle size, zeta potential, pH measurement, stability study and rheology study.
- 4 To assess the safety such as antimicrobial and irritancy test of the optimized formulation.

REFERENCES

- Abd Gani, S. S. 2010. Formation and Characteristics of Engkabang-Based Nanocosmeceuticals. PhD Thesis, Universiti Putra Malaysia.
- Ahmad Robin, W. 2004. Guidelines for the Preparation of Halal Food and Goods for the Muslim Consumers. pp 1–12.
- Akoh, C.; Min, D. 2002. In *Food lipids: chemistry, nutrition, and biochemistry*. New York: Marcel Dekker, Second Edition.
- Alserhan, B.A. 2010. On Islamic Branding: brands as good deeds. *J. Islam Marketing*, 1(2), pp 101-106.
- Anton, N.; Benoit, J.-P.; Saulnier, P. 2008. Design and Production of Nanoparticles Formulated from Nano-Emulsion Templates-a Review. *J. Control. release*, 128, 185–199.
- Arens, E.; Zhang, H. 2006. The Skin's Role in Human Thermoregulation and Comfort. In *Thermal and Moisture Transport in Fibrous Materials*; N. Pan and P. Gibson, eds.; Woodhead Publishing Ltd, pp 560-602.
- Armijo, R.; Orellana, M.; Mediona, E.; Coulson, A. H.; Sayre, J. W.; R., D. 1981. Epidemiology of Gastric Cancer in Chile: I. Case-Control Study. *Int. J. Epidemiol.*, 10, 53–56.
- Arosio, P. 2011. *Stability and Aggregation Kinetics of Colloidal Systems: Application to Polymer Colloids, Proteins and Peptides*, PhD Thesis, ETH Zurich.
- Association of Analytical Communities (AOAC). Official Methods of Analysis. 17th ed. Gaithersburg, Maryland, USA, AOAC International.
- Aubrun, O. S.; Simonnet, J. T., L'Alloret, F. 2004. Nanoemulsions: a new vehicle for skincare products. *Adv. Colloid Interface Sci.*, 108-109, 145-149.
- Azeem, A.; Khan, Z. I.; Aqil, M.; Ahmad, F. J.; Khar, R. K.; Talegaonkar, S. 2009. Microemulsions as a Surrogate Carrier for Dermal Drug Delivery. *Drug Dev. Ind. Pharm.*, 35(5):525-47.
- Baş, D.; Boyacı, İ. H. 2007. Modeling and Optimization I: Usability of Response Surface Methodology. *J. Food Eng.*, 78, 836–845.
- Bazaraa, W. A.; Hassan, E. E. 1996. Response Surface Optimization for the Continuous Glucose Isomerization Process. *J. Ind. Microbiol.*, 17, 100–103.

- Beresford, S. A. A. 1985. Is Nitrate in the Drinking Water Associated with the Risk of Cancer in the Urban UK. *Int. J. Epidemiol.*, 14, 57–63.
- Bfs magazine, Business For Sale Online Magazine. 2012. An Introduction to Malaysia's Cosmetic Industry. Retrieved 6 June 2014 from http://www.bfsmagazine.com.my/index.php?option=com_content&view=article&id=1546:an-introduction-to-malaysias-cosmetic-industry&catid=228:articles&Itemid=1524.
- Bonney, G. E.; Elston, R. C.; Correa, P.; Tannenbaum, S. R. Haenszel, W.; Zavala, D. E.; Fontham, E.; Zarama, G.; Gordillo, G.; Cuello, C. 1987. Genetic Etiology of Gastric Cancer: II. Segregation Analysis of Gastric pH, Nitrate and Nitrite. *Genet. Epidemiol.*, 4, 103–114.
- CDC 24/7. 2014b. Methicillin Resistant Staphylococcus Aureus (MRSA) Infections. Retrieved 26 September 2014 from <http://www.cdc.gov/mrsa/>.
- Cebrián, J.; Messeguer, A.; Facino R. M.; Garcia Antón J. M. 2005. New Anti-RNS and -RCS Products for Cosmetic Treatment. *Int. J. Cosmet. Sci.*, 27, 271-278.
- Centers for Disease Control and Prevention (CDC), USA. CDC 24/7. 2014a. Candidiasis. Retrieved 26 September 2014 from <http://www.cdc.gov/fungal/diseases/candidiasis/index.html>.
- Chan, S. W. 2006. Review of Scientific Research on Edible Bird's Nest. Retrieved 6 October 2014 from <http://www.hkfsta.com.hk/articles/special/article7.htm>.
- Chellamboli, C.; Perumalsamy, M. 2014. Application of Response Surface Methodology for Optimization of Growth and Lipids in Scenedesmus Abundans using Batch Culture System. *RSC Adv.*, 4, 22129–22140.
- Chiesa, M.; Garg, J.; Kang, Y.T.; Chen, G. 2008. Thermal Conductivity and Viscosity of Water-in-Oil Nanoemulsions. *Colloids Surf., A*, 326, 1-2, 67-72.
- Colombo, J. P.; Garcia-Rodenas, C.; Guesry, P. R.; Rey, J. 2003. Potential Effects of Supplementation with Amino Acids, Choline or Sialic Acid on Cognitive Development in Young Infants. *Acta Paediatr. Suppl.*, 92, 42–46.
- Colombo, A. L.; Guimarães, T.; Luis Fernando Aranha Camargo, L. F. A.; Richtmann, R.; de Queiroz-Telles, F.; Sallese, M. J. C.; da Cunha, C. A.; Yasudag, M. A. S.; Moretti, M. L.; Nuccii, M. 2013. Brazilian Guidelines for the Management of Candidiasis – A Joint Meeting Report of Three Medical Societies: Sociedade Brasileira de Infectologia, Sociedade Paulista de Infectologia and Sociedade Brasileira de Medicina Tropical. *Braz. J. Infect. Dis.*, 17(3), 283–312.

- Crowley, M. M. 2013. Solutions, Emulsions, Suspensions and Extracts. In *Essentials of Pharmeceutics*; Felton, L., Ed.; Pharmaceutical Press, United Kingdom, pp 435-460.
- Dickinson, E.; McClements, D. J. 1995. *Advances in Food Colloids*, Chapman & Hall, London, U.K.
- DSM, Department of Standards Malaysia. 2008. MS 2200:2008 Islamic Consumer Goods – Part 1 Cosmetic and Personal Care – general Guidelines, Ministry of Science and Innovation (MOSTI), Malaysia
- Department of Standards Malaysia (DSM). 2011. MS 2334:2001 – Edible Bird Nest (EBN) Spesification.
- Eastman. 1998. Sorbic Acid and Potassium Sorbate as Cosmetic Preservatives. pp. 2–16.
- Elibol, M. 2004. Optimization of Medium Composition for Actinorhodin Production by *Streptomyces Coelicolor* A3 (2) with Response Surface Methodology. *J. Proc. Biochem.*, 39, 1057–1062.
- Epstein, H.; Simion, F. A. 2001. Emulsion-based skincare products: Formulating and measuring their moisturizing benefit. In *Handbook of Cosmetic Science and Technology*; Barel, A.O., Paye, M., Maibach, H.I., Eds.; Marcel Decker, Inc. 270 Madison Avenue, New York, pp 511-529.
- Evison, J.; Dickinson, E.; Owusu-Apenten, R. K.; Williams, A. 1995. Food Macromolecules and Colloids. In Lorient, D. E.; Lorient, D., Eds., Royal Society of Chemistry, Cambridge, U.K., pp 235–243.
- Ezzine, M.; Ghorbel, M. H. 2006. Physiological and Biochemical Responses Resulting from Nitrite Accumulation in Tomato (*Lycopersicon Esculentum* Mill. Cv. Ibiza F1). *J. Plant Physiol.*, 163, 1032–1039.
- Farage, M. A.; Miller, K. W.; Elsner, P.; Maibach, H. I. 2008. Intrinsic and Extrinsic Factors in Skin Ageing: A Review. *Int J Cosmet Sci*, 30, 87–95.
- Gardouh, A.; Ghorab, M.; Abdel-Rahman, S. 2011. Effect of Viscosity, Method of Preparation and Homogenization Speed on Physical Characteristics of Solid Lipid Nanoparticles., 2, 996–1006.
- Goh, D.L.M.; Chua, K.Y.; Chew, F.T.; Seow, T.K.; Ou, K.L.; Yi, F.C.; Lee, B.W. 2001. Immunochemical Characterization of Edible Bird's Nest Allergens. *J. Allergy Clin. Immunol.*, 107(6), 1082–1088.
- Gunasekaran, S.; Mehmet Ak, M. 2003. *Cheese Rheology and Texture*; CRC Press LLC: Boca Raton, Florida, pp 31–112.

- Gupta, P. K.; Pandit, J. K.; Kumar, A., Swaroop, P.; Gupta, S. 2010. Pharmaceutical Nanotechnology Novel Nanoemulsion–high Energy Emulsification Preparation, Evaluation and Application. *T. Ph. Res.*, 3, 117-138.
- Haritha; Basha Syed, P.; Rao P, K.; Vedantham, C. 2013. A Brief Introduction to Methods of Preparation, Applications and Characterization of Nanoemulsion Drug Delivery Systems. *Indian J. Res. Pharm. Biotechnol.*, 1, 25–28.
- Harris, P. L.; Cuppett, S. L.; Bullerman, L. B. 1990. Optimization of Lipase Synthesis by *Pseudomonas Fluorescens* by Response Surface Methodology. *J. Food Prot.*, 53, 481–483.
- Harwansh, R. K.; Patra, K. Ch.; Pareta, S. K. 2011. Nanoemulsion as Potential Vehicles for Transdermal Delivery of Pure Phytopharmaceuticals and Poorly Soluble Drug. *Int. J. Drug Deliv.*, 3, 209-218.
- Hicks, R. M. 1982. Nitrosamines as Possible Etiological Agents in Bilharzial Bladder Cancer. pp 455–472.
- Hsu, J.; Arcot, J.; Alice Lee, N. 2009. Nitrate and Nitrite Quantification from Cured Meat and Vegetables and Their Estimated Dietary Intake in Australians. *Food Chem.*, 115, 334–339.
- Hsu, J.-P.; Nacu, A. 2003. Behavior of Soybean Oil-in-Water Emulsion Stabilized by Nonionic Surfactant. *J. Colloid Interface Sci.*, 259, 374–381.
- Human Skin Layers. Retrieved 3 November 2014 from <http://pixgood.com/human-skin-layers.html>.
- JAKIM, Jabatan Kemajuan Islam Malaysia. 2014. Hukum Memakan Sarang Burung Layang-layang. Retrieved 7 September 2014 from <http://www.e-fatwa.gov.my/fatwa-kebangsaan/hukum-memakan-sarang-burung-layang-layang>.
- Joglekar, A.; May, A. 1987. Product Excellence through Design of Experiments. *Cereal Foods World*, 32, 857–868.
- Jovanović, N.; Bouchard, A.; Hofland, G. W.; Witkamp, G.-J.; Crommelin, D. J. a; Jiskoot, W. 2006. Distinct Effects of Sucrose and Trehalose on Protein Stability during Supercritical Fluid Drying and Freeze-Drying. *Eur. J. Pharm. Sci.*, 27, 336–345.
- Juhasz, L.; Hill, M. J.; Nagy, G. 1980. Possible Relationship between Nitrate in Drinking Water and the Incidence of Stomach Cancer. In *N-Nitroso Compounds: Analysis, Formation and Occurrence.*; Walker, E. A.; Gastegnaro, M.; Griciute, L.; Borz, M., Eds.; International Agency for Research on Cancer (Lyon), pp 619–623.

- Kabalnov, A.S.; Pertzov, A.V.; Shchukin, E.D. 1987. Ostwald Ripening in Emulsions: I. Direct Observations of Ostwald Ripening in Emulsions. *J. Colloid Interface Sci.*, 118, 2, 590-597.
- Kathan, R. H.; Weeks, D. I. 1969. Structure Studies of Collocalia Mucoid I. Carbohydrate and Amino Acid Composition. *Arch. Biochem. Biophys.*, 134, 572–576.
- Keng, P. S.; Basri, M.; Ariff, A. B.; Abdul Rahman, M. B.; Abdul Rahman, R. N. Z.; Salleh, A. B. 2008. Scale-up Synthesis of Lipase-Catalyzed Palm Esters in Stirred-Tank Reactor. *Bioresour. Technol.*, 99, 6097–6104.
- Kerr, S. 2013. Halal Cosmetics Allow Muslim Women to Enjoy Beauty and Duty. Retrieved 7 June 2103 from <http://halalfocus.net/halal-cosmetics-allow-muslim-women-to-enjoy-beauty-and-duty/>.
- Kiran, K. R.; Manohar, B.; Karanth, N. G.; Divakar, S. 2000. Response Surface Methodological Study of Esterification of Lactic Acid with Palmitic Acid Catalysed by Immobilised Lipases from *Mucor Miehei* and Porcine Pancreas. *Eur. Food Res. Technol.*, 211, 130–135.
- Kong, Y. C.; Keung, W. M.; Yip, T. T.; Ko, K. M.; Tsao, S. W.; Ng, M. H. 1987. Evidence That Epidermal Growth Factor Is Present in Swiftlet's (Collocalia) Nest. *Comp. Biochem. Physiol. B.*, 87, 221–226.
- Kornfeld, R.; Kornfeld, S. 1980. Structure of Glycoproteins and Their Oligosaccharide Units. In *The Biochemistry of Glycoproteins and Proteoglycans*; J. Lennarz, W., Ed.; Springer US, pp 1–34.
- Kumar, D.; Jain, V. K.; Shanker, G.; Srivastava, A. 2003. Citric Acid Production by Solid State Fermentation Using Sugarcane Bagasse. *J. Process Biochem.*, 38, 1731–1738.
- Ligade, V.; Sreedhar, D.; Ajay, M.; Udupa, N. 2007. Nanotechnology in Cosmeceuticals: Benefits vs Risks. *Curr. Sci.*, 93, 597.
- Ma, F.; Liu, D. 2012. Sketch of the Edible Bird's Nest and Its Important Bioactivities. *Food Res. Int.*, 48, 559–567.
- Mannan, S.; Fakhru'l-Razi, A.; Alam, M. Z. 2007. Optimization of Process Parameters for the Bioconversion of Activated Sludge by *Penicillium Corylophilum*, Using Response Surface Methodology. *J. Environ. Sci.*, 19, 23–28.
- Marcone, M. F. 2005. Characterization of the Edible Bird's Nest the “Caviar of the East”. *J. Food Sci.*, 38, 1125–1134.
- Martínez-Pla, J. J.; Martín-Biosca, Y.; Sagrado, S.; Villanueva-Camañas, R. M.; Medina-Hernández, M. J. 2004. Evaluation of the pH Effect of Formulations on

- the Skin Permeability of Drugs by Biopartitioning Micellar Chromatography. *J. Chromatogr. A*, 1047, 255–262.
- Mason, T.; Wilking, J.; Meleson, K.; Chang, C.; Graves, S. 2006. Nanoemulsions: formation, structure, and physical properties. *J. Phys.: Condens. Matter*, 18, R635-R666.
- Massey, R. C. 1996. Methods for the Analysis of Nitrate and Nitrite in Food and Water. In *Nitrates and Nitrites in Food and Water*; Hill, M. J., Ed.; Woodhead Publishing: Cambridge, England, pp 13–32.
- Massey, R. C.; Key, P. E.; Mallett, A. K.; Rowland, I. R. 1988. An Investigation of the Endogenous Formation of Apparent Total N-Nitroso Compounds in Conventional Microflora and Germ-Free Rats. *Food Chem. Toxicol*, 26, 595–600.
- Masuda, M. 2001. Legislation in Japan. In *Handbook of Cosmetic Science and Technology*; Barel, A.O., Paye, M., Maibach, H.I., Eds.; Marcel Decker, Inc. 270 Madison Avenue, New York, pp 761-767.
- McClements, D.J. 2005. Food Emulsions: Principles, Practice, and Techniques. Second Edition Boca Raton, FL: CRC Press.
- McClements, D. J.; Weiss, J. 2005. Lipid Emulsions. In *Bailey's Industrial Oil and Fat Products*; Shahidi, F., Ed.; John Wiley & Sons, Inc., pp 457–502.
- McClements, D.J. 2010. Emulsion Design to Improve the Delivery of Functional Lipophilic Components. *Annu. Rev. Food Sci. Technol.*, 1, 241-69.
- Mei, Z.; Liu, S.; Wang, L.; Jiang, J.; Xu, J.; Sun, D. 2011. Preparation of Positively Charged Oil/water Nano-Emulsions with a Sub-PIT Method. *J. Colloid Interface Sci.*, 361, 565–572.
- Meinders, M.; Vliet, T.V. 2004. The Role of Interfacial Rheological Properties on Ostwald Ripening in Emulsions. *Adv. Colloid Interface Sci.*, 108-109, 119-126.
- MOH, Ministry of Health, Malaysia. 2012. *Standard Operating Procedure on the Control of Nitrite Level in Edible Bird's Nest*.
- Mohd Nazri, N. A. A.; Ahmat, N.; Abdullah, M.; Jaafar Sidik, N.; Tuan Johari, S. A. T. 2012. Antioxidant, Antimicrobial and Cytotoxic Activities of Resveratrol Oligomers of Shorea Macroptera Dyer. *Aust. J. Basic Appl. Sci.*, 6, 431–436.
- Motwani, S. K.; Chopra, S.; Talegaonkar, S.; Kohli, K.; Ahmad, F. J.; Khar, R. K. 2008. Chitosan-Sodium Alginate Nanoparticles as Submicroscopic Reservoirs for Ocular Delivery: Formulation, Optimisation and in Vitro Characterisation. *Eur. J. Pharm. Biopharm.*, 68, 513–525.

- Nanocomposix.com. 2012. NanoComposix Guidelines for Zeta Potential Analysis of Nanoparticles; San Diego, pp 1–6.
- Ng, M. H.; Chan, K. H.; Kong, Y. C. 1986. Potentiation of Mitogenic Response by Extracts of the Swiftlet's (Collocalia) Nest. *Biochem. Int.*, *13*, 521–531.
- Norhayati, M.; Azman, O.; Wan Nazaimoon, W. 2010. Preliminary Study of the Nutritional Content of Malaysian Edible Bird ' S Nest. *Mal. J. Nutr.*, *16*, 389–396.
- Ohshima, H.; Bartsch, H. 1981. Quantitative Estimation of Endogenous Nitrosation in Humans by Monitoring N-Nitrosoproline Excreted in the Urine. *Cancer Res.*, *41*, 3658–3662.
- Organisation for Economic Co-operation and Development (OECD). 2002. OECD 404: Guidelines for Testing of Chemicals, Acute Dermal Irritation/ Corrosion. Updated version, Adopted: 24 April 2002. Retrieved 1 September 2014 from <https://law.resource.org/pub/us/cfr/ibr/005/oecd.404.2002.pdf>.
- Pal, R. 2003. Viscous Behavior of Concentrated Emulsions of Two Immiscible Newtonian Fluid with Interfacial Tension. *J. Colloid Interface Sci.*, *263*, 296–305.
- Peng, L.; Yu-Long, Y.; Defa, L.; Sung Woo, K., Guoyao, W. 2007. Amino Acids and Immune Function. *Br. J. Nutr.*, *98*, 237–252.
- Ponto, T. 2010. Rejuvenating Potentials of Bird Nest in Aqueous Base for Cosmeceutical Application, Universiti Teknologi Mara.
- Puziah, H.; Shahab, N.; Masilamani, T.; Rozita, B.; Rohaidah, I. 2009. A Cosmetic Analysis in Compliance with the Legislative Requirements, Halal and Quality Control. *Malaysian J. Chem.*, *11*, 81–87.
- Qian, C.; McClements, D. J. 2011. Formation of Nanoemulsions Stabilized by Model Food-Grade Emulsifiers Using High-Pressure Homogenization: Factors Affecting Particle Size. *Food Hydrocoll.*, *25*, 1000–1008.
- Raissi, S.; Farsani, R. 2009. Statistical Process Optimization Through Multi-Response Surface Methodology. *World Acad. Sci. Eng. Technol.*, *3*, 251–255.
- Rao, P.; Jayaraman, K.; Lakshmanan, C. 1993. Production of Lipase by *Candida Rugosa* in Solid State Fermentation: 2. Medium Optimization and Effect of Aeration. *J. Proc Biochem*, *28*, 391–395.
- Rincón, F.; Martínez, B.; Delgado, J. M. 2003. Detection of Factors Influencing Nitrite Determination in Meat. *Meat Sci.*, *65*, 1421–1427.

- Rodrigues, R. C. L. B.; Felipe, M. G. A.; Silva, J. B. A.; Vitobol, M. 2003. Response Surface Methodology for Xylitol Production from Sugarcane Bagasse Hemicellulosic Hydrolyzate Using Controlled Vacuum Evaporation Process Variables. *J. Process Biochem.*, 38, 1231–1237.
- Rowland, I. R.; Granli, T.; Bøckman, O. C.; Key, P. E.; Massey, R. C. 1991. Endogenous N-Nitrosation in Man Assessed by Measurement of Apparent Total N-Nitroso Compounds in Faeces. *Carcinogenesis*, 12, 1395–1401.
- Ryad, A.; Lakhdar, K.; Majda, K.-S.; Samia, A.; Mark, A.; Corinne, A.-D.; Eric, G. 2010. Optimization of the Culture Medium Composition to Improve the Production of Hyoscyamine in Elicited *Datura Stramonium* L. Hairy Roots Using the Response Surface Methodology (RSM). *Int. J. Mol. Sci.*, 11, 4726–4740.
- Sagalowicz, L.; Leser, M.E. 2010. Delivery Systems for Liquid Food Products. *Curr. Opin. Colloid Interface Sci.*, 15, 61-72.
- Schramm, L.L. 2005. *Emulsions, Foams, and Suspensions*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, FRG, pp 4–399.
- Set, J. 2012. *Fast, Effective Evaluation of Edible Bird Nests Using the Handheld Agilent 4100 ExoScan FTIR. Application Note.*
- Shinoda, K.; Saito, H. 1968. The Effect of Temperature on the Phase Equilibria and the Types of Dispersions of the Ternary System Composed of Water, Cyclohexane, and Nonionic Surfactant. *J. Colloid Interface Sci.*, 26, 70–74.
- Shinoda, K.; Saito, H. 1969. The Stability of O/W Type Emulsions as Functions of Temperature and the HLB of Emulsifiers: The Emulsification by PIT-Method. *J. Colloid Interface Sci.*, 30, 258–263.
- Shokuhfar, A.; Khalili, S.M.R.; Ashenai Ghasemi, F.; Malekzadeh, K.; Raissi, S. 2008. Analysis and Optimization of Smart Hybrid Composite Plates Subjected to Low-velocity Impact using the Response Surface Methodology (RSM). *Thin Wall Struct.*, 46, 1204–1212.
- Siong, T.B. 2010. Edible Birdnest Cleaning Process. Swiftlets Nests Research. Retrieved 30 July 2014 from <http://swifletresearch.blogspot.com/2010/08/edible-birdnest-cleaning-process.html>
- Siong, T.B. 2009. What is 100% Pure Bird's Nest? Retrieved 27 October 2014 from <http://yongkangbirdnest.blogspot.com/2009/08/what-is-100-pure-birds-nest.html>
- Solans, C.; Izquierdo, P.; Nolla, J.; Azemar, N.; Garcíacelma, M. 2005. Nano-Emulsions. *Curr. Opin. Colloid Interface Sci.*, 10, 102–110.

- Solè, I.; Maestro, A.; Pey, C. M.; González, C.; Solans, C.; Gutiérrez, J. M. 2006. Nano-Emulsions Preparation by Low Energy Methods in an Ionic Surfactant System. *Colloids Surfaces A.*, 288, 138–143.
- Soo, E.; Salleh, A.; Basri, M.; Rahman, R.; Kamaruddin, K. 2004. Response Surface Methodological Study on Lipase-Catalyzed Synthesis of Amino Acid Surfactants. *J. Process Biochem.*, 39, 1511–1518.
- Tadros, T.; Izquierdo, P.; Esquena, J.; Solans, C. 2004. Formation and stability of nano-emulsions. *Adv. Colloid Interface Sci.*, 108-10, 303-318.
- Tadros, T. F. 2005. Microemulsions. In *Applied Surfactants: Principles and Applications*; Wiley-VCH Verlag GmbH & Co.: Weinheim.
- Talegaonkar, S.; Tariq, M.; Alabood, R. M. 2011. Design and Development of O/W Nanoemulsion for the Transdermal Delivery of Ondansetron. *Bull. Pharm. Res.*, 1(3), 18-30.
- Tang, S. Y.; Manickam, S.; Wei, T. K.; Nashiru, B. 2012. Formulation Development and Optimization of a Novel Cremophore EL-Based Nanoemulsion Using Ultrasound Cavitation. *Ultrason. Sonochem.*, 19, 330–345.
- Temporal, P. 2011. *Islamic Branding and Marketing: Creating a Global Islamic Business*. Singapore: John Wiley & Sons (Asia) Pte.Ltd.
- Thomassen, H. a; Wiersema, A. T.; de Bakker, M. a. ; de Knijff, P.; Hetebrij, E.; Povel, G. D. E. 2003. A New Phylogeny of Swiftlets (Aves: Apodidae) Based on Cytochrome-B DNA. *Mol. Phylogenet. Evol.*, 29, 86–93.
- Tilak, K. S.; Veeraiah, K.; Milton Prema Raju, J. 2007. Effects of Ammonia, Nitrite and Nitrate on Hemoglobin Content and Oxygen Consumption of Freshwater Fish, *Cyprinus Carpio* (Linnaeus). *J. Environ. Biol.*, 28(1) 45-47.
- Triveni, R.; Shamala, T.; Rastogi, N. 2001. Optimised Production and Utilisation of Exopolysaccharide from *Agrobacterium Radiobacter*. *J. Proc. Biochem.*, 36, 787–795.
- Troncoso, E.; Aguilera, J.; McClements, D. 2011. Development of Nanoemulsions by an Emulsification-Evaporation Technique. In *Proceedings of the 11th International Congress on Engineering and Food*; Taoukis, P. S.; Stoforos, N. G.; Karathanos, V. T.; Saravacos, G. D., Eds.; Athens, Greece, pp 929–930.
- Umbach, W. 1995. Cosmeceuticals-the future of cosmetics. *Cosmetics & Toiletries.*, 110(11), 33-40.
- US Commercial Service Hong Kong. 2013. *Cosmetics & Toiletries Market Overview 2013*. Retrieved 18 October 2014 from <http://business.usa.gov/export->

portal?hongkong/build/groups/public/@eg_hk/documents/webcontent/eg_hk_065324.pdf.

Utusan. 2011. Hukum Makan Sarang Burung. Retrieved 6 June 2014 from http://ww1.utusan.com.my/utusan/info.asp?y=2011&dt=0603&pub=Utusan_Malaysia&sec=Bicara_Agama&pg=ba_06.htm.

Walstra, P. 1996. Emulsion Stability. In *Encyclopedia of Emulsion Technology*; Becher, P. Ed; Marcel Dekker, New York, pp. 1-62.

Walters, C. L.; Smith, P. L. R.; Reed, P. I.; Haines, K.; House, F. R. 1982. N-Nitroso Compounds in Gastric Juice and Their Relationship to Gastrointestinal Disease. In *N-Nitroso Compounds: Occurrence and Biological Effects*; Bartsch, H.; O'Neill, I. K.; Castegnaro, M.; Okada, M.; Davis, W., Eds.; International Agency for Research on Cancer, Lyon, pp. 345–355.

Wang, C. C. 1921. The Composition of Chinese Edible Birds' Nests and the Nature of Their Proteins. *J. Biol. Chem.*, 49, 429–439.

Wei, W.; Wang, X.; Huang, Z. 2011. Bird Nest Health-Care Product Buccal Tablet. CN 102239984 A.

Weiss J.; Herrmann, N.; McClements, D.J. 1999. Ostwald Ripening of Hydrocarbon Emulsion Droplets in Surfactant Solutions. *Langmuir.*, 15, 6652-57.

Wooster, T.J.; Golding, M.; Sanguansri, P. 2008. Impact of Oil Type on Nanoemulsion Formation and Ostwald Ripening Stability. *Langmuir.*, 24, 12758-12765.

Yeoh, W. (August 1, 2011). Group: Getting Zero Nitrite Level in Bird's Nest Impossible. The Star Online. Retrieved 31 Desember 2014 from <http://www.thestar.com.my/story/?sec=nation&file=%2F2011%2F8%2F1%2Fnation%2F9213607>.

Zainab, H.; Sarojini, J.; Nur Hulwani, I.; Othman, H.; Boon-Beng, L.; Kamarudin, H. 2013a. A Rapid Technique to Determine Purity of Edible Bird Nest. *Adv. Environmental Biol.*, 7, 3758–3765.

Zainab, H.; Nur Hulwani, I.; Sarojini, J.; Kamarudin, H.; Othman, H.; Boon-Beng, L. 2013b. Nutritional Properties of Edible Bird Nest. *J. Asian Sci. Res.*, 3, 600–607.

Zainol, S.; Basri, M.; Basri, H.; Shamsuddin, A. F.; Abd Gani, S. S.; Karjiban, R. A.; Abdul-Malek, E. 2012. Formulation Optimization of a Palm-Based Nanoemulsion System Containing Levodopa. *Int. J. Mol. Sci.*, 13, 13049–13064.

Zhang, Y.; Liu, X.; McHale, C.; Li, R.; Zhang, L.; Wu, Y.; Ye, X.; Yang, X.; Ding, S. 2013. Bone Marrow Injury Induced via Oxidative Stress in Mice by Inhalation Exposure to Formaldehyde. *PLoS ONE*, 8(9), e74974. Retrieved 23 Desember 2014 from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3770590/>

Zhao, N.; Qin, Y.; Sun, Y.; Sun, J.; He, Z. 2011. Formulation and Preparation of a Nitrendipine Submicron Emulsion. *Asian J. Pharm. Sci.*, 6, 151–158.

