



UNIVERSITI PUTRA MALAYSIA

***PREDICTORS OF MALARIA AMONG PREGNANT WOMEN
ATTENDING ANTE-NATAL CLINIC IN GENERAL HOSPITAL
IN ZAMFARA STATE, NIGERIA***

KALLAMU HADIZA

FPSK(m) 2015 35



**PREDICTORS OF MALARIA AMONG PREGNANT WOMEN ATTENDING
ANTE-NATAL CLINIC IN GENERAL HOSPITAL
IN ZAMFARA STATE, NIGERIA**

By

KALLAMU HADIZA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

November 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

PREDICTORS OF MALARIA AMONG PREGNANT WOMEN ATTENDING ANTE-NATAL CLINIC IN GENERAL HOSPITAL IN ZAMFARA STATE, NIGERIA

By

KALLAMU HADIZA

November 2015

Chair : Assoc. Prof. Hejar Binti Abd Rahman, PhD
Faculty : Medicine and Health Sciences

Introduction: Malaria is a major health problem with global concern. It is one of the leading causes of morbidity and mortality in the tropics and subtropics. In 2014, there were 2.16 billion cases and one million deaths per year. Nigeria is included among the 45 countries that are endemic for malaria, and about 97% of the population were at risk especially children and pregnant women.

Objectives: To determine the predictors of malaria among pregnant women attending ante-natal clinic in general hospitals Zamfara State, Nigeria.

Methodology: A case control study was conducted among pregnant women attending ante-natal clinic in general hospital Zamfara State, Nigeria. A total of 522 pregnant women 261 cases and 261 controls were selected using multistage random sampling. Cases and controls in this study were defined as a pregnant woman attending ante-natal clinic from the selected general hospitals in Zamfara, confirmed with and without malaria respectively, using giemsa staining method based on their medical records. Face to face interview and self-administered pretested questionnaire in English and Hausa languages was used to obtain information based on their socio demographic characteristics, maternal history, knowledge, attitude and preventive practices regarding malaria from May to August 2014. The data was analysed using SPSS version 21, chi square test was used to determine the association between the predictors of malaria. Significant level (p) was set at 0.05.

Results: The overall response rate was 89.8%. Chi square results shows that pregnant women with older age ($\chi^2 = 17.835$, $df = 3$, $p < 0.001$), informal education ($\chi^2 = 166.619$, $df = 4$, $p < 0.001$), unemployment ($\chi^2 = 220.519$, $df = 2$, $p < 0.001$), monthly income < 5000 naira ($\chi^2 = 353.841$, $df = 2$, $p < 0.001$), first trimester ($\chi^2 = 27.754$, $df = 2$, $p < 0.001$). Other essential significant variables include low level of knowledge on malaria ($\chi^2 = 96.632$, $df = 1$, $p < 0.001$), negative attitude ($\chi^2 = 248.309$, $df = 1$, $p < 0.001$) and low level of preventive practices ($\chi^2 = 148.761$, $df = 1$, $p < 0.001$) are

significantly associated with malaria. Predictors of malaria include informal education (AOR = 8.340, 95% CI = 3.170, 21.947), unemployment (AOR = 8.437, 95% CI = 1.695, 42.007), monthly income < 1000 naira (AOR = 18.809, 95% CI = 3.829, 92.393), low knowledge (AOR = 5.363, 95% CI = 2.130, 13.501), negative attitude (AOR = 33.831, 95% CI = 12.749, 89.778) and no practice (AOR = 44.622, 95% CI = 3.829, 92.392).

Conclusion: This study has identified informal education, unemployment, low monthly income, low level of knowledge, negative attitude and poor preventive practices as predictors of malaria among pregnant women attending ante-natal clinic in general hospitals in Zamfara State, Nigeria. The findings in this study can be used by policy makers to plan how to tackle the risk factors of malaria among pregnant women in the State.

Keywords: Malaria, predictors, pregnancy, Zamfara State, Nigeria

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Sarjana Sains

PERAMAL MALARIA DALAM KALANGAN WANITA HAMIL YANG MENGHADIRI KLINIK ANTENATAL DI HOSPITAL BESAR DI ZAMFARA NEGERI, NIGERIA 2014

Oleh

KALLAMU HADIZA

November 2015

Pengerusi : Prof. Madya Hejar Binti Abd Rahman, PhD
Fakulti : Perubatan dan Sains Kesihatan

Pengenalan: Malaria merupakan masalah kesihatan utama dengan kebimbangan global. Ia merupakan salah satu penyakit berjangkit yang serius paling lazim di dunia, dengan kira-kira 250 juta kes dan satu juta kematian setiap tahun. Nigeria adalah termasuk di kalangan 45 negara yang endemik malaria, dan kira-kira 97% daripada populasi berisiko terutamanya kanak-kanak dan wanita hamil.

Objektif: Untuk menentukan peramal malaria dalam kalangan wanita hamil yang menghadiri klinik antenatal di Hospital besar Zamfara Negeri, Nigeria.

Metodologi: Satu kajian kes-kes kawalan telah dijalankan dalam kalangan wanita hamil yang menghadiri klinik antenatal di Hospital besar Zamfara Negeri, Nigeria. Seramai 522 wanita hamil 261 kes dan 261 kawalan telah dipilih menggunakan pensampelan rawak pelbagai peringkat. Kes dan kawalan dalam kajian ini ditakrifkan sebagai seorang wanita hamil yang menghadiri klinik antenatal dari hospital besar terpilih di Zamfara, disahkan dengan dan atampa malaria masing-masing menggunakan kaedah pewarnaan giemsa berdasarkan rekod perubatan mereka. Temuduga bersemuka dan soal selidik praujian tadbir sendiri dalam Bahasa Inggeris dan Bahasa Hausa telah digunakan untuk mendapatkan maklumat berdasarkan kepada ciri-ciri sosio demografi mereka, sejarah keibuan, pengetahuan, sikap dan amalan pencegahan mengenai malaria dari Mei hingga Ogos 2014. Data dianalisis dengan menggunakan SPSS versi 21, ujian khi kuasa dua telah digunakan untuk menentukan hubungan antara kumpulan kes dan kawalan, dan regresi logistik telah digunakan untuk menentukan peramal malaria. Aras signifikan (p) telah ditetapkan pada 0.05

Hasil: Kadar sambutan keseluruhan adalah 89.8%. Keputusan khi kuasa dua dengan malaria berbanding mereka yang lebih berusia ($F = 17.835$, $df = 3$, $p < 0.001$), pendidikan tidak formal ($F = 166.619$, $df = 4$, $p < 0.001$), pengangguran ($F = 220.519$, $df = 2$, $p < 0.001$), pendapatan bulanan < 5000 naira ($F = 353.841$, $df = 2$, $p < 0.001$), trimester pertama ($F = 27.754$, $df = 2$, $p < 0.001$). Pembolehubah signifikan penting yang lain termasuk tahap pengetahuan yang rendah mengenai malaria ($F = 96.632$, $df = 1$, $p < 0.001$), sikap negatif ($F = 248.309$, $df = 1$, $p < 0.001$) dan tahap

amalan pencegahan yang rendah ($\chi^2 = 148.761$, $df = 1$, $p < 0.001$) adalah berhubung secara signifikan dengan malaria. Peramal malaria termasuk pendidikan tidak formal (AOR = 8.340, 95% CI = 3.170, 21.947), pengangguran (AOR = 8.437, 95% CI = 1.695, 42.007), pendapatan bulanan <1000 naira (AOR = 18.809, 95% CI = 3.829, 92.393), pengetahuan rendah (AOR = 5.363, 95% CI = 2.130, 13.501), sikap negatif (AOR = 33.831, 95% CI = 12.749, 89.778) dan amalan miskin (AOR = 44.622, 95% CI = 3.829, 92.392).

Kesimpulan: Kajian ini telah mengenal pasti pendidikan tidak formal, pengangguran, pendapatan bulanan yang rendah, tahap rendah pengetahuan, sikap negative dan amalan pencegahan miskin sebagai peramal malaria dalam kalangan wanita hamil yang menghadiri klinik antenatal di Hospital besar di Zamfara Negeri, Nigeria. Penemuan dalam kajian ini boleh digunakan oleh pembuat dasar dalam merancang bagaimana untuk menangani faktor-faktor risiko malaria dalam kalangan wanita hamil di negeri ini.

Kata kunci: Malaria, peramal, kehamilan, Zamfara Negeri, Nigeria

ACKNOWLEDGEMENT

I wish to express my deepest appreciation to my supervisor Associate Professor (Dr) Hejar Binti Abdul Rahman whose despite her tight schedule was always available and helpful. Special thanks to my co-supervisor Dr Hayati Kadir @ Shahar for her warm support and guidance throughout the programme.

I cannot express my gratitude to my husband Dr. Usman Ismaila Gusau, without him, I would not have completed this programme. Special thanks to my student nurses and my siblings Fadalu A.A Gusau, Mubarak Kallamu, Bello Ismaila Iwucw'cpf" Dcdc" Cwf wou" hc o kn { "hqt" v j gkt "mkpf" cuukvcpeg" fwtkpi "v j g" fcvc" eqnngcvkqp0

Words cannot really express my gratitude to my husband Dr. Usman Ismaila Gusau, without him, I would not have completed this programme. Special thanks to my parents Arch. A. A. Gusau, Hajiya Rabi and Hajiya Ummu A. A. Gusau who laid the foundation, their support, prayers and the responsibility of taking care of the kids throughout the programme. Warmest thanks to my lovely kids Waleed, Amina and Rukayya Usman Ismaila for the sacrifice of bearing our absence.

Finally, my appreciation to my colleagues and friends Dr. Tayo Martins, Dr. Jamila U. Garba, Zubaida Mahmud, Safiyya Kalgo, Kubrah Usman, Hadiza Anka and all those who contributed one way or the other during the programme.

I certify that a Thesis Examination Committee has met on 18 November 2015 to conduct the final examination of Kallamu Hadiza on her thesis entitled "Predictors of Malaria among Pregnant Women Attending Ante-Natal Clinic in General Hospital in Zamfara State, Nigeria" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Ahmad Azuhairi Ariffin, PhD

Senior Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Huda binti Zainuddin, PhD

Senior Lecturer
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Mohamad Ikhsan bin Selamat, PhD

Senior Lecturer
Universiti Teknologi MARA
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 January 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Hejar Binti Abdul Rahman; M.D, Master Community Health

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairperson)

Hayati Kadir @ Shahar M.D., PhD

Senior Lecturer (Medical)
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Kallamu Hadiza, GS36517

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman
Supervisory
Committee:

Hejar Binti Abdul Rahman; M.D, Master Community Health

Signature: _____

Name of
Member of
Supervisory
Committee:

Hayati Kadir @ Shahar M.D., PhD

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem statements	2
1.3 Significance of the Study	2
1.4 Objectives	3
1.4.1 General Objectives	3
1.4.2 Specific Objectives	3
1.5 Research Hypotheses	3
2 LITERATURE REVIEW	4
2.1 Epidemiology of malaria worldwide	4
2.2 Epidemiology of malaria in Africa	5
2.3 Epidemiology of malaria in Nigeria	5
2.4 Aetiology of malaria	6
2.5 Mode of Transmission	6
2.6 Susceptibility and resistance	7
2.6.1 Susceptibility	7
2.6.2 Resistance	7
2.7 Life Cycle of Malaria Parasite	7
2.8 Clinical Signs and Symptoms	8
2.9 Diagnosis of malaria	9
2.10 Treatment of malaria	9
2.11 Prevention and control of malaria	10
2.11.1 Primary prevention of malaria	10
2.11.1.1 Insecticide-treated mosquito nets (ITNs)	10
2.11.1.2 Indoor residual spraying (IRS)	11
2.11.1.3 Intermittent preventive treatment in pregnancy	11
2.11.2 Secondary prevention of malaria	12
2.11.2.1 Early diagnosis	12
2.11.2.2 Prompt treatment	12
2.11.3 Tertiary prevention of malaria	13
2.11.3.1 Pharmaceutical and commodity management	13
2.11.3.2 Advocacy, communication, and social mobilization	13
2.12 The association between malaria and pregnancy	14
2.12.1 Malaria in Pregnancy	14
2.12.2 Natural Immunity against Malaria in Pregnancy	15
2.12.3 Malaria in Pregnancy with HIV	16

2.13	Risk factors of malaria during pregnancy	16
2.13.1	Socio demographic characteristics	16
2.13.1.1	Age	16
2.13.1.2	Educational status	16
2.13.1.3	Monthly income	17
2.13.1.4	Gravidity	17
2.13.1.5	Trimester	18
2.13.2	Knowledge of malaria	18
2.13.3	Attitude towards malaria	19
2.13.4	Practice regarding malaria	19
2.13.4.1	Insecticide Treated Nets (ITNs)	19
2.13.4.2	Intermittent Preventive Treatment (IPT)	20
2.14	Conceptual Framework	21
3	METHODOLOGY	23
3.1	Study Location	23
3.2	Study Duration	24
3.3	Study Design	24
3.4	Study Population	24
3.5	Selection Criteria	25
3.5.1	Case group	25
3.5.1.1	Inclusion Criteria for Cases	26
3.5.1.2	Exclusion Criteria for Cases	26
3.5.2	Control group	26
3.5.2.1	Inclusion Criteria for Control	26
3.5.2.2	Exclusion Criteria for Control	26
3.6	Sampling frame	26
3.6.1	Case group	26
3.6.2	Control group	26
3.7	Sampling unit	26
3.8	Sample size estimation	26
3.9	Sampling method	27
3.10	Instrument of the study	27
3.10.1	Questionnaire	27
3.10.2	Components of questionnaire	28
3.10.2.1	Socio demographic characteristics	30
3.10.2.2	Maternal history	30
3.10.2.3	Knowledge	30
3.10.2.4	Attitude	30
3.10.2.5	Preventive practices	31
3.11	Data Collection	31
3.12	Variables of the study	31
3.12.1	Dependent Variable	31
3.12.2	Independent Variables	31
3.13	Operational definition of terms	32
3.13.1	Cases	32
3.13.2	Controls	32
3.13.3	Socio demographic characteristic	32
3.13.4	Gravidity	32
3.13.5	Primigravida	32
3.13.6	Gravida 2	32

3.13.7	Multigravida	32
3.13.8	Trimester	32
3.13.9	Maternal history	33
3.14	Quality control	33
3.14.1	Validity	33
3.14.1.1	Content validity	33
3.14.1.2	Face validity	33
3.14.2	Reliability	33
3.15	Data analysis	34
3.16	Ethical considerations	34
4	RESULTS	35
4.1	Response rate	35
4.2	Test of normality	35
4.2.1	Test of normality for knowledge	36
4.2.2	Test of normality for attitude	36
4.2.3	Test of normality for practice	36
4.3	Socio-demographic characteristics of the pregnant women	37
4.4	Maternal history of the pregnant women	38
4.5	Knowledge on Malaria	39
4.5.1	Knowledge of malaria based on causes, symptoms, breeding sites and people at high risk	39
4.5.2	Knowledge on mode of transmission and prevention of malaria	41
4.5.3	Knowledge on effects, seasons and diagnosis of malaria	43
4.5.4	Level of knowledge of the pregnant women	44
4.6	Attitude regarding Malaria	46
4.6.1	Attitude regarding malaria on causes and breeding sites	46
4.6.2	Attitude regarding malaria on mode of transmission and seasons	47
4.6.3	Attitude regarding malaria on treatment and control	48
4.6.4	Level of attitude of the respondents	50
4.7	Practices towards Malaria	51
4.7.1	Preventive practices towards malaria	51
4.7.2	Level of practices of the pregnant women	52
4.8	Association between socio demographic characteristics of the respondents and malaria	52
4.9	Association between maternal history of the pregnant women and malaria	53
4.10	Association between level of knowledge of the pregnant women and malaria	54
4.11	Association between attitude of the pregnant women and malaria	55
4.12	Association between practices of the respondents and malaria	56
4.13	Risk factors for malaria	56
4.14	Multiple logistic regression analysis showing the Predictors of malaria	57
4.15	Model based on Log. Regression	59
5	DISCUSSION	60
5.1	Socio demographic characteristics of the respondents	60
5.2	Maternal history of the respondents	61
5.3	Knowledge on malaria	62

5.4	Attitude towards malaria	63
5.5	Practices related to malaria	64
5.6	Factors associated with malaria	64
5.7	Predictors of malaria	65
6	SUMMARY, CONCLUSION AND RECOMMENDATION	67
6.1	Summary and conclusion	67
6.2	Strengths and limitations of the study	67
6.2.1	Strength of the study	67
6.2.2	Limitations of the study	67
6.3	Recommendations	68
	REFERENCES	69
	APPENDICES	80
	BIODATA OF STUDENT	104
	PUBLICATION	105

LIST OF TABLES

Table	Page
3.1: Average attendance of ante-natal clinic ANC and deliveries for the selected hospitals from May ó July 2014	25
3.2: Knowledge range of scores	30
3.3: Attitude range of scores	31
3.4: Reliability test results	34
4.1: Response rate from the selected General Hospitals in Zamfara State, Nigeria	35
4.2: Socio-demographic characteristics of the pregnant women (N=522)	38
4.3: Maternal history of the pregnant women (N=522)	38
4.4: Knowledge regarding causes and symptoms of malaria, breeding sites and people at high risk (N=522)	40
4.5: Knowledge on mode of transmission and prevention of malaria	42
4.6: Knowledge on effects, seasons and diagnosis of malaria	43
4.7: Level of knowledge of the respondents (N=522)	45
4.8: Attitude regarding malaria on causes and breeding sites (N=522)	46
4.9: Attitude regarding malaria on mode of transmission and seasons	47
4.10: Attitude regarding malaria on treatment and control (N=522)	49
4.11: Level of attitude of the pregnant women (N=522)	51
4.12: Practices towards malaria (N=522)	52
4.13: Level of practices of the pregnant women (N=522)	52
4.14: Association between socio demographic characteristics of the respondents and malaria (N=522)	53
4.15: Association between maternal history of the pregnant women and malaria (N=522)	54
4.16: Level of knowledge of the pregnant women and malaria (N=522)	55
4.17: Level of attitude of the pregnant women (N=522)	56
4.18: Level of practices of the respondents	56
4.19: Logistic regression analysis showing the risk factors associated with malaria among pregnant women	57
4.20: Multiple logistic regression showing adjusted odds ratio (AOR) of predictors of malaria	59

LIST OF FIGURES

Figure	Page
2.1: Global areas of elevated risk of malaria	4
2.2: Life cycle of mosquito	6
2.3: Life cycle of malaria parasite	8
2.4: Areas where symptoms of malaria occur	9
2.5: The conceptual framework of factors associated with malaria among pregnant women attending ANC in Zamfara.	22
3.1: Map of Nigeria showing Zamfara State	24
3.2: Map of Zamfara State showing local government areas	25
3.3: Flow chart showing multi-stage sampling technique of pregnant women attending ANC in Zamfara State, Nigeria	29
4.1: Test of normality for knowledge	36
4.2: Test of normality for attitude	36
4.3: Test of normality for practice	37

LIST OF ABBREVIATIONS

<	Less than
>	Greater than
Ö	Less than or equal to
×	Greater than or equal to
AOR	Adjusted odds ratio
CI	Confidence interval
COR	Crude odds ratio
DF	Degree of freedom
F	Frequency
FMOH	Federal Ministry of Health
IPT	Intermittent Preventive Treatment
IRS	Insecticide Residual Spray
ITNs	Insecticide Treated Nets
LLINs	Long Lasting Insecticide Nets
MDGs	Millennium Development Goals
N	Total number
NDHS	National Demographic and Health Survey
NMCP	National Malaria Control Programme
NMCSP	National Malaria Control Strategic Plan
NPC	National Population Commission
OR	Odds ratio
RBM	Roll Back Malaria
SP	Sulfadoxine pyrimethamine/Fansidar
WHO	World Health Organisation

CHAPTER 1

INTRODUCTION

1.1 Background

Malaria is a major health problem, and has attracted global concern. There were approximately 198 million cases and 584,000 deaths in 2013 (WHO, 2014), and it was estimated that more than 80% of the cases were in Sub-Saharan Africa. Approximately one death occurs every 30 seconds with 90% of the mortality rate occurring in Sub-Saharan Africa, and 90% of malaria deaths are of children less than five years of age and pregnant women (Tillotson, 2012).

Control of malaria still remains a challenge in Africa, as evidenced by the 163 million estimated cases and 528,000 deaths in 2013. Nigeria and the Democratic Republic of the Congo together accounted for 39% and 34% of the global total of estimated malaria deaths and cases respectively in 2013 (WHO, 2014). Nigeria, which is included among the 45 countries that are endemic for malaria, has a population of over 170 million, and 97% of the population are at risk particularly pregnant women and children. The statistics indicate that Nigeria alone accounts for 45% of the prevalence in the African continent (Agomo, Oyibo, Anorlu, & Agomo, 2009; Aregawi, Cibulskis, Otten, & Williams, 2009; Duffy & Fried, 2005). Malaria in pregnancy is a serious health problem both in pregnant women and her foetus with 11% maternal deaths annually in Nigeria (Nzeako, Nduka, & Origie, 2013; World Health Organization, 2012).

The Nigerian climate makes malaria transmission suitable throughout the country, due to geographic location of the country; only 3% of the populace who live at an altitude ranging from 1,200 to 1,400 metres in the Southern Jos, Plateau State are at relatively low risk of malaria. Similarly, it has been estimated that about 140 million people are living in areas of high malaria transmission due to large population of Nigeria (Polsa, Spens, Soneye, & Antai, 2011). About 30 million women in malaria-endemic areas of Africa become pregnant each year, and are at risk of infection with *Plasmodium falciparum*. The prevalence of malaria infection in the north-western part of Nigeria, Sokoto State is 27.29%, as malaria is endemic in Nigeria with seasonal variation in different geographic regions of the country; it has been described as moderately high (Abdullahi et al., 2009; Steketee, Nahlen, Parise, & Menendez, 2001).

Pregnancy complications have been associated with increased incidence and severity of malaria. In Sub-Saharan Africa, the effects of the disease are anaemia, spontaneous abortion, prematurity and stillbirths are effects of the disease (Okpere, Enabudoso, & Osemwenkha, 2010). The increased susceptibility of pregnant women to malaria is as a result of decreased immunity caused by pregnancy which makes them more vulnerable to anaemia, still birth, placental parasitisation and increasing the risk of illness leading to death. Maternal malaria also affects the unborn babies by increasing the risk of spontaneous abortion, stillbirth, premature delivery and low birth weight, and is a leading cause of child mortality (World Health Organization, 2010).

Malaria is usually linked with poverty and may also be a major burden to economic development (Aregawi et al., 2009). The prevalence of the disease is higher in Sub-Saharan Africa than in many other regions of the world, more than 75% of the cases in this region are due to *P. falciparum*, while in most other countries the disease transmission is as a result of the other less virulent plasmodial species. It has however been reported that the majority of the mortality recorded as a result of malaria is caused by *P. falciparum* (Aregawi et al., 2009).

The increased risk of malarial infection could be related to illiteracy, low educational status, unemployment, low income and gravidity of the pregnant women. The other risk factors are environment, such as the presence of unclean gutters, residence in swamps that are favourable conditions for the breeding of Anopheles mosquitoes, poor knowledge, attitude and preventive practices towards the prevention and control measures of malaria (Amuta, Houmsou, Wama, & Ameh, 2014; Bawa, Auta, & Liadi, 2014; Oyefabi, Sambo, & Sabitu, 2015).

1.2 Problem statements

Malaria still remains an important public health concern globally; about 3.3 billion people were at risk of malaria infection in 2013. It has been indicated by the World Malaria Reports that Africa bears the heaviest burden and the highest risk of malaria infection. Africa accounted for about 82% and 90% of the reported malaria cases and deaths respectively, with pregnant women and children below five years suffering the most (WHO, 2014).

High prevalence rates of malaria among pregnant women have been reported by many studies in different parts of Nigeria, (namely Kogi, Katsina, Kano, Benue, Osun and Rivers States) ranging from 30.0% to 72.5% (Adefioye, Adeyeba, Hassan, & Oyeniran, 2007; Amuta, Houmsou, Wama, & Ameh, 2014; Bawa, Auta, & Liadi, 2014; Gajida, Iliyasu, & Zoakah, 2010; Mofolorunsho, Audu, & Omatola, 2014; Nzeako, Nduka, & Origie, 2013). More than 90% of the Nigerian population were at risk of malaria and at least 50% of the total population experience an incidence of malaria yearly. The disease also affects the general population, beyond the effects on pregnant women and children (Federal Ministry of Health, 2004; Roll Back Malaria, 2013).

Each year up to 10,000 maternal deaths were recorded as a result of maternal malaria and it contributes to high maternal morbidity rates; that is, severe anaemia, fever, and placental parasitaemia especially in first time mothers (Falade, Tongo, Ogunkunle, & Qtkocfgiwp."4232-"Ucxcig."Ou{c o dq|c." Ikgu."FøCnguucpftq." ("Dtcdkp."4229+. It has also been attributed to between 75,000 to 200,000 infant deaths annually (Steketee et al., 2001).

1.3 Significance of the Study

The significance of the study is to identify the predictors of malaria among pregnant women, and highlight areas for possible management modification and further research based on the findings of current predictors of malaria among pregnant women in Zamfara State. Identifying the predictors and implementing prevention of malaria during pregnancy is one of the major interventions in helping to reduce maternal and infant mortality and morbidity.

Since no research has been reported so far on the predictors of malaria among pregnant women in Zamfara State, this study will provide more information. Hence, information that will be gathered from the study will be used by the Ministry of Health, Zamfara State, to improve programmes dealing with prevention services against malaria in the State.

1.4 Objectives

1.4.1 General Objectives

The general objective is to determine the predictors of malaria among pregnant women attending ante-natal clinic in general hospitals Zamfara State.

1.4.2 Specific Objectives

- i. To determine the socio-demographic factors (such as age, ethnicity, religion, marital status, educational status, occupation and monthly income) of pregnant women among cases and controls.
- ii. To determine the maternal history (such as gravidity and trimester) of pregnant women among cases and controls.
- iii. To determine the level of knowledge (such as causes, signs and symptoms, breeding sites, people at risk, mode of transmission, prevention and control, effects, seasonal variation and diagnosis), attitude (such as mode of transmission, treatment, prevention and control) and preventive practices (using LLIN, repellent spray, long sleeves, mosquito coil and good hygiene) on malaria in pregnant women among cases and controls.
- iv. To determine the association between socio-demographic factors, maternal history, knowledge, attitude and practices of pregnant women and malaria.
- v. To determine the predictors of malaria in pregnant women among cases and controls.

1.5 Research Hypotheses

- i. There is a significant association between the socio-demographic factors of cases as compared to controls with regards to malaria in pregnancy.
- ii. There is a significant association between the maternal histories of cases compared to controls with regards to malaria in pregnancy.
- iii. There is a significant association between the levels of knowledge of cases compared to controls with regards to malaria in pregnancy.
- iv. There is a significant association between the levels of attitude of cases compared to controls with regards to malaria in pregnancy.
- v. There is a significant association between the levels of practice of cases compared to controls with regards to malaria in pregnancy.
- vi. Socio-demographic factors, maternal history, level of knowledge, attitude and practice are risk factors of malaria among pregnant women.
- vii. Socio-demographic factors, maternal history, level of knowledge, attitude and practice are predictors of malaria among pregnant women.

REFERENCES

- Abba, K., Deeks, J. J., Olliaro, P., Naing, C., Jackson, S. M., Takwoingi, Y., Garner, P. (2011). Rapid diagnostic tests for diagnosing uncomplicated *P. falciparum* malaria in endemic countries. *Cochrane Database Syst Rev*, Issue 12. Art. No.: CD011431. doi: 10.1002/14651858.CD011431.
- Abdulla, S. M. K. (2000). Malaria control strategies in the Kilombero valley, Tanzania. University of Basel, faculty of science PhD thesis
- Abdullahi, M. (2009). Malaria in Sokoto, North Western Nigeria. *African Journal of Biotechnology*, 8(24), 7101-7105
- Adefioye, O., Adeyeba, O., Hassan, W., & Oyeniran, O. (2007). Prevalence of malaria parasite infection among pregnant women in Osogbo, Southwest, Nigeria. *American-Eurasian Journal of Scientific Research*, 2(1), 43-45.
- Ademiluyi, I. A., & Aluko-Arowolo, S. O. (2009). Infrastructural distribution of healthcare services in Nigeria: An overview. *Journal of Geography and Regional Planning*, 2(5), 104-110.
- Adetokunbo, T., Akinola, O., Shittu, L., Ottun, T., Bankole, M., Akinola, R., Okunribido, A. (2009). Prevalence of malaria parasitaemia in the booking antenatal (ANC) patients at the Lagos State University Teaching Hospital. *African Journal of Biotechnology*, 8(15), 3628-3631.
- Adeyemo, M. O., Oluwatosin, O. A., Amodu, O. K., & Yekinni, O. T. (2014). Home management and prevention of malaria among under-five children: experiences of mothers in Egbedore Local Government Area (LGA), Osun State Nigeria. *African Journal of Biomedical Research*, 17(2), 83-91.
- Agomo, C. O., & Oyibo, W. A. (2013). Factors associated with risk of malaria infection among pregnant women in Lagos, Nigeria. *Infect Dis Poverty*, 2(1), 19. doi:10.1186/2049-9957-2-19.
- Agomo, C. O., Oyibo, W. A., Anorlu, R. I., & Agomo, P. U. (2009). Prevalence of malaria in pregnant women in Lagos, South-west Nigeria. *The Korean Journal of Parasitology*, 47(2), 179-183.
- Ahorlu, C. K., Dunyo, S. K., Afari, E. A., Koram, K. A., & Nkrumah, F. K. (1997). Malaria related beliefs and behaviour in Southern Ghana: Implications for treatment, prevention and control. *Tropical Medicine & International Health*, 2(5), 488-499.
- Akaba, G., Otubu, J., Agida, E., & Onafowokan, O. (2013). Knowledge and utilization of malaria preventive measures among pregnant women at a Nigerian Journal of Clinical Practice, 16(2), 201-206.

- Akinleye, S. O., Falade, C. O., & Ajayi, I. O. (2009). Knowledge and utilization of intermittent preventive treatment for malaria among pregnant women attending antenatal clinics in primary health care centers in rural Southwest, Nigeria: A cross-sectional study. *BMC Pregnancy and Childbirth*, 9, 28-2393-9-28.
- Alaku, I., Abdullahi, A., & Kana, H. (2015). Epidemiology of malaria parasites infection among pregnant women in some part of Nasarawa State, Nigeria. *Developing Country Studies*, 5(2), 30-33.
- Amuta, E., Houmsou, R., Wama, E., & Ameh, M. (2014). Malarial infection among antenatal and maternity clinics attendees at the federal medical centre, Makurdi, Benue State, Nigeria. *Infectious Disease Reports*, 6:5050 doi:10.4081/idr.2014.5050
- Anchang-Kimbi, J. K., Achidi, E. A., Nkegoum, B., Sverremark-Ekström, E., & Troye-Blomberg, M. (2009). Diagnostic comparison of malaria infection in peripheral blood, placental blood and placental biopsies in Cameroonian parturient women. *Malaria Journal*, 8(1), 126. doi:10.1186/1475-2875-8-126.
- Ankomah, A., Adebayo, S. B., Arogundade, E. D., Anyanti, J., Nwokolo, E., Ladipo, O., & Meremikwu, M. M. (2012). Determinants of insecticide-treated net ownership and utilization among pregnant women in Nigeria. *BMC Public Health*, 12, 105-2458-12-105.
- Aregawi, M., Cibulskis, R. E., Otten, M., & Williams, R. (2009). *World malaria report 2009* World Health Organization.
- Babalola, M. (2013). An examination of the association between malaria knowledge and bed net use of pregnant women receiving antenatal care at federal medical centre, Abeokuta, Nigeria. Thesis, Georgia State University.
- Dcvk."L0." Ngiguug." O0." (" Ogfj kp." I 0" *4235+0" Eq o o w p kv {øu" mpq y ng f i g." c w k w f g u" and practices about tuberculosis in itang special district, Gambella Region, South Western Ethiopia. *BMC Public Health*, 13(1), 1-9.
- Bawa, J., Auta, T., & Liadi, S. (2014). Prevalence of malaria: Knowledge, attitude and cultural practices of pregnant women in Katsina metropolis, Nigeria. *European Scientific Journal*, 10(21), 148-167.
- Beck, S., Mockenhaupt, F. P., Bienzle, U., Eggelte, T. A., Thompson, W. N., & Stark, K. (2001). Multiplicity of *Plasmodium falciparum* infection in pregnancy. *The American Journal of Tropical Medicine and Hygiene*, 65(5), 631-636.
- Beeson, J. G., Rogerson, S. J., Cooke, B. M., Reeder, J. C., Chai, W., Lawson, A. M., Brown, G. V. (2000). Adhesion of plasmodium falciparum-infected erythrocytes to hyaluronic acid in placental malaria. *Nature Medicine*, 6(1), 86-90.

- Beeson, J., & Duffy, P. (2005). The immunology and pathogenesis of malaria during pregnancy. *Immunology and immunopathogenesis of malaria* (pp. 187-227) Springer.
- Beyene, H. B., Telele, N. F., & Mekuria, A. H. (2015). Knowledge, attitude and practice on malaria and associated factors among residents in Pawe district, North West Ethiopia: A cross-sectional study. *Science*, 3(3), 303-309.
- Bouyou-Akotet, M. K., Adegnika, A. A., Agnandji, S. T., Ngou-Milama, E., Kombila, M., Kremsner, P. G., & Mavoungou, E. (2005). Cortisol and susceptibility to malaria during pregnancy. *Microbes and Infection*, 7(11), 1217-1223.
- Breman, J. G., Alilio, M. S., Mills, A., Ter Kuile, F. O., Parise, M. E., Verhoeff, F. H., Rogerson, S. J. (2004). The burden of co-infection with human immunodeficiency virus type 1 and malaria in pregnant women in Sub-Saharan Africa. *The American journal of tropical medicine and hygiene*, 71(2 suppl), 41-54.
- Campbell, G. L., Marfin, A. A., Lanciotti, R. S., & Gubler, D. J. (2002). West Nile virus. *The Lancet Infectious Diseases*, 2(9), 519-529.
- Centers for Disease Control and Prevention (CDC). (2012). Impact of malaria. *Centers for Disease Control and Prevention CDC*.
- Dellicour, S., Tatem, A. J., Guerra, C. A., Snow, R. W., & Ter Kuile, F. O. (2010). Quantifying the number of pregnancies at risk of malaria in 2007: A demographic study. *PLoS Medicine*, 7(1), e1000221.
- Desai, M., ter Kuile, F. O., Nosten, F., McGready, R., Asamo, K., Brabin, B., & Newman, R. D. (2007). Epidemiology and burden of malaria in pregnancy. *The Lancet Infectious Diseases*, 7(2), 93-104.
- DHS, M. (2013). Malaria indicator survey. *Data Processing*, 5, 3. Retrieved from www.rbm.who.int
- Dockrill, P. (2015). World's first malaria vaccine approved by European regulators. Retrieved from www.sciencealert.com
- Doctor, H. V., Olatunji, A., Findley, S. E., Afenyadu, G. Y., Abdulwahab, A., & Jumare, A. (2012). Maternal mortality in Northern Nigeria: Findings of a health and demographic surveillance system in Zamfara State, Nigeria. *Tropical Doctor*, 42(3), 140-143.
- Doumbo, O. K., Thera, M. A., Kone, A. K., Raza, A., Tempest, L. J., Lyke, K. E., Rowe, J. A. (2009). High levels of *Plasmodium falciparum* rosetting in all clinical forms of severe malaria in African children. *The American Journal of Tropical Medicine and Hygiene*, 81(6), 987-993.
- Duffy, P., & Fried, M. (2005). Malaria in the pregnant woman. *Malaria: Drugs, disease and post-genomic biology* (pp. 169-200) Springer.

- Eisele, T. P., Keating, J., Littrell, M., Larsen, D., & Macintyre, K. (2009). Assessment of insecticide-treated bednet use among children and pregnant women across 15 countries using standardized national surveys. *The American Journal of Tropical Medicine and Hygiene*, 80(2), 209-214.
- Enato, E. F., Okhamafe, A. O., & Okpere, E. E. (2007). A survey of knowledge, attitude and practice of malaria management among pregnant women from two health care facilities in Nigeria. *Acta Obstetrica Et Gynecologica Scandinavica*, 86(1), 33-36.
- Eng, J. L. V., Thwing, J., Wolkon, A., Kulkarni, M. A., Many, A., Erskine, M., Slutsker, L. (2010). Research assessing bed net use and non-use after long-lasting insecticidal net distribution: A simple framework to guide programmatic strategies. 9:133 <http://www.malariajournal.com>
- Eze, U. I., Eferakeya, A. E., Oparah, A. C., & Enato, E. F. (2007). Assessment of prescription profile of pregnant women visiting antenatal clinics. *Pharmacy Practice*, 5(3), 135-139.
- Ezeigbo, O., Osuagwu, M., Ibegbulem, Z., & Agomoh, N. (2015). Evaluation of the knowledge, attitude and practice of the use of insecticide-treated nets (ITNs) in Aba, Nigeria. *British Journal of Medicine and Medical Research*, 5(1), 57-64.
- Falade, C. O., Tongo, O. O., Ogunkunle, O. O., & Orimadegun, A. E. (2010). Effects of malaria in pregnancy on newborn anthropometry. *The Journal of Infection in Developing Countries*, 4(07), 448-453.
- Falade, C. O., Dada-Adegbola, H. O., Ogunkunle, O. O., Oguike, M. C., Nash, O., & Ademowo, O. G. (2014). Evaluation of the comparative efficacy and safety of artemether-lumefantrine, artesunate-amodiaquine and artesunate-amodiaquine-chlorpheniramine (artemoclo) for the treatment of acute uncomplicated malaria in Nigerian children. *Medical Principles and Practice: International Journal of the Kuwait University, Health Science Centre*, 23(3), 204-211.
- Federal Ministry of Health. (2004). Maternal mortality situation and determinants in Nigeria. *Section 2*, 1-5.
- Fischer, P. R. (2003). Malaria and newborns. *Journal of Tropical Pediatrics*, 49(3), 132-135.
- Gajida, A., Iliyasu, Z., & Zoakah, A. (2010). Malaria among antenatal clients attending primary health care facilities in Kano State, Nigeria. *Annals of African Medicine*, 9(3), 188-93 doi: 10.4103/1596-3519.68352.
- Gamble, C., Ekwaru, J. P., & ter Kuile, F. O. (2006). Insecticide-treated nets for preventing malaria in pregnancy. *Cochrane Database Syst Rev*, Issue 2. Art. No.: CD003755. doi: 10.1002/14651858.CD003755.pub2.

- Garner, P., & Gülmezoglu, A. (2007). Drugs for preventing malaria in pregnant women (Review). Issue 4. Art. No.: CD000169. doi: 10.1002/14651858.CD000169.pub2.
- Gharoro, E. P., & Igbafe, A. (2000). Antenatal care: Some characteristics of the booking visit in a major teaching hospital in the developing world. *Medical Science Monitor*, 6(3), 519-522.
- Guyatt, H. L., & Snow, R. W. (2004). Impact of malaria during pregnancy on low birth weight in Sub-saharan Africa. *Clinical Microbiology Reviews*, 17(4), 760-769.
- Hamer, D. H., Mwanakasale, V., MacLeod, W. B., Chalwe, V., Mukwamataba, D., Champo, D., Mulele, C. K. (2007). Two-dose versus monthly intermittent preventive treatment of malaria with sulfadoxine-pyrimethamine in HIV-seropositive pregnant Zambian women. *Journal of Infectious Diseases*, 196(11), 1585-1594.
- Hezelgrave, N. L., Whitty, C. J., Shennan, A. H., & Chappell, L. C. (2011). Advising on travel during pregnancy. *BMJ*, 342:d2506.
- Hosmer, D. W., & Lemeshow, S. (2000). Introduction to the logistic regression model. *Applied Logistic Regression, Second Edition*, 1-30.
- Idowu, O. A., Mafiana, C. F., & Sotiloye, D. (2008). Traditional birth home attendance and its implications for malaria control during pregnancy in Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102(7), 679-684.
- Iriemenam, N., Dosunmu, A., Oyibo, W., & Fagbenro-Beyioku, A. (2011). Knowledge, Attitude, Perception of malaria and Evaluation of malaria parasitaemia among pregnant women attending antenatal care clinic in metropolitan Lagos, Nigeria. *Journal of Vector Borne Disease* 3(1),6-7.
- Isah Ibrahim, G. (2015). Zamfara APGA lauds INEC for credible elections. Retrieved from <http://www.nrguardiannews.com>
- Jesse Uneke, C. (2012). Malaria during pregnancy: Incidence, manifestations, therapy, and prevention. *Current Women's Health Reviews*, 8(4), 326-336.
- Jimoh, A., Sofola, O., Petu, A., & Okorosobo, T. (2007). Cost effectiveness and resource allocation. *Open Access journal*, 5, 6 doi:10.1186/1478-7547-5-6
- Kanhutu, K., & Torda, A. (2011). Travel and pregnancy: An infectious diseases perspective. *Obstetric Medicine*, 4(2), 53-58.
- Kayentao, K., Garner, P., van Eijk, A. M., Naidoo, I., Roper, C., Mulokozi, A., Doumbo, O. K. (2013). Intermittent preventive therapy for malaria during pregnancy using 2 vs 3 or more doses of sulfadoxine-pyrimethamine and risk of low birth weight in Africa: Systematic review and meta-analysis. *Jama*, 309(6), 594-604.

- Kilian, A., Koenker, H., Baba, E., Onyefunafoa, E. O., Selby, R. A., Lokko, K., & Lynch, M. (2013). Universal coverage with insecticide-treated nets-applying the revised indicators for ownership and use to the Nigeria 2010 Malaria Indicator Survey data. *Malar J*, 12(1), 314.
- Kovacs, S. D., Rijken, M. J., & Stergachis, A. (2015). Treating Severe Malaria in Pregnancy: A Review of the Evidence. *Drug safety*, 38(2), 165-181.
- Lengeler, C. (2004). Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev*, Issue 2. Art. No.: CD000363. doi: 10.1002/14651858.CD000363.pub2.
- Lorenzetti, L. (2015). World's first malaria vaccine approved by European Regulators. Retrieved from www.fortune.com
- Malhotra, I., Dent, A., Mungai, P., Wamachi, A., Ouma, J. H., Narum, D. L., King, C. L. (2009). Can prenatal malaria exposure produce an immune tolerant phenotype?: A prospective birth cohort study in Kenya. *PLoS Medicine*, 6(7), e1000116.
- Marchant, T., Nathan, R., Jones, C., Mponda, H., Bruce, J., Sedekia, Y., Hanson, K. (2008). Individual, facility and policy level influences on national coverage estimates for intermittent preventive treatment of malaria in pregnancy in Tanzania. *Malar J*, 7, 260. doi:10.1186/1475-2875-7-260.
- McGready, R., Davison, B. B., Stepniewska, K., Cho, T., Shee, H., Brockman, A., Meshnick, S. R. (2004). The effects of *Plasmodium falciparum* and *P. vivax* infections on placental histopathology in an area of low malaria transmission. *The American Journal of Tropical Medicine and Hygiene*, 70(4), 398-407.
- Menendez, C., Fleming, A., & Alonso, P. (2000). Malaria-related anaemia. *Parasitology Today*, 16(11), 469-476.
- Meritnation. (2015). Life cycle of mosquito in malaria. Retrieved from www.meritnation.com
- Mofolorunsho, C., Audu, H., & Omatola, A. (2014). Prevalence of malaria among pregnant women attending a healthcare facility in Lokoja, North-Central, Nigeria. *Asian Journal of Pharmaceutical & Health Sciences*, 4(1), 936.
- Mutagonda, R. F. (2012). A study of pregnant women and health workers knowledge on malaria prevention and treatment guidelines during pregnancy (Doctoral dissertation, Muhimbili University of Health and Allied Sciences).
- Nankwanga, H. A. A., & Gorette, N. (2008). Adherence to intermittent preventive treatment for malaria in pregnancy. *African Journal of Midwifery and Women's Health*, 2(3), 131-141.
- National Population Commission (2012). National malaria control programme, and ICF international. 2012. *Nigeria Malaria Indicator Survey 2010*.

- Nosten, F., McGready, R., & Mutabingwa, T. (2007). Case management of malaria in pregnancy. *The Lancet Infectious Diseases*, 7(2), 118-125.
- Nwagha, U. I., Ugwu, V. O., Nwagha, T. U., & Anyaehie, B. U. (2008). Asymptomatic plasmodium parasitaemia in pregnant Nigerian women: Almost a decade after roll back malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 103(1), 16-20.
- Nwizu, E., Iliyasu, Z., Ibrahim, S., & Galadanci, H. (2011). Socio-demographic and maternal factors in anaemia in pregnancy at booking in Kano, Northern Nigeria. *African Journal of Reproductive Health*, 15(4), 33-41.
- Nzeako, S. O., Nduka, F. O., & Origie, O. A. (2013). Prevalence of malaria in pregnant women attending ante natal care at university of Portharcourt primary health care centre Aluu, Portharcourt, Rivers State, Nigeria. *International Journal of Scientific Research in Environmental Sciences*, 1(10), 268.
- Okafor, U., Oguonu, T., & Onah, H. (2006). Risk factors associated with congenital malaria in Enugu, South Eastern Nigeria. *Journal of Obstetrics & Gynecology*, 26(7), 612-616.
- Okpere, E., Enabudoso, E., & Osemwenkha, A. (2010). Malaria in pregnancy. *Nigerian Medical Journal*, 51(3), 109.
- Oladeinde, B. H., Omoregie, R., Odia, I., & Oladeinde, O. B. (2012). Prevalence of malaria and anemia among pregnant women attending a traditional birth home in Benin city, Nigeria. *Oman Medical Journal*, 27(3), 232-236.
- Omoti, C. E., Ojide, C. K., Lofor, P. V., Eze, E., & Eze, J. C. (2013). Prevalence of parasitemia and associated immunodeficiency among HIV-malaria co-infected adult patients with highly active antiretroviral therapy. *Asian Pacific Journal of Tropical Medicine*, 6(2), 126-130.
- Onoka, C. A., Hanson, K., & Onwujekwe, O. E. (2012). Low coverage of intermittent preventive treatment for malaria in pregnancy in Nigeria: Demand-side influences. *Malar J*, 11(82), S1-S8.
- Qp ywmc."G0"E0*4228+0"Cpqv jgt"nqqm"cv"vjg"ko rcev"qh"Pki gtkcøu"itqy kpi"rqrwncvkqp" qp" vjg" Eqwpvt {øu" fgxgnqr o gpv0 *African Population Studies/Etude De La Population Africaine*, 21(1), 1-18.
- Onyenekwe, C. C., Meludu, S. C., Dioka, C. E., & Salimonu, L. S. (2002). Prevalence of asymptomatic malaria parasitaemia amongst pregnant women. *Indian Journal of Malariology*, 39(3-4), 60-65.
- Opiyo, P., Mukabana, W. R., Kiche, I., Mathenge, E., Killeen, G. F., & Fillinger, U. (2007). An exploratory study of community factors relevant for participatory malaria control on Rusinga Island, Western Kenya. *Malaria Journal*, 6, 48. doi:10.1186/1475-2875-6-48.

- Ordinioha, B. (2012). The use and misuse of mass distributed free insecticide-treated bed nets in a semi-urban community in Rivers State, Nigeria. *Annals of African Medicine*, 11(3), 163-168.
- Oyefabi, A., Sambo, M., & Sabitu, K. (2015). Effect of primary health care workers training on the knowledge and utilization of intermittent preventive therapy for malaria in pregnancy in Zaria, Nigeria. *Journal of Medicine in the Tropics*, 17(1), 4-11.
- Peat, J., & Barton, B. (2005). Continuous variables: Descriptive statistics. *Medical statistics: A guide to data analysis & critical appraisal* (pp. 31-32)
- Perkins, D. J., Were, T., Davenport, G. C., Kempaiah, P., Hittner, J. B., & Ong'echa, J. M. (2011). Severe malarial anemia: Innate immunity and pathogenesis. *International Journal of Biological Sciences*, 7(9), 1427-1442.
- Perlmann, P., & Troye-Blomberg, M. (2000). Malaria blood-stage infection and its control by the immune system. *Folia Biologica*, 46(6), 210-218.
- Pluess, B., Tanser, F. C., Lengeler, C., & Sharp, B. L. (2010). Indoor residual spraying for preventing malaria. *The Cochrane Library*. doi: 10.1002/14651858.CD006657
- Polsa, P., Spens, K., Soneye, A., & Antai, I. (2011). Comparing the perceived quality of private and public health services in Nigeria. *Journal of Management Policy and Practice*, 12(7), 18-26.
- President's Malaria Initiative. (2014). *President's Malaria Initiative Nigeria, Malaria Operational Plan FY 2014*. Nigeria: USAID and CDC.
- President's Malaria Initiative. (2015). *President's Malaria Initiative Nigeria, Malaria Operational Plan FY 2015*. Nigeria: USAID and CDC.
- Radeva Petrova, D., Kayentao, K., ter Kuile, F. O., Sinclair, D., & Garner, P. (2014). Drugs for preventing malaria in pregnant women in endemic areas: Any drug regimen versus placebo or no treatment. *The Cochrane Library*. Issue 10. Art. No.: CD000169. doi: 10.1002/14651858.CD000169.pub3.
- Rai, N., & Abraham, J. (2012). Different clinical features of malaria. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 2(12), 28-31.
- Rai, R. K., Singh, P. K., & Singh, L. (2012). Utilization of maternal health care services among married adolescent women: Insights from the Nigeria Demographic and Health Survey, 2008. *Women's Health Issues*, 22(4), e407-e414.
- Ribera, J. M., Hausmann-Muela, S., D'Alessandro, U., & Grietens, K. P. (2007). Malaria in pregnancy: What can the social sciences contribute? *PLoS Medicine*, 4(4), e92. doi:10.1371/journal.pmed.0040092.

- Rich, S. M., Leendertz, F. H., Xu, G., LeBreton, M., Djoko, C. F., Aminake, M. N., Rosenthal, B. M. (2009). The origin of malignant malaria. *Proceedings of the National Academy of Sciences*, 106(35), 14902-14907.
- Rogerson, S. J. (2010). Malaria in pregnancy and the newborn. *Hot topics in infection and immunity in children VI* (pp. 139-152) Springer.
- Roll Back Malaria. (2013). Roll back malaria (RBM) partnership programme- the goal frame work for coordinated action against malaria. Retrieved from www.rollbackmalaria.org
- Sabin, L. L., Rizal, A., Brooks, M. I., Singh, M. P., Tuchman, J., Wylie, B. J., Hamer, D. H. (2010). Attitudes, knowledge, and practices regarding malaria prevention and treatment among pregnant women in Eastern India. *The American Journal of Tropical Medicine and Hygiene*, 82(6), 1010-1016.
- Sangaré, L. R., Stergachis, A., Brentlinger, P. E., Richardson, B. A., Staedke, S. G., Kiwuwa, M. S., & Weiss, N. S. (2010). Determinants of use of intermittent preventive treatment of malaria in pregnancy: Jinja, Uganda. *PLoS One*, 5(11), e15066. doi:10.1371/journal.pone.0015066.
- Ucxcig." G0." Ou{codq|c." M0." Ikgu." U0." F0Cnguucpftq." W0." ("Dtcdkp." D0" *4229+0" Maternal anaemia as an indicator for monitoring malaria control in pregnancy in Sub Saharan Africa. *BJOG: An International Journal of Obstetrics & Gynaecology*, 114(10), 1222-1231.
- Schantz-Dunn, J., & Nour, N. M. (2009). Malaria and pregnancy: A global health perspective. *Reviews in Obstetrics and Gynecology*, 2(3), 186-192.
- Schlesselman, J., & Stolley, P. (1982). Sources of bias. *Case-Control Studies Design, Conduct, Analysis*. New York: Oxford, 124-143.
- Sharling, L., Enevold, A., Sowa, K. M., Staalsoe, T., & Arnot, D. E. (2004). Antibodies from malaria-exposed pregnant women recognize trypsin resistant epitopes on the surface of plasmodium falciparum-infected erythrocytes selected for adhesion to chondroitin sulphate A. *Malaria Journal*, 3, 31. doi:10.1186/1475-2875-3-31.
- Singh, R., Musa, J., Singh, S., & Ebere, U. V. (2014). Knowledge, attitude and practices on malaria among the rural communities in Aliero, Northern Nigeria. *Journal of Family Medicine and Primary Care*, 3(1), 39-44. doi: 10.4103/2249-4863.130271.
- Steketee, R. W., Nahlen, B. L., Parise, M. E., & Menendez, C. (2001). The burden of malaria in pregnancy in malaria-endemic areas. *The American Journal of Tropical Medicine and Hygiene*, 64(1 suppl), 28-35.
- Suresh, K., & Chandrashekhara, S. (2012). Sample size estimation and power analysis for clinical research studies. *Journal of Human Reproductive Sciences*, 5(1), 7613. <http://doi.org/10.4103/0974-1208.97779>.

- Vcmgo."G0" P0." (" F0Cnguucpf tq."W0"*4235+0" Ocncctk"kp"Rtgi pcpe{0 *Mediterranean Journal of Hematology and Infectious Diseases*, 5(1), e2013010. <http://doi.org/10.4084/MJHID.2013.010>.
- Taylor, G. M., Murphy, E., Hopkins, R., Rutland, P., & Chistov, Y. (2007). First report of mycobacterium bovis DNA in human remains from the iron age. *Microbiology*, 153(4), 1243-1249.
- Ter Kuile, F. O., Van Eijk, A. M., & Filler, S. J. (2007). Effect of sulfadoxine-pyrimethamine resistance on the efficacy of intermittent preventive therapy for malaria control during pregnancy. *JAMA: The Journal of the American Medical Association*, 297(23), 2603-2616.
- Thwing, J., Eisele, T. P., & Steketee, R. W. (2011). Protective efficacy of malaria case management and intermittent preventive treatment for preventing malaria mortality in children: A systematic review for the lives saved tool. *BMC Public Health*, 11(3), 1-9.
- Tillotson, G. S. (2012). Foreword: 10-year special edition. *Expert Review of Anti-Infective Therapy*, 10(11), 1219-1220.
- Umaru, M. L., & Uyaiabasi, G. N. (2015). Prevalence of malaria in patients attending the General Hospital Makarfi, Makarfi Kaduna State, North-western Nigeria. *American Journal of Infectious Diseases and Microbiology*, 3(1), 1-5.
- Warrell, D. (1997). Cerebral malaria: Clinical pathophysiology and treatment features. *Annals of Tropical Medicine and Parasitology*, 91(7), 875-884.
- Wikipedia, the free encyclopedia (2014). Zamfara State. Retrieved from www.en.wikipedia.org/wiki/zamfara_state
- World Health Organization. (2008). World Malaria Report 2008. *World Health Organization Geneva*. Available <http://www.who.int/malaria/publications/atoz/9789241563697/en/>
- World Health Organization. (2010). Malaria fact sheet no. 94. *World Health Organization Geneva*. Available <http://www.who.int/mediacentre/factsheets/fs094/en/>
- World Health Organization. (2012). World Malaria Report 2010. 2010. *World Health Organization Geneva*. Available http://www.who.int/malaria/world_malaria_report_2010/en/index.html
- World Health Organization. (2014). World Malaria Report 2013. *World Health Organization Geneva*. Available http://www.who.int/malaria/publications/world_malaria_report_2014/report/en/

Yosaatmadja, F., Andrews, K. T., Duffy, M. F., Brown, G. V., Beeson, J. G., & Rogerson, S. J. (2008). Characterization of VAR2CSA-deficient *Plasmodium falciparum*-infected erythrocytes selected for adhesion to the BeWo placental cell line. *Malaria Journal*, 7(1), 51. doi:10.1186/1475-2875-7-51.

Zamfara State Government. (2010). Map of Zamfara State. Retrieved from www.nigeriagallery.com/Nigeria/States_Nigeria/Zamfara_State.html

Zamfara State, N. (2000). Approved recurrent and capital estimates. Zamfara State. Retrieved from https://books.google.com.my/books/about/Approved_Recurrent_and_Capital_Estimates.html

Zinga, M. M. (2013). Malaria parasitaemia and associated preventive factors among pregnant women in Nyamagana and Misungwi districts, Mwanza region, 2012 (Doctoral dissertation, Muhimbili University of Health and Allied Sciences).