

UNIVERSITI PUTRA MALAYSIA

PARTIAL PURIFICATION AND CHARACTERIZATION OF GLUTATHIONE S-TRANSFERASE FROM THE LIVERS OF MALAYSIAN CATFISH (CLARIAS BATRACHUS) AND MALAYSIAN RED TILAPIA (OREOCHROMIS SP.)

CHIN CHEE MENG.

FBSB 2005 32



PARTIAL PURIFICATION AND CHARACTERIZATION OF GLUTATHIONE S-TRANSFERASE FROM THE LIVERS OF MALAYSIAN CATFISH (CLARIAS BATRACHUS) AND MALAYSIAN RED TILAPIA (OREOCHROMIS SP.)



Ву

CHIN CHEE MENG

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

July 2005



UPM

C)

Dedicated to my parents



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science

PARTIAL PURIFICATION AND CHARACTERIZATION OF GLUTATHIONE S-TRANSFERASE FROM THE LIVERS OF MALAYSIAN CATFISH (CLARIAS BATRACHUS) AND MALAYSIAN RED TILAPIA (OREOCHROMIS SP.)

By

CHIN CHEE MENG

July 2005

Chairman : Professor Nor Aripin Shamaan, PhD

Faculty : Biotechnology and Biomolecular Sciences

The present study is concerned mainly with partial purification and characterization of cytosolic GST from livers of Malaysian catfish and Malaysian red tilapia as both species are of economic and commercial importance in Malaysia being the major freshwater fishes consumed. This study hopes to establish the patterns of cytosolic glutathione S-transferase isoenzymes in the catfish and red tilapia. This may be useful to further achieve an understanding toward this enzyme in view of using it as a tool in environmental monitoring. The hepatic GST enzyme from catfish and red tilapia was partially purified 15X and 27X respectively in comparison to the ultra-centrifuged cytosolic fraction by affinity chromatography. Specific GST activity of 12.69 unit/mg protein and 33.42 unit/mg protein was obtained from livers of catfish and red tilapia using 1-chloro-2,4-dinitrobenzene (CDNB) as a substrate. Isolation of GST isoenzymes from affinity purified fractions was achieved by preparative isoelectric-focusing. Two isoenzymes; one major isoenzyme designated Ci1 and one minor isoenzyme

designated Ci2 were isolated from catfish liver with an apparent pl of 6.20 and 8.73 respectively. One isoenzyme designated Ti1 was isolated from red tilapia liver with an apparent pl of 9.14. SDS-PAGE analysis suggests that the isolated isoenzymes appear to be homodimeric in nature with subunit molecular weight of 29.7 ±1.7 kDa (Ci1), 27.7 ±1.3 kDa (Ci2) and 29.9 ±0.9 kDa (Ti1). As has been found for most GSTs, highest catalytic activity was obtained with CDNB. With the exception of the isoenzyme from cytosol and affinity purified fractions of red tilapia, none of the catfish fractions displayed enzyme activities towards 1,2dichloro-4-nitrobenzene (DCNB) and ethacrynic acid (EA). Therefore, both catfish and tilapia possess hepatic glutathione S-transferase activity, indicating that they capable of conjugating endogenous xenobiotic are or metabolites/compounds as a result of foreign exposure or oxidative metabolism with glutathione, thereby making it a useful tool as a effective biomarker of aquatic contamination.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

PURIFIKASI SEPARA DAN PENCIRIAN ENZIM GLUTATHIONE S-TRANSFERASE DARIPADA HATI IKAN KELI MALAYSIA (CLARIAS BATRACHUS) DAN IKAN TILAPIA MERAH MALAYSIA (OREOCHROMIS SP.)

Oleh

CHIN CHEE MENG

July 2005

Pengerusi : Profesor Nor Aripin Shamaan, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Kajian ini mengenai purifikasi separa dan pencirian GST sitosol daripada ikan keli Malaysia dan ikan tilapia merah Malaysia disebabkan kedua-dua spesis tersebut mempunyai kepentingan ekonomi dan komersial di Malaysia memandangkan ia adalah ikan air tawar yang sering dimakan. Kajian ini diharapkan dapat menentukan corak isoenzim sitosol GST dalam hati ikan keli dan tilapia merah. Ini mungkin berguna untuk mencapai pemahaman ke atas enzim ini dimana ia boleh dijadikan sebagai alat untuk memantau persekitaran. Enzim GST hepatik daripada ikan keli dan tilapia merah telah disepara-tulen sebanyak 15X dan 27X masing-masing dibandingkan dengan fraksi sitosol ultracentrifuged oleh kromatografi afiniti. Aktiviti spesifik enzim yang diperolehi adalah 12.69 unit/mg dan 33.42 unit/mg daripada hati ikan keli dan tilapia merah dengan menggunakan 1-chloro-2,4-dinitrobenzene (CDNB) sebagai substrat. Pemencilan isoenzim GST daripada fraksi yang ditulenkan dengan kromatografi afiniti dicapai dengan menggunakan "preparative"



isoelectric-focusing". Dua isoenzim: satu isoenzim major Ci1 dan satu isoenzim minor Ci2 dipencilkan daripada hati ikan keli dengan nilai pl 6.20 dan 8.73 masing-masing. Satu isoenzim Ti1 dipencilkan daripada hati ikan tilapia merah dengan nilai pl 9.14. Analisis SDS-PAGE mencadangkan isoenzim-isoenzim yang dipencilkan mempunyai struktur homodimer dengan berat molekul subunit 29.7 ±1.7 kDa (Ci1), 27.7 ±1.3 kDa (Ci2) dan 29.9 ±0.9 kDa (Ti1). Seperti yang ditemui dengan kebanyakan GST yang lain, aktiviti katalitik yang tertinggi diperolehi dengan CDNB. Dengan pengecualian isoenzim daripada sitosol dan fraksi yang ditulen oleh kromatografi afiniti daripada ikan tilapia merah, tiada fraksi ikan keli yang menunjukkan aktiviti dengan 1,2-dichloro-4-nitrobenzene (DCNB) dan ethacrynic acid (EA). Oleh itu, kedua-dua ikan keli dan tilapia mempunyai aktiviti glutathione S-transferase, menunjukkan bahawa mereka mampu mengkonjugasikan metabolik/bahan endogenus atau xenobiotik akibat daripada terdedah kepada persekitaran atau metabolisma oksidatif dengan glutathione, oleh itu membuatkannya alat yang berguna sebagai petanda biologi untuk pencemaran akuatik.



ACKNOWLEDGEMENTS

First and foremost I would like to express my heartfelt gratitude and appreciation to my supervisor, Prof. Nor Aripin Shamaan for his advice, guidance, ideas and critical discussions throughout the course of this project. He was always available for consultation even after office hours and has never failed to give an encouraging word when in need. Special thanks are extended to the other members of the Supervisory Committee, Dr. Yunus and Encik Ismail Omar for their ideas, advice and guidance given whenever sought.

I would also like to thank Encik Ibrahim and Encik Shamsudin for their help in preparing the chemicals and equipments needed during the course of the project. Not forgetting also to my lab members; Eddie Chia, Thong Chuan, Che Wan, Lan, Kak Neni, and Nina for their cooperation, companionship in the lab till the wee hours of morning and their help in supplying chemicals and lab equipments. A million thanks to them!

I wish to thank the Ministry of Science and Technology for the IRPA grant that funded this project and also for the PASCA scholarship given to me for two years.

A big thank you is also expressed to my parents for their encouragement and financial support to me throughout my study.



TABLE OF CONTENTS

		Page
DEDICATIO ABSTRACT ABSTRAK ACKNOWLE APPROVAL DECLARAT LIST OF TA LIST OF FIG LIST OF AB	N EDGEMENT ION BLES BURES BREVIATIONS	ii iii vii viii x xiii xiv xvi
CHAPTER		
-	INTRODUCTION Objectives	19 22
	LITERATURE REVIEW The Enzymes of Detoxification Glutathione Mercapturic acid Glutathione S-Transferase The Discovery of Glutathione S-Transferase Isoenzymes of Glutathione S-Transferase Cellular location of Glutathione S-Transferase Nomenclature The Final Classification The Substrates of Glutathione S-Transferase Assay Methods Spectrophotometric Method Titrimetric Method Colorimetric Method Radioisotope Method Thiolysis Method Purification Studies Properties of Glutathione S-Transferase in Fish The Structure of Glutathione S-Transferase	22 23 24 26 26 33 46 37 42 43 48 49 49 51 55 60 66 8
	Inhibition of Glutathione S-Transferase Activity Glutathione S-Transferase as Biomarkers The Multifunctional Nature of Glutathione S-Transferase Evolution of Glutathione S-Transferase	71 72 73 79

111	METHODOLOGY	81
	Tissue Samples	81
	Determination of the Enzymatic Activity	81
	Measurement of Protein Concentration	83
	Preparation of the Cytosol	83
	Cytosol Purification by Glutathione-Agarose Affinity	
	Chromatography	84
	Isoelectric-focusing	85
	Denaturing SDS-Polyacrylamide Gel Electrophoresis	88
	Determination of Substrate Specificity	89
IV	RESULTS	90
	Preparation of Cytosol	90
	Purification of Cytosol by Affinity Chromatography	91
	Effects of High Concentration of Salt	95
	Isoelectric-focusing	98
	Repeat of the Purification Run	104
	SDS-PAGE Analysis	109
	Substrate Specificity	115
	Summary of Purification and Characterization of	
	Glutathione S-Transferase From the Livers of Catfish	
	and Red Tilapia	117
v	DISCUSSION	119
		400
VI	CONCLUSION	132
BIBILIO	BRAPHY	134
APPEND		159
BIODAT	A OF THE AUTHOR	160



 Θ

LIST OF TABLES

Table		Page
1	Human members of the MAPEG family	29
2	Classification of rat glutathione S-transferases	41
3	Conditions for spectrophotometric assays	50
4	Quaternary structure of rat GST	
5	Formation of the light and dense pH gradient	87
6	Preparation of catfish cytosol	90
7	Preparation of red tilapia cytosol	90
8	Purification of catfish cytosol by affinity chromatography	91
9	Purification of red tilapia cytosol by affinity chromatography	
10	Effect of high salt concentration on the purification	
11	Purification of catfish cytosol by affinity chromatography (repeat experiment)	104
12	Purification of red tilapia cytosol by affinity chromatography (repeat experiment)	104
13	GST activities of isolated isoenzymes and isolated electro-focused peaks from affinity purified samples and cytosol fractions from catfish and red Tilapia toward different substrates	
14	Summary of purification procedures and quantitative data of hepatic glutathione S-transferases from The Malaysian Catfish (<i>Clarius batrachus</i>) and Malaysian Red Tilapia (<i>Oreochromis sp.</i>)	118



 \bigcirc

LIST OF FIGURES

Figure		Page
1	The mercapturic acid pathway for urinary excretion of glutathione conjugate	27
2	Affinity-chromatographic purification of catfish livers on glutathione-agarose matrix	92
3	Affinity-chromatographic purification of red tilapia livers on glutathione-agarose matrix	93
4	Affinity-chromatographic purification of catfish livers on glutathione-agarose matrix using 1M NaCl eluting buffer	97
5	Isoelectric-focusing profile of affinity purified samples from catfish liver	100
6	Isoelectric-focusing profile of affinity purified samples from red tilapia liver	101
.7	Isoelectric-focusing profile of cytosol samples from catfish liver	102
8	Is <mark>oelectric-focusing profile of cytosol s</mark> amples from red tilapia liver	103
9	Affinity-chromatographic purification (repeat experiment) of catfish liver on glutathione-agarose matrix	106
10	Affinity-chromatographic purification (repeat experiment) of red tilapia liver on glutathione-agarose matrix	107
11	Isoelectric-focusing profile of affinity purified samples (repeat experiment) from catfish liver	108
() 12	SDS-PAGE analysis of electrofocused samples from affinity purified pooled fractions of catfish and red tilapia livers	110
13	SDS-PAGE analysis of electrofocused samples from cytosolic fractions of catfish liver	111



14	SDS-PAGE analysis of electrofocused samples from cytosolic fractions of red tilapia liver	112
15	SDS-PAGE analysis of electrofocused samples from affinity purified pooled fractions (first run and repeat experiment) of catfish and (repeat experiment) of red tilapia livers	114





 \bigcirc

LIST OF ABBREVIATIONS

	α	alpha
	β	beta
	Br	bromine
	BSA	bovine serum albumin
	CDNB	1-chloro-2,4-dinitrobenzene
	CI	chlorine
	C-terminal	carboxyl-terminal
	Cys	cysteine
	Δε	molar extinction coefficient
	DCNB	1,2-dichloro-4-nitrobenzene
	DDT	dichlorodiphenyltrichloroethane
	DNA	deoxyribonucleic acid
	3	epsilon
	E.C.	Enzyme Commission
	EA	ethacrynic acid
	EDTA	ethylenediaminetetraacetic acid
	EST	expressed sequence tag
	GDP	gross domestic product
	Glu	glutamate
	Gly	glycine
	GS ⁻	glutathione thiolate ion
	GSH	reduced glutathione





G-site	glutathione binding site	
GST	glutathione S-transferase	
HPLC	high performance liquid chromatography	
H-site	hydrophobic binding site	
I	iodine	
KCI	potassium chloride	
kDa	kilo Dalton	
Μ	molar	
MAPEG	Membrane Associated Proteins in Eicosanoid and Glutathione metabolism	
mol	mole	
мров	Malaysian Palm Oil Board	
mRNA	messenger RNA	
MVV	molecular weight	
NaCl	natrium chloride	
N-terminal	amino-terminal	
θ	theta	
ρ	rho	
PhIP	2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine	
RNA	ribonucleic acid	
SDS-PAGE	PAGE sodium dodecyl sulfate polyacrylamide gel electrophoresis	
-SH	sulfurhydryl group	
π	pi	
TEMED	N, N, N', N'-tetramethylethylenediamine	



G

μ	mu
U	unit
UPM	Universiti Putra Malaysia
Y	gamma





CHAPTER I

INTRODUCTION

The Malaysian fisheries sector plays an important role in providing food and as a source of protein to the general public. In 2000, it contributed about 1.60% to the national Gross Domestic Product (GDP)^a and provided direct employment to 82,000 fishermen and 22,000 fish culturists. With the recent episodes of Nipah virus infecting pigs and the bird flu attack in chickens, more people are turning to other sources of proteins such as fish. Therefore, the Department of Fisheries is taking several measures in order to provide a sustained production of fish in the country. Although deep sea fishing still remains as the easiest method for fish production, however water pollution and over-fishing are taking its toll on the fish population in the wild causing shortages and resulting in the increasing prices of fish. Therefore, the real potential lies in the aquaculture industry in order to achieve a sustainable growth in fish production.

Freshwater aquaculture contributes a significant percentage (30.2%) to the total aquaculture production in 2000. Some of the major freshwater species cultured were freshwater Malaysian Catfish (*Clarius batrachus*) and Red Tilapia (*Oreochromis sp.*).

^a National GDP RM 339.42 billion (Statistics Handbook, Statistics Department Malaysia, 2001



The catfish are demersal creatures with the ability to survive in muddy and polluted water. Generally, they are found in rivers and swampy areas. They are omnivorous in nature surviving on water insects, worms, shrimps, snails and rotting materials such as plants. They have a special breathing system which allows them to survive for short periods of time on land. In Malaysia the Malaysian catfish is usually mated with the African catfish (*Clarias gariepinus*) as the African catfish displays higher growth rate and can attain a size of 1 kg in 6 months. The red tilapia can be found in freshwater, saltwater and also brackish water bodies. They are also omnivorous, surviving on a diet of worms, plants and small fish (Basir and Daud, 1994).

As the two species represent an economic importance in Malaysia, maximum yield during the breeding and production are of utmost importance to maximize profits. Therefore, there is a need for a monitoring system to monitor the state of the fish habitat and the stress levels of the fish (i.e overcrowding, parasite infections).

Glutathione S-Transferases (GSTs, E.C. 2.5.1.18) are a multigene family of multifunctional enzymes which can be found in bacteria, yeasts, higher plants, mollusks, crustaceans, insects, fish, amphibians and mammals (Jakoby, 1978; Mannervik, 1985b). They catalyze the nucleophilic attack of the sulfur atom of glutathione on compounds possessing an electrophilic center (Hayes and Pulford, 1995). Therefore, GSTs play a key role in the cellular detoxification process in by conjugating a broad spectrum of endobiotic and xenobiotic electrophilic substrates (as a result of



20

environmental exposure or oxidative metabolism) with intracellular glutathione into more readily excretable products (Mannervik and Danielson, 1988; Hayes and Pulford, 1995; Armstrong, 1997).

Certain GST isoenzymes display selective response towards pollutants and high activity towards certain substrates, making it a useful tool as an effective biomarker of aquatic contamination (Egaas et al., 1999; Gallagher et al., 1996).





Objectives

The objectives of the present study are to:-

- 1. Partially purified the cytosolic glutathione S-transferase (E.C. 2.5.1.18) from the livers of Malaysian catfish (*Clarius batrachus*) and red tilapia (*Oreochromis sp.*)
- 2. Partially characterized the cytosolic glutathione S-transferase from the livers of Malaysian catfish (*Clarius batrachus*) and red tilapia (*Oreochromis sp.*)

Ultimately, this study hopes to establish the patterns of the cytosolic glutathione S-transferases (i.e. the dimeric structures, specific activities, pl values, molecular weight and substrate specificity) in the catfish and red tilapia. This may be useful to further achieve an understanding toward this enzyme in view of using it as a tool in environmental monitoring.



CHAPTER II

LITERATURE REVIEW

The Enzymes of Detoxification

All living organism are continuously subjected to foreign external (xenobiotics) compounds; naturally occurring or man-made such as chemicals in the air, water, food additions and drugs or endogenous toxicants produced as a result of the metabolism of hormones, steroids and oxidative metabolism. Most of these compounds are lipophilic and might be excreted slowly from the body. Therefore, the toxicity of the compounds is increased as a result of the accumulation in the tissues of the organism. Given these exposures, the ability to detoxify is important to the overall health and the survivability of the organism.

The process of detoxification through which the non-polar (lipophilic) toxins are converted to polar (hydrophilic) compounds that can be readily excreted by the organism occurs in two classical steps known as Phase 1 and Phase 2. Each of the phases involved the so-called detoxification enzymes that have broad specificity towards a wide variety of toxins faced by the organism. The two phases are interconnected; Phase 1 reactions normally generate intermediate end-products that are toxic to the tissues. Phase 2 reactions conjugates the end-products of Phase 1 reactions to yield final products that are eliminated from the organism. Phase 1 reactions are characterized by the cytochrome P450 supergene family of isoenzymes that catalyzed the addition



of a reactive group to the substrate (toxins). However the product yields a compound that is more toxic compared to the parent compound. Furthermore, the reactions of Phase 1 enzymes generate very reactive damaging free radicals. Therefore it is necessary that this compound is further metabolized by the Phase 2 enzymes to avoid damage of the organism's proteins and genes. Phase 2 reactions consist of conjugating enzymes that normally adds a water-soluble molecule to the active site of the Phase 1 metabolite to facilitate its removal in urine or bile. These conjugates are usually less toxic and more water-soluble than their precursors. However, not all toxins undergo detoxification using the Phase 1 to Phase 2 route. Occasionally the parent toxins undergo detoxification directly through the Phase 2 route bypassing the Phase 1 route. Of all the Phase 2 enzymes, glutathione S-transferases (GSTs E.C. 2.5.1.18), the most abundant protein in the cytosolic fraction of the liver (Booth et al., 1961; Wilce and Parker, 1994), is the most widely studied and major Phase 2 detoxification enzyme catalyzing the conjugation of cellular nucleophile glutathione with a wide range of endogenous or xenobiotic hydrophobic molecules (Armstrong, 1997; Hayes and Pulford, 1995; Mannervik and Danielson, 1988).

Glutathione

Reduced glutathione (GSH), a tripeptide consisting of γ -glutamyl-cysteinylglycine can be conjugated in the initial step of mercapturic acid synthesis (George, 1994). Its cysteinyl residue provides a nucleophilic thiol important for the detoxication of electrophilic metabolites. Its net negative charge and overall hydrophilicity increases the aqueous solubility of the lipophilic

