



UNIVERSITI PUTRA MALAYSIA

***PRECIPITATION TREND ANALYSIS FOR THE LANGAT RIVER BASIN,
SELANGOR, MALAYSIA***

NARGES PALIZDAN

FK 2014 75



**PRECIPITATION TREND ANALYSIS FOR THE LANGAT RIVER BASIN,
SELANGOR, MALAYSIA**

By

NARGES PALIZDAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

April 2014

COPYRIGHT

All material contained within the thesis, including without limitation text, logos icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of nay material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia

Copyright © Universiti Putra Malaysia



© COPYRIGHT UPM

DEDICATED

TO

My Mother

A strong and gentle soul who thought me to trust in Allah, believe in hard work and that so much could be done with little

My Father

For earning and honest living for us and for supporting and encouraging me to believe in myself

My Parents-in-law

For being my guardian, their support and encouragement

My Husband

Without whom none of my success would be possible

And finally my lovely Siblings

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement of the degree of Master of Science

**PRECIPITATION TREND ANALYSIS FOR THE LANGAT RIVER BASIN,
SELANGOR, MALAYSIA**

By

NARGES PALIZDAN

April 2014

Chairman: Professor Ir. Lee Teang Shui, PhD

Faculty: Engineering

Almost all the hydrological and meteorological variables such as rainfall and temperature etc are affected by global climate change. Precipitation is one significant climate factor that affects the state of the environment and contributes significantly towards the natural hydrological process. Its temporal fluctuation is significant both in terms of the scientific knowledge and practical applications, such as in water resources planning and management. It is necessary to evaluate the regional precipitation trend to improve water management strategies for a given region. A study was conducted to analyse the annual and seasonal precipitation trends of 30 stations in the Langat River Basin, located in the state of Selangor, Malaysia, at both the stationary and on regional scales. On an at site scale the rainfall trends were detected using the Mann-Kendall (MK) test, the Linear Regression analysis for the period 1982-2011. The lag-1 approach was utilized at the 95% significance level to test the serial correlation of the data series. Then the autocorrelated time series were pre-whitened. On the annual scale, it was found that most of the stations in the basin were characterized with insignificant precipitation trends. The significant trends were found only at the four stations, namely stations 44301, 44305, 44320 and 2719001. The results of the seasonal precipitation trend analysis showed that most of the stations, during the North East Monsoon (NEM) and Inter Monsoon 1 (INT1) seasons, and half of the stations in South West Monsoon (SWM) season, experienced insignificant positive trends. To the contrary, for the Inter Monsoon 2 (INT2) season, the majority of the stations showed negative trends. It was found that during NEM season the station 44301, for INT1 the stations 44301, 2719001 and 3118069 were established as having significant changes, while in SWM, the station 2917001 and during INT2 the stations 2615131 and 44301 showed significant trends.

In the regional study, the regional annual and seasonal precipitation trends for the period 1982-2011 were examined at the 95% level of significance, using the Regional Average Mann-Kendall (RAMK) test and the Regional Average Mann-Kendall coupled with bootstrap (RAMK-bootstrap) method. In order to identify the homogeneous regions respectively for the annual and seasonal scales, firstly, at-site mean total annual and separately at site mean total seasonal precipitation were

spatialized into 5 km × 5 km grids using the inverse distance weighting (IDW) algorithm. Next, the optimum number of homogeneous regions (clusters) is computed using the Silhouette Coefficient approach. Next, the homogeneous regions were then formed using the K-mean Clustering method. The applied homogeneous region analysis method in the present study is a new approach. From the annual scale perspective, all three regions showed positive trends. However, the application of the two methods at this scale showed a significant trend only in the region AC1. The region AC2 experienced a significant positive trend using only the RAMK test. On a seasonal scale, all regions showed insignificant trends, except the regions I1C1 and I1C2 in the Inter Monsoon 1 (INT1) season which experienced significant upward trends. In addition, it was proved that the significance of trends has been affected by the existence of serial and spatial correlations.

We also aimed to determine the most dominant periodic components that affect the trends in each homogeneous region for the period 1982-2011. The combination of wavelet analysis and Mann-Kendall test at regional scale was used for the first time in this study. In each region the areal precipitation series were computed using the Thiessen polygon method. In the method using wavelet transformation, the mother wavelet type and the number of decomposition levels were assessed based on two criteria. The first one is the Mean Relative Error (MRE) between the wavelet approximation component and the original data set and the second one is the relative error of the MK Z-value (e_r) between the approximation series and the original data series. The Discrete Wavelet Transform (DWT) coupled with the Mann-Kendall (MK) test and the sequential MK analysis were utilized in order to find out which time scale are affecting the trends observed in each homogeneous region. On annual scale it was found that DWT1 in region AC1 and AC2 and DWT2 in region AC3 were the periodic modes responsible for trends. On seasonal scale, in regions NC1, I1C1, I1C2, I1C3, SC1, SC2, I2C1 and I2C2, DWT1, in region NC2, DWT2 and DWT3, in region I2C3, DWT2 were the most influential periodicity for trends, respectively.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**ANALISIS KECENDERUNGAN HUJAN BAGI KAWASAN TADAHAN
SUNGAI LANGAT, SELANGOR, MALAYSIA**

Oleh

NARGES PALIZDAN

April 2014

Pengerusi: Professor Ir. Lee Teang Shui, PhD

Fakulti: Kejuruteraan

Hampir semua perubahan hidrologi dan meteorologi seperti hujan, suhu dan lain-lain dipengaruhi oleh perubahan iklim global. Hujan adalah satu faktor iklim yang ketara yang menjejaskan tahap persekitaran alam sekitar dan banyak menyumbang kepada proses semula jadi hidrologi. Turun naik semasanya adalah penting dari segi ilmu saintifik dan penggunaan praktikal seperti dalam bidang pengurusan dan perancangan sumber air. Keperluan untuk menilai trend taburan hujan adalah bagi meningkatkan strategi pengurusan air bagi sesebuah kawasan. Satu kajian telah dijalankan untuk menganalisa trend taburan hujan mengikut tahun dan musim bagi 30 stesen di kawasan tadahan Sungai Langat, Selangor, Malaysia, pada kedua-dua skala pegun dan kawasan tersebut. Pada skala di tapak, trend taburan hujan dikesan menggunakan Ujian Mann-Kendall (MK), Penaksir Cerun Sen dan Analisis Linear Regression bagi tempoh 1982-2011. Pendekatan Lag-1 diguna pada tahap kepentingan 95% untuk menguji korelasi bersiri pada siri data. Kemudian, auto-korelasi masa bersiri di pra-putihkan. Pada skala tahunan, kebanyakan stesen dalam kawasan tadahan disifatkan sebagai tidak cenderung kepada corak taburan hujan. Trend kecenderungan ketara ditemui hanya di empat stesen, iaitu stesen 44301, 44305, 44320 and 2719001. Keputusan analisis bagi corak taburan hujan bermusim menunjukkan kebanyakan stesen pada musim Monsun Timur Laut (NEM) dan Pertengahan Monsun 1 (INT1), dan sebahagian daripada stesen di musim Monsun Barat Daya (SWM) mengalami trend tidak cenderung positif. Sebaliknya, bagi musim Pertengahan Monsun 2 (INT2), majoriti stesen menunjukkan trend kecenderungan negatif. Didapati semasa musim NEM bagi stesen 44301 dan INT1 bagi stesen 44301, 2719001 dan 3118069 mengalami kecenderungan perubahan yang ketara, manakala pada SWM, stesen 2917001 dan INT2 di stesen 2615131 dan 44301 menunjukkan trend yang berkecenderungan ketara.

Dalam kajian kawasan serantau, trend taburan hujan tahunan dan bermusim bagi tahun 1982-2011 diperiksa pada tahap bermakna 95%, menggunakan ujian Regional Average Mann-Kendall (RAMK) dan Regional Average Mann-Kendall digabungkan dengan bootstrap (RAMK-bootstrap) method. Demi mengenal pasti kawasan homogen masing-masing bagi skala tahunan dan bermusim, pertamanya, jumlah

purata di kawasan tahunan dan jumlah purata kawasan bermusim secara berasingan di spatialkan kepada grid 5 km x 5 km menggunakan algoritma Inverse Distance Weighting (IDW). Kemudian, nombor optimum bagi kawasan homogenus (gugusan) dihitung menggunakan pendekatan Silhouette Coefficient. Selepas itu, kawasan homogenus dibentuk menggunakan kaedah K-mean Clustering. Daripada perspektif skala tahunan, ketiga-tiga kawasan menunjukkan trend yang positif. Walau bagaimanapun, penggunaan kedua-dua kaedah tersebut pada skala menunjukkan trend yang berkecenderungan ketara pada kawasan AC1. Kawasan AC2 mengalami trend kecenderungan ketara yang positif dengan hanya menggunakan ujian RAMK. Pada skala bermusim, semua kawasan menunjukkan trend kecenderungan tidak ketara, kecuali kawasan I1C1 dan I1C2 pada musim Pertengahan Monsun 1 (INT1) yang mengalami trend kecenderungan ketara menaik. Sebagai tambahan, terbukti bahawa trend kecenderungan ketara telah terjejas hasil daripada kewujudan korelasi bersiri dan spatial.

Kami juga berazam untuk menentukan komponen berkala yang paling dominan yang menjejaskan trend setiap kawasan homogenus bagi tahun 1982-2011. Di setiap kawasan bahagian hujan bersiri dikira menggunakan kaedah poligon Thiessen. Dalam kaedah menggunakan transformasi gelombang, gelombang jenis induk dan bilangan tahap penguraian dinilai berdasarkan dua kriteria. Kriteria pertama adalah Ralat Purata Relatif (MRE) antara komponen gelombang anggaran dan set data asal dan yang kedua adalah ralat relatif MK nilai-Z (e_r) antara siri anggaran dan siri data asal. Discrete Wavelet Transform (DWT) digandingkan dengan ujian Mann-Kendall (MK) dan analisis berurutan MK digunakan untuk mengetahui mana skala masa yang terkesan dengan trend yang diperhatikan di setiap kawasan yang homogenus. Pada skala tahunan didapati DWT1 di kawasan AC1 dan AC2 dan DWT2 di kawasan AC3 adalah trend mod berkala. Pada skala bermusim, di kawasan NC1, I1C1, I1C2, I1C3, SC1, SC2, I2C1 dan I2C2, DWT1 di kawasan NC2, DWT2 dan DWT3, di kawasan I2C3, DWT2 masing-masing merupakan trend jangka masa yang paling berpengaruh.

ACKNOWLEDGEMENTS

First of all, I would like to thank God, the almighty, for having made everything possible by giving me strength and courage to do this work.

I would like to express the deepest appreciation to my committee chair Professor Lee Teang Shui, for all the assistance and advice. He continually and persuasively conveyed a spirit of adventure in regard to research and scholarship, and an excitement in regard to teaching. Without his supervision and constant help this dissertation would not have been possible.

I would like to thank my committee members, Dr. Abdul Halim Ghazali and Dr. Huang Yuk Feng whose work demonstrated to me that concern for global affairs supported by an “engagement” in comparative literature and modern technology, should always transcend academia and provide a quest for our times.

Finally, to my caring, loving, and supportive husband, Yashar: my deepest gratitude. Your encouragement when the times got rough are much appreciated and duly noted.

I certify that a Thesis Examination Committee has met on 16 April 2014 to conduct the final examination of Narges Palizdan on her thesis entitled "Precipitation Trend Analysis for The Langat River Basin, Selangor, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Hasfalina binti Che Man, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Badronnisa binti Yusuf, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Aimrun Wayayok, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Chau Kwok-Wing, PhD

Professor
Hong Kong Polytechnic University
Hong Kong
(External Examiner)



NORITAH OMAR, PhD

Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 23 June 2014

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Lee Teang Shui, PhD

Professor.Ir
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Abdul Halim Bin Ghazali, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Huang Yuk Feng, PhD

Assistant Professor
Faculty of Engineering and Science
Universiti Tunku Abdul Rahman
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

Declaration by Graduate Student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003(Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: _____

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 202-2013) are adhered to:

Signature _____
Name of _____
Chairman of _____
Supervisory _____
Committee: _____

Signature _____
Name of _____
Member of _____
Supervisory _____
Committee: _____

Signature _____
Name of _____
Member of _____
Supervisory _____
Committee: _____

Signature _____
Name of _____
Member of _____
Supervisory _____
Committee: _____




TABLE OF CONTENTS

	Page
ABSTRACT	iii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xvii
LIST OF APPENDICES	xix
LIST OF ABBREVIATIONS	xx
LIST OF NOTATIONS	xxi
CHAPTER	
1	
INTRODUCTION	
1.1 Background	1
1.2 Problem Statement	3
1.3 Objectives	4
1.4 Significance of the Study	4
2	
LITERATURE REVIEW	
2.1 General	5
2.2 Common Methods to detect trend	5
2.3 Parametric and Non-Parametric Statistical Test	6
2.4 Non-Parametric methods	6
2.4.1 Sen's Slope Estimator	6
2.4.2 Spearman's rank correlation	7
2.4.3 Sen's T-test	7
2.4.4 Mann-Kendall Test	8
2.4.5 Sequential Mann-Kendall Test	8
2.4.6 Regional Average Mann-Kendall Test	9
2.5 Understanding resampling	9
2.6 Methods to investigate the serial correlation issue	10
2.6.1 Pre-whitening	10
2.6.2 Modified Mann Kendall Test for Autocorrelated data	11
2.6.3 Seasonal Mann-Kendall Test	11
2.7 Parametric Methods	14
2.7.1 Regression Analysis	14
2.8 Moving Average Method	16
2.9 Wavelet Analysis	17
2.10 Summary	18
3	
METHODOLOGY	
3.1 General	20
3.2 Study Area	20

	3.3 Data	21
	3.4 Homogeneity	24
	3.5 Normality	24
	3.6 Linear regression analysis	25
	3.7 Mann-Kendall (MK) Test	26
	3.8 Lag-1 serial correlation	27
	3.9 Sen's Slope Estimator	28
	3.10 Inverse Distance Weighting (IDW) Algorithm	29
	3.11 Silhouette coefficient	30
	3.12 K-mean Clustering Method	31
	3.13 Regional Average Mann-Kendall Test	32
	3.14 Bootstrap Method	33
	3.15 Thessien Polygon	33
	3.16 Wavelet analysis	34
	3.16.1 Discrete Wavelet Transform (DWT)	34
	3.17 Sequential Mann-Kendall test	36
4	RESULTS AND DISCUSSION	
	4.1 Results of Normality and Homogeneity Test	41
	4.2 Stationary Analysis	41
	4.2.1 Annual Trends	43
	4.2.2 Seasonal Trends	45
	4.3 Regional Analysis	49
	4.3.1 Homogenous Regions	49
	4.3.2 Regional trend test	57
	4.3.3 The effect of serial correlation and cross correlation	64
	4.4 Wavelet analysis	65
	4.4.1 Annual perspective	67
	4.4.2 Seasonal perspective	70
	4.4.3 Variation of sequential Mann-Kendall values of the original time series and the DWT models	73
5	CONCLUSIONS AND RECOMMENDATIONS	82
	5.1 Conclusion	82
	5.2 Recommendations	84
	REFERENCES	85
	APPENDICES	92
	BIODATA OF STUDENT	143
	PUBLICATIONS	144

LIST OF TABLES

Table	Page
2.1 The sample schema of data	7
3.1 The coordinates, elevation, name and codes of the selected stations	23
4.1 The results of homogeneity and normality test	42
4.2 The result of lag-1 serial correlation test and the magnitude of r-lag1 for annual and seasonal data series at each station	43
4.3 Values of Z in MK test, Qmed in Sen's slope estimator and b in linear regression analysis for annual data series (1982-2011)	45
4.4 Values of Z in MK test, Qmed in Sen's slope estimator and b in Linear Regression analysis for seasonal data series in NEM and INT1 seasons (1982-2011)	47
4.5 Values of Z in MK test, Qmed in Sen's slope estimator and b in Linear Regression analysis for seasonal data series in SWM and INT2 seasons (1982-2011)	48
4.6 Area and mean total precipitation of the homogeneous regions	57
4.7 The results of the RAMK test and the RAMK-bootstrap method in the Langat River Basin	58
4.8 Results of MK test at the selected stations in annual, NEM and INT1 scales	59
4.9 Results of MK test at the selected stations in SWM and INT2 scales	63
4.10 The amount of mean cross correlation and number of the stations with significant autocorrelation in each region	64

4.11	Amount of Mean Relative Error (MRE) and MK Z-value relative error in region AC1	67
4.12	Amount of Mean Relative Error (MRE) and MK Z-value relative error in region AC2	68
4.13	Amount of Mean Relative Error (MRE) and MK Z-value relative error in region AC3	68
4.14	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total Mean Square Error (MSE) between the sequential value of original series and different DWT models of the region AC1 precipitation series	69
4.15	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region AC2 precipitation series	69
4.16	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region AC3 precipitation series	70
4.17	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region NC2 precipitation series	71
4.18	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region I1C3 precipitation series	72
4.19	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region I2C1 precipitation series	72
4.20	Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region I2C2 precipitation series	72

4.21 Mann–Kendall (MK) test results of different DWT models and the Correlation (Co) and total mean Square error (MSE) between the sequential value of original series and different DWT models of the region I2C3 precipitation series

72



LIST OF FIGURES

Figure	Page
3.1 The Langat River Basin and the distribution of the selected stations	22
3.2 The K-Mean clustering method	31
3.3 The schematic of decomposition process	35
3.4 Stationary analysis flow chart	37
3.5 Homogeneous region analysis flow chart	38
3.6 Regional analysis flow chart	39
3.7 Wavelet analysis flow chart	40
4.1 The topographic homogeneous region map	50
4.2 The annual homogenous region map	52
4.3 The NEM homogenous region map	53
4.4 The INT1 homogenous region map	54
4.5 The SWM homogenous region map	55
4.6 The INT2 homogenous region map	56
4.7 The empirical bootstrap cdf curve for the region NC1 in the NEM season	61
4.8 The comparison chart for the results of the RAMK test and the RAMK–bootstrap method	65

4.9	The Sequential Mann – Kendall value test on (---) original precipitation series and (—) DWT models of AC1 region	74
4.10	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of AC2 region	75
4.11	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of AC3 region	76
4.12	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of NC2 region	77
4.13	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of I1C3 region	78
4.14	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of I2C1 region	79
4.15	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of I2C2 region	80
4.16	The Sequential Mann–Kendall value test on (---) original precipitation series and (—) DWT models of I2C3 region	81

LIST OF APPENDICES

Appendix	Page
A: The empirical bootstrap cdf curve	92
B: Amount of Mean Relative Error (MRE) and MK Z-value relative error	104
C: Original rainfall data (mm)	109
D: Sample Matlab codes for the analysis of the trends	139

LIST OF ABBREVIATIONS

cdf	Cumulative Distribution Function
Co	Correlation
CWT	Continues Wavelet Transform
db	Daubechies wavelet
DID	Department of Irrigation and Drainage
DWT	Discrete Wavelet Transform
EMA	Exponential Moving Average
ESS	Effective Sample Size
IDW	Inverse Distance Weighting
INT1	Inter Monsoon 1
INT2	Inter Monsoon 2
MAP	Mean Annual Precipitation
MK	Mann-Kendall
MMD	Malaysian Meteorology Department
MRE	Mean Relative Error
MSE	Mean Square Error
NEM	North East Monsoon
OLS	Ordinary Least Square
PMW	Pettitt Mann-Whitney
RAMK	Regional Average Mann-Kendall
RAMK-bootstrap	Regional Average Mann-Kendall coupled with bootstrap
SMA	Simple Moving Average
STFT	Short –Time Fourier Transformation
SWM	South West Monsoon

LIST OF NOTATIONS

n_a	Number of “a” in Run test
n_b	Number of “b” in Run test
U	Sum of series “a” and “b”
n	Number of data
b	Slope in Linear regression
x	Sample data
t	time
L_{XY}	Sum of deviation square of variable X
U	Regression sum of square in regression analysis
Q	Error sum of square in regression analysis
S	S statistic in MK test
Var(S)	Variance of S in MK test
Z	Standard normal variable
r_1	Lag-1 serial correlation coefficient
Y_t	Pre-whitened series
Q_i	Slope in Sen’s Slope Estimator
C_α	Confidence interval in Sen’s Slope Estimator
W	Weight function of IDW
P	Precipitation value at unmeasured point
S(i)	The silhouette coefficient at point i
SC	Silhouette coefficient
\bar{S}_m	Regional Average Kendall’s S in RAMK
m	Number of stations in RAMK
H_0	Null hypothesis

H_1	Alternative hypothesis
ψ	Mother wavelet
S_0	Dilation step
γ_0	Location variable
$u(t)$	Sequential value of MK test



CHAPTER 1

INTRODUCTION

1.1 Background

Precipitation is the resultant production in condensation of atmospheric water vapor that falls under gravity. It can also be defined as water in various forms that fall back to the earth from the atmosphere. There are different forms of precipitation; these include rain, snow, sleet, hail and mist. Precipitation is the most important factor that has a direct impact on hydrology cycle. The average amount of precipitation on the whole earth has been reported at 1050 mm, and this amount changes in different regions of the earth (Pidwirny, 2006). It can be zero in some deserts such as Sudan Desert in Africa to more than 12 meter in some places such as the Himalayan Heights. In each of these regions, there is a specific distribution of precipitation over time. For instance, the distribution of precipitation in wet areas is in a good order and the rain falls almost throughout the whole year. However, in arid and semi-arid areas this distribution can be unbalanced and it is possible that approximately 50 percent of annual precipitation occurs in just one event.

Water is the most valuable material on the earth and fresh water is the reason for human survival, where upon precipitation is the main source. There is no doubt that precipitation is one significant climate factor that affects the environmental and natural processes. Precipitation is an essential component of the hydrological cycle and any change in its distribution may have a significant effect on the hydrosphere, biosphere and the society at large. In addition, changes in the pattern of precipitation may lead to floods and droughts in different areas (Ahani et al., 2012). Precipitation plays an important role in the agricultural production; the significance of precipitation can be obviously seen when applied in dry land agriculture. This is because in dry land agriculture it is expected that the entire moisture required for plant is provided through precipitation, either through irrigation facilities or through direct rain fed options.

Changes in the hydro-meteorological series can take place in many different ways. A change can occur abruptly (step changes) or gradually (trend) or in more complex forms. In general, trend analysis is used to obtain information and in studying whether it is possible to detect a pattern or trend from the information. According to Rana et al. (2011) information about the trends of rainfall is important because it is connected to the water related issues of a region, its associated problems and environment and water management purposes. This information will be the most valuable to researchers studying climate change and its effects on water management. The detection of rainfall trends has been one of the main activities for both hydrologists and climatologists in pursuing climate change studies and research (Rana et al., 2011). Furthermore, studying the climate change needs the information of trends in different indices because climate change is continuously changing.

Climate is the pattern or cycle of weather conditions such as temperature, wind, rain, snowfall, humidity and clouds including extreme or occasional ones, over a large area and averaged over many years. Climate parameters can be affected by some factors such as place, the type and the area of green cover like forest and grassland, local land feature like mountains and human activities (burning fossil fuel, forming or cutting down forest). Climate change is a considerable change in weather patterns in the world over a long period of time. There are many factors that cause climate change including those oceanic processes like oceanic circulation, variation in solar system, plate tectonics and human activities which have increased global warming due to carbon dioxide emission etc.

In recent years concerns about the impact of climate change have increased. Therefore, researchers have utilized various methods and techniques to identify trends and shifts in hydrological series at different scales and places (Caloiero et al., 2011). Trends in hydro-climate data can be investigated at individual sites and regional scales (Sadri et al., 2009). Regional trend analysis has some advantages over at-site analysis. For instance, according to Yue and Hashino (2003) a regional trend analysis provides a better representation of a time series over a wide area. Also, it is more suitable to study the effect of global phenomena such as climate change. There are different methods for detecting trends such as parametric and non-parametric methods. Some researchers have utilized parametric tests to analyze trends in hydrological series. For example Longobardi and Villani (2009) applied the t-test (parametric method) for trend analysis in annual and seasonal rainfall time series in the Mediterranean area. Caloiero et al. (2011) used a parametric method for trend detection in annual and seasonal precipitation in Calabria (southern Italy). The hydrologic components such as precipitation are not constant and they change spatially and temporally. Therefore even over small areas, the parametric methods could not always be appropriate for detecting the precipitation trend. As such many researchers have turned to non-parametric tests such as the Mann-Kendall, the Spearman, the Sen's slope estimator, and so on.

The Mann-Kendall (MK) test has been utilized by many researchers because it has several advantages. For instance, when the data series are non-normal (skewness), the MK test is a suitable choice (Basistha et al., 2009). Another advantage of the MK test is that it has a low sensitivity to inhomogeneous time series (Ahani et al., 2012). Another non-parametric method is the Sen's Slope Estimator. This method estimates the slope if a linear trend exists in time series. Ahani et al. (2012) applied the MK test and Sen's slope estimator to investigate the trend in precipitation volume in different regions of Fars province, Iran. They concluded that the Sen's slope estimator was in a good agreement with MK test. Although the MK test is very popular for analyzing trend, it is sensitive to serial correlation. It has been suggested that before applying the MK test, it is necessary to test the serial correlation of data series (Ahani et al., 2012; Kumar et al., 2009; Matalas and Langbein, 1962; Oguntunde et al., 2011).

Besides these two approaches, the wavelet analysis can also be used to investigate trend. The wavelet transform is a strong mathematical tool that provides a time-frequency representation of an analyzed signal in the time domain (Partal and Küçük, 2006). There are two kinds of wavelet transform function, namely the continuous wavelet transform (CWT) and the discrete wavelet transform (DWT). The advantage of the CWT is that all potential scales or frequencies are analyzed, detected and extracted. However it suffers from edge effects. Meanwhile, the advantage of the DWT is that the transform and reconstruction of the signals of the scales available is perfect, but it is restricted to a discrete numbers of scales (Adamowski et al., 2009). Adamowski et al. (2009) applied the CWT to analyze the recorded monthly minimum stream flow series for five different eco-zones in Canada. It was said that the CWT method is more efficient than spectral analysis when reconstructing highly variable time series from a few wave bands. Partal and Kahya (2006) used the DWT to determine the possible trends in annual total precipitation series in the Marmara region (Turkey). They found that the trend of hydro meteorological variables may be better understood using the Wavelet approach.

1.2 Problem Statement

In recent years, detection of the precipitation trends has attracted wide spread interest. This is not only because of the concerns about the global warming, the rising demand for water and economic growth but also due to the role of water in natural hazards such as floods, droughts and severe erosions (López-Moreno et al., 2009; Oguntunde et al., 2011). Furthermore changing in the pattern of precipitation affects the quality and quantity of water resources. Therefore it is important to know how the climate change affects the water resources, the society and the nature. Studying the climate change is linked directly with trends in various climatic variables because climate is continuously changing.

The climate of Malaysia is equatorial (hot and humid) throughout the year. It has been mentioned that the climate change is likely to have a significant impact on Malaysia, such as increasing sea levels, rainfall and flood risks. Therefore, it is necessary to determine the trends of climate variables to understand the effect of climate change on them. Since in Malaysia, a few studies such as Suhaila et al. (2010) and Tangang et al. (2006) have been performed to investigate the trends of climatic variables, in the present study the annual and seasonal precipitation trends at the Langat River Basin, Malaysia will be investigated. The reservoirs located within the Langat River Basin are the Semenyih dam and Langat dam that provide water for about 1.2 million people in the Langat and Klang valleys. Furthermore, there are 8 water treatment plants in the basin that provide clean water for the users. Beside of providing fresh water, the reservoirs are also important to control floods. The Langat reservoir is also used to generate hydroelectric power. Any change in the pattern of rainfall affects the quantity of water in the basin. Therefore, studying the trend of precipitation is essential for the study area.

As it was mentioned the wavelet transform is a strong method for detection trend. However most of researches have used the parametric and non-parametric tests. The parametric methods are easy to apply but they are restricted to the normal distributed series. On the other hand, the non-parametric methods do not have any assumption about the distribution of the series. They are not also sensitive to non-homogenous data series and the existence of outliers in the series. The wavelet transform can assess both the trend and the periods which are mainly responsible for trend in the measurement series. So in this study, beside the parametric and non-parametric tests, the wavelet transform will be applied.

1.3 Objectives

The main objective of this study is to detect the annual and seasonal precipitation trends in the Langat River Basin.

The following are the specific objectives:

1. To determine the annual and seasonal precipitation trends at both at-site and regional scales.
2. To identify the homogenous rainfall regions in the Langat River Basin.
3. To find the most influential periodicity for annual and seasonal precipitation trends in each homogenous region using Discrete Wavelet Transform (DWT).

1.4 Significance of the Study

As mentioned earlier, in recent years concerns about the impacts of climate change have increased. Various environmental variables including rainfall is affected by the global climate change phenomenon. Therefore, researchers have used different methods to detect trends in hydrology series. Changes in the amount of precipitation directly influence water resources, agriculture, hydrology and ecosystems (Cannarozzo et al., 2006). Furthermore, precipitation has an important role in contributing to natural hazards such as floods, droughts and severe erosions. Since the at-site analysis do not have clear conclusions, the regional-scale analysis is also needed which might be more suitable than at-site studies for determination the impact of climate change (Renard et al., 2008). Thus studying and establishing the trends of rainfall can contribute significantly to the study of climate change scenario. While precipitation trends help to certain extent with interpreting climate change responses, the usefulness of the trend analysis as well as the magnitude over time can be seen when water managers need to plan water resources project and their related management well. One example is the management of water resources hydro dams for electricity production and assessment of long time water availability.

REFERENCES

- Adamowski, K. and Bocci, C. 2001. Geostatistical regional trend detection in river flow data. *Hydrological Processes* 15: 3331–3341.
- Adamowski, K. and Bougadis, J. 2003. Detection of trends in annual extreme rainfall. *Hydrological Processes* 17: 3547–3560.
- Adamowski, K., Prokoph, A. and Adamowski, J. 2009. Development of a new method of wavelet aided trend detection and estimation. *Hydrological Processes* 23: 2686–2696.
- Ahani, H., Kherad, M., Kousari, M.R., Rezaeian-Zadeh, M., Karampour, M.A., Ejraee, F. and Kamali, S. 2012. An investigation of trends in precipitation volume for the last three decades in different regions of Fars province, Iran. *Theoretical and Applied Climatology* 109: 361–382.
- Andreo, B., Jiménez, P., Durán, J.J., Carrasco, F., Vadillo, I. and Mangin, a. 2006. Climatic and hydrological variations during the last 117–166 years in the south of the Iberian Peninsula, from spectral and correlation analyses and continuous wavelet analyses. *Journal of Hydrology* 324: 24–39.
- Basistha, A., Arya, D. and Goel, N. 2009. Analysis of historical changes in rainfall in the Indian Himalayas. *International Journal of Climatology* 29: 555–572.
- Birsan, M.V., Molnar, P., Burlando, P. and Pfaundler, M. 2005. Stream flow trends in Switzerland. *Journal of Hydrology* 314: 312–329.
- Bonaccorso, B., Cancelliere, A. and Rossi, G. 2005. Detecting trends of extreme rainfall series in Sicily. *Advances in Geosciences* 2: 7–11.
- Buishand, T. and Kempen, G. 1988. Trend and seasonal variation of precipitation chemistry data in the Netherlands. *Atmospheric Environment* 22: 339–348.
- Burn, D.H., Sharif, M. and Zhang, K., 2010. Detection of trends in hydrological extremes for Canadian watersheds. *Hydrological processes* 24: 1781–1790.
- Caloiero, T., Coscarelli, R., Ferrari, E. and Mancini, M. 2011. Trend detection of annual and seasonal rainfall in Calabria (Southern Italy). *International Journal of Climatology* 31: 44–56.
- Cannarozzo, M., Noto, L.V. and Viola, F. 2006. Spatial distribution of rainfall trends in Sicily (1921–2000). *Physics and Chemistry of the Earth, Parts A/B/C.* 31: 1201–1211.

- Chaouche, K., Neppel, L., Dieulin, C., Pujol, N., Ladouche, B., Martin, E., Salas, D. and Caballero, Y. 2010. Analyses of precipitation, temperature and evapotranspiration in a French Mediterranean region in the context of climate change. *Comptes Rendus Geoscience* 342: 234–243.
- Cheung, W., Senay, G. and Singh, A. 2008. Trends and spatial distribution of annual and seasonal rainfall in Ethiopia. *International journal of climatology* 13: 1723–1734.
- Chou, C. 2007. Applying multi-resolution analysis to differential hydrological grey models with dual series. *Journal of Hydrology* 332: 174–186.
- Cluis, D., Langlois, C., Van Coillie, R. and Laberge, C. 1990. Development of a software package for trend detection in temporal series: application to water and industrial effluent quality data for the St. Lawrence River, in *Statistical Methods for the Assessment of Point Source Pollution*. Springer. 329–341.
- Corder, G.W. and Foreman, D.I. 2009. *Nonparametric Statistics for Non-Statisticians: A Step-by-Step Approach*. Wiley New Jersey.
- Cunderlik, J. and Burn, D. 2004. Linkages between regional trends in monthly maximum flows and selected climatic variables. *Journal of Hydrologic Engineering* 9: 246–256.
- Dai, A., Trenberth, K.E. and Karl, T.R. 1998. Global variations in droughts and wet spells: 1900-1995. *Geophysical Research Letters* 25: 3367–3370.
- Dai, J. and Schieber, F. 2008. Resampling : The New Statistics The “ Sampling Distribution ” is the foundation of Statistical Inference.. <http://sunburst.usd.edu/~schieber/psyc770/resampling/Resampling-Colloquium.pdf>, Heimstra Labs Colloq.
- Daubechies, I. 1992. *Ten Lectures on Wavelets*. Society for Industrial and Applied Mathematics.
- De Lima, M.I.P., Carvalho, S.C.P. and de Lima, J.L.M.P. 2010. Investigating annual and monthly trends in precipitation structure: an overview across Portugal. *Natural Hazards and Earth System Sciences* 10: 2429–2440.
- Dietz, E.J. and Killeen, T.J. 1981. A nonparametric multivariate test for monotone trend with pharmaceutical applications. *Journal of the American Statistical Association* 76: 169–174.
- Douglas, E., Vogel, R., Kroll, C., 2000. Trends in floods and low flows in the United States: impact of spatial correlation. *Journal of Hydrology* 240: 90–105.

- Gabor, D. 1946. Theory of communication. Part 1: The analysis of information. Electrical Engineers. Part III Radio Commun. Journal of the Institution 93: 429–441.
- Golberg, M.A. and Cho, H.A. 2004. Introduction to regression analysis, illustrate. ed. WIT Press.
- Hamed, K. and Rao, A.R. 1998. A modified Mann-Kendall trend test for autocorrelated data. Journal of Hydrology 204: 182–196.
- Hamed, K.H. 2008. Trend detection in hydrologic data: The Mann--Kendall trend test under the scaling hypothesis. Journal of Hydrology 349: 350–363.
- Helsel, D.R. and Hirsch, R.M. 1992. Statistical methods in water resources. Elsevier, Amsterdam.
- Hirsch, R.M. and Slack, J.R. 1984. A nonparametric trend test for seasonal data with serial dependence. Water Resources Research 20: 727–732.
- Hirsch, R.M., Slack, J.R. and Smith, R.A. 1982. Techniques of trend analysis for monthly water quality data. Water Resources Research 18: 107–121.
- Huang, Y., Wang, Y. and Zhang, L. 2008. Long-term trend of chemical composition of wet atmospheric precipitation during 1986–2006 at Shenzhen City, China. Atmospheric Environment 42: 3740–3750.
- Huntington, T.G. 2006. Evidence for intensification of the global water cycle: review and synthesis. Journal of Hydrology 319: 83–95.
- Intergovernmental Panel on Climate Change (IPCC). 2001. Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, edited by J. T. Houghton et al., 881 pp., Cambridge Univ. Press, New York.
- Jiang, T., Su, B. and Hartmann, H. 2007. Temporal and spatial trends of precipitation and river flow in the Yangtze River Basin, 1961–2000. Geomorphology 85: 143–154.
- Juahir, H. 2009. Water quality data analysis and modeling of the Langat river basin. University Malaya.
- Kang, S. and Lin, H. 2007. Wavelet analysis of hydrological and water quality signals in an agricultural watershed. Journal of Hydrology 338: 1–14.
- Kardi, T. 2007. K-Means clustering tutorials. <http://people.revoledu.com/kardi/tutorial/kMean/>.

- Karpouzou, D., Kavalieratou, S. and Babajimopoulos, C. 2010. Trend analysis of precipitation data in Pieria Region (Greece). *European Water* 30: 31–40.
- Kendall, M.G. 1975. *Rank correlation methods*: Charles Griffin. Oxford, London, UK.
- Kim, S. 2004. Wavelet analysis of precipitation variability in northern California, U.S.A. *KSCE Journal of Civil Engineering* 8: 471–477.
- Kisi, O. and Cimen, M. 2011. A wavelet-support vector machine conjunction model for monthly stream flow forecasting. *Journal of Hydrology* 399: 132–140.
- Kulkarni, A. and von Storch, H. 1995. Monte Carlo experiments on the effect of serial correlation on the Mann-Kendall test of trend. *Meteorologische Zeitschrift* 4: 82–85.
- Kumar, S., Merwade, V., Kam, J. and Thurner, K. 2009. Streamflow trends in Indiana: Effects of long term persistence, precipitation and subsurface drains. *Journal of Hydrology* 374: 171–183.
- Kumar, V. and Jain, S.K. 2010. Trends in seasonal and annual rainfall and rainy days in Kashmir Valley in the last century. *Quaternary International* 212: 64–69.
- Kundzewicz, Z.W. and Robson, A.J. 2004. Change detection in hydrological records—a review of the methodology/Revue méthodologique de la détection de changements dans les chroniques hydrologiques. *Hydrological Sciences Journal* 49: 7–19.
- Lanzante, J.R. 1996. Resistant, robust and non-Parametric techniques for the analysis of climate data: theory and examples, including applications to historical radiosonde station data. *International Journal of Climatology* 16: 1197–1226.
- Liu, Q., Yang, Z. and Cui, B. 2008. Spatial and temporal variability of annual precipitation during 1961–2006 in Yellow River Basin, China. *Journal of Hydrology* 361: 330–338.
- Longobardi, A. and Villani, P. 2009. Trend analysis of annual and seasonal rainfall time series in the Mediterranean area. *International journal of Climatology* 30: 1538–1546.
- López-Moreno, J.I., Vicente-Serrano, S.M., Angulo-Martínez, M., Beguería, S. and Kenawy, a. 2009. Trends in daily precipitation on the northeastern Iberian Peninsula, 1955-2006. *International journal of Climatology* 30: 1026–1041.
- Lu, G.Y. and Wong, D.W. 2008. An adaptive inverse-distance weighting spatial interpolation technique. *Computers & Geosciences* 34: 1044–1055.

- Mallat, S. 1989. A theory for multiresolution signal decomposition: the wavelet representation. *Pattern Analysis and Machine Intelligence, IEEE Transactions on* 11: 674–693.
- Mann, H.B. 1945. Nonparametric tests against trend. *Econometrica: Journal of the Econometric Society* 245–259.
- Matalas, N. and Langbein, W. 1962. Information content of the mean. *Journal of Geophysical Research* 57: 3441–3448.
- McCuen, R.H. 2005. *Hydrologic analysis and design*, 3, illustr. ed. Pearson Prentice Hall, Michigan.
- Miao, L., Jun, X. and Dejuan, M. 2012. Long-term trend analysis of seasonal precipitation for Beijing, China. *Journal of Resources and Ecology* 3: 64–72.
- Misiti, M. and Misiti, Y. 1996. *Wavelet toolbox*, MA. ed, The MathWorks, Inc, Natick.
- Modarres, R., da Silva, V. and de P.R. 2007. Rainfall trends in arid and semi-arid regions of Iran. *Journal of Arid Environments* 70: 344–355.
- Moh'd B, A.Z. and al Rawi, M. 2008. An efficient approach for computing silhouette coefficients. *Journal of Computer Science* 4: 252.
- Nalley, D. 2012. *Analyzing trends in temperature, precipitation and streamflow data over southern ontario and quebec using the discrete wavelet transform*. McGill University.
- Nalley, D., Adamowski, J. and Khalil, B. 2012. Using discrete wavelet transforms to analyze trends in streamflow and precipitation in Quebec and Ontario (1954–2008). *Journal of Hydrology* 475: 204–228.
- Novotny, E. V. and Stefan, H.G. 2007. Stream flow in Minnesota: Indicator of climate change. *Journal of Hydrology* 334: 319–333.
- Oguntunde, P.G., Abiodun, B.J. and Lischeid, G. 2011. Rainfall trends in Nigeria, 1901–2000. *Journal of Hydrology* 411:207–218.
- Partal, T. 2010. Wavelet transform-based analysis of periodicities and trends of Sakarya basin (Turkey) streamflow data. *River Research and Applications* 26: 695–711.
- Partal, T. and Kahya, E. 2006. Trend analysis in Turkish precipitation data. *Hydrological processes* 20: 2011–2026.

- Partal, T. and Küçük, M. 2006. Long-term trend analysis using discrete wavelet components of annual precipitations measurements in Marmara region (Turkey). *Physics and Chemistry of the Earth, Parts A/B/C.* 31: 1189–1200.
- Percival, D.B. 2008. Analysis of geophysical time series using discrete wavelet transforms: An overview, in: *Nonlinear Time Series Analysis in the Geosciences.* Springer, Berlin. 61–79.
- Pidwirny, M. 2006. Global distribution of precipitation. <http://www.eoearth.org/view/article/153027>.
- Prokoph, A. and Patterson, R.T. 2004. Application of wavelet and regression analysis in assessing temporal and geographic climate variability: Eastern Ontario, Canada as a case study. *Atmosphere-Ocean* 42: 201–212.
- Rana, A., Uvo, C.B., Bengtsson, L. and Parth Sarthi, P. 2011. Trend analysis for rainfall in Delhi and Mumbai, India. *Climate dynamics* 38: 45–56.
- Renard, B., Lang, M., Bois, P., Dupeyrat, A., Mestre, O., Niel, H., Sauquet, E., Prudhomme, C., Parey, S., Paquet, E., Neppel, L. and Gailhard, J. 2008. Regional methods for trend detection: Assessing field significance and regional consistency. *Water Resources Research* 44: W08419.
- Robson, A.J. 2004. Change detection in hydrological records — a review of the methodology. *Hydrological Sciences Journal* 49: 7–20 .
- Sadri, S., Madsen, H., Mikkelsen, P.S. and Burn, D.H. 2009. Analysis of extreme rainfall trends in Denmark, in: *proceedings of the 33rd iahr congress: water engineering for a sustainable environment. international association of hydraulic engineering & research (IAHR), Vancouver, Canada, 9-14 August .1731–1738.*
- Santos, C. and Galvão, C. 2001. Matsuyama city rainfall data analysis using wavelet transform. *Annual Journal of Hydraulic Engineering-JSCE* 45: 211–216.
- Sen, P.K. 1968. Estimates of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association* 63: 1379–1389.
- Serrano, A., Mateos, V.L. and Garcia, J.A. 1999. Trend analysis of monthly precipitation over the Iberian peninsula for the period 1921–1995. *Physics and Chemistry of the Earth, Part B Hydrology, Oceans and Atmosphere* 24: 85–90.
- Shaaban, A. J. B. 2013. Impact of climate change on Malaysia, National Hydraulic Research Institute of Malaysia, Ministry of Natural Resources & Environment.

- Sneyers, R. 1990. On the statistical analysis of series of observations. Secretariat of the world meteorological organization, Geneva.
- Suhaila, J., Deni, S., Zin, W. and Jemain, A. 2010. Trends in peninsular Malaysia rainfall data during the Southwest Monsoon and Northeast Monsoon Seasons: 1975–2004. *Sains Malaysiana* 39: 533–542.
- Svensson, C., Kundzewicz, W.Z. and Maurer, T. 2005. Trend detection in river flow series: 2. Flood and low-flow index series/ Détection de tendance dans des séries de débit fluvial: 2. Séries d'indices de crue et d'étiage. *Hydrological Sciences Journal* 50.
- Tabari, H. and Marofi, S. 2010. Changes of Pan Evaporation in the West of Iran. *Water resources management* 25: 97–111.
- Tabari, H. and Talaei, P.H. 2011. Temporal variability of precipitation over Iran: 1966–2005. *Journal of Hydrology* 396: 313–320.
- Tomé, A.R. and Miranda, P.M.A. 2005. Continuous partial trends and low-frequency oscillations of time series. *Nonlinear Processes in Geophysics* 12: 451–460.
- Torrence, C. and Compo, G. 1998. A practical guide to wavelet analysis. *Bulletin of the American Meteorological society* 79: 61–78.
- Van Belle, G. and Hughes, J.P. 1984. Nonparametric tests for trend in water quality. *Water resources research* 20: 127–136.
- World Meteorological Organization. 1989. Calculation of monthly and annual 30-year standard normals. WCDP, No. 10, WMO-TD/No. 341. Geneva.
- Xu, J., Chen, Y., Li, W., Ji, M., Dong, S. and Hong, Y. 2009. Wavelet analysis and nonparametric test for climate change in Tarim River Basin of Xinjiang during 1959–2006. *Chinese Geographical Science* 19: 306–313.
- Yue, S. and Hashino, M. 2003. Long term trends of annual and monthly precipitation in Japan. *JAWRA Journal of the American Water Resources Association* 39: 587–596.
- Yue, S. and Wang, C.Y. 2002. Regional streamflow trend detection with consideration of both temporal and spatial correlation. *International Journal of Climatology* 22: 933–946.
- Zhang, X., Vincent, L. a., Hogg, W.D. and Niitsoo, A. 2000. Temperature and precipitation trends in Canada during the 20th century. *Atmosphere-Ocean* 38: 395–429.