



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF SALINITY ON YIELD AND POSTHARVEST QUALITY
OF TOMATO (LYCOPERSICON ESCULENTUM MILL.)**

ABDUL-RAQEEB ALI AHMED AL-ERYANI.

FP 2004 8

**EFFECTS OF SALINITY ON YIELD AND POSTHARVEST QUALITY OF
TOMATO (*Lycopersicon esculentum* Mill.)**

By

ABDUL- RAQEEB ALI AHMED AL- ERYANI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Agricultural Science**

February 2004



Special dedicated

To

My beloved father and mother

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements of the degree of Master of Agricultural Science

**EFFECTS OF SALINITY ON YIELD AND POSTHARVEST QUALITY OF
TOMATO (*Lycopersicon esculentum* Mill.)**

By

ABDUL- RAQEEB ALI AHMED AL- ERYANI

February 2004

Chairman: Associate Professor Dr. Mohd. Razi Ismail, Ph.D.

Faculty: Agriculture

The effects of salinity on the yield and chemical composition of tomato (*Lycopersicon esculentum* Mill.) fruits produced in soilless culture under protected environment were investigated. Two experiments were carried out at Hydroponic Complex, Crop Science Department, Faculty of Agriculture, Universiti Putra Malaysia.

Increasing salinity (EC) from 2 mScm⁻¹ (control) to 6 mScm⁻¹ linearly reduced total yield, size, firmness and water content of tomato fruits, and dry weights of roots and shoots of plants. The incidence of blossom end rot (BER) was higher at high salinity level as a consequence of deficiency of Ca content, which was found to decrease with increasing salinity levels. On the contrary, high salinity conditions resulted in increasing total soluble solids, carbohydrates (fructose, glucose, sucrose), titratable acidity and ascorbic acid (Vitamin C) concentrations and dry matter content of tomato fruits. The redness (a*) values gradually increased with increasing salinity level from ECs 3 to 6 mScm⁻¹, whereas the lightness (L*) and yellowness (b*) values decreased. These observations indicated that it is possible to obtain a good quality tomato fruits with acceptable yield reduction at EC 4.5 mScm⁻¹.

Salinity affected both shelf life of tomato fruits stored at ambient temperature (21°C) or in cool condition (15°C) with relative humidity (RH) between 48-66% and 91-92%, respectively. There was a negative relationship between salinity and fruit shelf life, probably due to an increase in polygalacturonase activity, which enhances softening and hence causes shorter shelf life.

In another experiment, both high (6 mScm⁻¹) and moderate (3 mScm⁻¹) salinities were applied at different growth stages of plants development. Saline irrigation at EC 3 mScm⁻¹ during late developmental stages (onset of ripening) improved the quality of the fruits with acceptable yield reduction (fresh weight, number and size of fruits). In general, maintaining the proper ECs 3 and 4.5 mScm⁻¹ applied at onset of ripening and flowering stages, respectively resulted to in an acceptable yield reductions and high quality products.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Pertanian

**KESAN KEMASINAN KE ATAS HASIL DAN
KUALITI PASCA TUAI TOMATO (*Lycopersicon esculentum* Mill.)**

Oleh

ABDUL-RAQEEB ALI AHMED AL-ERYANI

February 2004

Pengerusi: Profesor Madya Dr. Mohd. Razi Ismail, Ph.D.

Fakulti: Pertanian

Kajian ke atas kesan kemasinan terhadap hasil dan komposisi kimia buah tomato (*Lycopersicon esculentum* Mill.) telah dijalankan menggunakan media tanpa tanah di bawah persekitaran terkawal. Dua eksperimen telah dijalankan di Unit Hidroponik, Jabatan Sains Tanaman, Fakulti Pertanian, Universiti Putra Malaysia.

Peningkatan kemasinan (EC) dari 2.0 mScm^{-1} (kawalan) kepada 6.0 mScm^{-1} mengurangkan jumlah hasil, saiz, kekerasan dan kandungan air buah tomato, berat kering akar dan pucuk pokok tomato. Walaubagaimana pun, kejadian reput pangkal buah adalah tinggi pada tahap kemasinan yang tinggi disebabkan oleh kandungan kalsium yang di dapati berkurangan dengan meningkatnya paras kemasinan. Tetapi, keadaan kemasinan yang tinggi di dapati meningkatkan jumlah bahan terlarut, karbohidrat (fruktos, glukos dan sukros), keasidan titrat dan asid askorbik (vitamin C) serta kandungan bahan kering buah tomato. Nilai kemerahan (a^*) meningkat seiringan dengan peningkatan kemasinan dari EC 3.0 ke 6.0 mScm^{-1} , manakala nilai kejelasan

(L*) dan kekuningan (b*) menurun. Pemerhatian ini menunjukkan kemungkinan untuk mendapatkan kualiti buah tomato yang baik dengan pengurangan hasil yang boleh diterima pada EC 4.5 mScm⁻¹.

Kemasinan memberi kesan ke atas jangkahayat buah tomato yang disimpan pada suhu bilik (21 °C) dan bilik sejuk (15 °C) dengan kelembapan relatif (RH) masing – masing 48-66% dan 91-92%. Perhubungan negatif ditunjukkan di antara kemasinan dengan jangkahayat buah tomato, berkemungkinan disebabkan oleh peningkatan aktiviti polygalacturonase, yang meningkatkan kelembutan isi buah dan menyebabkan jangkahayat yang pendek.

Dalam eksperimen yang berasingan, kedua-dua kadar kemasinan tinggi (6.0 mScm⁻¹) dan sederhana (3.0 mScm⁻¹) telah diberikan pada peringkat tumbesaran yang berbeza. Pengairan dengan kadar kemasinan 3.0 mScm⁻¹ pada akhir peringkat pembentukan buah (permulaan kemasakan) meningkatkan kualiti buah walaupun hasilnya berkurangan. Pada amnya, pengekalan kadar kemasinan 3 dan 4.5 mScm⁻¹, pada peringkat permulaan kemasakan buah dan peringkat pembungaan boleh menghasilkan buah yang berkualiti tinggi dengan pengurangan hasil yang boleh diterima.

ACKNOWLEDGEMENTS

In the name of Allah the Beneficial and the Compassionate

Praise be to Allah SWT, upon His permission I could complete this thesis smoothly. Contributions of individuals and institution for the successful completion of this thesis are also deeply acknowledged.

I would like to express my sincere thank and profound appreciation to the chairman of my supervisory committee, Associate Professor Dr. Mohd Razi Ismail, Department of Crop Science, Faculty of Agriculture for his supervision, kind support beyond supervisory duties and for being so patient and understanding. I am also indebted to Associate Professor Dr. Mahmud Tengku Muda Mohamed, my co-supervisor for his time and advice, as well as for his encouragement and friendship. Special thanks to Associate Professor Dr. Abdul Ghani Yunus as a member of my supervisory committee for his critical comments and suggestion during the course of this project.

Completion of this research work is also much owes to my wife, for her continuous encouragement and my beloved sons Mohammed and Ryan and my beloved daughter Shuhd who provided the comfort and for being the constant source of inspiration for me to complete this research.

My special appreciation and gratitude goes to my beloved father and mother, I pray to Allah to bless my parents who guided and supported me to achieve the highest level of

education. I also wish to express my sincere gratitude to my brothers Dr. Abdul Rahman and Amin and my sisters for their constant encouragement, love and ever ready to give helping hands.

My deepest thanks to my beloved father-in-law Al Haj Ahmed Ali Al Eryani and my mother-in-law and their sons Nabil, Rafiq and Maher and other family members for their kind consideration and all the valuable supports. I am indebted to them for their support to me and my family in completing my study.

My appreciations and honest thanks are due to all Lecturers and staff members in Sana'a University, Faculty of Agriculture, Department of Horticultural Science in particular Dr. Ahmed Hamzah, Dr. Mane'e, Dr. Mehasen, Dr. Al ward, Dr. Hassan and Dr. Mansor as well as for their encouragement and friendship.

Finally, most profound thanks go to the Malaysian government and people represented by Universiti Putra Malaysia, for giving me this opportunity to study in their prestigious and reputed institute.

DECLARATION

I here by declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



ABDUL- RAQEEB ALI AHMED AL- ERYANI

Date: 05 APR 2004

TABLE OF CONTENTS

	Page
DEDICATION	2
ABSTRACT	3
ABSTRAK	5
ACKNOWLEDGMENTS	7
APPROVAL	9
DECLARATION	11
LIST OF FIGURES	15

CHAPTER

1	INTRODUCTION	17
2	LITERATURE REVIEW	19
	2.1 Tomato	19
	2.2 Tomato Handling and Postharvest Process	19
	2.3 Storage Requirements	21
	2.4 Tomato Storage	22
	2.5 Salinity and Soil Salinity	23
	2.6 Salinity and Other Environmental Factors	25
	2.7 Salinity Mineral Nutrient Relations in Horticultural Crops	25
	2.8 Responses and Salt Tolerance of Crops to Soil Salinity	27
	2.9 Effect of Salinity on Growth Performance	28
	2.9.1 Effect on Shoot Development	28
	2.9.2 Effect on Root Development	30
	2.9.3 Effect of salinity on flowering quality and quantity of pollen	31
	2.10 Effect of Salinity on Fruit Growth	32
	2.11 Effect of Salinity on Yield	33
	2.11.1 Fruit Number and Weight	33
	2.11.2 Fruit Size and Dry Matter Content	34
	2.12 Effect of Salinity on Fruit Quality	35
	2.12.1 Fruit Composition	35
	2.12.2 Fruit Firmness	37
	2.12.3 Fruit Skin Colour	37
	2.12.4 Fruit Shelf Life and Blossom End Rot	38
	2.13 Effect of Salinity on Carbohydrates Functioning	38
	2.14 Tomato Cuticle	39
	2.14.1 Cuticle Anatomy Description	39
	2.14.2 Importance of Wax	39
	2.14.3 Changes of Cuticular Waxes During Fruit Ripening	39

3	GENERAL MATERIALS AND METHODS	41
3. 1	Establishment of Planting Materials	41
3. 2	Treatments and Design	42
3. 3	Fruits Sampling	43
3. 4	Fruit Parameters Determination	43
3. 4. 1	Fruit Weight, Number and Size Determination	43
3. 4. 2	Fruit Firmness Determination	44
3. 4. 3	Fruit Total Soluble Solids (TSS) Determination	44
3. 4. 4	Total Titratable Acidity	45
3. 5	Statistical Analysis	46
 4	 Effects of Increasing Salinity in the Root Environment on Yield, Quality, Wax Deposition and Storability of Tomatoes (<i>Lycopersicon esculentum</i> L.) Grown in Coconut Coir Dust Culture	
4. 1	Introduction	47
4. 2	Materials and Methods	50
4. 2. 1	Plant Materials and Treatments	50
4. 2. 2	Fresh and Dry Weight of Shoots an Roots	50
4. 2. 3	Fruit Weight, Number and Size Determination	66
4. 2. 4	Fruit Moisture and Dry Matter Content Determinations	51
4. 2. 5	Fruit Quality Parameters Determination	52
4. 2. 5. 1	Firmness, Titratable Acidity and TSS	52
4. 2. 5. 2	Ascorbic Acid Determination	52
4. 2. 5. 3	Skin Colour Determination	54
4. 2. 5. 4	Sugar Determination	54
4. 2. 5. 5	Cuticle Wax Measurement	55
4. 2. 5. 6	Blossom End Rot	58
4. 2. 5. 7	Ca Content	58
4. 2. 5. 8	Shelf Life	58
4. 3	Results and Discussion	59
4. 3. 1	Total Yield	59
4. 3. 2	Number of fruits	60
4. 3. 3	Fruit Size	62
4. 3. 4	Fruit Firmness	63
4. 3. 5	Titrratable Acidity	64
4. 3. 6	Ascorbic Acid	65
4. 3. 7	Total Soluble Solids (TSS)	67
4. 3. 8	Water Content	68
4. 3. 9	Dry Matter Content	70
4. 3. 10	Dry weight of Roots and Shoots	71
4. 3. 11	Sugar Content	75
4. 3. 12	Calcium Content	78

	4. 3. 13 Blossom End Rot	79
	4. 3. 14 Shelf life	81
	4. 3. 15 Skin Colour	84
	4. 3. 16 Wax deposition	86
	4. 3. 17 The relationship between Yield and Quality (TSS)	86
5	Effect of Regulating Salinity at Different Phenological Stages of Plants in the Root Environment on Yield and Quality of Tomato (<i>Lycopersicon esculentum</i> L.) Fruits	
	5. 1 Introduction	89
	5. 2 Materials and Methods	90
	5. 2. 1 Plant Materials and Treatments	90
	5. 2. 2 Fresh Weight, Number and Size Determination	91
	5. 2. 3 Fruit Quality Parameters Determination	91
	5. 3 Result and Discussion	91
	5. 3. 1 Fruit Number	91
	5. 3. 2 Total Fruit Yield	92
	5. 3. 3 Fruit Firmness	95
	5. 3. 4 Fruit Size	95
	5. 3. 5 Titratable Acidity	98
	5. 3. 6 Total Soluble Solids (TSS)	98
	5. 3. 7 Skin Colour	101
	5. 3. 8 Sugar Contents	104
	5. 3. 9 The relationship between Sugars Contents and Yield	107
6	GENERAL DISCUSSION AND CONCLUSION	109
	REFERENCES	114
		103
	APPENDICES	
	Appendix A	125
	Appendix B	127
	BIODATA OF THE AUTHOR	140

LIST OF FIGURES

Figure		Page
3. 1	Typical daily temperature in the greenhouse at the location of the experiments.	42
4. 3. 1	Effect of different salinity level on total fruit yield of tomato plants (g/plant).	60
4. 3. 2	Effect of different salinity levels on fruit number of tomato plants.	61
4. 3. 3	Effect of different salinity levels on size (cm) of tomato fruits.	63
4. 3. 4	Effect of different salinity levels on firmness (kg.f) of tomato fruits.	64
4. 3. 5	Effect of different salinity levels on titratable acidity content of tomato fruits.	65
4. 3. 6	Effect of different salinity levels on ascorbic acid content of tomato fruits.	66
4. 3. 7	Effect of different salinity level on total soluble solid content of tomato fruits.	67
4. 3. 8	Effect of different salinity levels on water content of tomato fruits.	69
4. 3. 9	Effect of different salinity levels on fruit dry matter content of tomato fruits.	71
4. 3. 10	Effect of different salinity levels on plants dry matter content of (A) roots and (B) shoots of tomato.	74
4. 3. 11	Effect of different salinity levels on sugar contents (A) fructose, (B) glucose and (C) sucrose of tomato fruits.	77
4. 3. 12	Effect of different salinity levels on (A) Ca content and (B) blossom end rot of tomato fruits.	80
4. 3. 13	Effect of different salinity levels on shelf life of tomato fruits stored under room condition (A) and cool condition (B).	83
4. 3. 14	Effect of different salinity levels on (A) lightness (L* values), (B) redness (a* values) and (C) yellowness (b* values) of colour indicators of tomato fruits.	85

4. 3. 15	Effect of different levels of salinity on tomato fruit skin cuticle wax.	87
4. 3. 16	Effect of different salinity levels on yield and quality (TSS) of tomato fruits.	88
4. 3. 17	The Relationship between yield and quality (TSS) obtained under salinity levels.	88
5. 3. 1	Effect of salinity imposed at different phenological stages and concentrations on (A) fruit number and (B) total fruit yield.	94
5. 3. 2	Effect of salinity imposed at different phenological stages and concentrations on firmness (A) and size (B) of tomato fruits.	97
5. 3. 3	Effect of salinity imposed at different phenological stages and concentrations on (A) the contents of titratable acidity and (B) total soluble solids of tomato fruits.	100
5. 3. 4	Effect of salinity imposed at different phenological stages and concentrations on (A) lightness, (B) a* values and (C) b* values of skin colour of tomato fruits.	103
5. 3. 5	Effect of salinity imposed at different phenological stages and concentrations on (A) fructose, (B) glucose and (C) sucrose contents of tomato fruits.	106
5. 3. 6	The relationship between quality as sugars contents {fructose (A), glucose (B) and sucrose(C)} and total fruit yield related to salinity imposed at different phenological stages and concentrations.	108

CHAPTER 1

INTRODUCTION

In drought areas where water resources are limited, poor quality water is increasingly being used, although such irrigation water is one of the major sources of salinity, which may result in crop yield reductions and soil degradation (Oster, 1994). In countries like Yemen, drought and salinity are prevalent especially in coastal areas and ground water that contains high concentrations of NaCl is being used by farmers for irrigation. However, the use of such water leads to long-term environmental problems such as permanent soil salinization and soil degradation. Likewise, the usage of saline water induces an effect that results in reduced crop yield and quality. Salinity reduces the relative growth rate of plants and restricts leaf expansion.

Tomato is a widely distributed annual vegetable crop which is consumed fresh, cooked or processed such as canning, juice, pulp, paste or a variety of sauces (FAO, 1995). In Malaysia, tomato is mostly cultivated in highlands such as Cameron Highlands. Local farmers successfully cultivate high temperature resistant tomato variety. Presently, cultivation of tomato under protected conditions is being attempted and problems such as salinity were prevalent either due to the use of nutrient solutions or environmental contamination of land by sources of salinity.

In today's world market, fruit flavour is of paramount importance to guarantee consumer satisfaction. Tomatoes grown under saline conditions produce fruits with a higher content of sugars and organic acids, which contributes to improved market quality of the

fruit. However, significant increase in total solids (TS) and total soluble solids (TSS) of fruits produced by salinized plants can compensate for the reduction of fresh yield when the yield is expressed on a dry weight basis (Ehert and Ho, 1986; Mizrahi *et al.*, 1988; Gao *et al.*, 1996).

Tomato plants which grow under salinity conditions may have a good quality characteristics that may improve the taste without reducing other quality aspects. However, tomato fruits developed under salinity conditions tend to have tough skin and enhanced wax layer of the skin that gives good appearance and reduces water loss. These characteristics may help to maintain longer shelf life during handling and under storage. Therefore, regulating the salinity level may be of much benefits for exporting, industrial processing or fresh marketing of the fruits. Though a few work had been done on the effect of various salinity levels on post-harvest quality and storability of tomato, not much works were being done on the imposition of salinity at different phenological stages of the plants. Hence, the objectives of this study are to determine the effects of different levels of salinity on yield, postharvest quality and storability of tomato and improvement of yield and quality by manipulating salinity levels at different phenological stages of plant.

CHAPTER 2

LITERATURE REVIEW

2. 1 Tomato

Tomato belongs to the genus *Lycopersicon* and *esculentum* is the species mostly grown for its edible fruit. The genus *Lycopersicon* of the family Solanaceae is believed to have originated from Latin America between Mexico and Peru. The tomato crop is adapted to a wide range of climates from the tropics to within a few degrees of the Arctic Circle. However, its broad adaptation, production is concentrated in a few warm and rather dry areas: more than 30% of world production comes from countries around the Mediterranean Sea and about 20% from California (Taylor, 1986; Papadopoulos, 1991).

2. 2 Tomato Handling and Postharvest Process

Successful postharvest handling of vegetable crops requires careful coordination and integration of the various steps from harvest operations to consumer in order to maintain the initial product quality. Horticultural quality refers to those characteristics which are associated with consumers for each commodity that are dependent upon the particular end use, such as sweetness in strawberries and melons, tenderness in snap beans and sweet corn, and crispness in carrots and celery. Quality also refers to freedom from defects such as blemishes, mechanical injury, physiological disorders, decay and water loss. It is important to keep in mind those quality losses for fresh vegetables are

cumulative: each incident of mishandling reduces final quality at consumer level (Sargent *et al.*, 1995).

Several factors that reduce quality during postharvest handling, including:

- Harvest at the incorrect maturity stage
- Careless handling at harvest and during packaging and shipping
- Poor sanitation
- Delays to cooling or sub-optimal cooling
- Shipping/storage above or below optimal temperature
- Lack of proper relative humidity
- For some commodities, exposure to ethylene gas.

Numerous technologies and procedures can significantly reduce quality losses during handling. These include the use of drying, curing, temperature conditioning, disinfestation (for exports/imports), ethylene treatment, application of surface coatings, sanitation treatments, controlled atmosphere storage, shipping and modified atmosphere packaging. Two of the most important means for maintaining vegetable quality during postharvest handling are minimizing mechanical injury and managing temperature. Proper handling and temperature management will significantly reduce losses due to decay and accelerated senescence. With vegetables typically being handled several times from harvest to retail level, it is critical that personnel at each step be properly trained and supervised (Sargent *et al.*, 1995).

Fruit ethylene production is associated with colour development and this natural hormone is also involved in fruit softening. Ethylene application can be used to accelerate colour development and fruit softening when provided at or slightly beyond

the mature green stage. Ethylene gas (100-150 ppm) is usually applied to the fruit in a sealed ripening room. The best ripening response to ethylene occurs at a temperature of 20-21° C and 85-90% RH for 12-24 h (Sargent *et al.*, 1995).

Fruit deterioration because of excessive softening is a major reason for marketable losses. Rough handling, poorly designed containers and exposure to hot and dry conditions also contribute to significant losses (Sargent *et al.*, 1995).

At many fresh market tomato-packaging facilities, fruit are washed in chlorinated water to remove dirt and to limit postharvest diseases. When cold washwater is used, postharvest decay often increases because of contaminated water entering the fruit through the stem scar. However, when the washwater is equal to or warmer than the fruit temperature, contaminated water is not drawn into the fruit (Rubatzky and Yamaguchi, 1997).

Fresh market fruit are graded for uniform size and quality before packaging; in some facilities this is accomplished mechanically. Occasionally, fruit are waxed to reduce moisture loss and improve appearance (Grierson and Kader, 1986; Rubatzky and Yamaguchi, 1997).

2. 3 Storage Requirements

Horticultural crops may be grouped into two broad categories based on sensitivity to storage temperatures. However, the degree of chilling sensitivity for the lowest safe

storage temperature is crop specific. Those crops, which are chilling sensitive, should be held at temperatures generally above 50°F (10°C). Storage below this threshold will give rise to a physiological disorder known as chilling injury. Chilling injury symptoms are characterized by development of sunken lesions on the skin, increased susceptibility to decay, increased shrivel and incomplete ripening (poor flavour, texture, aroma and colour). Those crops that are not sensitive to chilling injury may be stored at temperatures as low as 32°F (0°C). The extent of chilling symptoms is also dependent on the length of exposure to low temperatures. Short exposure times will result in less injury than longer exposure to chilling temperatures (Sargent *et al.*, 1995).

2. 4 Tomato Storage

Tomatoes can be stored successfully for several weeks, but recommended storage temperatures differ with stage of fruit maturation. When mature green fruit are stored, temperatures should be between 13° C and 18° C and 85-95% RH. At these temperatures, chilling damage does not occur, but colour development is slow. The optimum temperature for ripen mature green fruit is between 18° C and 21° C; below 13° C, fruit will not develop a dark red colour. Mature green fruit have been stored for 6 weeks at 13° C in 3% oxygen, 97% nitrogen atmosphere, and upon ripening, there was no noticeable flavour or other quality impairment (Rubatzky and Yamaguchi, 1997).

Red fruit have a shorter shelf life at room temperature but can tolerate storage at lower temperatures than mature green fruit. Firm red fruit can be held at 7-10° C for several days without significant quality losses. For ripe fruit, temperatures less than 7° C will cause chilling damage and the fruit loses firmness, flavour and shelf life. Chilling injury

is cumulative and increases with length and level of low temperature. Red fruit can be stored as long as 3 weeks at 0-1.5° C in acceptable condition. However, fruit should be used within a day or two following removal from storage because flavour and textural quality becomes unacceptable. The usual recommendation for red fruit to maintain quality is to avoid low temperature exposure (Rubatzky and Yamaguchi, 1997).

2. 5 Salinity and Soil Salinity

Over the course of history, thriving civilizations declined in part due to their inability to sustain food production on lands that had been salinized. It is estimated that 10 million hectares are now being lost every year as a result of salinity and water logging (El-Haddad *et al.*, 1998).

Excessive use of water for irrigation due to inefficient irrigation distribution systems, poor on-farm management practices and inappropriate management of drainage water causes many of these problems. Inefficient on-farm irrigation practices cause local salinity problems. These problems increase as a result of poor on-farm drainage. Excessive irrigation increases salt loading in water tables and downstream aquifers, which causes regional salinization. Lack of these local and regional drainage systems result in lands being put out of agricultural production (El-Haddad *et al.*, 1998).

Salinity is a threat to the health and productivity of many catchments, and to the rural and urban communities that live on them. It is affecting rural landholders, urban developments, infrastructure (roads and bridges), water users and the environment. In some places, salinity is a natural phenomenon but in others, increasing salinity caused by

rising water tables is often the result of particular land use practices, such as over-clearing, urban development, river regulation, irrigation or the cultivation of crops. Salinity from rising watertables is grouped into dryland, irrigation and urban salinity. River salinity and industrial salinity can be observed as related phenomena. Dryland salinity is the build up of salt in the soil surface in non-irrigated areas, usually as a result of a rising watertable. Irrigation salinity is caused by over irrigation, inefficient water use and poor drainage (NSW Department of Land and Water Conservation, 1998).

Soil salinity is a measurement of the total amount of soluble salt in soil. As salinity levels increase, plants extract water less easily from soil, aggravating water stress conditions. High soil salinity can also cause nutrient imbalances, result in the accumulation of elements toxic to plants, and reduce water infiltration if the level of one salt element-sodium is high. In many areas, soil salinity is the factor limiting plant growth. Salt-affected plants are stunted with dark green leaves, which in some cases are thicker and more succulent than normal. In woody species, high soil salinity may lead to leaf burn and defoliation. High salinity causes alfalfa yield to decrease while the leaf-to-stem ratio increases, influencing forage quality. Grasses also appear dark green and stunted with leaf burn symptoms (Kotuby-Amacher *et al.*, 2000).

In most instances, dryland salinity is caused by saline groundwater seeping to the surface of the land. Its impacts are on soil quality and water resources, depleting their utility and environmental value. Salinity is usually evident after water tables reach within 1.5 or two meters of the surface when shallow saline groundwater, drawn up by capillary action, is further concentrated by evaporation (Ailie, 2000).