



UNIVERSITI PUTRA MALAYSIA

**BROODSTOCK NUTRITION OF AUSTRALIAN RED CLAW
CRAYFISH *CHERAX QUADRICARINATUS* (VON MARTENS)**

LADAN ASGARI.

FP 2004 13

**BROODSTOCK NUTRITION OF AUSTRALIAN RED CLAW CRAYFISH
CHERAX QUADRICARINATUS (VON MARTENS)**

By

LADAN ASGARI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy**

January 2004



Dedicated

To my family for their support..

...and to all enthusiasts of this field.



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**BROODSTOCK NUTRITION OF AUSTRALIAN RED CLAW CRAYFISH
CHERAX QUADRICARINATUS (VON MARTENS)**

By

LADAN ASGARI

January 2004

Chairman: Che Roos Saad, Ph.D.

Faculty: Agriculture

A series of experiments were conducted to develop least cost feed formulations for red claw crayfish *Cherax quadricarinatus* broodstock based on the gonad, egg protein and essential amino acids content . All diets were developed using least cost feedstuffs and a linear interactive and discrete optimiser software (LINDO).

Prior to evaluation of different formulated diets, the protein and amino acid contents of red claw crayfish broodstock tissues, eggs and newly hatched juvenile under a culture condition were analysed. Increases in ovarian and embryonic proteins content from 10.48 to 65.29 % were observed with the maturation stages of red claw crayfish *C. quadricarinatus*. These results emphasized the critical role of protein as a main structural component in maturation. However, amino acid compositions of gonad and embryonic

tissue of *C. quadricarinatus* were generally remained constant with the possible exception of histidine.

A comparative study on performance of five dietary protein levels (30-50%) on red claw crayfish was carried out. Based on spawning rate, fecundity, hatchability and egg size, a range of 40-45% crude protein and 16.72 kJ g^{-1} ($400 \text{ kcal } 100 \text{ g}^{-1}$) energy were the best/optimal for red claw crayfish broodstock. Essential amino acids index (EAAI) of all test diets (developed based on available least cost ingredients for red claw crayfish broodstock) were within the best range (0.960-0.996) recommended for crustaceans. Therefore, all test diets were best in terms of protein and essential amino acids sources. In term of biochemical composition, an increase in dietary protein level was also increased the protein content in eggs and muscle. A positive relationship between dietary protein level and protein contents of eggs and muscle of red claw broodstock was found.

A following study later showed that a diet providing 40% protein and 16.72 kJ g^{-1} ($400 \text{ kcal } 100 \text{ g}^{-1}$) energy or 45% protein and $14.65 - 16.72 \text{ kJ g}^{-1}$ ($350 - 400 \text{ Kcal } 100 \text{ g}^{-1}$) energy could support a higher eggs production and hatchability in red claw broodstock. In this study, at a constant dietary protein level, broodstock performance increased with an increase in dietary energy level. Whereas, feeding red claw crayfish broodstock with lower energy diets resulted in a noticeable reduction in spawning rate and a delay of 2-3 weeks in the time of onset of spawning. The extremely low performance of red claw crayfish broodstock fed with low dietary energy content suggested the important role of energy in its reproduction and metabolism.

This study also demonstrated that different protein-energy ratio diets developed based on least cost ingredients were highly digestible for red claw crayfish broodstock. This results indicated that protein digestibility coefficients obtained for all formulated diets using external marker (Cr_2O_3) and internal markers (crude fibre and ash) were very similar to one-another. This suggested that crude fibre and ash ratio techniques alongside of Cr_2O_3 as a standard marker can also be used as reliable digestibility markers in red claw crayfish nutrition study.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMAKANAN INDUK UDANG AUSTRALIAN *CHERAX*
QUADRICARINATUS (VON MARTENS) DARI**

Oleh

LADAN ASGARI

Januari 2004

Pengerusi: Che Roos Saad, Ph.D.

Fakulti : Pertanian

Satu siri eksperimen telah dijalankan untuk menghasilkan rumusan makanan berkos rendah berdasarkan protein gonad, telur dan kandungan asid amino perlu pada induk udang *Cherax quadricarinatus*. Semua makanan dihasil menggunakan bahan-bahan kos rendah dan dirumus dengan perisian interaktif linear dan pengoptima diskrit (LINDO).

Sebelum menilai rumusan diet yang berlainan, satu kajian telah dijalankan ke atas kandungan protein dan asid amino bagi tisu induk udang, telur dan juwana yang baru menetas semasa pengkulturan. Satu peningkatan di dalam kandungan protein ovari dan embrio dari 10.48 kepada 65.29 % didapati dengan peningkatan peringkat kematangan induk udang *C. quadricarinatus*. Keputusan ini menekankan peranan protein sebagai satu komponen struktural di dalam kematangan. Walaubagaimanapun komposisi asid amino

pada gonad dan tisu embrio *C. quadricarinatus* secara amnya adalah sama kecuali asid amino histidina.

Satu kajian bandingan ke atas prestasi lima peringkat (30-50%) protein diet berkos rendah telah dijalankan. Berdasarkan kadar pembiakan, fekunditi, penetasan dan saiz telur, satu julat 40-45% protein kasar dan 16.72 kJ g^{-1} ($400 \text{ kcal } 100\text{g}^{-1}$) tenaga telah didapati sebagai julat optima di dalam perumusan diet berkos rendah untuk induk udang. Indeks asid amino perlu (EAAI) dalam semuadiet berdasarkan kepada bahan-bahan berkos rendah yang ada untuk induk udang berada di dalam julat yang terbaik (0.960-0.996) yang dicadangkan untuk krustasea. Oleh itu, diet yang dirumus adalah terbaik dari segi sumber protein dan asid amino perlu. Dari segi komposisi biokimia, secara amnya, peningkatan takat protein dalam akan diet meningkatkan kandungan protein dalam telur dan otot. Terdapat hubungan positif di antara kandungan dalam diet dan kandungan protein telur serta otot induk udang.

Kajian turut menunjukkan bahawa diet yang mengandungi 40% protein dan 16.72 kJ g^{-1} ($400 \text{ kcal } 100\text{g}^{-1}$) tenaga atau 45% protein dan $14.62 - 16.72 \text{ kJ g}^{-1}$ ($350-400 \text{ Kcal } 100\text{g}^{-1}$) tenaga boleh membantu pengeluaran telur dan penetasan yang lebih tinggi oleh induk udang. dalam kajian ini, pada takat protein diet yang tetap, prestasi induk turut meningkat dengan meningkatnya takat tenaga dalam diet. Pemberian makanan dengan diet bertenaga rendah kepada induk udang menghasilkan penurunan kadar pembiakan dan melambatkan permulaan pembiakan selama 2-3 minggu. Prestasi rendah oleh induk udang yang diberi makan diet dengan kandungan tenaga rendah

telah menunjukkan penting tenaga di dalam pembiakan dan metabolismenya.

Kajian ini juga menunjukkan nisbah tenaga protein yang berbeza di dalam diet yang dirumuskan dari bahan-bahan berkong rendah adalah sangat mudah dihadamkan oleh induk udang. Kajian seterusnya juga menunjukkan bahawa koefisien kadar penghadaman untuk semua diet yang dirumus dengan menggunakan penanda luar (Cr_2O_3) dan penanda dalaman (gentian kasar dan abu) adalah sama ($P>0.05$). Keputusan ini mencadangkan di samping Cr_2O_3 sebagai penanda piawai, teknik nisbah gentian kasar dan abu turut boleh digunakan sebagai penanda penghadaman untuk kajian makanan udang.

ACKNOWLEDGEMENTS

First of all, I would like to express my deepest gratitude, appreciation and thanks to Dr. Che Roos Saad, chairman of my supervisory committee, for his kind supervision, assistance and encouragement during the period of my doctoral studies. I am also deeply indebted to him for provision of the necessary funding, laboratory and hatchery facilities and the friendly environment during my study. My sincere thanks are also extended to Assoc. Prof. Dr. Abd. Razak Alimon whose assistance and support were essential factors in commencement and completion of my Ph.D. degree.

I would also like to thank Assoc. Prof. Dr. Mohd. Salleh Kamarudin who provided me with an ample opportunity to explore research ideas in his well-equipped laboratory. I am also gratefully acknowledging his critical reading and suggestions of my Ph.D. thesis. I gratefully acknowledge constructive advice of my co-supervisor Dr. Hishamuddin Omar. Staff of Faculty of Agriculture and Institute of Bioscience at Universiti Putra Malaysia receive my appreciation for their support during my studies in Malaysia as well.

I would also like to express my sincerest appreciation and thanks to:

- Mr. Ismail from Food Analysis Laboratory of Malaysian Veterinary Department, Petaling Jaya, Kuala Lumpur who sincerely trained and helped me in amino acids analysis;
- The staff of Aquatic Resources Technology Laboratory, Faculty of Agriculture, UPM especially Mr. Jasni for their valuable assistance;

- Dr. Paymon Roustaian, Dr. Hamid Rezai and Dr. Farshad Shishechian for their pure friendship and support;
- My lab mates in Aquatic Biotechnology Laboratory, Dr. Annie Christianus, Mannuel, Emil, Dr. Reza Rafiee and Carina for their sincere friendship and assistance during the conduct of this study;
- My mother and my father for their love, understanding, support, principle guide and encouragement ;
- My sister Yasmin for her. everyday moral and spiritual support, encouragement, guidance and her concern during my study; Mr. Kambiz Shamsi for his critical review of my thesis and his editorial assistance, my only brother Alireza for his kind and lovely support
-
- My husband, Dr. Samad, for his support and encouragements. It was with his company which enabled me to overcome every obstacle in the way of my study;

And above all, to the GOD almighty for making this study possible.

TABLE OF CONTENT

	page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxi
 CHAPTER	
 I INTRODUCTION	 1
The Background of Study	1
Statement of Problem	3
The Significance of Study	5
Objectives	7
 II LITERATURE REVIEW	 8
The Biology, Culture and Nutrition of Red Claw Crayfish <i>Cherax quadricarinatus</i>	9
Taxonomy	9
Distribution	10
Ecology	10
Reproduction	11
Hatchery	12
Egg Incubation	13
Morphology of Eggs	14
Fecundity	14
Water Quality for Red Claw Crayfish	15
Nutrition of Crayfish	16
Natural Food	16
Formulated Food	17
The Role of Protein Compounds in Crustaceans	18
The Role of Protein Compounds in Crustacean's Broodstock	20
The Role of Protein Compounds in Fish Broodstock	22
Least Cost Feed Formulation	24
Digestion	26
 III GENERAL METHODOLOGY	 30
Location of Study	30
Experimental Crayfish	30
Fibreglass Experimental Tank	31
Biological Filter	31
Least Cost Feed Formulations	31
Food Preparation	35

Crude Protein	35
Lipids	36
Moisture Determination	37
Crude Fibre	37
Ash Determination	38
Gross Energy Determination	39
Amino Acid Analysis	40
Preparation of Glassware	40
Determination of Amino Acid	40
Acid Hydrolysis	41
Performic Acid Oxidation	41
Alkaline Hydrolysis	42
Start up and operation	42
Chromic Oxide	43
Water Quality Monitoring	43
Statistical Analyses	44
 IV	
PROTEIN AND AMINO ACID CONTENTS OF BROODSTOCK TISSUES, EGGS AND NEWLY HATCHED JUVENILE OF CULTURED RED CLAW CRAYFISH, <i>CHERAX QUADRICARINATUS</i>	45
Introduction	45
Materials and Methods	46
Results	47
Discussion	53
Conclusion	56
 V	
EFFECTS OF DIETARY PROTEIN LEVEL ON SPAWNING PERFORMANCE AND EGG PROTEIN COMPOSITION OF RED CLAW CRAYFISH <i>CHERAX QUADRICARINATUS</i> BROODSTOCK	57
Introduction	57
Materials and Methods	59
Test Diets	59
Experimental Animals	61
Experimental Tanks	61
Experimental Procedure	61
Water Quality	62
Chemical Analysis	63
Evaluation	63
Statistical Analysis	64
Results	64
Discussion	76
Conclusion	82
 VI	
EFFECTS OF DIETARY PROTEIN ENERGY RATIO ON REPRODUCTION PERFORMANCE AND BIOCHEMICAL COMPOSITION OF EGGS OF RED CLAW CRAYFISH <i>CHERAX QUADRICARINATUS</i> BROODSTOCK	83
Introduction	83
Materials and Methods	85
Diets	85
Experimental Animals	89

Experimental Tanks	89
Experimental Procedure	89
Water Quality	90
Chemical Analyses	90
Feed Evaluation	91
Statistical Analysis	91
Results	92
Discussion	103
Conclusion	108
 VII APPARENT NUTRIENT DIGESTIBILITY OF LEAST COST DIETS BY AUSTRALIAN RED CLAW CRAYFISH, <i>CHERAX QUADRICARINATUS</i> USING INTERNAL AND EXTERNAL MARKERS	 109
Introduction	109
Materials and Methods	111
Experimental Diet	111
Fibreglass Experimental Tanks	112
Experimental Animals	112
General Design	112
Biochemical Analysis and Calculation	113
Chromic Oxide	114
Statistical Analyses	115
Results	115
Discussion	126
Conclusion	131
 VII GENERAL DISCUSSION	 132
REFFRENCES	137
APPENDICES	160
BIODATA OF THE AUTHOR	170

LIST OF TABLES

Table	page
4.1: Protein contents of gonad of <i>C. quadricarinatus</i> at different ovarian stages	48
4.2: Morphological characteristics and percentage of protein for various stages of <i>C. quadricarinatus</i> embryonic development.	49
4.3: The amino acid compositions of gonad, egg and newly hatched juvenile of red claw crayfish (g 100 g ⁻¹ sample)	50
4.4: Essential amino acid (A/E) ratios of gonad, egg and newly hatched juvenile of red claw crayfish <i>C. quadricarinatus</i> .	52
5.1: Composition and proximate composition of the experimental diets for red claw crayfish <i>C. quadricarinatus</i> (g 100 g ⁻¹) expressed in percent dry weight and cost of the diets	60
5.2: Essential amino acid (EAA) composition of different diets (g 100g ⁻¹ feed) expressed in dry weight	66
5.3: Effect of different protein level on fecundity, hatchability, spawning and egg quality of crayfish broodstock <i>C. quadricarinatus</i>	67
5.4: Results of multiple regression analysis between spawning rate, fecundity, egg size and hatchability of red claw broodstock (Dependent variables) and dietary nutritional factors.	69
5.5: Biochemical composition of egg, gonad and muscle of red claw crayfish <i>C. quadricarinatus</i> fed with different protein level diets	72
5.6: EAA composition of egg (g 100 g ⁻¹ sample) with different protein levels diets	73
5.7: EAA composition of larvae (g 100 g ⁻¹ sample) of red claw crayfish broodstock fed on diet with different protein levels	74
5.8: EAA composition of muscle (g 100 g ⁻¹ sample) of red claw crayfish broodstock fed on diet with different protein levels	75

6.1: Composition and proximate composition of the experimental diets for red claw crayfish <i>C. quadricarinatus</i> (g 100 g ⁻¹) expressed in percent dry weight and cost of the diets	86
6.2: Essential amino acid (EAA) composition of different diets (g 100 g ⁻¹ feed) expressed in dry weight	88
6.3: Effect of different protein (energy) level on fecundity, hatchability, spawning and egg quality of crayfish broodstock <i>C. quadricarinatus</i>	93
6.4: Results of two-way ANOVA and mean multicomparisons (Tukey test) that were performed to calculate the effects of experimental diets variables by protein and energy level on red claw crayfish broodstock performance	94
6.5: Results of multiple regression analysis between spawning rate, fecundity, egg size and hatchability of red claw broodstock (Dependent variables) and dietary nutritional factors	96
6.6: Biochemical composition of egg, gonad and muscle of red claw crayfish <i>C. quadricarinatus</i> fed with different protein (energy) level diets	97
6.7: EAA composition of eggs (g 100 g ⁻¹ sample) with different protein (energy) level diets	98
6.8: EAA composition of Larvae (g 100 g ⁻¹ sample) with different protein (energy) level diets	99
6.9: EAA composition of muscle (g 100 g ⁻¹ sample) with different protein (energy) level diets	100
6.10: Results of two-way ANOVA and mean multicomparisons (Tukey test) that were performed to	101
7.1: Composition and proximate composition of the experimental diets in red claw crayfish <i>C. quadricarinatus</i> (g 100 g ⁻¹) expressed in percent dry weight	118
7.2: Composition and proximate composition of the experimental diets for red claw crayfish <i>C. quadricarinatus</i> (g 100 g ⁻¹) expressed in percent dry weight, which used in experiment trail 2	119
7.3: Proximate composition of red claw crayfish faeces fed with different protein (energy) level diets	120
7.4: Protein digestibility coefficients (\pm SE) of different diets using different external and internal markers	121

- 7.5: Comparative protein digestibility coefficients (\pm SE) in different dietary protein (energy) level using several digestibility markers 121
- 7.6: Lipid digestibility coefficients (\pm SE) using different external and internal markers 122
- 7.7: Energy digestibility coefficients (\pm SE) using different external and internal markers. 122
- 7.8: Results of two-way ANOVA that were performed to calculate the interaction effect of dietary protein and energy content on digestibility of dietary protein, lipid and digestible energy in red claw crayfish broodstock using different external and internal marker 125

LIST OF FIGURES

Figure	page
4.1:Protein content of various stages of gonads and embryo	49
5.1:Relationship between dietary protein level and a) egg protein content, b) gonad protein content and c) muscle protein content	70
7.1:Relationship between dietary fibre content and a) Protein digestibility coefficient b) Lipid digestibility coefficient and c) Energy digestibility coefficient	123
7.2:Relationship between a) Dietary energy content and Energy digestibility coefficient b) Dietary lipid content and lipid digestibility coefficient	124

LIST OF ABBREVIATIONS

AABA	L- α -Amino- η -Butric Acid
AND	Apparent Nutrient Digestibility
ANOVA	Analysis of Variance
AOAC	Association of Official Analytical Chemist
AQB	Aquabind
Arg	Arginine
BHT	Butylated hydroxy toluene
BM	Bone meal
BW	Body weight
CAP	Calcium Propionate
CF	Crude fibre
CH ₃ OH	Methanol
CMC	Carboxy methyl α cellulose
COP	Copra meal
COT	Cottonseed meal
CP	Crude protein
DE	Digestible energy
DMD	Dry Matter Digestibility
DMRT	Duncan Multiple Range Test
DO	Dissolved oxygen
DP	Digestible protein
DW	Dry weight
EAA	Essential amino acid
EAAI	Essential amino acid index

FM	Fish meal
ha	hectare
HCL	Hydrochloric acid
His	Histidine
Ileu	Iso leucine
Kcal	Kilo calorie
Leu	Leucine
LINDO	Linear interactive and discrete optimiser
LV	Lipovitellin
Lys	Lysine
µm	Microgram
Meth	Methionine
MN	Mineral
MT	Metric tonne
N	Normality
NaOH	Sodium hydroxide
NRC	National research council
P/E	Protein / Energy
P: E ratio	Protein: Energy ratio
P30	Protein 30 percent
Phe	Phenylalanine
PO	Palm oil
ppm	Part per million
RB	Rice bran
- RM	Ringgit Malaysia
SAS	Statistical Analysis System

SBM	Soybean meal
SD	Standard deviation
SE	Standard error
SEAFDEC	Southeast Asian Fisheries Development Centre
SEM	Standard error of mean
SM	Shrimp meal
TAP	Tapioca flour
Thr	Threonine
TRT	Treatment
Trypt	Tryptophan
UPM	Universiti Putra Malaysia
Val	Valine
VC	Vitamin C
VG	Vitelloenin
VITP	Vitamin premix
VP	Vitellin, egg yolk protein

CHAPTER I

INTRODUCTION

The Background of Study

The total harvest of the world capture fishery reached a maximum level in 1989 (New, 1991) and has recently entered a period of decline (Garcia and Newton, 1994). The decline has been brought by multiple factors including increasing demand, development of wide ranging fishing and processing vessels (Larkin, 1991), environmental degradation and climate change (Donaldson, 1996). Thus, aquaculture production of various fish and shellfish species has been developed as an alternative for fishery industries in many countries, and further development is expected as the demand for fishery products continues to increase and supply from natural sources becomes more limited (Gatlin, 1996). Aquaculture is increasingly recognized as a sustainable means of producing high quality aquatic food for human consumption. Production has increased rapidly during the last decade and especially over the last half decade to the extent that aquaculture accounts for over 20% of the world production of aquatic foods of marine and freshwater origin (Donaldson, 1996).

Among the aquaculture industries, crayfish farming is one of the few industries developed in the United States, European countries and Australia in the latter half of this century (Brown, 1995). Total world production of freshwater crayfish is estimated to be around 70,000 to 100,000 metric tons (MT) per annum (Huner, 1989). Although, the single most important species