



**UNIVERSITI PUTRA MALAYSIA**

**ELECTROCHEMICAL CHARACTERISATION AND APPLICATION OF  
SOLID ELECTRODES CHEMICALLY MODIFIED WITH C60-  
FULLERENE, TETRATHIAFULVALENE AND SILICON  
POLYOXOTUNGSTATE**

**GOH JOO KHENG.**

**FS 2005 9**



**ELECTROCHEMICAL CHARACTERISATION AND APPLICATION OF  
SOLID ELECTRODES CHEMICALLY MODIFIED WITH C<sub>60</sub>-FULLERENE,  
TETRATHIAFULVALENE AND SILICON POLYOXOTUNGSTATE**

**By**

**GOH JOO KHENG**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia in  
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**November 2005**



Dedicated to my beloved parents, Shawn and Kelvin Choi



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**ELECTROCHEMICAL CHARACTERISATION AND APPLICATION OF  
SOLID ELECTRODES CHEMICALLY MODIFIED WITH C<sub>60</sub>-  
FULLERENE, TETRATHIAFULVALENE AND SILICON  
POLYOXOTUNGSTATE**

By

**GOH JOO KHENG**

**November 2005**

**Chairman : Associate Professor Tan Wee Tee, PhD**

**Faculty : Science**

The modified glassy carbon electrodes by adhered microcrystals of fullerene (C<sub>60</sub>), tetrathiafulvalene (TTF) and  $\alpha$ -Keggin silicon polyoxotungstate, [SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> has been studied extensively in characterization and application of these modified electrode in mediating some selected compounds such as cysteine, methionine and nitrite ions in the presence of aqueous electrolytes.

The use of a glassy carbon electrode modified by adhered microcrystals of fullerene (C<sub>60</sub>), mediates the oxidation of cysteine in the presence of aqueous potassium containing electrolytes. Under conditions of cyclic voltammetry, the potential for the oxidation of cysteine was lowered by approximately 100 mV and



current was enhanced significantly relative to the situation prevailing when a bare glassy carbon electrode was used. Additional mediation occurs when the potential range covered includes that of  $C_{60}/C_{60}^{n-}$  redox couples. The sensitivity under condition of cyclic voltammetry was significantly dependent on pH, temperature and  $C_{60}$  dosage. Excellent analytical and recovery data were obtained with vitamin pill, cassamino acid (hydrolyzed casein) and for a range of beverages.

The usefulness of a  $C_{60}$ -fullerene modified gold (Au) electrode was extended in mediating the oxidation of methionine in the presence of potassium ions electrolyte. During cyclic voltammetry, an oxidation peak of methionine appearing at +1000 mV vs. Ag/AgCl was observed. The oxidation current of methionine was enhanced by about 2 times using a modified gold electrode. The current enhancement was significantly dependent on pH, temperature and  $C_{60}$  dosage. The variation of scan rate study shows that the system undergoes diffusion-controlled process.

Then, diffusion coefficient and rate constant of methionine were determined using hydrodynamic method (rotating disk electrode) with values of  $1.11 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$  and  $0.0026 \text{ cm s}^{-1}$  respectively for unmodified electrode while the values of diffusion coefficient and rate constant of methionine using  $C_{60}$  modified Au electrode were  $5.67 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$  and  $0.0021 \text{ cm s}^{-1}$  respectively. Excellent analytical and recovery data were obtained with root beer beverage.



Tetrathiafulvalene modified glassy carbon electrode has been characterized and been applied in the determination of L-cysteine at the potential of +800 mV vs. Ag/AgCl. It has shown to catalyze the oxidation peak of cysteine in the alkaline buffer solution under cyclic voltammetry conditions. The sensitivity of tetrathiafulvalene modified electrode was found to be dependent on pH and dosages of varying concentration of TTF in acetonitrile. Excellent analytical and recovery rates also has been obtained using vitamin pills and root beer syrup either with known amount of cysteine or added deliberately into the samples.

The voltammetric reduction of  $\alpha$  form of Keggin silicon polyoxotungstate anions,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> has been carried out under acidic conditions. The cyclic voltammograms obtained from the solution voltammetry of the Keggin polyoxotungstate anions,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> showed cyclic voltammogram of a two reversible, one-electron process and a reversible two-electron process in the potential range of +200 mV to -1000 mV (vs. Ag/AgCl). However, only the first reduction process was undertaken in this study.

The use of Keggin silicon polyoxotungstate anion,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> was successfully applied to the synthesis of ruthenium bipyridine silicon polyoxotungstate, [Ru(bipy)<sub>3</sub>]<sub>2</sub>[ $\alpha$ -SiW<sub>12</sub>O<sub>40</sub>] via reduction of  $\alpha$ -[TBA]<sub>4</sub>[SiW<sub>12</sub>O<sub>40</sub>] solvent cast onto the glassy electrode in sulphuric acid. The reduction of nitrite has shown to be catalyzed by [Ru(bipy)<sub>3</sub>]<sub>2</sub>[ $\alpha$ -SiW<sub>12</sub>O<sub>40</sub>] modified electrode at lower pH of aqueous sulphuric acid media. Bulk electrolysis technique was employed to

reduce  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> to form a blue solution of  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup>. The reduction of  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> to  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup> was measured spectrophotometrically and there was no stable complex formation between  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup> and reduction product of NO<sub>2</sub><sup>-</sup> species in the acidic solution. Therefore, the mechanism involving the catalytic effect on  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> in the presence of nitrite suggested that nitrous acid, HNO<sub>2</sub> has been reduced to a product that possibly containing NO. Beta,  $\beta$  form of Keggin [SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> was found to be less active as compared to the  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup>.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**ELEKTROKIMIA PENCIRIAN DAN APLIKASI ELEKTROD PEPEJAL  
MODIFIKASI KIMIA DENGAN C<sub>60</sub>-FULLERENE,  
TETRATHIAFULVALENE DAN SILIKON POLYOXOTUNGSTAT**

Oleh

**GOH JOO KHENG**

**November 2005**

**Pengerusi : Profesor Madya Tan Wee Tee, PhD**

**Fakulti : Sains**

Modifikasi elektrod karbon kaca melalui pelekatan mikrokrystal fullerene (C<sub>60</sub>), tetrathiafulvalene (TTF) dan  $\alpha$ -Keggin silikon polyoxotungstat,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> telah di kaji dengan terperinci dalam pencirian and aplikasi elektrod-elektrod ini sebagai perantaraan sebatian terpilih, contohnya, sistina, metionina dan ion nitrit dalam elektrolit akues.

Penggunaan modifikasi elektrod karbon kaca melalui pelekatan mikrokrystal fullerene C<sub>60</sub> telah meningkatkan pengoksidaan sistina dalam kehadiran elektrolit akues yang mengandungi ion kalium. Dalam keadaan voltametri siklik, potensi bagi pengoksidaan sistina telah direndahkan kira-kira 100 mV dan arus ditingkatkan relatif kepada situasi di mana elektrod karbon kaca tanpa modifikasi digunakan. Pertambahan mediasi telah berlaku apabila skala potensi meliputi



proses redoks  $C_{60}/C_{60}^{n-}$ . Sensitiviti di bawah keadaan siklik voltametri adalah bergantung kepada pH, suhu, dan dosej  $C_{60}$ . Data analisis dan perolehan semula yang baik diperolehi menggunakan pil vitamin, asid cassamino (casein yang dihidrolisis) dan juga beberapa jenis minuman.

$C_{60}$ -fullerene terlekat pada elektrod emas juga digunakan bagi mediasi pengoksidaan metionina dengan kehadiran ion kalium dalam elektrolit akues. Puncak oksidasi methionine diperhatikan muncul pada potensi +1000 mV (vs. Ag/AgCl) semasa siklik voltametri. Arus pengoksidaan methionine telah ditingkatkan sebanyak 2 kali dengan penggunaan modifikasi elektrod emas. Peningkatan arus ini adalah bergantung kepada pH, suhu, dan dosej  $C_{60}$ . Kadar imbasan yang berbeza menunjukkan sistem ini melalui proses terkawal resapan.

Justeru itu, pemalar penyerapan and pemalar kadar methionine telah diperolehi menggunakan cara hidrodinamik (elektrod cakera putar) dengan nilai masing-masing  $1.11 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$  dan  $0.0026 \text{ cm s}^{-1}$  untuk elektrod tanpa modifikasi. Manakala, pemalar penyerapan and pemalar kadar metionina untuk  $C_{60}$  modifikasi elektrod emas adalah masing-masing  $5.67 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$  and  $0.0021 \text{ cm s}^{-1}$ . Data analisis dan perolehan semula yang baik didapati menggunakan minuman root bir.

Modifikasi elektrod dengan tetrathiafulvalene telah dicirikan dan juga diaplikasikan dalam penentuan kehadiran sistina pada potensi +800 mV vs. Ag/AgCl. Ia menunjukkan katalisis puncak pengoksidaan sistina dalam penimbal alkali elektrolit di bawah keadaan siklik voltametri. Sensitiviti tetrathiafulvalene modifit elektrod didapati bergantung kepada pH, dosej TTF/ACN, dan kepekatan TTF/ACN. Data analisis dan perolehan semula yang baik telah dicapai dengan penggunaan pil vitamin and root bir sirup samada dengan kepekatan sistina yang diketahui ataupun yang ditambahkan kepada sampel.

Voltametri bagi penurunan bentuk Keggin silikon polyoxotungstat anion,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> telah diperolehi di bawah keadaan berasid. Siklik voltamogram yang diperolehi daripada larutan voltametri Keggin polyoxomethalate anion,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> menunjukkan dua proses berbalik satu elektron dan satu proses berbalik dua elektron pada potensi antara +200 mV to -1000 mV (vs. Ag/AgCl). Walau bagaimanapun, hanya puncak penurunan yang pertama di ambil kira dalam projek ini.

Penggunaan jenis Keggin silikon polyoxotungstat anion,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> telah berjaya diaplikasikan kepada sintesis [Ru(bipy)<sub>3</sub>]<sub>2</sub>[ $\alpha$ -SiW<sub>12</sub>O<sub>40</sub>] melalui penurunan dalam asid sulfurik  $\alpha$ -[TBA]<sub>4</sub>[SiW<sub>12</sub>O<sub>40</sub>] yang dilekat atas elektrod karbon kaca menggunakan cara pengewapan pelarut. Penurunan nitrit, NO<sub>2</sub><sup>-</sup> telah dikatalis oleh penurunan ruthenium bipyridine silikon polyoxotungstat, [Ru(bipy)<sub>3</sub>]<sub>2</sub>[ $\alpha$ -SiW<sub>12</sub>O<sub>40</sub>] modifikasi elektrod pada pH rendah asid sulfuric.

Teknik elektrolisis pukal telah digunakan untuk menurunkan  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> kepada larutan biru,  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup>. Penurunan  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> to  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup> telah dianalisis dengan spektrofotometer dan didapati tiada kompleks yang stabil terbentuk antara  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>5-</sup> dan produk penurunan NO<sub>2</sub><sup>-</sup> spesies dalam larutan  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup>. Dengan demikian, mekanisma melibatkan kesan katalisis  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> dalam kehadiran NO<sub>2</sub><sup>-</sup> telah dicadangkan bahawa asid nitrous, HNO<sub>2</sub> telah diturunkan ke suatu produk yang mengandungi nitric oxida, NO. Bentuk beta,  $\beta$  Keggin [SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup> telah didapati kurang elektrokatalitik aktif dalam kehadiran nitrit berbanding dengan  $\alpha$ -[SiW<sub>12</sub>O<sub>40</sub>]<sup>4-</sup>.

## ACKNOWLEDGEMENTS

I wish to express my most sincere thanks and deepest appreciation to my supervisor, Associate Professor Dr. Tan Wee Tee for his continuous guidance, constant encouragement, invaluable advices and valuable comments throughout the course of this project. I wish to extend my sincere thanks to my supervisory committee members Professor Dr. Zulkarnain Zainal and Professor Dr. Anuar Kassim for their support, advices and guidance.

I would like to convey many thanks and deepest gratitude to Professor Alan M. Bond and Dr. Jie Zhang for their constant support, valuable advices, comments and guidance throughout my research attachment in School of Chemistry, Clayton Campus, Monash University, Australia for three months commencing October 2004. Also thanks to the staff of School of Chemistry, Clayton Campus, Monash University, Mr. Victor De Guzman and Mrs. Sally Duck for their help spectrophotometry measurements.

I wish to express many thanks and appreciation to my colleagues, Ei Bee, Lai Peng and Zidan for their support and encouragement throughout the research project. Besides, I would also like to acknowledge the financial support provided by the IRPA programme.



Finally, I wish to convey my deepest appreciation and gratefulness to my parents, brother, and friends for their constant moral support, encouragement and assistance during the course of my research. Many thanks and appreciations are extended to Kelvin Choi for his continuous encouragement and invaluable help that have contributed towards the success of this project.



## TABLE OF CONTENTS

	Page
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	vii
<b>ACKNOWLEDGEMENTS</b>	xi
<b>APPROVAL</b>	xiii
<b>DECLARATION</b>	xv
<b>LIST OF TABLES</b>	xxii
<b>LIST OF FIGURES</b>	xxiv
<b>LIST OF ABBREVIATION OF TERMS</b>	xxxv

CHAPTER	Page
<b>1 INTRODUCTION</b>	
1.1 Chemically Modified Electrode	1
1.2 Electroanalysis	2
1.3 Solid State Electrochemistry	4
1.4 Fundamental of Voltammetry Techniques	7
1.4.1 Linear Sweep Voltammetry	7
1.4.2 Cyclic Voltammetry	9
1.5 Fundamental of Chronoamperometry	13
1.6 Fundamental of Chronocoulometry	15
1.7 Buckminsterfullerene, C <sub>60</sub>	16
1.8 Tetrathiafulvalene	19
1.9 Silicon Polyoxometalate	21
1.10 L-cysteine	23
1.11 Methionine	25
1.12 Objectives	27
<b>2 LITERATURE REVIEWS</b>	
2.1 Solid Phase Voltammetry	28
2.2 Reviews on Buckminsterfullerene, C <sub>60</sub>	38
2.3 Reviews on Tetrathiafulvalene, TTF	51
2.4 Reviews on Polyoxometalates	56
2.5 Reviews on Cysteine and Methionine	59
<b>3 METHODOLOGY</b>	
3.1 Materials and Reagents	71
3.2 Instrumentation and Apparatus	72



3.2.1	BAS 100W and CV 50W Voltammetric Analyzer	72
3.2.2	Working Electrode (WE)	72
3.2.3	Counter Electrode (CE)	73
3.2.4	Reference Electrode (RE)	74
3.2.5	Voltammetric Cell	75
3.2.6	Ultra Violet Visible, UV-Vis and Liquid Chromatography Mass Spectrophotometer, LCMS	76
3.3	Procedures for Preparation of Chemically Modified Electrodes	77
3.3.1	Mechanically Attachment (MA)	77
3.3.2	Solvent Casting (SC)	78
3.3.2.1	Preparation of Fullerene-C <sub>60</sub> Modified Electrode via Solvent Casting	78
3.3.2.2	Preparation of TTF Modified Electrode via Solvent Casting	79
3.3.2.3	Preparation of Ruthenium Bipyridine Silicon Polyoxotungstate, [Ru(bipy) <sub>3</sub> ] <sub>2</sub> [α-SiW <sub>12</sub> O <sub>40</sub> ] Modified Electrode	79
3.3.3	General Procedures for Electrochemical Measurement	80
3.4	Procedures for Data Collections on Characterization of Fullerene-C <sub>60</sub> Modified Electrode	81
3.4.1	Cyclic Voltammograms of C <sub>60</sub> Adhered to Electrode Surface via Mechanical Attachment and Solvent Casting	81
3.4.2	Effect of the Nature of Anions and Cations in the Electrolytes	81
3.4.3	Effect of Varying Scan Rate	82
3.4.4	Chronocoulometry of C <sub>60</sub>	82
3.4.5	Scan Reversal Studies of C <sub>60</sub>	83
3.5	Procedures for Data Collection on Characterization of C <sub>60</sub> -Modified Electrode on Cysteine: Scanning Potential Range I: C <sub>60</sub> Electroinactive	83
3.5.1	Enhancement Study on Cysteine	83
3.6	Procedures for Data Collections on Application of C <sub>60</sub> -Modified Electrode on Cysteine: Scanning Potential Range II: C <sub>60</sub> Electroactive	84
3.6.1	Enhancement Study on Cysteine	84
3.6.2	Effect of Potential Cycling	85
3.6.3	Effect of Varying C <sub>60</sub> Dosages	85
3.6.4	Calibration Graph	85

3.6.5	Cysteine Determination / Recovery Experiment	86
3.7	Procedures for Data Collections on Application of C <sub>60</sub> -Modified Gold Electrode on Methionine: Scanning Potential Range: C <sub>60</sub> Electroinactive	87
3.7.1	Enhancement Study on Methionine	87
3.7.2	Effect of Varying pH	88
3.7.3	Effect of Varying Temperature	88
3.7.4	Effect of Varying C <sub>60</sub> Dosage	88
3.7.5	Effect of Varying Scan Rate	89
3.7.6	Hydrodynamic Voltammetry using Rotating Disk Electrode (RDE)	89
3.7.7	Calibration Graph	89
3.7.8	Methionine Determination / Recovery Experiment	90
3.8	Procedures for Data Collections on Characterization of Tetrathiafulvalene (TTF) Modified Electrode	90
3.8.1	Cyclic Voltammogram of TTF Adhered to Electrode Surface via Mechanical Attachment and Solvent Casting	90
3.8.2	Effect of Different Electrodes Materials	91
3.8.3	Effect of Varying Scan Rate	91
3.8.4	Effect of Varying Electrode Sizes	92
3.8.5	Scan Reversal Studies of TTF	92
3.8.6	Cyclic Voltammogram of TTF-Modified Electrode in Different Electrolyte	92
3.9	Procedures for Data Collections on Application of Tetrathiafulvalene (TTF)-Modified Electrode	93
3.9.1	Catalytic Current of L-cysteine in Various pH of Supporting Electrolyte	93
3.9.2	Effect of Varying Concentration of TTF Dissolved in Acetonitrile	93
3.9.3	Effect of Varying Amount of TTF in Acetonitrile Deposited on Electrode	94
3.9.4	Effect of Varying Concentration of L-cysteine	94
3.9.5	Cysteine Determination / Recovery Experiment	94
3.9.6	Electrochemical Quartz Crystals Microbalance, EQCM	95
3.9.7	Liquid Chromatography Mass Spectrometer (LCMS)	95
3.10	Procedures for Data Collections on Potassium Silicon Polyoxotungstate, $\alpha$ -K <sub>4</sub> SiW <sub>12</sub> O <sub>40</sub>	96



3.10.1	Solution Voltammetry of $\alpha$ -Potassium Silicon Polyoxotungstate, $\alpha$ -K <sub>4</sub> SiW <sub>12</sub> O <sub>40</sub>	96
3.10.2	Catalytic Effect in the Presence of Nitrite Ion, NO <sub>2</sub> <sup>-</sup>	96
3.10.3	UV Spectrophotometry	97
3.10.4	Data Simulation of Varying Scan Rate and Effect of Varying Concentration of Nitrite Ion, NO <sub>2</sub> <sup>-</sup>	98
3.10.5	Ruthenium Bipyridine Silicon Polyoxotungstate Modified Glassy Carbon Electrode	98
3.10.6	Effect of Varying Concentration of Nitrite Ion, NO <sub>2</sub> <sup>-</sup>	99
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	
4.1	Characterization of Fullerene-C <sub>60</sub> Modified Electrode	100
4.1.1	Cyclic Voltammograms of C <sub>60</sub> Adhered to Electrode Surface via Mechanical Attachment and Solvent Casting	101
4.1.2	Effect of the Nature of Anions and Cations in the Electrolytes	109
4.1.3	Effect of Varying Scan Rate	114
4.1.4	Chronocoulometry of C <sub>60</sub>	116
4.1.5	Scan Reversal Studies of C <sub>60</sub>	117
4.2	Application of C <sub>60</sub> -Modified Electrode on Cysteine Scanning Potential Range I: C <sub>60</sub> Electroinactive	120
4.2.1	Enhancement Study on Cysteine	120
4.3	Application of C <sub>60</sub> -Modified Electrode on Cysteine Scanning Potential Range II: C <sub>60</sub> Electroactive	123
4.3.1	Enhancement Study on Cysteine	124
4.3.2	Effect of Potential Cycling	126
4.3.3	Effect of Varying C <sub>60</sub> Dosage	128
4.3.4	Calibration Graph	129
4.3.5	Cysteine Determination / Recovery Experiment	130
4.4	Application of C <sub>60</sub> -Modified Electrode on Methionine Scanning Potential Range: C <sub>60</sub> Electroinactive	134
4.4.1	Enhancement Study on Methionine	135
4.4.2	Effect of Varying pH	137
4.4.3	Effect of Varying Temperature	138
4.4.4	Effect of Varying C <sub>60</sub> Dosage	141
4.4.5	Effect of Varying Scan Rate	142
4.4.6	Hydrodynamic Voltammetry using Rotating Disk Electrode (RDE)	143
4.4.7	Calibration Graph	146

4.4.8	Methionine Determination / Recovery Experiment	147
4.5	Characterization of Tetrathiafulvalene (TTF) Modified Electrode	150
4.5.1	Cyclic Voltammogram of TTF Adhered onto Electrode Surface via Mechanical Attachment and Solvent Casting	151
4.5.2	Effect of Different Electrode Materials	153
4.5.3	Effect of Varying Scan Rate	156
4.5.4	Effect of Varying Electrode Sizes	159
4.5.5	Scan Reversal Studies of TTF	162
4.5.6	Cyclic Voltammogram of TTF-Modified Electrode in Different Electrolyte	165
4.6	Application of Tetrathiafulvalene (TTF)-Modified Electrode	170
4.6.1	Catalytic Current of L-cysteine in Various pH Supporting Electrolyte	170
4.6.2	Effect of Varying Concentration of TTF Dissolved in Acetonitrile	174
4.6.3	Effect of Varying Amount of TTF Dissolved in Acetonitrile Deposited on Electrode	174
4.6.4	Effect of Varying Concentration of L-cysteine	175
4.6.5	Cysteine Determination / Recovery Experiment	176
4.6.6	Electrochemical Quartz Crystals Microbalance, EQCM	179
4.6.7	Liquid Chromatography Mass Spectrometer (LCMS)	181
4.7	Potassium Silicon Polyoxotungstate, $\alpha$ -K <sub>4</sub> SiW <sub>12</sub> O <sub>40</sub>	183
4.7.1	Solution Voltammetry of $\alpha$ -Potassium Silicon Polyoxotungstate, $\alpha$ -K <sub>4</sub> SiW <sub>12</sub> O <sub>40</sub>	183
4.7.2	Catalytic Effect in the Presence of Nitrite	185
4.7.3	UV Spectrophotometry	188
4.7.4	Data Simulation of Varying Scan Rate and the Effect of Varying Concentration of Nitrite Ion, NO <sub>2</sub> <sup>-</sup>	191
4.7.5	Ruthenium Bipyridine Silicon Polyoxotungstate Modified Glassy Carbon Electrode	194
4.7.6	Effect of Varying Concentration of Nitrite Ion, NO <sub>2</sub> <sup>-</sup>	198

5	<b>CONCLUSION</b>	200
	<b>RECOMMENDATIONS FOR FURTHER STUDY</b>	203
	<b>REFERENCES</b>	204
	<b>APPENDIX</b>	214
	<b>BIODATA OF THE AUTHOR</b>	240

## LISTS OF TABLES

Table	Page
4.1 Reduction peak potential $E_{pc}$ (ii) of $C_{60}$ solvent cast onto 3 mm diameter GC electrode in 0.1 M alkali-metal salts in water with scan rate of 100 mV	110
4.2 Comparisons of crystal, hydrated ion radii, ionic conductivities, and hydration enthalpies for various first class alkali-metal cations (Burgess, 1988 and Conway, 1981)	112
4.3 Reproducibility of oxidative current of cysteine detected at $C_{60}$ /toluene coated GC electrode under specified conditions in 0.1 M $KH_2PO_4$ electrolyte. Parameters were as in Figure 4.1, current was obtained at a potential of 0.5 V.	131
4.4(a) Recovery data for L-cysteine spiked at known concentration into various samples prepared in 0.1 M $KH_2PO_4$ buffer at 7.0 using 5 $\mu$ L of a saturated $C_{60}/CH_2Cl_2$ modified GC electrode. Data was obtained using the calibration curve and case (II) method. Current was obtained at 1000 mV. Concentration of L-Cysteine added to sample = 0.20 mM	132
4.4(b) Recovery data for L-cysteine spiked at known concentrations into various samples prepared 0.1 M $KH_2PO_4$ buffer at pH 7.0 using a 5 $\mu$ L of a saturated $C_{60}/CH_2Cl_2$ modified GC electrode. Data was obtained using the calibration curve and case (II) method. Current was obtained at 1000 mV. Concentration of L-Cysteine added to sample = 0.50 mM	133
4.5 Recovery data for methionine spiked at 0.1 mM into 0.1 M $KNO_3$ at pH 6.0 using a 5 $\mu$ L of a saturated $C_{60}/CH_2Cl_2$ modified Au electrode. Data was obtained using the calibration curve. Current was obtained at 1010 mV.	148
4.6 Recovery data for methionine spiked at 0.05 mM into 0.1 M $KNO_3$ at pH 6.0 using a 5 $\mu$ L of a saturated $C_{60}/CH_2Cl_2$ modified Au electrode. Data was obtained using the calibration curve. Current was obtained at 1010 mV.	149

- 4.7 Recovery data for L-cysteine spiked at 0.2 mM into two samples prepared 0.1 M Borax at pH 10.0 using a 2  $\mu$ L of 0.2 mM TTF /ACN modified GC electrode. Data was obtained using the calibration curve (Figure 4.45). Current was obtained at 100 mV.

178



## LIST OF FIGURES

<b>Figures</b>	<b>Pages</b>	
1.1	Linear Sweep Voltammetry	8
1.2	Potential-time behavior at the working electrode following imposition of a triangular waveform	12
1.3	A typical cyclic voltammogram showing the method of extrapolation baseline and determining peak current	12
1.4	Chronoamperogram (current-time)	14
1.5	Chronocoulogram (charge-time)	16
1.6	Structure of Buckminsterfullerene, C <sub>60</sub>	18
1.7	Structure of Tetrathiafulvalene (TTF)	19
1.8	Structure of Keggin polyoxotungstate	22
1.9	Structure of L-cysteine	25
1.10	Structure of methionine	26
2.1	An electrochemistry model of solid ferrocene (Bond <i>et al.</i> , 1991)	32
2.2	A scheme of possible reactions of solid compounds in abrasive stripping voltammetry (Scholz <i>et al.</i> , 1992).	34
2.3	A schematic diagram of nucleation process proposed by Bond <i>et al.</i> (1996) occurs at oxidation of TTF and reduction of TTFBr when attached to an electrode in an aqueous electrolyte.	35
2.4	Schematic form of representation of the mechanism proposed to account for the electrochemical conversion of solid TCNQ to surface-attached microcrystals during the initial stages of redox cycling experiments at a glassy carbon electrode (Suárez <i>et al.</i> , 1999).	37

