



UNIVERSITI PUTRA MALAYSIA

**ASSESSMENT OF PUTATIVE BACTERIA AS PROBIOTICS IN
JUVENILE FRESHWATER PRAWN, MACROBRACHIUM
ROSENBERGII (DE MAN)**

MEHRAN AVAKH KEYSAMI.

FP 2006 23



**ASSESSMENT OF PUTATIVE BACTERIA AS PROBIOTICS IN JUVENILE
FRESHWATER PRAWN, *MACROBRACHIUM ROSENBERGII* (DE MAN)**

By

MEHRAN AVAKH KEYSAMI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

June 2006



DEDICATION

To my most beloved wife, Roya,
for all her understanding, patience and support during all difficulties and for
her technical help during my study

To my lovely daughter, Romina,
for making every thing worthwhile

To my dearest parents,
for their true love, constant trust, principle guide and encouragement since
my childhood

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

ASSESSMENT OF PUTATIVE BACTERIA AS PROBIOTICS IN JUVENILE FRESHWATER PRAWN, *Macrobrachium rosenbergii* (de Man)

By

MEHRAN AVAKH KEYSAMI

June 2006

Chairman: Associate Professor Che Roos Saad, PhD

Faculty : Agriculture

A study focused on the use of putative bacteria as probiotics to reduce nutritional and disease problems in aquaculture industry was carried out. This study was conducted in 8 experiments to investigate juvenile *M. rosenbergii* putative bacterial flora as probiotics, for enhancement of growth and survival of juvenile *M. rosenbergii* during November 2002 to August 2005 at University Putra Malaysia.

Bacterial counts and the putative bacterial flora in juvenile *M. rosenbergii* cultivation (0.1-9.0 g), were investigated and the isolated strains were classified to genera. Total viable cell counts were $96 \pm 2.6 \times 10^4$ CFU/ml for culture water, $3100 \pm 169.0 \times 10^4$ CFU/ml for culture tank sediment, $2100 \pm 143.0 \times 10^4$ CFU/ml for prawn body surface, $14000 \pm 4.6 \times 10^4$ CFU/g for prawn digestive tracts and $5.6 \pm 2.8 \times 10^4$ CFU/g for prawn abdominal muscle. Altogether, 12 genera were identified from water, sediment and different organs of prawn. The most frequently isolated ones were *Bacillus*, *Aeromonas* and *Pseudomonas*.



The three groups of putative bacteria (*Bacillus subtilis*, *Micrococcus luteus* and *Corynebacterium ammoniagenes*), collectively showed antagonistic activity against the six pathogens in antibacterial activity test. *B. subtilis* showed greatest inhibition of pathogens in well and disc diffusion assays and cross-streak method. Addition of 10 ml of cell free supernatant of *B. subtilis* to pathogens resulted in complete suppression of it within 12 h. Maximum growth and maximum zone of inhibition against *A. hydrophila* was observed at 30 °C and at pH 8 with 1% NaCl, respectively.

A ranking index (RI) was calculated to screen potential aquaculture probiont. The RI is based on the doubling time (t_d) and lag period (λ) obtained from the growth profile of each bacterium. Three candidates (*B. subtilis*, *M. luteus* and *C. ammoniagenes*) and two pathogens (*A. hydrophila* and *Pseudomonas aeruginosa*) were grown in TSB broth and absorbance recorded at each 0.5 h during 24 h period. The *B. subtilis* grew equally as well as the pathogens and candidate *B. subtilis* had a faster specific growth rate ($\mu=0.034$) than the other candidates, *M. luteus* (0.031) and *C. ammoniagenes* (0.032) respectively.

Sensitivity of juvenile prawns to two putative bacteria strains (*B. subtilis* and *A. hydrophila*) was examined using *in vitro* and *in vivo* pathogenicity challenge test. *In vitro* challenge test was done on juvenile prawn (0.1 g) using the immersion method. An *in vivo* challenge experiment was carried out using the feeding trial method. The result of *in vitro* and *in vivo*

pathogenicity tests indicated that the survival of juvenile (0.1 and 0.45 g) was not affected by the presence of *B. subtilis* in fresh water but, *A. hydrophila* infected juvenile *M. rosenbergii* and cause about 78.67% mortality and exhibited disease clinical signs.

A feeding trial was carried out to investigate the potential probiotic properties of *B. subtilis* and its suitable level in juvenile *M. rosenbergii*. Putative *B. subtilis* bacterium was added to commercial prawn feed as a probiotic. Six types of diets were prepared by mixing the prawn feed with *Bacillus* form in the ratio of 1:1, 2:1, 3:1, 4:1 and 5:1, with an unsupplemented control. After 60 days, the prawns fed diet at level 3:1, showed a higher mean weight gain (593.45g) or 170.29% increase in growth over control. Clearly, feed treated with *B. subtilis* appeared to enhance growth and survival of *M. rosenbergii* at a ratio of 3:1.

Another feeding trial was carried out to investigate the suitable methods of administration of *B. subtilis* to the commercial feed in juvenile *M. rosenbergii* culture. Four types of diets were prepared by mixing the prawn feed to *Bacillus* form in the ratio of 3:1 with an unsupplemented as a control. The different methods used were blending, soaking, spraying and bathing. After 60 days, the prawns fed diet under soaking method treated group, showed a higher mean weight gain (1.49 g) or 142.28% increase in growth over control.

To investigate the potential probiotic-ability of *B. subtilis* to combat the freshwater prawn diseases problem, *B. subtilis* bacterium was added to

prawn feed as a probiotic at a level of 10^{10} CFU/g feed by soaking method. Sixty days after the start of the *B. subtilis* feeding, the prawns were challenged by bath exposure to *A. hydrophila* (10^7 CFU/ml). After 28 days of post challenge, *B. subtilis* treated group had 88.33% survival, whereas the control group had only 20.83% survival, with an unhealthy external appearance.

After then a feeding experiment was conducted for juvenile *M. rosenbergii* (0.037 g) using three types of diets prepared with the incorporation of putative *B. subtilis*, non-putative *B. subtilis* and a commercial probiotic at level of 10^{10} (CFU/g) with an unsupplemented control. After 60 days the prawns fed putative *B. subtilis* added diets showed a higher survival, mean weight gain (0.663 ± 0.017 g) with 0.267 g increase in growth over control and best FCR value (2.33 ± 0.02 g).

Finally, the probiotic-ability of the *B. subtilis* was evident from properties consisting dominant isolated one, greatest antagonistic activity against pathogenic agents, highest RI, non-pathogenic to the host, growth enhancement, health improvement and more effectiveness than non-putative and commercial probiotics. This study demonstrated that among putative bacterial flora of juvenile *M. rosenbergii*, *B. subtilis* can be a promising probiotic for this prawn.

Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia sebagai memenuhi syarat untuk mendapatkan ijazah Doktor Falasafah.

PENILAIAN POTENSI BAKTERIA SEBAGAI PROBIOTIK DALAM UDANG GALAH JUVENIL, *Macrobrachium rosenbergii* (de Man)

Oleh

MEHRAN AVAKH KEYSAMI

Jun 2006

Pengerusi : Professor Madya Che Roos Saad, PhD

Fakulti : Pertanian

Satu kajian yang difokuskan kepada penggunaan probiotik untuk menyelesaikan masalah pemakanan dan penyakit dalam industri akuakultur telah dilakukan. Dalam kajian ini, sebanyak lapan eksperimen dilakukan untuk mengenalpasti flora bakteria daripada udang juvenil *M. rosenbergii* sebagai probiotik untuk meningkatkan pertumbuhan dan kemandirian udang untuk hidup, dari November 2002 hingga Ogos 2005 di Universiti Putra Malaysia.

Pengiraan bakteria dan flora bakteria putatif dalam ternakan udang juvenil *M. rosenbergii* (0.1 - 9.0 g) telah ditentukan dan jenis udang yang diasingkan kemudiannya dikelaskan kepada genera. Jumlah pengiraan sel hidup adalah sebanyak $96 \pm 2.6 \times 10^4$ CFU/ml bagi air kultur, $3100 \pm 169.0 \times 10^4$ CFU/ml bagi mendakan dalam tangki kultur, $2100 \pm 143.0 \times 10^4$ CFU/ml bagi permukaan badan udang, $14,000 \pm 4.6 \times 10^4$ CFU/g bagi salur pencernaan udang dan sebanyak $5.6 \pm 2.8 \times 10^4$ CFU/g bagi otot abdomen pada udang. Secara keseluruhannya, terdapat 12 genera telah dikesan daripada air, mendakan

dan organ-organ udang yang berbeza. Bakteria yang paling kerap diasingkan adalah daripada jenis *Bacillus*, *Aeromonas* dan *Pseudomonas*.

Sebanyak tiga kumpulan bakteria (*Bacillus subtilis*, *Micrococcus luteus* dan *Corynebacterium ammoniagenes*), telah menunjukkan aktiviti antagonis terhadap enam jenis patogen dalam ujian aktiviti antibakteria. *B. Subtilis* telah menunjukkan perencatan patogen yang tertinggi dalam kaedah telaga dan assei resapan cakera serta kaedah coretan lintang. Penambahan sebanyak 10 ml supernatan *B. subtilis* tanpa sel kepada patogen menyebabkan perencatan patogen dalam tempoh 12 jam. Pertumbuhan maksimum dan zon maksimum perencatan terhadap *A. hydrophila* diperhatikan pada 30 °C dan pada pH 8 dengan 1% NaCl, masing-masing.

Indeks kedudukan (RI) telah dikira sebagai saringan kepada probiont akuakultur yang berpotensi. RI dikira berdasarkan masa penggandaan (td) dan tempoh lanjut (λ) yang diperolehi daripada profil pertumbuhan bagi setiap bakterium. Tiga jenis bakteria (*B. subtilis*, *C. ammoniagenes* dan *M. luteus*) dan 2 jenis patogen (*A. hydrophila* dan *P. aeruginosa*) telah dikulturkan dalam agar TSB dan bacaan penyerapan (absorbans) direkodkan setiap setengah jam dalam tempoh masa 24 jam. Bakteria jenis *B. subtilis* membiak sama banyak dengan patogen-patogen dan mempunyai kadar pertumbuhan spesifik lebih tinggi ($\mu=0.034$) berbanding spesis bakteria yang lain iaitu *M. luteus* (0.031) dan *C. ammoniagenes* (0.032) masing-masing.

Tahap sensitiviti udang juvenil kepada dua jenis bakteria putatif (*B. subtilis* dan *A. hydrophila*) telah ditentukan menggunakan ujian cabaran patogenisiti secara *in vitro* dan *in vivo*. Ujian *in vitro* telah dijalankan ke atas udang juvenil (0.1 g) menggunakan kaedah rendaman. Sementara ujian *in vivo* pula telah dijalankan menggunakan kaedah percubaan pemakanan. Hasil kedua-dua ujian menunjukkan bahawa kadar kebolehan udang juvenil untuk hidup (0.1 dan 0.45g) tidak dipengaruhi oleh kehadiran *Bacillus* dalam air tawar tetapi, *A. hydrophila* menyebabkan jangkitan kepada udang juvenil *M. rosenbergii* dan menyebabkan sebanyak 78.67% kematian dan menunjukkan ciri-ciri klinikal penyakit-penyakit tertentu.

Satu percubaan pemakanan telah dijalankan untuk menentukan potensi bakteria *B. subtilis* sebagai probiotik dan tahap yang sesuai dalam udang juvenil *M. rosenbergii*. Bakteria *B. subtilis* putatif telah ditambahkan ke dalam makanan udang komersil sebagai probiotik. Sebanyak enam jenis diet telah disediakan dengan mencampurkan makanan udang kepada *Bacillus* dengan nisbah 1:1, 2:1, 3:1, 4:1 dan 5:1, dan satu kumpulan kawalan. Selepas 60 hari, udang yang diberi makan diet dengan nisbah 3:1, telah menunjukkan peningkatan berat yang lebih tinggi (593.45 g) atau peningkatan pertumbuhan sebanyak 170.29% berbanding kumpulan kawalan. Dengan ini, didapati probiotik yang dirawat dengan *B. subtilis* dapat meningkatkan kadar pertumbuhan dan kebolehan udang juvenil *M. rosenbergii* untuk hidup pada nisbah 3:1.

Satu lagi percubaan pemakanan juga telah dijalankan untuk mengenalpasti kaedah yang paling sesuai untuk pemberian *B. subtilis* dalam makanan komersil untuk pengkulturan udang juvenil *M. rosenbergii*. Empat jenis diet telah disediakan dengan mencampurkan makanan udang dan bakteria *Bacillus* dengan nisbah 3:1 dengan satu kumpulan kawalan (tanpa campuran bakteria). Kaedah-kaedah yang dilakukan ialah penggaulan, rendaman dan penyemburan. Selepas 60 hari, kumpulan udang yang diberi makan diet yang disediakan dengan kaedah rendaman telah menunjukkan min berat tertinggi (1.49 g) atau peningkatan pertumbuhan sebanyak 142.28% berbanding kumpulan kawalan.

Untuk mengenalpasti potensi probiotik *B. subtilis* bagi mengatasi penyakit udang air tawar, bakteria *B. subtilis* telah ditambahkan kepada makanan udang sebagai probiotik pada tahap 10^{10} CFU/g makanan udang menggunakan kaedah rendaman. Enam puluh hari selepas pemberian makanan rawatan, kumpulan udang tersebut telah didedahkan kepada *A. hydrophila* (10^7 CFU/ml) dalam rendaman air. Selepas 28 hari, kumpulan udang yang dirawat dengan probiotik mempunyai 88.33% kemandirian untuk hidup, sementara kumpulan kawalan hanya mempunyai 20.83% kemandirian untuk hidup, dengan rupa luaran yang tidak sihat (sempurna).

Kajian selanjutnya telah dijalankan ke atas kumpulan udang juvenil *M. rosenbergii* (0.037g) dengan memberi makan tiga jenis diet yang disediakan menggunakan *B. subtilis* putatif, *B. subtilis* bukan-putatif dan probiotik komersil pada tahap 10^{10} (CFU/g) berbanding kumpulan kawalan. Selepas

60 hari, udang yang diberi makan diet bercampur *B. subtilis* putatif menunjukkan min pertambahan berat udang yang lebih tinggi ($0.663 \pm 0.017\text{g}$) dan peningkatan pertumbuhan sebanyak 0.267 g berbanding kumpulan kawalan, di samping mempunyai nilai FCR yang terbaik (2.33 ± 0.02).

Sebagai kesimpulannya, potensi probiotik bagi bakteria *B. subtilis* telah dibuktikan berdasarkan bahan dominan yang diasingkan, kadar aktiviti antagonistik yang tinggi terhadap ejen patogenik, nilai RI yang tinggi, tidak mendatangkan penyakit kepada hos, peningkatan pertumbuhan dan kesihatan serta lebih efektif berbanding bakteria bukan putatif dan probiotik komersil. Kajian ini menunjukkan bahawa bakteria *B. subtilis* berpotensi sebagai probiotik dikalangan flora-flora bakteria juvenil *M. rosenbergii* yang lain.

ACKNOWLEDGEMENTS

First and foremost, I want to thank Associate Professor Dr. Che Roos Saad, chairman of my advisory committee, for providing me with a wonderful opportunity to complete my doctoral studies under his exceptional guidance. Other than the provision of the necessary funding, this work would not have been possible without his patience, constant encouragement, guidance and knowledge. Through frequent meetings and his open door policy, Dr. Che Roos made an immense contribution to this dissertation and my academic growth, as well as my professional and personal life.

My sincerest appreciation is also extended to Associate Professor Dr. Kamaruzaman Sijam who is a member of my supervisory committee and provided me some laboratory facilities and cooperation during the course of the study. I would like also to extend my most sincere gratitude and thanks to Associate Professor Dr. Hassan Mohd Daud and Professor Dr. Razak Alimon who are also members of my supervisory committee, for their constructive suggestions and guidance during the study period. I am also grateful for their willingness to serve on my committee, provide me assistance whenever required, involvement in my qualifying examination, and for reviewing this dissertation.

I would also like to extend my appreciation and thanks to:

- The Government of Malaysia for providing the research grant for this program.

- My laboratory mates in Microbiology Laboratory, Mr. Antario Dikin, and Eshetu Derso for their technical assistance during the conduct of this study;
- The staff of Aquatic Resources Technology and Microbiology laboratory, Faculty of Agriculture, UPM especially Mr. Jasni and Mr. Khir for their valuable assistance;
- My family for their love, understanding, encouragement and moral support towards this achievement;
- Above all, to GOD almighty for making this study possible.

I certify that an Examination Committee has met on 12th June 2006 to conduct the final examination of Mehran Avakh Keysami on his Doctor of Philosophy thesis entitled "Assessment of Putative Bacteria as Probiotics in Juvenile Freshwater Prawn, *Macrobrachium rosenbergii* (de Man)" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Mihdzar Bin Abdul Kadir, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Mustafa Kamal Abdul Satar, PhD


Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Annie Christianus, PhD

Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Faizah Mohd Shaharom, PhD

Professor
Faculty of Science and Technology
Kolej Universiti Sains dan Teknologi Malaysia
(External Examiner)



HASANAH MOHD GHAZALI, PhD
Professor / Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : 11 JUL 2006

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Che Roos Saad, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Hassan Mohd Daud, PhD


Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Kamaruzaman Sijam, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Abdul Razak Alimon, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)



AINI IDERIS, PhD
Professor / Dean
School of Graduate Studies
Universiti Putra Malaysia

Date 10 AUG 2006



DECLARATION

I hereby declare that this thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

MEHRAN AVAKH KEYSAMI

Date: 10/7/06



Keysami

TABLE OF CONTENTS

		Page
DEDICATION		ii
ABSTRACT		iii
ABSTRAK		vii
ACKNOWLEDGEMENTS		xii
APPROVAL		xiv
DECLARATION		xvi
LIST OF TABLES		xxii
LIST OF FIGURES		xxvii
LIST OF ABBREVIATIONS		xxx
CHAPTER		
1	INTRODUCTION	1
1.1	Background of the Study	1
1.2	Statement of Problem	9
1.3	Significance of the Study	12
1.4	Objectives	14
2	LITERATURE REVIEW	16
2.1	Biology of Fresh Water Prawn	16
2.1.1	Taxonomy and Life-history of <i>M. rosenbergii</i>	16
2.1.2	Growth	17
2.2	Hatchery	19
2.2.1	Broodstock, Incubation and Hatching	19
2.2.2	Larvae Culture	22
2.2.3	Feeds and Feeding	23
2.2.4	Separation of Larvae and Post larvae	24
2.3	Nursery	25
2.3.1	Post larvae Culture	25
2.3.2	Size-grading of Nursery Populations	28
2.4	Grow out	28
2.4.1	Extensive	29
2.4.2	Semi-intensive	29
2.4.3	Intensive	34
2.4.4	Polyculture	35
2.5	Diseases	36
2.6	Harvesting	37
2.7	Probiotics	37
2.7.1	Definition of Probiotics	37
2.7.2	Selection of Probiotics Bacteria	39
2.7.3	Possible Mode of Action	40
2.7.4	Production of Inhibitory Compounds	41
2.7.5	Competition for Chemicals	44
2.7.6	Competition for Adhesion Sites	48
2.7.7	Enhancement of the Immune Response	50

2.7.8	Improvement of Water Quality	51
2.7.9	Interaction with Phytoplankton	52
3	GENERAL METHODOLOGY	54
3.1	Location of the Study	54
3.2	Experimental Prawns	55
3.3	<i>Artemia</i> Preparation	56
3.4	Experimental Set-up	56
3.5	Identification of Bacterial Species Using Biolog Kit	57
3.6	Bacterial Enumeration	58
3.7	Growth Measurement	58
3.8	Proximate Analyses of Experimental Diets and Prawns	59
3.8.1	Crude Protein	59
3.8.2	Lipids	60
3.8.3	Crude Fibre	61
3.8.4	Moisture Determination	62
3.8.5	Ash Determination	62
3.9	Water Quality Management	63
3.10	Statistical Analysis	63
4	ISOLATION AND IDENTIFICATION OF PUTATIVE BACTERIAL FLORA IN JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> (DE MAN)	64
4.1	Introduction	64
4.2	Materials and Methods	66
4.2.1	Sample Collection	66
4.2.2	Isolation and Identification of Putative Bacteria	69
4.2.3	Statistical Analysis	69
4.3	Results	70
4.4	Discussion	76
5	ASSESSMENT OF PUTATIVE BACTERIAL FLORA IN JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> AS PROBIOTICS USING ANTIBACTERIAL ACTIVITY TEST	81
5.1	Introduction	81
5.2	Materials and Methods	83
5.2.1	Study of Inhibition by Well-diffusion, Disc-diffusion and Cross-streak Methods	84
5.2.2	Effect of Cell free Supernatant of Candidates on Washed cells of Pathogens in Sterile Normal Saline	85
5.2.3	Influence of Growth Conditions on Production of Inhibitor	86
5.3	Results	86
5.4	Discussion	96

6	COMPARISON OF PROBIOTIC-ABILITY FOR THREE PUTATIVE BACTERIA IN JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> BASED ON IN VITRO BACTERIA GROWTH CHARACTERISTICS	98
	6.1 Introduction	98
	6.2 Materials and Methods	99
	6.2.1 Bacterial Strains	99
	6.2.2 Determination of Bacterial Population Density	100
	6.2.3 Estimation of Bacterial Growth Parameters	101
	6.3 Results	105
	6.4 Discussion	112
7	USE OF PATHOGENICITY CHALLENGE TEST FOR ASSESSMENT OF PUTATIVE <i>BACILLUS SUBTILIS</i> AS PROBIOTIC IN JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> (<i>DE MAN</i>)	115
	7.1 Introduction	115
	7.2 Materials and Methods	116
	7.2.1 Preparation of Putative Bacterial Strains	116
	7.2.2 <i>In vitro</i> Testing Pathogenicity of <i>Bacillus subtilis</i> in Juvenile Prawns	116
	7.2.3 <i>In vivo</i> Testing Pathogenicity of <i>Bacillus subtilis</i> in Juvenile Prawns	117
	7.2.4 Water Quality Parameters	118
	7.2.5 Bacteriological Study	118
	7.2.6 Statistical Analysis	119
	7.3 Results	119
	7.4 Discussion	127
8	RESPONSE OF JUVENILE <i>MACROBRACHIUM ROSENBERGII</i>, TO DIFFERENT LEVELS OF <i>BACILLUS SUBTILIS</i> AS PROBIOTICS	132
	8.1 Introduction	132
	8.2 Materials and Methods	133
	8.2.1 Preperation of Experimental Prawns and Design	133
	8.2.2 Diet Preparation	135
	8.2.3 Growth Analysis	136
	8.2.4 Proximate Analysis	137
	8.2.5 Water Quality Parameters	138
	8.2.6 Bacteriological Study	138
	8.2.7 Statistical Analysis	139
	8.3 Results	139
	8.4 Discussion	149

9	SUITABLE METHODS FOR THE INCORPORATION OF <i>BACILLUS SUBTILIS</i> IN TO THE FEED TO ENHANCE GROWTH AND SURVIVAL OF JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> (DE MAN)	152
9.1	Introduction	152
9.2	Materials and methods	153
9.2.1	Preparation of Experimental Prawns and Design	153
9.2.2	Diet Preparation	154
9.2.3	Growth Analysis	157
9.2.4	Proximate Analysis	158
9.2.5	Water Quality Parameters	159
9.2.6	Bacteriological Study	159
9.2.7	Statistical Analysis	159
9.3	Results	160
9.4	Discussion	172
10	PROTECTION OF JUVENILE <i>MACROBRACHIUM ROSENBERGII</i> FROM <i>AEROMONAS HYDROPHILA</i> INFECTION BY <i>BACILLUS SUBTILIS</i>	176
10.1	Introduction	176
10.2	Materials and Methods	177
10.2.1	Preparation of Experimental Prawns and Design	177
10.2.2	Diet Preparation	179
10.2.3	Pathogen Challenge Test	179
10.2.4	Growth Analysis	180
10.2.5	Water Quality Parameters	181
10.2.6	Bacteriological Study	181
10.2.7	Statistical Analysis	181
10.3	Results	182
10.4	Discussion	191
11	EFFECTIVENESS OF PUTATIVE, NON-PUTATIVE AND COMMERCIAL PROBIOTIC BACTERIA ON THE GROWTH AND SURVIVAL OF JUVENILE <i>MACROBRACHIUM ROSENBERGII</i>	193
11.1	Introduction	193
11.2	Materials and Methods	193
11.2.1	Preparation of Experimental Prawns and Design	193
11.2.2	Diet Preparation	195
11.2.3	Growth Analysis	197
11.2.4	Proximate Analysis	197
11.2.5	Water Quality Parameters	198
11.2.6	Bacteriological Study	198
11.3	Results	199
11.4	Discussion	211

12	GENERAL DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS	
12.1	General Discussion and Conclusions	214
12.2	Recommendations	220
	REFERENCES	223
	APPENDICES	250
	BIODATA OF AUTHOR	271

LIST OF TABLES

Table		Page
1.1	Top 15 countries farming freshwater prawn, <i>Macrobrachium rosenbergii</i> in 2001 (FAO, 2003).	4
2.1	Weight-dependent feeding rates for semi-intensive pond grow -out of <i>Macrobrachium rosenbergii</i> .	32
4.1	The physico-chemical characteristics of prawn culture water and weight of juvenile at the time of sampling (mean± SD; n=3).	70
4.2	Bacterial count on different selective media (mean±SD; n=3) observed during <i>M. rosenbergii</i> culture (CFU×10 ⁴ /ml for prawn culture water ,sediment and prawn body surface; CFU×10 ⁴ /g for digestive tract and muscle).	71
4.3	Main groups of isolates categorized based on similar results of biochemical tests and Biolog.	73
4.4	Biochemical characteristics of putative bacterial flora in juvenile <i>M. rosenbergii</i> .	74
4.5	Relative frequency of bacteria isolated from culture water, sediment, prawn body surface, digestive tract and body muscle.	76
5.1	Antagonism between candidate probionts and pathogenic bacteria grown on TSA.	89
5.2	Effect of cell free supernatant of <i>B. subtilis</i> , <i>C. ammoniagenes</i> and <i>M. luteus</i> on washed cells of <i>A. hydrophila</i> .	91
5.3	Effect of pH on the growth of <i>B. subtilis</i> and production of antibacterial component.	93
5.4	Effect of temperature and salt concentration on the growth of <i>B. subtilis</i> and production of antibacterial component.	93
5.5	Effect of duration of incubation on the production of antibacterial component.	94

6.1	The absorbance (600nm) and cell density of three candidate probionts and two pathogens.	106
6.2	Measured absorbance of three candidate probionts and two pathogens grown in TSB during 14 hours (Mean + SD).	109
6.3	Log measured cell density (CFU/ml) of three candidate probionts and two pathogens grown in TSB during 14 hours.	110
6.4	Growth parameters of three candidate probionts and two pathogens grown in TSB media.	112
7.1	<i>In vitro</i> effectiveness of <i>B. subtilis</i> and <i>A. hydrophila</i> culture on mortality of juvenile Prawns during 96 h.	120
7.2	<i>In vivo</i> effectiveness of <i>B. subtilis</i> and <i>A. hydrophila</i> culture on mortality of juvenile prawns during 28 days of pathogenicity challenge test (Mean Mean \pm SD).	122
7.3	Water quality of control and treated tanks during 28 days of pathogenicity challenge test (Mean \pm SD).	124
7.4	Log total bacteria, <i>Bacillus</i> and gram-negative bacteria count (CFU/ml) in prawns gut during 28 days of <i>A. hydrophila</i> challenge test.	126
7.5	Log total bacteria, <i>Bacillus</i> and gram-negative bacteria count (CFU/ml) in prawn rearing tank water during 28 days of <i>A. hydrophila</i> challenge test.	126
8.1	Experimental design to determine the level of <i>B. subtilis</i> for incorporation into the prawn feed.	135
8.2	Water quality of control and treated tanks during feeding trial (Mean \pm SD).	140
8.3	Biochemical composition of diets (DM% basis, Mean \pm SD).	141
8.4	Biochemical composition of whole body tissues (DM % basis, Mean \pm SD).	141

8.5	Growth-related performance of <i>M. rosenbergii</i> by using diets incorporated with <i>B. subtilis</i> at different levels (Mean \pm SD).	144
8.6	Log total viable bacteria, <i>Bacillus</i> and gram-negative bacteria count (CFU/g) in prawn guts (hepatopancreas and intestine) during 60 days of feeding diets incorporated with <i>B. subtilis</i> at different levels.	147
8.7	Log total viable bacteria, <i>Bacillus</i> and gram-negative bacteria count (CFU/g) in prawn feces during 60 days of feeding diets incorporated with <i>B. subtilis</i> at different levels.	147
9.1	Experimental design to determine method of <i>B. subtilis</i> administration into the prawn feed.	154
9.2	Water quality of control and treated tanks during feeding diets prepared using different methods of <i>B. subtilis</i> administration into the prawn feed (Mean \pm SD).	161
9.3	Biochemical composition of diets prepared using different methods of <i>B. subtilis</i> administration into the prawn feed (DM % basis, Mean \pm SD).	163
9.4	Biochemical composition of whole body tissues of prawns fed diets prepared using different methods of <i>B. subtilis</i> administration into the prawn feed (DM % basis, Mean \pm SD).	164
9.5	Growth-related performance of <i>M. rosenbergii</i> by using diets prepared using different methods of <i>B. subtilis</i> administration into the prawn feed (Mean \pm SD).	167
9.6	Log total viable cells of bacteria, <i>Bacillus</i> and gram-negative bacteria count (CFU/g) in prawn guts (hepatopancreas and intestine) during 60 days of feeding diets prepared using different methods of <i>B. subtilis</i> administration into the prawn feed.	170