



UNIVERSITI PUTRA MALAYSIA

**ANTIMICROBIAL AND WOUND HEALING ACTIVITIES OF LOCAL
EUPHORBIA HIRTA , ACALYPHA INDICA AND .PHYLLANTHUS
NIRURI**

REEZAL BIN ISHAK

FPSK(M) 2004 1

**ANTIMICROBIAL AND WOUND HEALING ACTIVITIES OF LOCAL
EUPHORBIA HIRTA, *ACALYPHA INDICA* AND *PHYLLANTHUS NIRURI***

By

REEZAL BIN ISHAK

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Master of Science**

January 2004



DEDICATION

**“This research project is dedicated to those who have endure a great deal of pain
and joy for just being around me”**

To my parents – Thank you for believing in me

To my brothers – Thanks for keeping me tough

To Elysha Nur Ismail – Thank you for your patience

“Your sacrifices and supports gave strength to me”

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**ANTIMICROBIAL AND WOUND HEALING ACTIVITIES OF LOCAL
EUPHORBIA HIRTA, *ACALYPHA INDICA* AND *PHYLLANTHUS NIRURI***

By

REEZAL BIN ISHAK

January 2004

Chairman: Associate Professor Muhammad Nazrul Hakim Abdullah, Ph.D.

Faculty: Medicine and Health Sciences

Herbal plants are well known for their capabilities in treating diseases for centuries. The World Health Organisation (WHO) has estimated that 80% of the world's population use botanical medicines for their primary health care needs. Some of the medicinal plants under the *Euphorbiaceae* family are *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri*. These plants are known as traditional remedy in many rural areas throughout the world and have been used for various purposes. This study focuses on the antimicrobial activity of aqueous and ethanol extracts of these plants against selected bacteria and fungi, and its wound healing properties on surgically incised mice. The antimicrobial test using the disc diffusion method at 20, 40 and 80 mg/ml extract-impregnated disc concentrations showed significant results ($P < 0.05$) of *Euphorbia hirta* aqueous leaves and barks extracts against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, and *Salmonella enteritidis* with inhibition zones ranged from 8.33 mm to 19.67 mm. The inhibition zones for *Acalypha indica* and *Phyllanthus niruri* aqueous leaves extracts were measured between 8.33 mm to 11.0



mm and 7.33 mm to 12.33 mm respectively, but no inhibition zones were seen with the bark extracts of both plants. Only the ethanol extracts of the leaves showed antibacterial activity. The results for *Euphorbia hirta* was the most significant ($P < 0.001$) with inhibition zones ranging from 7.0 mm to 13.33 mm. Ethanol leaves extracts of *Acalypha indica* and *Phyllanthus niruri* had inhibition zones between 7.0 mm to 9.67 mm and 7.0 mm to 9.67 mm respectively. The results obtained showed comparable effects to commercial antibiotics of chloramphenicol, ampicilin, penicillin G, erythromycin, tetracycline and enrofloxacin as some of the plant extracts exhibit almost equal and even bigger inhibition zones. Antifungal activity were detected only with the *Euphorbia hirta* aqueous leaves extracts against *Candida tropicalis* with mean inhibition zones of 8.67 mm at 80 mg/ml discs concentrations. The aqueous extracts of the leaves were then tested for its wound healing properties on surgically incised mice. The extracts were applied topically over the incision and inflammatory and cicatrization activities evaluated according to a designated scoring table. *Acalypha indica* showed the most significant results ($P < 0.05$) with better reduction of inflammation and increased cicatrization activities on the wound compared to *Euphorbia hirta* and *Phyllanthus niruri* treated mice. In comparison to the acriflavine treated mice, the results demonstrated an equally good epithelial regeneration of the skins and showed to be better than the control group (no treatment).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**AKTIVITI ANTIMIKROB DAN PENYEMBUHAN LUKA OLEH
EUPHORBIA HIRTA, *ACALYPHA INDICA* DAN *PHYLLANTHUS NIRURI*
TEMPATAN**

Oleh

REEZAL BIN ISHAK

Januari 2004

Pengerusi: Profesor Madya Muhammad Nazrul Hakim Abdullah, Ph.D.

Fakulti: Perubatan dan Sains Kesihatan

Tumbuhan-tumbuhan herba sangat dikenali kerana kebolehannya untuk merawat pelbagai penyakit sejak dahulu lagi. Pertubuhan Kesihatan Sedunia (WHO) menganggarkan sebanyak 80% populasi dunia menggunakan ubatan-ubatan herba untuk keperluan kesihatan. Sebahagian daripada tumbuhan perubatan di bawah famili *Euphorbiaceae* adalah *Euphorbia hirta*, *Acalypha indica* dan *Phyllanthus niruri*. Pokok-pokok ini telah diketahui sebagai ubatan tradisional di kawasan-kawasan kampung di merata dunia dan digunakan untuk pelbagai tujuan. Penyelidikan ini memfokuskan kepada aktiviti antimikrob oleh ekstrak akues dan etanol pokok-pokok ini terhadap bakteria dan kulat yang dipilih, serta kebolehannya menyembuhkan luka pembedahan pada mencit. Ujian antimikrob menggunakan kaedah penyebaran cakera pada konsentrasi ekstrak 20, 40 dan 80 mg/ml menunjukkan keputusan yang signifikan ($P < 0.05$) oleh ekstrak akues daripada daun dan batang *Euphorbia hirta* terhadap *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, dan *Salmonella enteritidis* dengan zon perencatan berjarak dari 8.33 mm ke 19.67 mm. Ukuran zon



perencatan oleh ekstrak daun akues dari *Acalypha indica* dan *Phyllanthus niruri*, masing-masing adalah di antara 8.33 mm ke 11.0 mm dan 7.33 mm ke 12.33 mm, tetapi tiada zon perencatan dapat dilihat dengan ekstrak batang dari kedua-dua pokok. Hanya ekstrak daun ethanol yang menunjukkan aktiviti antimikrob. Ekstrak bagi *Euphorbia hirta* menunjukkan keputusan yang paling signifikan ($P < 0.001$) dengan zon perencatan berjarak dari 7.0 mm ke 13.33 mm. Ekstrak ethanol dari *Acalypha indica* dan *Phyllanthus niruri* dilihat dengan zon perencatan di antara 7.0 mm ke 9.67 mm and 7.0 mm ke 9.67 mm. Keputusan yang diperolehi ini menunjukkan kesan-kesan yang setanding dengan ubatan antibiotik komersial seperti chloramphenicol, ampicilin, penicillin G, erythromycin, tetracycline dan enrofloxacin menunjukkan kesan yang setanding, di mana sesetengah ekstrak pokok-pokok ini mempamerkan zon perencatan yang hampir sama dan juga yang lebih besar. Aktiviti antikulat pula hanya dilihat dengan ekstrak akues dari daun *Euphorbia hirta* terhadap *Candida tropicalis* dengan purata zon perencatan sebanyak 8.67 mm pada 80 mg/ml konsentrasi ekstrak. Ekstrak akues dari daun pokok-pokok ini kemudiannya diuji untuk keupayaannya menyembuhkan luka pembedahan yang telah dilakukan pada mencit. Ekstrak-ekstrak ini telah diberikan secara aplikasi luaran pada luka tersebut dan aktiviti inflamasi dan pembentukan semula kulitnya dinilai mengikut carta penilaian yang telah direka. Ekstrak *Acalypha indica* menunjukkan keputusan yang paling signifikan ($P < 0.05$) dengan pengurangan inflamasi dan peningkatan pembentukan semula kulit pada luka berbanding dengan rawatan oleh *Euphorbia hirta* and *Phyllanthus niruri* pada mencit. Perbandingan keputusan dengan rawatan acriflavine mempamerkan pertumbuhan semula epitelial yang setanding dan lebih baik daripada kumpulan kawalan (tiada rawatan).

ACKNOWLEDGEMENTS

In the name of Allah, Most gracious, Most merciful

All gratifications are referred to Allah

I would like to take this opportunity to express my highest gratitude and thanks to my supervisor, Associate Professor Dr. Muhd Nazrul Hakim Abdullah, for his time, support, advises, encouragement and constant guidance throughout the completion of my study. And most of all, for giving me chance to improve myself to be a better person in real life.

I also like to express my appreciation to my supervisory committee for advises, help and suggestion during the period of my study. My sincere thanks go to Associate Professor Dr. Abdul Rahim Mutalib for his ideas, help and providing his Bacteriology Laboratory. I am also grateful to Dr. Wan Nordin Wan Mahmud for his advises and permission to use the Histology Laboratory for my research project. A word of thanks and appreciation to Prof. Dato' Dr. Abd. Salam Abdullah and Associate Prof. Dr. Abd. Manan Mat Jais for their interest and advises during the progress of my research.

I wish to express my gratitude to the staff of Bacteriology Laboratory, En. Hajaraih and En. Jefri for their helping hands; to Puan Sapiah at Histology Laboratory for her guidance; and also to the staff at Physiology Laboratory, especially En. Kufli for sharing his experience.



I wish to express my thanks to my friends, Azmahani Abdullah, Nor Shahida Abdul Rahman, and Solihah Mohd Hassan for their help and co-operation in making this project a success. To Elysha Nur Ismail, I would like to give my millions of thanks for your patience, help and continuous support throughout this course.

Finally, to my beloved family, my parents, Ishak b. Mohd. Ali and Tialan bt. Kamari, for their prayers, loves, support, understanding and patience for the last 25 years. To my elder brother Dr. Azlan Iskandar b. Ishak, who inspire me to improve myself from day to day; and also to my youngest brother, Azree Ishak, who makes me push forward to improve myself day by day.



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVALS	ix
DECLARATION	xi
LIST OF FIGURES	xv
LIST OF TABLES	xvii
LIST OF ABBREVIATIONS	xviii

CHAPTER

1	INTRODUCTION	1
	The Importance of Herbal Plants	1
	<i>Euphorbiaceae</i>	2
	<i>Euphorbia hirta</i>	2
	<i>Acalypha indica</i>	3
	<i>Phyllanthus niruri</i>	3
	Justification of the Study	4
	Objectives of the Study	4
2	LITERATURE REVIEW	5
	Herbal Medication	5
	<i>Euphorbia hirta</i>	6
	Traditional Uses	6
	Previous Studies	8
	Chemical Constituents	12
	<i>Acalypha indica</i>	14
	Traditional Uses	14
	Previous Studies	16
	Chemical Constituents	18
	<i>Phyllanthus niruri</i>	20
	Traditional Uses	20
	Previous Studies	22
	Chemical Constituents	25
	Bacteria	29
	Bacterial Infections	30
	Gram Positive Bacteria	32
	<i>Staphylococcus aureus</i>	32
	<i>Bacillus subtilis</i>	32
	Gram Negative Bacteria	33
	<i>Escherichia coli</i>	33
	<i>Salmonella enteritidis</i>	33
	Fungi	34
	Fungal Infections	35
	Moulds	36
	<i>Microsporium canis</i>	36



	<i>Aspergillus fumigatus</i>	36
Yeast		37
	<i>Candida albicans</i>	37
	<i>Candida tropicalis</i>	37
Skin Wound Healing		38
	Repair of Skin Wounds and Healing Process	38
	Previous studies	39
3	<i>IN VITRO</i> ANTIBACTERIAL ACTIVITY OF <i>EUPHORBIA HIRTA, ACALYPHA INDICA</i> AND <i>PHYLLANTHUS NIRURI</i>	42
	Introduction	42
	Methodology	43
	Preparation of Plant Extracts	43
	Extract-Impregnated Discs Preparation	44
	Media Preparation for Bacteria Culture	45
	Bacteria Preparation for the Sensitivity Test	45
	Biochemical Test	46
	Disc Diffusion Method (Kirby-Bauer Method) – Sensitivity Test	46
	Results	47
	Comparison on the Aqueous Extracts of <i>Euphorbia hirta, Acalypha indica</i> and <i>Phyllanthus</i> <i>niruri</i> Antibacterial Activity	47
	Comparisons on the Ethanol Extracts of <i>Euphorbia</i> <i>hirta, Acalypha indica</i> and <i>Phyllanthus niruri</i> Antibacterial Activity	68
	Discussions	81
4	<i>IN VITRO</i> ANTIFUNGAL ACTIVITY OF <i>EUPHORBIA HIRTA, ACALYPHA INDICA</i> AND <i>PHYLLANTHUS NIRURI</i>	90
	Introduction	90
	Methodology	91
	Preparation of Plant Extracts	91
	Extract-Impregnated Discs Preparation	91
	Standard Antifungal Discs Preparation	92
	Media Preparation for Fungi Culture	92
	Fungi Preparation for the Sensitivity Test	93
	Fungi Identification	93
	Disc Diffusion Method (Kirby-Bauer Method) – Sensitivity Test	94
	Results	95
	Morphological Observation of the Selected Fungi	95
	Comparisons on the Aqueous and Ethanol Extracts of <i>Euphorbia hirta, Acalypha indica</i> and <i>Phyllanthus niruri</i> Antifungal Activity	98
	Discussions	104

5	WOUND HEALING ACTIVITY OF <i>EUPHORBIA</i> <i>HIRTA</i>, <i>ACALYPHA INDICA</i> AND <i>PHYLLANTHUS</i> <i>NIRURI</i> IN MICE	106
	Introduction	106
	Methodology	107
	Preparation of the Plant Extracts	107
	Experimental Animals	107
	Preparation of the Skin Samples for Histological Study	109
	Evaluation of the Wound Healing Activity	110
	Results	112
	Observation of the Wound Healing Process at 3, 6 and 9 Days of Experiment	112
	Discussions	145
6	GENERAL DISCUSSIONS AND CONCLUSIONS	152
	REFERENCES	155
	APPENDICES	162
	BIODATA OF THE AUTHOR	169



LIST OF FIGURES

Figure		Page
1	<i>Euphorbia hirta</i> .	13
2	<i>Acalypha indica</i> .	19
3	<i>Phyllanthus niruri</i> .	28
4	Sensitivity test of <i>E. hirta</i> aqueous leaves and barks extracts against <i>E. coli</i> .	52
5	Sensitivity test of <i>E. hirta</i> aqueous leaves and barks extracts against <i>S. enteritidis</i> .	53
6	Sensitivity test of <i>E. hirta</i> aqueous leaves and barks extracts against <i>S. aureus</i> .	54
7	Sensitivity test of <i>E. hirta</i> aqueous leaves and barks extracts against <i>B. subtilis</i> .	55
8	Sensitivity test of <i>A. indica</i> aqueous leaves extracts against <i>E. coli</i> and <i>S. enteritidis</i> .	56
9	Sensitivity test of <i>A. indica</i> aqueous leaves and barks extracts against <i>S. aureus</i> .	57
10	Sensitivity test of <i>A. indica</i> aqueous leaves extracts against <i>B. subtilis</i> .	58
11	Sensitivity test of <i>P. niruri</i> aqueous leaves extracts against <i>S. aureus</i> and <i>B. subtilis</i> .	59
12	Mean inhibition zones of <i>E. hirta</i> aqueous leaves extracts antibacterial activities at different concentrations.	61
13	Mean inhibition zones of <i>E. hirta</i> aqueous barks extracts antibacterial activities at different concentrations.	62
14	Mean inhibition zones of <i>A. indica</i> aqueous leaves extracts antibacterial activities at different concentrations.	64
15	Mean inhibition zones of <i>A. indica</i> aqueous barks extracts antibacterial activities at different concentrations.	65
16	Mean inhibition zones of <i>P. niruri</i> aqueous leaves extracts antibacterial activities at different concentrations.	67
17	Sensitivity test of <i>E. hirta</i> ethanol leaves extracts against <i>E. coli</i> and <i>B. subtilis</i> .	71

Figure		Page
18	Sensitivity test of <i>E. hirta</i> ethanol leaves extracts against <i>S. aureus</i> and <i>B. subtilis</i> .	72
19	Sensitivity test of <i>A. indica</i> ethanol leaves extracts against <i>S. aureus</i> and <i>B. subtilis</i> .	73
20	Sensitivity test of <i>P. niruri</i> ethanol leaves extracts against <i>S. aureus</i> and <i>B. subtilis</i> .	74
21	Mean inhibition zones of <i>E. hirta</i> ethanol leaves extracts antibacterial activities at different concentrations.	76
22	Mean inhibition zones of <i>A. indica</i> ethanol leaves extracts antibacterial activities at different concentrations.	78
23	Mean inhibition zones of <i>P. niruri</i> ethanol leaves extracts antibacterial activities at different concentrations.	80
24	<i>Microsporium canis</i> .	96
25	<i>Aspergillus fumigatus</i> .	96
26	<i>Candida albicans</i> .	97
27	<i>Candida tropicalis</i> .	97
28	Sensitivity test of <i>E. hirta</i> aqueous leaves extracts against <i>Candida tropicalis</i> .	99
29	Mean inhibition zones of <i>E. hirta</i> aqueous leaves extracts antifungal activities at different concentrations.	101
30	Wound healing process after day 3 of control group (40x & 100x magnification).	115
31	Wound healing process after day 3 of control group (200x & 400x magnification).	116
32	Wound healing process after day 6 of control group (40x & 100x magnification).	117
33	Wound healing process after day 6 of control group (200x & 400x magnification).	118
34	Wound healing process after day 9 of control group (40x & 100x magnification).	119
35	Wound healing process after day 9 of control group (200x & 400x magnification).	120

Figure		Page
36	Wound healing process after day 3 of acriflavine treatment (40x & 100x magnification).	121
37	Wound healing process after day 3 of acriflavine treatment (200x & 400x magnification).	122
38	Wound healing process after day 6 of acriflavine treatment (40x & 100x magnification).	123
39	Wound healing process after day 6 of acriflavine treatment (200x & 400x magnification).	124
40	Wound healing process after day 9 of acriflavine treatment (40x & 100x magnification).	125
41	Wound healing process after day 9 of acriflavine treatment (200x & 400x magnification).	126
42	Wound healing process after day 3 of <i>E. hirta</i> aqueous leaves extracts treatment (40x & 100x magnification).	127
43	Wound healing process after day 3 of <i>E. hirta</i> aqueous leaves extracts treatment (200x & 400x magnification).	128
44	Wound healing process after day 6 of <i>E. hirta</i> aqueous leaves extracts treatment (40x & 100x magnification).	129
45	Wound healing process after day 6 of <i>E. hirta</i> aqueous leaves extracts treatment (200x & 400x magnification).	130
46	Wound healing process after day 9 of <i>E. hirta</i> aqueous leaves extracts treatment (40x & 100x magnification).	131
47	Wound healing process after day 9 of <i>E. hirta</i> aqueous leaves extracts treatment (200x & 400x magnification).	132
48	Wound healing process after day 3 of <i>A. indica</i> aqueous leaves extracts treatment (40x & 100x magnification).	133
49	Wound healing process after day 3 of <i>A. indica</i> aqueous leaves extracts treatment (200x & 400x magnification).	134
50	Wound healing process after day 6 of <i>A. indica</i> aqueous leaves extracts treatment (40x & 100x magnification).	135
51	Wound healing process after day 6 of <i>A. indica</i> aqueous leaves extracts treatment (200x & 400x magnification).	136

Figure		Page
52	Wound healing process after day 9 of <i>A. indica</i> aqueous leaves extracts treatment (40x & 100x magnification).	137
53	Wound healing process after day 9 of <i>A. indica</i> aqueous leaves extracts treatment (200x & 400x magnification).	138
54	Wound healing process after day 3 of <i>P. niruri</i> aqueous leaves extracts treatment (40x & 100x magnification).	139
55	Wound healing process after day 3 of <i>P. niruri</i> aqueous leaves extracts treatment (200x & 400x magnification).	140
56	Wound healing process after day 6 of <i>P. niruri</i> aqueous leaves extracts treatment (40x & 100x magnification).	141
57	Wound healing process after day 6 of <i>P. niruri</i> aqueous leaves extracts treatment (200x & 400x magnification).	142
58	Wound healing process after day 9 of <i>P. niruri</i> aqueous leaves extracts treatment (40x & 100x magnification).	143
59	Wound healing process after day 9 of <i>P. niruri</i> aqueous leaves extracts treatment (200x & 400x magnification).	144
60	Mean value on inflammation effects based on the attributed score.	150
61	Mean value on cicatrisation effects based on the attributed score.	151

LIST OF TABLES

Table		Page
1	Mean inhibition zones \pm S.D of <i>Euphorbia hirta</i> aqueous extracts antibacterial activities.	60
2	Mean inhibition zones \pm S.D of <i>Acalypha indica</i> aqueous extracts antibacterial activities.	63
3	Mean inhibition zones \pm S.D of <i>Phyllanthus niruri</i> aqueous extracts antibacterial activities.	66
4	Mean inhibition zones \pm S.D of <i>Euphorbia hirta</i> ethanol extracts antibacterial activities.	75
5	Mean inhibition zones \pm S.D of <i>Acalypha indica</i> ethanol extracts antibacterial activities.	77
6	Mean inhibition zones \pm S.D of <i>Phyllanthus niruri</i> ethanol extracts antibacterial activities.	79
7	Mean inhibition zones \pm S.D of <i>Euphorbia hirta</i> aqueous extracts antifungal activities.	100
8	Mean inhibition zones \pm S.D of <i>Acalypha indica</i> aqueous extracts antifungal activities.	102
9	Mean inhibition zones \pm S.D of <i>Phyllanthus niruri</i> aqueous extracts antifungal activities.	103
10	Attributed score to value anti-inflammatory and cicatrisation activities.	111

LIST OF ABBREVIATIONS

AIDS	Acquired Immunodeficiency Syndrome
ATCC	American Types Cell Culture
CNS	Central nervous system
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
IBS	Institute of Bioscience
MARDI	Malaysian Agricultural and Research Development Institute
μl	Micro-litre
MIC	Minimum inhibitory concentration
mg	Milligram
ml	Millilitre
mm	Millimetre
MR/VP	Methyl Red/ Voges-Proskauer
s.d	Standard deviation
UPM	Universiti Putra Malaysia

CHAPTER 1

INTRODUCTION

The Importance of Herbal Plants

Herbal plants have been known for its capabilities in healing or treating diseases for centuries. The beginning of the nineteenth century heralded an era, in which the active secondary metabolite of medicinal plants began to be purified, with which such pure constituents were then introduced into therapy. Treatments using herbal medicines are generally safer due to its natural constituents (Perry, 1980).

In tropical countries, modern medicines are not available to most of the rural populations. The World Health Organisation (WHO) has estimated that 80% of the world's population use botanical medicines for their primary health care needs (Farnsworth *et al.*, 1985). Many common health problems in tropical rural communities including respiratory infections, diarrhoea, fungal infections, diabetes, and malaria are treated by traditional botanical medicines. Numerous tropical medicinal plant species have shown *in vivo* and *in vitro* biological activities against these health problems (Carlson *et al.*, 1997).



Euphorbiaceae

Euphorbiaceae is a family of medicinal plants often with milky juice. These plants can be seen with either alternate leaves or opposite and glands which located at the barks of the plants or at the base of the leaf-blade. It has numerous flowers which are usually small or minute. The fruit is in a capsular formation and attached together with the seeds laterally. Plants in this family are mostly found in large numbers in less expected places such as waste ground and along the drain. Some of the useful plants in this family are *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri*. These plants are known as traditional healer in many rural areas throughout the world. It has been used for different purposes in different places (Kirtikar *et al.*, 1975).

Euphorbia hirta

Euphorbia hirta is also popularly known as *Euphorbia pilulifera* or locally called “ara tanah” or “susun nabi”. It is a small annual herb common to the tropical countries and belongs to the same family as the tic, tapioca and the rubber tree. This plant can easily be found in waste places, roadsides, gardens and rice fields. It can attain a height of 30 cm, growing erects or close to the ground (Blomquist and Oosting, 1940). *E. hirta* has a long history of usage in the treatment of various ailments. It is locally used in Africa and Australia. The plant has been used in almost every part of the world particularly Africa (Khan *et al.*, 1980).



Acalypha indica

Acalypha indica is an annual erect plant that can grow up to 30-75 cm high. It is also known as “rumpit lis-lis” or “kucing galak” in Malaysia. This plant has numerous branches and flowers on their soft bark. It grows widely in fields and bushes, but preferably on a sandy and open area. *Acalypha indica* is mostly found in warm tropical areas such as Peninsular Malaysia, Thailand and Java, Indonesia (Said, 2000). This plant has been used as traditional medication for quite sometime but not widely used due to the little information known about it. Usually, the leaves of this plant are used to treat several skin infections such as scabies and ringworm (Kirtikar *et al.*, 1975).

Phyllanthus niruri

Phyllanthus niruri is a short-lived usually erect little-branched annual herb, often completely green in colour including its flowers. It has simple and very small leaves with the seeds attached along the branches below the leaves. Locally, it is known as “dukung anak” and also referred as *Phyllanthus amarus*. This plant can be found in the waste ground and is native to tropical countries (Seaforth *et al.*, 1985). *Phyllanthus niruri* is known to be used to treat various types of ailments; almost every part of the plants can be use as medication. The plant is much used as diuretic and other problems of the genito-urinary tract. The fruit is useful for wounds, scabies and ringworm remedies (Kirtikar *et al.*, 1975).



Justification of the Study

In general, this study focuses on the effects of *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri* as antimicrobial and wound healing agents. With alarming situation of microbial resistance against several antibacterial and antifungal drugs, alternative medicine such as medicinal plants has become a very popular source for research in recent years. Another problem face nowadays is with prolonged or incomplete wound healing process. Vast efforts are being made to discover natural agents including from medicinal plants that can expedite healing and thereby reduce other severe complications.

Objectives of the Study

- 1) To evaluate the antibacterial activities of the aqueous and ethanol extracts of *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri* against selected gram positive and gram negative bacteria.
- 2) To evaluate the antifungal activities of the aqueous and ethanol extracts of *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri* against selected moulds and yeasts.
- 3) To observe the wound healing activity of the aqueous extracts of the leaves of *Euphorbia hirta*, *Acalypha indica* and *Phyllanthus niruri* on surgical incision in mice.