

UNIVERSITI PUTRA MALAYSIA

EFFECT OF EDUCATIONAL INTERVENTION ON THE INTENTION TO PROVIDE FAMILY-CENTRED CARE AMONG SELECTED HOSPITALS' PAEDIATRIC NURSES IN TEHRAN, IRAN

FOROUZAN ROSTAMI

FPSK(p) 2015 17



EFFECT OF EDUCATIONAL INTERVENTION ON THE INTENTION TO PROVIDE FAMILY-CENTRED CARE AMONG SELECTED HOSPITALS' PAEDIATRIC NURSES IN TEHRAN, IRAN

By

FOROUZAN ROSTAMI

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of Philosophy

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

Dedicated to the omnipresent God, my dear Lord for all the gifts that were given to us, my family for their love and endless support, and my great supervisor Professor Syed Tajuddin Syed Hassan for his valuable guidance and help.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

EFFECT OF EDUCATIONAL INTERVENTION ON THE INTENTION TO PROVIDE FAMILY-CENTRED CARE AMONG SELECTED HOSPITALS' PAEDIATRIC NURSES IN TEHRAN, IRAN

By

FOROUZAN ROSTAMI

August 2015

Chairman: Syed Tajuddin Syed Hassan, PhD Faculty: Medicine and Health Sciences

Family-centred care (FCC), which has become the cornerstone for paediatric nursing practice, supports the trust of child and family health. Families are recognised as an essential part of a specialised childcare during illness and they are expected to be skilful in childcare. In Iran, studies have shown that for parents to be involved in the care for hospitalized children there are obstacles that need to be discussed and investigated. There are huge gaps between effective health services and real world practice. Moreover, there is no common agreement between health care workers about provision of FCC in Iran. Hence, The main goal of this study was to examine, the effect of an educational intervention on the intention to provide family-centred care among paediatric nurses in Iran.

The research design of the study was quasi-experimental. A total of 200 paediatric nurses in two groups (100 intervention and 100 control), aged 25-55 from hospitals affiliated to Shahid Beheshti University (1 control and 3 intervention hospitals) in Tehran, Iran. Four hospitals were selected randomly based on the purpose of choosing hospitals that have paediatric wards for two groups (intervention and control groups). Proportionate sampling technique was employed. The educational intervention (twoday education classes) was conducted during the first month, and the effects on dependent variables of the study were assessed almost immediately after the intervention and three months later. In the intervention group, nurses were provided with presentations and internet-based education concerning FCC, in addition to booklets and pamphlets. The control group received one educational pamphlet about nutrition in children. To evaluate the effect of the intervention, data were collected at baseline, immediately following, and three months after the intervention, for both groups. Researcher-guided self-administered questionnaires were given to both groups prior to the intervention, immediately after the intervention, and three months after the intervention.

Descriptive and multivariate statistics (repeated measures ANOVA) were used for analysing the data. The mean age of participants was 36.1 years (SD = 9.1), and they were female (100%) without any specific training about FCC (93.8%). After the

intervention, there was a significant increase in the mean score of attitude [mean i.e. M (pre) = 3.35%, M (post) = 3.97, P < 0.001)], subjective norms [M (pre) = 3.72, M (post) = 4.21, P < 0.001)], perceived behaviour control [M (pre) = 3.36, M (post) = 3.75, P < 0.001)], and intention [M (pre) = 3.56, M (post) = 4.37, P < 0.001)], over the 3-month follow-up in the intervention group. The data provided preliminary support for effectiveness of the Theory of Planned Behaviour in producing constructs that significantly increased attitude, subjective norms, perceived behaviour control, and intention towards FCC in the intervention group.

The multiple linear regression models showed that changes in attitude, subjective norms, and perceived behaviour control scores were the predictors of the change in intention towards FCC. The most significant predictor was attitude ($\beta = 0.393$, t = 5.914, p < 0.01), followed by perceived behaviour control ($\beta = 0.320$, t = 4.815, p < 0.01) and subjective norms ($\beta = 0.172$, t = 2.184, p < 0.01). The last model was related to the follow-up test and the results showed the same pattern where attitude still showed the highest impact on intention ($\beta = 0.319$, t = 4.889, p < 0.01) followed by perceived behaviour control ($\beta = 0.309$, t = 5.174, p < 0.01) and subjective norms ($\beta = 0.264$, t = 4.138, p < 0.01).

The results provide evidence in support of the thesis that education can enhance attitude, subjective norms, perceived behaviour control, and intention to provide FCC within an educated group such as paediatric nurses, who represent a large portion of educated nurses in Iran. The study suggests that health care professionals can focus especially on teaching young nurses and parents to promote paediatric health through FCC.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KESAN INTERVENSI PENDIDIKAN TERHADAP NIAT UNTUK MENYEDIAKAN PENJAGAAN BERPUSATKAN KELUARGA DI HOSPITALS TERPILIH DI KALANGAN JURURAWAT PAEDIATRIK DI TEHRAN, IRAN

Oleh

FOROUZAN ROSTAMI

Ogos 2015

Pengerusi: Syed Tajuddin Syed Hassan, PhD Fakulti: Perubatan dan Sains Kesihatan

Penjagaan Berpusatkan Keluarga ("Family-Centred Care"- FCC), yang telah menjadi asas bagi amalan kejururawatan paediatrik, menyokong kepercayaan kanak-kanak dan kesihatan keluarga. Keluarga diakui sebagai bahagian yang penting bagi penjagaan kanak-kanak yang khusus ketika menghidapi penyakit dan keluarga juga diharap perlu mahir dalam aspek penjagaan kanak-kanak. Di Iran, kajian telah menunjukkan bahawa bagi ibu bapa melibatkan diri dalam penjagaan kanak-kanak yang dirawat di hospital, banyak halangan yang perlu dibincang dan diselidiki. Terdapat jurang yang besar antara perkhidmatan kesihatan yang efektif dan amalan dunia sebenar. Tambahan pula, tidak terdapat persetujuan umum antara pekerja penjagaan kesihatan, tentang penyediaan Penjagaan Berpusatkan Keluarga (FCC) di Iran. Justeru, tujuan utama kajian ini adalah untuk meneliti, menganalisis, dan menghuraikan kesan intervensi pendidikan terhadap niat untuk menyediakan penjagaan berpusatkan keluarga dalam kalangan jururawat paediatrik di Iran.

Pensampelan secara berkadar digunakan. Reka bentuk kajian ini ialah separa eksperimen. Sebanyak 200 jururawat paediatrik dalam dua kumpulan (100 kumpulan intervensi dan 100 kumpulan terkawal), berumur 25-55 tahun dari hospital yang bergabung dengan Universiti Shahid Beheshti (1 hospital terkawal dan 3 hospital intervensi) di Tehran (Iran), telah dipilih. Empat hospital dipilih secara rawak berdasarkan hospital yang mempunyai wad paediatrik. Intervensi Pendidikan (dua hari kelas pendidikan) diadakan dalam bulan pertama, dan dinilai hampir sebaik sahaja selepas intervensi, serta tiga bulan kemudiannya. Dalam kumpulan intervensi, jururawat telah diberikan pembentangan slaid, dan pendidikan berasaskan komputer tentang FCC di samping beberapa buku kecil dan risalah yang berkaitan. Kumpulan terkawal menerima satu buku kecil pendidikan tentang nutrisi kanak-kanak. Bagi menilai kesan intervensi, data dikumpul sebelum intervensi, dan sebaik sahaja selepas intervensi, dan pada tiga bulan selepas intervensi, bagi kedua-dua kumpulan. Bagi setiap kumpulan, soal selidik secara swaguna bimbingan penyelidik telah diberikan, sebelum intervensi, sebaik sahaja selepas intervensi, dan tiga bulan selepas intervensi.

Statistik deskriptif dan multivariat (Pengukuran Berulang ANOVA) telah digunakan bagi menganalisis data menggunakan SPSS versi 21. Purata umur responden ialah 36.1 tahun (SD = 9.1) dan mereka ialah perempuan (100%) dan tanpa latihan yang khusus tentang FCC (93.8%). Selepas intervensi, terdapat peningkatan yang signifikan dalam purata skor tentang sikap [min iaitu M(pra)= 3.35%, M(pasca)= 3.97, P<0.001)], norma subjektif [M(pra)= 3.72, M(pasca)= 4.21, P<0.001)], kawalan tingkah laku tercerap [M(pra)= 3.36, M(pasca)= 3.75, P<0.001)], dan niat [M(pra)= 3.56, M(pasca)= 4.37, P<0.001)], dalam tiga bulan janka masa susulan, dalam kumpulan intervensi. Data menyokong secara awalan, tentang keberkesanan Teori Tingkah Laku Terancang dalam menghasilkan beberapa konstruk yang signifikan, yang meningkatkan sikap, norma subjektif, kawalan tingkah laku tercerap, dan niat terhadap FCC, dalam kumpulan intervensi.

Model regresi linear berganda menunjukkan bahawa perubahan sikap, norma subjektif, dan skor tingkah laku terkawal tercerap, merupakan prediktor perubahan FCC dari segi niat. Prediktor yang paling signifikan ialah sikap (β = 0.393, t= 5.914, p<0.01), diikuti oleh kawalan tingkah laku tercerap (β = 0.320, t= 4.815, p<0.01) dan norma subjektif (β = 0.172, t= 2.184, p<0.01). Ujian susulan selepas tiga bulan intervensi juga menunjukkan corak yang sama, iaitu aspek sikap masih merupakan pemberi impak yang paling tinggi terhadap niat (β =0.319, t=4.889, p<0.01), diikuti oleh tingkah laku terkawal (β =0.309, t=5.174, p<0.01) dan norma subjektif (β =0.264, t=4.138, p<0.01).

Hasil kajian memberikan bukti yang menyokong tesis tentang pendidikan dapat mempertingkatkan sikap ("attitude"-ATT), norma subjektif ("subjective norms"-SN), tingkah laku terkawal tercerap ("perceived behavior control"-PBC), dan niat ("intention"-INT) bagi menyediakan FCC dalam kumpulan yang berpendidikan seperti jururawat paediatrik, yang mewakili sebahagian besar jururawat terdidik di Iran. Kajian ini juga mencadangkan supaya golongan profesional penjagaan kesihatan dapat memberikan tumpuan, terutamanya dari segi mendidik jururawat dan ibu bapa muda untuk meningkatkan kesihatan paediatrik, melalui FCC.

ACKNOWLEDGEMENTS

First and foremost, all praises to Allah the Almighty for giving me the strength and utmost courage to complete this thesis. It is by his wish that this humble work has been completed.

I would like to sincerely express my deepest gratitude to my supervisor, Professor Dr. Syed Tajuddin bin Syed Hassan, and my supervisory committee members, Professor Dr, Turiman bin Suandi, Dr, Suriani, and Associate Professor, Dr, Farideh Yaghmaie for their dedication, patience and invaluable comments that made writing this thesis possible. I am also grateful to the officials in Shahid Beheshti University of Iran for their assistance and advice that gave me impetus to complete this thesis on time.

My thanks also go to the Dean and Deputy Dean of Research of Graduate Studies, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for all their support to this study. My sincere thanks also go to all coordinators in Shahid Beheshti and International Branch of Shahid Beheshti University in Iran and all nurses in the selected hospitals (Shohada Tajrish, Mofid, Moddaress, and Taleghani) for their cooperation through the study.

I am also grateful to Chaloos Islamic Azad University, Iran for giving me this opportunity to pursue my studies. I would also like to thank my colleagues for their kind assistance.

Last but not least, a big thank you to my father and mother and two brothers for all their love and support and their constant encouragement, throughout my life and this study.

I certify that a Thesis Examination Committee has met on 10 August 2015 to conduct the final examination of Forouzan Rostami on her thesis entitled "Effect of Educational Intervention on the Intention to Provide Family-Centred Care among Selected Hospitals' Paediatric Nurses in Tehran, Iran" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Zailina binti Hashim, PhD

Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

Latifah binti A Latiff, PhD

Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Internal Examiner)

Khatijah binti Abdullah, PhD

Professor University Malaysia Malaysia (External Examiner)

Rama C Nair, PhD

Professor University of Ottawa Canada (External Examiner)

Common of the co

ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date: 22 September 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Syed Tajuddin bin Syed Hassan, PhD

Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

Turiman bin Suandi, PhD

Professor Faculty of Educational studies Universiti Putra Malaysia (Member)

Suriani binti Ismail, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Farideh Yaghmaei, PhD

Associate Professor
Faculty of Nursing
Islamic Azad University, Zanjan Branch
Iran
(External Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:	Date:	
Nama and Matria No : Foro	uzon Poetami GS23508	

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature:		Signature:	
Name of		Name of	
Chairman of		Member of	
Supervisory	Syed Tajuddin bin Syed	Supervisory	Turiman bin Suandi, PhD
Committee:	Hassan, PhD	Committee:	
Signature:		Signature:	
Name of		Name of	
Member of		Member of	
Supervisory	Suriani binti Ismail, PhD	Supervisory	Farideh Yaghmaei, PhD
Committee:		Committee:	

TABLE OF CONTENTS

			Page
ΑŦ	BSRTA	CT	i
	STRA		iii
		WLEDGEMENT	V
	PROV		vi
		RATION	viii
		TABLES	xiii
		FIGURES	xiv
		ABBREVIATION	XV
CI	HAPTI	ER .	
1	INT	RODUCTION	1
1	1.1	Background	1
	1.2		4
	1.3		6
	1.4	1	6
	1.5		6
	1.6	Research Hypothesis	7
	1.0	Research Trypodiesis	,
2	LIT	ERATURE REVIEW	8
	2.1		8
	2.2	Family-Centred Care	9
		2.2.1 Issues and Relevance	12
		2.2.2 ElementI: Two-way and Respectful Communication	14
		2.2.3 Element II: Partnership and Collaboration	15
		2.2.4 Element III: Caring and Relationship Building	16
		2.2.5 Advantages of FCC	17
		2.2.6 Disadvantages of FCC	18
	2.3	Role of Paediatric Nurse in FCC	19
	2.4	Theory of Planned Behaviour Related to FCC	19
		2.4.1 Attitude	19
		2.4.2 Subjective Norms (SN)	22
		2.4.3 Perceived Behavioural Control (PBC)	22
		2.4.4 Behavioural Intention	24
	2.5	Transformative Learning Theory Foundation for Developing FCC	28
	2.6	Barriers to the Implementation and Practice of FCC	30
	2.7	Conceptual Framework for the Family-Centred Care	32
3	MET	THODOLOGY	34
	3.1	Introduction	34
	3.2	Study Location and Duration	34
	3.3	Study Design	34
	3.4	Sampling	34
		3.4.1 Study Population	34
		3.4.2 Sampling Unit	35
		3.4.3 Inclusion Criteria for Nurses	35
		3.4.4 Exclusion Criteria for Nurses	36

		3.4.5	Sample Size	36
		3.4.6		37
		3.4.7	Sampling Technique	40
	3.5	Admir	nistration Approval	40
	3.6	Data C	Collection	40
		3.6.1	Data Collection Method	40
	3.7	Instrui	ments	42
		3.7.1	Questionnaire Development	42
		3.7.2		47
		3.7.3	·	47
		3.7.4	Content Validity	48
			Pilot Study	49
			Reliability of the Questionnaire	50
	3.8		Educational Intervention	50
		3.8.1	Educational Material	50
		3.8.2	Implementation of the Intervention	51
	3.9	Variat	oles and Operational Definitions	52
		3.9.1	Independent Variables	52
		3.9.2	Dependent Variables	53
	3.10	Ethics	Consideration	54
	3.11	Data F	Procurement and Sequence of Analysis	54
	3.12		ality Test and Outliers	54
	3.13		Analysis	55
	3.14		ng Data	57
1	RESU	JLTS		59
	4.1	Introd	uction	59
	4.2	Demo	graphic Characteristics of the Respondents	59
	4.3	Relation	onship between Socio-Demographic Variables and Attitude,	62
		Subjec	ctive Norms, Perceived Behavior Control and Intention	
		4.3.1	Relationship between Socio-demographic Characteristics	62
			and Attitude	
		4.3.2	Relationship between Socio-demographic Characteristics	62
			and Subjective Norms	
		4.3.3	Relationship between Socio-demographic Characteristics	65
			and Perceived Behaviour Control	
		4.3.4	Relationship between Socio-demographic Characteristics	67
			and Intention	
	4.4		hanges in Values of TPB Variables between Intervention and	69
		Contro	ol Groups Comparing Baseline, Immediately after	
		Interve	ention and Follow-up Test	
		4.4.1	The Differences in Scores of Attitude between Intervention	69
			and Control Group Comparing Baseline, Immediately after	
			Intervention and Follow-up Test	
		4.4.2	The Differences in Scores of Subjective Norms between	71
			Intervention and Control Group Comparing Baseline,	
			Immediately after Intervention and Follow-up Test	
		4.4.3	The Differences in Scores of Perceived Behaviour Control	73
			between Intervention and Control Group Comparing	
			Baseline, Immediately after Intervention and Follow-up	

		4.4.4	The Differences in Scores of Behavioural Intention between Intervention and Control Groups Comparing Baseline, Immediately after Intervention and Follow-up Tests	75
	4.5	Interve	iation between TPB Variables at Baseline, Immediately after ention and Follow-up Test in Intervention and Control	77
		Group 4.5.1	Correlation between Attitude, Subjective Norms and Perceived Behaviour Control at Baseline, Immediately after Intervention and Follow-up Tests in the Intervention Group	77
		4.5.2	Correlation between Attitude, Subjective Norms and Perceived Behaviour Control at Baseline, Immediately after Intervention and Follow-up Tests in the Control	79
	4.6		Group of Attitude, Subjective Norms and Perceived Behavior of on Intention	81
	4.7	Summ		84
5		USSIO		85
	5.1	Overv		85
	5.2		of Socio-demographic Characteristics on Intention	85
	5.3		of Intervention on Changes in Attitude	87
	5.4		of Intervention on Changes in Subjective Norms	89
	5.5	Effect Contro	of Intervention on Changes in Perceived Behavioural	90
	5.6		ffect of Attitude, Subjective Norms, and Perceived Behaviour of on Intention towards Family-Centred Care	91
	5.7	Summ		93
6	CON	CLUSI	ON, LIMITATIONS AND RECOMMENDATIONS FOR	94
	FUT	URE RI	ESEARCH	
	6.1	Summ	nary and Conclusion	94
	6.2		ations of the Study	94
	6.3	Streng	ths of the Study	95
	6.4	Implic	cations	96
		6.4.1	Implications to Nursing Theories	96
		6.4.2	Implications to Nursing Practice	96
		6.4.3	Implications to Health System Organization	96
		6.4.4	Implications to Education	96
	6.5	Recon	nmendations for Future Research	97
RE	FERE	NCES		98
	PEND			118
			TUDENT	216
			CATIONS	217

LIST OF TABLES

Table]	Page
3.1	Information Used in Sample Size Calculation	37
3.2	Paediatric Nurses in the Selected Hospitals	38
3.3	Sampling Framework	39
3.4	Content of open-ended questions to elicit commonly held attitude, subjective norms, perceived behavior and intention to provide family-centered care see also Appendix (D. E. F. G)	46
3.5	Cronbach's Alpha Value of the Instruments	49
3.6 3.7	Normality Test of Variables in Both Intervention and Control Groups Hypotheses and Analysis	55 58
4.1	Comparison between Intervention and Control Group for Respondents' Background and Characteristics	61
4.2	Criteria for Interpreting Strength of Relationship between Two Variables	62
4.3	Correlation between Socio-Demographic Characteristics and Attitude	63
4.4	Correlation between Socio-Demographic Characteristics and Subjective Norms	64
4.5	Correlation between Socio-Demographic Characteristics and Perceived Behaviour Control	66
4.6	Correlation between Socio-Demographic Characteristics and Intention.	68
4.7	Comparison of Changes in the Attitude between Control and Intervention According to Time and Its Graph	70
4.8	Comparison of Changes in Subjective Norms between Control and Intervention Group According to Time and Its Graph	72
4.9	Comparison of Changes in Perceived Behavioural Control between Control and Intervention Groups According to Time and Its Graph (1) Baseline, (2) Immediately after Intervention, (3) Follow Up	74
4.10	Comparison of changes in Behavioural Intention between Control and Intervention Groups According to Time and its Graph (1) Baseline, (2) Immediately after Intervention, (3) Follow up (3 Months after Intervention)	76
4.11		78
4.12	Follow-up Test in the Intervention Group (n=99) Correlation between Attitude, Subjective Norms and Perceived Behaviour Control at Baseline, Immediately after Intervention, and	80
4.13	Follow up Test, in Control Group (n=97) Correlation between Intention and Attitude, Subjective Norms and Perceived Behaviour Control	81
4.14	Model Summary for Multiple Linear Regression in Baseline, Immediately after Intervention, and Follow-Up Test	82
4.15	Regression Coefficients in Baseline, Immediately after Intervention, and	83

LIST OF FIGURES

Figure	e	Page
1.1	Theory of Planned Behaviour (Ajzen & Fishbein, 1985)	3
2.1	Conceptual Framework for the Family-Centred Care of Pediatric Nurses	33
3.1	Flow chart of study design and outcome evaluation (number of hospitals: control; intervention)	35
3.2	The Flow Chart of Enrolment and Data Collection	41
3.3	Flow Chart of the Ouestionnaire Design	44

LIST OF ABBREVIATIONS

FCC Family-Centred Care PCC Patient-Centred Care

PFCC Patient and Family-Centred Care

IPFCC Institute of Patient and Family-Centred Care

PICU Paediatric Intensive Care Unit TPB Theory of Planned Behaviour TRA Theory of Reasoned Action

ATT Attitude

SN Subjective Norms

PBC Perceived Behaviour Control
WHO World Health Organization
MPOC Measure of Process of Care
CSHCN Children with Special Health Care
CCNFI Critical Care Family Needs Inventory

FPOR Family Presence during Paediatric Cardiopulmonary Resuscitation

IMCI Integrated Management of Children Illness

UNICEF United Nations International Children and Emergency Fund

ANOVA Analysis of Variance

RMANOVA Repeated Measures Analysis of Variances

SD Standard Deviation ICU Intensive Care Unit

NICU Neonatal Intensive Care Unit
AAP American Academy of Paediatrics

IPFCC Institute of Patient and Family-Centred Care

UNEPSA Union of National European Paediatric Societies and Associations

NHS National Health Services

CHAPTER 1

INTRODUCTION

1.1 Background

During the past 20 years, the pattern of mortality in children has dramatically changed as life expectancy in neonatal and infantile ages has increased (Hockenberry & Wilson, 2009). These changes were due to advances in medical technology, control and management of diseases, in-time diagnosis and progress in public health as well as preventive measures. Significant decrease in mortality rate has increased the number of children with special health care needs and with chronic conditions, who are in need of more medical services than ordinary medical conditions (Hockenberry & Wilson, 2009).

Family is defined as a unit made up of child admitted to hospital and his or her caretakers who are bound together by emotional, biological, economic or other ties (Maijala & Astedt-Kurki, 2009). The family role is pivotal to the child's sense of security and trust in hospital (Hopia *et al.*, 2005; Lehto, 2004).

When a child develops severe or chronic diseases, the family has to readjust their daily activities and commitments due to the need to care for the child. Daily activities of the family previously planned to meet its needs are converted to satisfy the present needs of the child (Ahmann, 1994). In addition to the influence of disease on the child and the family, the health care system is also affected, leading to decreased quality of life and increased medical costs for society (McCarthy *et al.*, 2002). Nevertheless, the family is the basic unit of society and it is considered the primary unit of care (Harrison, 2010). It seems that family-centred care (FCC) would be a central part of child nursing (Hutchfield, 1999). Traditionally, the focus of nursing is on the provision of care to individual patients. However, every patient is also a member of a family. Family nursing is a practice and a science that has been developed over the past 20 years as a way of approaching and working with families. All nursing, according to this way of thinking, should be done in the context of the family (Hanson, 2005).

During the past 40 years, great advances have been made in medical science and technology to satisfy the physical needs of a sick child; however, psychological needs of the child and the family have received less attention (Saunders *et al.*, 2003). Studies regarding the needs of families with hospitalised children show that their important needs include respect and support as well as cooperation in the health planning of their children (Conway *et al.*, 2006; Kuo et al., 2012; Wells, 2011).

These three concepts are the components of FCC (Pettoello-Mantovani *et al.*, 2009). Furthermore, in families' points of view, nurses, compared to other health professionals, have to attend much more to their needs, respect the activities and capabilities of them in providing care more and support in the best possible way (Balling & McCubbin, 2001).

According to reports of the Institute for Family-Centred Care in the USA, FCC can be implemented in any nursing area. For example, in neonatal wards, parents reported

feeling less anxious after the adoption of FCC. Also, FCC leads to shorter hospital stay and fewer re-hospitalisations (Conway *et al.*, 2006).

A number of authors have compared traditional medical approaches with patient-centred care. Today, patient-centred care is widely acknowledged as a core value in family medicine (Hudon *et al.*, 2011). Health care workers should use principals of FCC during the admission process of children in the hospitals (Bowden & Greenberg, 2007). The concept of FCC was described by the Association for the Care of Children's Health (ACCH) in 1987 (Harrison, 2010) as including the following eight equally important elements:

- 1. recognition that the family is the constant in the child's life whereas the service systems and health care workers within those systems change.
- 2. facilitation of parent and professional collaboration at all levels of health care.
- 3. recognition of family strengths and individuality and respect for different methods of coping and adaptation.
- 4. sharing unbiased and complete information with parents about their child's care on an ongoing basis in an appropriate and supportive manner.
- 5. encouragement and facilitation of parents-to-parents support.
- 6. understanding and incorporating the developmental needs of infants, children, and adolescents and their families into the health care delivery system.
- 7. implementation of appropriate policies and programmes that are comprehensive and provide emotional support to meet the needs of families.
- 8. assurance that the design of health care delivery systems is flexible, accessible, and responsive to family needs.

Researchers have utilised the theory of planned behaviour (TPB) in the area of providing FCC by nurses as a framework to understand stages of behavioural changes and their determining factors. It is a valid theory in understanding human behaviour (Mathew *et al.*, 2007). TPB is an extension of the Theory of Reasoned Action that also measures self-efficacy beliefs, also known as perceived behavioural control (Bandura, 1977). The TPB was developed and introduced by Fishbein and Ajzen in (1985). Thus, the TPB can predict behaviour by three factors; attitude toward the behaviour (ATT), subjective norms (SN) and perceived behavioural control (PBC). According to the TPB (Fishbein & Ajzen, 1975), intention can be determined by the attitude toward performing the behaviour and subjective norms and perceived behaviour control.

Attitude can predict the intention of an individual to perform behaviour about an objective. Fishbein and Ajzen (1975) state that attitude can determine a person's overall intentions. Attitude influences the general level of person's intention and the relationship between attitude and intention can be determined if the intention is measured. High correlation between attitude and intention is expected when attitude and intention are measured at the same time and on the same target. Thus, the measuring of intention can predict positive or negative behaviour.

Fishbein and Ajzen (1980) argue that high influence of subjective norms in a person can provide a greater intention to perform behaviour. Assessment and measurement of subjective norms for predicting and understanding intentions are necessary. In some situations, both the attitude and subjective norms are not equally weighted for the behavioural intention. A person may have a positive attitude towards performing a specific behaviour but his or her subjective norms suggest that he or she should not perform it or vice versa (Ajzen & Fishbein, 1980). The perceived behavioural control

reflects the perception of internal and external constraints of behaviour (Ajzen, 1985). Thus, a person's perception of control over behaviour that the person is able to control their behavioural performance is defined as perceived behavioural control.

The effects of attitude (A) and subjective norms (SN) and perceived behavioural control (PBC) are mediated by behavioural intention (BI). Intention is viewed as being related to the corresponding behaviour. Behavioural intention in individuals is a subjective probability that they will perform some behaviour and this is regarded as a function of attitude and the subjective norms. The best predictor for behaviour is the intention to perform the behaviour, thus, the measure of intention will lead to accurate prediction of behaviour. Furthermore, accurate prediction of behaviour depends on the strength of the intention-behaviour relationship (Fishbein & Ajzen, 1975).

Explanation of behaviour is considered in the PBC. This theory provides a comprehensive framework that has been used to explain many of the findings in behavioural change. Ajzen and Fishbein (1980) state that measurement of the variables of the TPB for understanding the relationship among variables is necessary. These measurements include: 1) attitude toward behaviour, 2) subjective norms, 3) perceived behaviour control, 4) intention to perform behaviour, and 5) behaviour.

The Theory of Planned Behaviour was likely to be most applicable in developed countries such as Canada and USA. However, it is essential to test this theory in developing countries such as Iran. This is especially important with respect to the concept of family-centred care. The Theory of Planned Behaviour was applicable through different approaches in developed countries such as Spain (Gonzalez et al., 2012), Australia (Kam et al., 2010), USA (Matthew et al., 2007), UK (Arnold et al., 2005), and Europe (Topa, 2010). There is thus a need to test FCC (based on TPB) in developing countries such as Iran. The hospitals in developed countries are well resourced, while many in developing countries, such as Iran, function with limited resources. Nurses in developed countries, but this might be culturally inappropriate (Shields & Nixon, 2004).




Figure 1.1: Theory of Planned Behaviour (Ajzen & Fishbein, 1985)

1.2 Problem Statement

The organization of paediatric care shows profound differences and divergent problems throughout the world, particularly between economically advantaged and challenged countries (Pettoello-Mantovani *et al.*, 2009). However, significant differences also exist among the economically developed countries. For instance, in the US, the concepts of family-oriented and family-centred care emerged and developed to become consolidated parts of its health system over the past fifty years. However, it has been observed that the US stands apart from most similarly developed countries in having a coherent set of public policies that would create a social context supportive of families, traditional or otherwise (American Academic of Paediatrics, 2003).

In Iran, the mortality rate among children has decreased considerably from 87 to 51 deaths per 1000 live births (UNICEF, 2012). In absolute terms, this means that the number of children who die each year has decreased from 12 to 6.9 million on a global basis. In spite of this significant achievement, more than 19,000 children still die every day, and most of these deaths are caused by preventable and treatable infectious diseases and by the deficiencies in family-centred practices. About 80 per cent of healthcare in developing countries occurs at home, and the majority of children die at home without being seen by a healthcare worker. It has been hypothesized in a report on equity and access to quality care that as many as 40% of children's deaths could be prevented with improved family and community care by providing access to solid knowledge, support, and basic provisions. In Iran, which has a total population of 74 million people, the infant mortality rate is 41(deaths/1,000 live births).

The statistics in Iran showed that 6.5% of neonates were of low birth weight among which 75% were premature and needed to be hospitalized in neonatal intensive care unit (NICU) (Ramezani *et al.*, 2014). This experience threatens not only neonates' health, but also family members' health and brings about a vast array of problems (Farahani *et al.*, 2014). This causes a delay in parental-neonatal attachment (especially for mothers) and upsets mothers' attachment process toward their children. In the Middle East, health care has undergone some changes with an increased stress on FCC, which is defined as a health care that recognizes the need for a partnership among healthcare workers, patients, and patients' families in a suitable and holistic manner. This partnership should ensure that the decisions that are made respect the wants, needs, and preferences of the patients and their families (Aliabadi *et al.*, 2011).

The Iranian tradition has maintained a strong family bond and parental responsibility for the care of children. Parental contribution to hospital care is a symbol of the strength of the Iranian tradition. The National Health Services in Iran have begun a movement for improving health care services in different settings based on consumer needs (Corlett and Twycross, 2006). The role of FCC in pediatric nursing, common understanding between the nursing staff and the child's parents can lead to providing higher quality of medical attention. However, nurses and parents have different perceptions of stressors in the child's admission to a hospital in Iran (Hassan Tehrani *et al.*, 2012).

In recent years, health care professionals have accepted that appropriate treatment should be done involving relationship and interaction between the family and the medical team at the hospital. This is one of the fundamental components of care and the philosophy of FCC (Khosravian *et al.*, 2013). Although improving the quality of

nursing care and higher patient and nurse satisfaction have been reported as the desired effects of family involvement in the nursing care of patients, neglecting the needs of the patients' families and imposing them more responsibilities than their capabilities can have adverse effects such as feelings of inadequacy and inability (Aein *et al.*, 2009). Few studies have assessed the status of participation of the patients' families as well as the views of families and nurses concerning the role of families in caring for their relatives. In particular, there is not enough reliable evidence on the status of participation of the families in the care of their patients. Family participation is a key indicator of family-centered care in Iran (Molazem *et al.*, 2011; Zarei *et al.*, 2009).

Interaction and relation of the families with the hospital treatment team is one of the essential components of care, but this need was not properly met. The effective presence of the nurse can balance the role of the family beside the patient. In this regard, training the personnel for better and more understanding of the families and involving them in the care based on their abilities according to the philosophy of family-centered care has been suggested (Khosravian *et al.*, 2013).

Fakhr-Movahedi *et al.* (2011) studied the nurse-patient relationship in Iran and showed that factors such as organization, culture and tradition, expectations from nurses, and heavy workloads can limit the time allocated to communicate with patients and reduce the effectiveness of communication. In addition, although nurses' communication behaviour including friendly personality, kindness, quick response, and timely attending to the needs of patients and allocating sufficient time to provide care could improve communication with the patient, heavy workload and shortage of nurses are common problems prevalent in the training hospitals of Iran, which could potentially affect this behaviour (Zamanzadeh *et al.*, 2014).

Despite substantial investments in researchers, dissemination and advocacy, a huge gap exists between what is known about effective health services and what is done in real world practice and there is not a consensus among health care workers about the provision of FCC in Iran. These barriers may be due to lack of knowledge and skills, lack of support, law, nurses' powerlessness, nurse-patient relationship, and recognizing patient needs that could be influential in the provision of FCC in Iran. It seems that participation of parents in care of their hospitalized children needs more discussion and research in Iran.

Most of the studies have considered the viewpoints of the parents, disregarding the viewpoints of the nurses in Iran. A small number of studies have researched about provision of FCC on TPB by nurses. However, there is no definitive study based on the concept, analyzing in depth the influence of ATT, SN, and PBC, and the interactions. As a result, this study focuses on ATT, SN and PBC of nurses related to the implementation of family-centered care in pediatric wards. The present study was done in an Iranian context in an effort to address the following questions:

- Is there a relationship between TPB elements and nurses' intention to provide ECC?
- Is there an intervention impact on the nurses' ATT, SN, PBC and intention to provide FCC?

1.3 Scope and Significance

Community health scholars contend that the integrity of community-based nursing is based on FCC (Ayers, Bruno, & Langford, 1999) and state that the individual client is part of a general unit as family. Therefore, the needs of this unit should be systematically assessed and met (Potter & Perry, 2009).

During the recent decade, family and patients' satisfaction with the health care system has gained increasing attention as a valid and reliable criterion to measure quality of care. This also applies in nursing and medical interventions for children. Due to the role of children in the health of society and future generation, their parents are regarded as their respondents and representatives. Parents as advocates of children in health care system are suitable indicators to reveal the views of children. In spite of extensive support for FCC, many hospitals still do not apply the principles of this philosophy persistently. Additionally, in organizational cultures of hospitals, there are barriers and limitations for including viewpoints of patients and families and their involvement. Central to these restrictions are the lack of understanding of their needs and knowledge concerning FCC (Conway et al., 2006).

FCC is a new phenomenon or innovative approach to planning and evaluation is based on mutual understanding and beneficial partnership among health care professionals, patients and families (Johnson *et al.*, 2008). The central concepts of FCC include dignity, exchanging information, participation and cooperation or collaboration (Johnson *et al.*, 2008). The core of this definition of FCC is the relationship between families and health professionals and understanding that families are considered full partners in the provision of health care to children (MacKay & Gregory, 2011). The family has a pivotal and critical role in the development and nursing of their child (Harrison, 2010). Although FCC is approved as an interesting idea in many paediatric institutions, biased attitudes of health care workers toward parents and families have hindered its complete and successful provision (Depompei, Whitword, & Beam, 1994).

Perceptions of parents and staff about the care of hospitalized children have been explored in developed countries; however, little research has examined these perceptions in developing countries. Assumptions about family-centred care are often based on Western values, with little evidence of how cultural constructs may affect care delivery in developing nations (Sheild & Nixon, 2004).

1.4 Research Question

Can an educational intervention on paediatric nurses improve their attitude, subjective norms, perceived behaviour control, and intention towards provision of family-centred care?

1.5 Research Objectives

General Objective

To determine the effect of an educational intervention on the intention to provide family- centred care services among nurses in Iran based on the Theory of Planned Behaviour.

Specific Objectives

- 1. To determine the relationship (Pearson product-moment correlation coefficient r, and linear multiple regression) between respondents' (nurses) socio-demographic characteristics and attitude, subjective norms, perceived behaviour control, and intention in the provision of family-centred care.
- 2. To compare the nurses' attitude about the provision of family-centred care before and after the intervention between the intervention and control group.
- 3. To compare the nurses' subjective norms related to the provision of family-centred care before and after the intervention between the intervention and control group.
- 4. To compare the nurses' perceived behaviour control related to the provision of family-centred care before and after the intervention between the intervention and control group.
- 5. To compare the nurses' intention to the provision of family-centred care before and after the intervention between the intervention and control group.
- 6. To determine the effect of an educational intervention on the intention of nurses to provide family-centred care during the study.

1.6 Research Hypothesis

- Socio-demographic characteristics of nurses (age, income, training and work experience in paediatric ward) are significantly associated with attitude, subjective norms, perceived behaviour control and intention in the provision of family-centred care behaviour by nurses.
 - 1. There are significant differences, between nurses' attitude towards the provision of family-centred care, comparing intervention and control groups.
 - 2. There are significant differences, between subjective norms related to the provision of family-centred care by nurses, comparing intervention and control groups.
 - 3. There are significant differences, between perceived behaviour control related to the provision of family-centred care by nurses, comparing intervention and control groups.
 - 4. Nurses' attitude, subjective norms and perceived behaviour control significantly predict the intention of the provision of family-centred care.

REFERENCES

- Abbaszadeh, A., Sabeghi, H., Heidary, A., & Borhani, F. (2012). Assessment of the Effect of Continuing Education Program on Nurse's Knowledge, Attitude and Performance about Documentation. *Evidence Based Care*, 2(1), 75–83.
- Abraham, M., Moretz, J. g. (2012). Implementing patient-and Family-Centered Care: Part 1 Understanding the challenges, Pediatric Nursing, 38(1), 44-48.
- Aday, L. A., & Cornelius, L. J. (2006). *Designing and conducting health surveys: a comprehensive guide* (3rd ed.). San Francisco, CA: John Wiley & Sons.
- Aein, F., Alhani, F., Mohammadi, E., & Kazemnejad, A. (2009). Parental participation and mismanagement: a qualitative study of child care in Iran. *Nursing & Health Sciences*, 11(3), 221–227.
- Ahmann, E. (1994). Family-centered care: Shifting orientation. *Pediatric Nursing*, 20, 113–116.
- Ahmann, E., & Johnson, B. H. (2000). Family-centered care: Facing the new millennium. *Pediatric Nursing*, 26(1), 87–90.
- Aliabadi T, Bastani F, Haghani H. (2011). Effect of Mothers' Participation in Preterm Infants' Care in NICU on Readmission Rates. *Journal of Hayat*, 17(2),71-77.
- American Heritage Dictionary of the English Language. (2015). 5th edn. Houghton Mifflin Horcourt Publishing Company.
- American Association of Colleges of Nursing (2013). 2012-2013 Enrollment and graduations in baccalaureate and graduate programs in nursing. Washington, DC.
- American Academy of Pediatrics. (2003). Preamble to Patient-Centered Medical Home Joint Principles. Every child deserves a medical home. *Pediatrics*, 112: 691-696.
- Arnold, J., Loan-Clark, J., Coombs, C. and Wilkinson, A. (2005). How Well can the Theroty of Planned Behaviour Account for Occupational Intentions? The Business School, Loughborough University, Ashby Road, Loughborough.
- Ajzen, I. (1985). From intentions to actions: A theory of planned behavior. In J. Kuhl & J. Beckmann (Eds.), *Action-control: From cognition to behavior*, 11–39. Heidelberg, Germany: Springer.
- Ajzen, I. (1991). The theory of planned behavior. *Organizational Behavior and Human Decision Processes*, *50*, 179–211.

- Ajzen, I. (2002). Perceived behavioral control, self-efficacy, locus of control, and the theory of planned behavior. *Journal of Applied Social Psychology*, 32, 665 683.
- Ajzen, I. (2005). *Attitudes, personality, and behavior* (2nd ed.). Milton-Keynes, England: Open University Press / McGraw-Hill.
- Ajzen, I. (2010). Icek Ajzen, Professor of psychology, University of Massachusetts. Retrieved from http://www.people.umass.edu/aizen/faq.html on September 23, 2010. American
- Ajzen, I. (2011). The theory of planned behaviour: reactions and reflections. *Psychology & Health*, 26(9), 1113–1127.
- Ajzen, I., & Fishbein, M. (1980). *Understanding attitudes and predicting social*. Englewood Cliffs, NJ: Prentice-Hall.
- Ajzen, I., & Fishbein, M. (1985). The Prediction of Behavior from Attitudinal and Normative Variables. *Journal of Experimental Social Psychology*, 466–488.
- Aminoroaia, M., Mashhadi, M., Maracy, M. R., & Attari, A. (2014). Efficacy of purposeful educational workshop on nursing care. *Journal of Education and Health Promotion*, 3, 82.
- Armitage, C. J., & Conner, M. (2001). Social Cognitive Determinants of Blood Donation. *Journal of Applied Social Psychology*, *31*(7), 1431–1457.
- Ayers, M., Bruno, A. A., & Langford, R. W. (1999). Community-based nursing care: Making the transition. St. Louis, MO: C. V. Mosby.
- Azocar, F., Cuffel, B., Goldman, W., & McCarter, L. (2003). The impact of evidence-based guideline dissemination for the assessment and treatment of major depression in a managed behavioral health care organization. *The Journal of Behavioral Health Services & Research*, 30(1), 109–118.
- Baier, R. (2012). Parents' and nurses' perceptions of patient-and-family centered care and the impact of the nurse-patient relationship (Master Thesis). University of Victoria. Retrieved from http://dspace.library.uvic.ca:8080/handle/1828/4146
- Balling, K., & McCubbin, M. (2001). Hospitalized children with chronic illness: Parental caregiving needs and valuing parental expertise. *Journal of Pediatric Nursing*, 16(2), 110–119.
- Bamm, E. L, Rosenbaum, P. (2008). Family-centered theory: Origins, development, barriers, and supports to implementation in rehabilitation medicine. *Archives of Physical Medicine and Rehabilitation*, 89(8), 1618–1624.
- Bandura, A. (1977). Social learning theory. Englewood Cliffs , *NJ*: Prentice Hall, 38(23). 247

- Barglin, G. M. (2013). Can a theory based on educational intervention change nurses' knowledge and attitudes concerning cancer pain management? A quasi-experimental design. *BMC Health Services Research*, *13*, 328.
- Bauer, M. S. (2002). A review of quantitative studies of adherence to mental health clinical practice guidelines. *Harvard Review of Psychiatry*, 10(3), 138–153.
- Beduz, M. A. (2008). A framework to guide the implementation of evidence based practice changes in nursing. Unpublished manuscript, school of nursing, McMaster University, Hamilton, Ontario.
- Beduz, M. A. (2010). *The Role of Attitude, Subjective norms, Perceived Behaviour Control, and Context in Nurses' Behavioural Intentions* (PhD dissertation). Mc master University, Hamilton, Ontario.
- Bell, J. M. (2013). Family nursing is more than family centered care. Journal of family nursing. 19(4), 411-417.
- Bellin, M. H., Osteen, P., Heffernan, C., Levy, J. M., & Snyder-Vogel, M. E. (2011). Parent and health care professional perspectives on family-centered care for children with special health care needs: are we on the same page? *Health & Social Work*, 36(4), 281–290.
- Berg, C. (2002). Influences on Swedish schoolchildren's dietary selection: focus on fat and fibre at breakfast. *Food & Nutrition Research*, 46(4), 194–196.
- Berry, R.S.Y. (1999). Collecting data by in-depth interviewing. Paper presented at the British Educational Research Association Annual Conference, University of Sussex at Brighton, September 2 5 1999
- Bernaix, L. W. (2000). Nurses' attitudes, subjective norms, and behavioral intentions toward support of breastfeeding mothers. *Journal of Human Lactation*, *16*(3), 201–209.
- Boler M, Zembylas, M. (2003). Discomforting Truths: The Emotional Terrain of Understanding Differences. In: Tryfonas P, editor. *Pedagogies of Differences: Rethinking Education for Social Justice*. New York: Routledge, 110–136.
- Boudreaux, E. D., Francis, J. L., & Loyacano, T. (2002). Family presence during invasive procedures and resuscitations in the emergency department: a critical review and suggestions for future research. *Annals of Emergency Medicine*, 40(2), 193–205.
- Bowden, V. R., & Greenberg, C. S. (2003). *Pediatric Nursing Procedures* (1st ed.). Philadelphia PA: Lippincott Williams-Wilkins.
- Bowden, V. R., & Greenberg, C. S. (2007). *Pediatric nursing procedures*. London, England: Lippincott Williams & Wilkins.

- Brady, D., & Hyde, A. (2002). Certificate-trained staff nurses' perceptions of the changes in nursing education in Ireland from certificate to diploma level. *Journal of Continuing Education in Nursing*, 33(5), 231–237.
- Bragadottir, H., & Reed, D. (2002). Psychometric instrument evaluation: the pediatric family satisfaction questionnaire. *Pediatric Nursing*, 28(5), 475–482.
- Brown, J., & Ritchie, J. A. (1990). Nurses' perceptions of parent and nurse roles in caring for hospitalized children. *Children's Health Care*, 19(1), 28–36.
- Bruce, B., Letourneau, N., Ritchie, J., Larocque, S., Dennis, C., & Elliott, M. R. (2002). A multisite study of health professionals' perceptions and practices of family-centered care. *Journal of Family Nursing*, 8(4), 408–429.
- Bruce, B., & Ritchie, J. (1997). Nurses Practices and perceptions of family centered care. *Journal of Pediatric Nursing*, 12(4), 214–222.
- Burns, N., & Grove, S.K.(2007). Understanding Nursing Research, building an Evidence-Based Practice . 4 edition. St. Louis, Missouri: Saunders.
- Callery, P. & Smith, L. (1991). A study of role negotiation between nurse and the parents of hospitalized children. *Journal of Advanced Nursing*, 16, 772-781
- Campbell, P. H., & Sawyer, L. B. (2007). Supporting learning opportunities in natural settings through participation-based services. *Journal of Early Intervention*, 29(4), 287–305.
- Cambridge University. (2015). Cambridge Advanced. Learners Dictionary & Thesaurus, Cambridge University Press.
- Casper, E. (2007). The theory of planned behavior applied to continuing education for mental health professionals. *Psychiatric Services*, *58*(10), 1324–1329.
- Cassista, J., Payne-Gagnon, J., Martel, B., & Gangnon, M. P. (2014). Applying Theory to understand Intervention Mapping Approach. *Nursing Research and Practice*, 8.
- Caty, S., Larocque, S., & Koren, I. (2000). Family-centered care in Ontario general hospitals: the views of pediatric nurses. *Canadian Journal of Nursing Leadership*, *14*(2), 10–18.
- Caty, S., Larocque, S.,& Koren, I. (2001). Family-centered care in Ontario general hospitals: The views of pediatric nurses. *Canadian Journal of Leadership*, 14(2), 10-18.
- Campinha-Bacote, J. (2011). Delivering patient-centered care in the midst of a cultural conflict: the role of cultural competence. *The online Journal of Issues Nursings*, 16(2): 5.

- Cranton P.(1994). Understanding and Promoting Transformative Learning, A Guide for Educators of Adults. San Francisco: Jossey-Bass Publishers.
- Chisholm A, Askham J. (2006).What do you think of your doctor? A review of questionnaires for gathering patients' feedback on their doctor [document on the Internet]. Picker Institute Europe; [cited 2008 Sep 14].Retrievedfromhttp://www.engage.hscni.net/library/What%20do%20you% 20think.pdf
- Christine, R., Kathleek, S., Abel, J. J., & Virginia, M. (2007). Attitudes Toward and Beliefs about Family Presence: A survey of Healthcare Providers, Patients' Families, and Patients. *American Journal of Critical Care*, 16(3), 270–282.
- Christianson, C. E, McBride, R. B, Vari, R. C, Olson, L, Wilson, H. D. (2007). From traditional to patient-centered learning: curriculum change as an intervention for changing institutional culture and promoting professionalism in undergraduate medical education. *Academic Medicine*, 82(11):1079–1088.
- Cohen, J., Cohen, P., West, S. G., & Aiken, L. S. (2003). Applied multiple regression/correlation analysis for the behavioral sciences (3rd ed.). Mahwah, NJ: Lawrence Erlbaum Associates.
- Cohen, L., Manion, L., & Morrison, K. (2007). *Research Methods in Education* (6th ed.). USA & Canada: Routledge.
- Coker, T. R., Rodriguez, M. A., & Flores, G. (2010). Family-centered care for US children with special health care needs: Who gets it and why? *Pediatrics*, 125(6), 1159–1167.
- Cole, J. C. (2008). How to deal with missing data: Conceptual overview and details for implementing two modern methods. In J. W. Osborne (Ed.), *Best practices in quantitative methods* (pp. 214–238). Newberry Park, CA: SAGE publications.
- Conway, J., Johnson, B., Edgman-Levitan, S., Schlucter, J., Ford, D., Sodomka, P., & Simmons, L. (2006). Partnering with patients and families to design a patient-and family-centered health care system: a roadmap for the future: a work in progress. Bethesda, MD: Institute for Family-Centered Care.
- Corlett, J., Twyeross, A. (2006). Negotiation of parental roles within family-centered care: a review of the research. *Journal of Cliical Nursing*, 15, 1308-1316.
- Cote, F., Gangon, J., Houme, P.K., Abdeljelil, A.B. & Gangon, M.P. (2011). Using the Theory of Planned Behaviour to predict nurses' intention to integrate research evidence into clinical descision-making. *Journal of Advanced Nursing*, 28, 2289-2298.
- Coyne, I., O'Neill, C., Murphy, M., Costello, T., & O'Shea, R. (2011). What does family-centred care mean to nurses and how do they think it could be enhanced in practice. *Journal of Advanced Nursing*, 67(12), 2561–2573.

- Coyne, I., Murphy, M., Costello, T., O'Neill, C.& Donellan, C. (2013). A survey of nurses' practices and perceptions of family-centered care in Ireland. *Journal of family Nursing*, 19(4), 469-488.
- Dadgari, A., Kasaeian, A., Dadvar, L., Kalatejari, M., Rezaee, M., & Mirzaee, M. (2012). Effects of Education on Nurses' Knowledge and Skill in Drug Dosage Calculation. *Knowledge and Health*, 7(1), 20–26.
- Davey N. (2006). *Unquiet Understanding: Gadamer's Philosophical Hermeneutics*. Albany, NY: State University of New York Press.
- Davies, B. L., & Hodnett, E. (2002). Labor Support: Nurses' Self-Efficacy and Views About Factors Influencing Implementation. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*, 31(1), 48–56.
- Dewberry, C. (2004). Statistical methods for organizational research: Theory and practice. Psychology Press.
- Depompei, K., Whitword, K., & Beam, P. (1994). One institution's efforts to implement family-centered care. *Pediatric Nursing*, 20, 119–121.
- De Rond, M., De Wit, R., & Van Dam, F. (2001). The implementation of a Pain Monitoring Programme for nurses in daily clinical practice: results of a follow-up study in five hospitals. *Journal of Advanced Nursing*, 35(4), 590–598.
- Desimone, R.L., Werner, J.M., Harrison, D. M. (2002). *Human Resource Development*. Orlando, FL: Harcourt. Inc.
- DeVellis, R. (2003). *Scale development: Theory and applications*. Newberry Park, CA: Sage Publications.
- DiCenso, A., Guyatt, G., & Ciliska, D. (2005). Evidence-based nursing: A guide to clinical practice. Hamilton, Ontario, Canada: *Elsevier Health Sciences*.
- Di Lorio, C. K. (2005). Measurement in health behavior: methods for research and evaluation. San Francisco, CA: Jossey-Bass.p; 218-219.
- Donabedia., A. (2003). *An Introduction to Quality Assurance in Health Care*. New York: Oxford University Press.
- Du Gas, B. W. (1983). *Introduction to patient care: a comprehensive approach to nursing* 4th ed. Philadelphia, PA: W.B. Saunders Company.
- Dunst, C. J., Trivette, C. M., & Hamby, D. W. (2007). Meta-analysis of family-centered helpgiving practices research. *Mental Retardation and Developmental Disabilities Research Reviews*, 13(4), 370–378.

- Dwyer, T., & Mosel Williams, L. (2002). Nurses' behaviour regarding CPR and the theories of reasoned action and planned behaviour. *Resuscitation*, 52(1), 85– 90.
- Eccles, M. P., Johnston, M., Hrisos, S., Francis, J., Grimshaw, J., Steen, N., & Kaner, E. F. (2007). Translating clinicians' beliefs into implementation interventions (TRACII): a protocol for an intervention modelling experiment to change clinicians' intentions to implement evidence-based practice. *Implementation Science*, 2(1), 27.
- Ekstedt, M., Stenberg, U., Olsson, M., & Ruland, C.M. (2014). Health Care Professionals' Perspectives of the experiences of family care givers during inpatient cancer care. *Journal of Family Nursing*, 20(4), 462-486.
- Ellis, J. A., McCleary, L., Blouin, R., Dube, K., Rowley, B., MacNeil, M., & Cooke, C. (2007). Implementing best practice pain management in a pediatric hospital. *Journal for Specialists in Pediatric Nursing*, 12(4), 264–277.
- Elliott, S. A., Brown, J. S. (2002). What are we doing to waiting list controls? *Behaviour Research and Therapy Journal*, 40(9), 1047–52.
- Epstein, R. M., Fiscella, K., Lesser, C. S., & Stange, K. C. (2010). Why the nation needs a policy push on patient-centered health care. *Health Affairs* (Millwood), 29(8), 1489–1495.
- Espezel, H. J. E., & Canam, C. J. (2003). Parent–nurse interactions: care of hospitalized children. *Journal of Advanced Nursing*, 44(1), 34–41.
- E-skills UK. (2004). Work Experience, http://www.e-skills.com/cgi-bin/wms.pl/430.
- Fakhr-Movahedi A, Salsali M, Negharandeh R, Rahnavard Z. (2011). A qualitative content analysis of nurse-patient communication in Iranian nursing. *International Nursing Review Journal*, 58:171-80.
- Farahani, M. A., Gaeeni, M., Mohammadi, N., Seyedfatemi, N. (2014). Giving information to family members of patients in the intensive care unit: Iranian nurses' ethical approaches. *Journal of Medical Ethical and History of Medicine*, 7:9.
- Farrell, M. E., Joseph, D. H., & Schwartz-Barcott, D. (2005). Visiting hours in the ICU: finding the balance among patient, visitor and staff needs. *Nursing Forum*, 40(1), 18–28.
- Federal Research Division. Library of Congress.(2008). *Ethical Groups and Languages of Iran*. Retrieved from:http://lcweb2.loc.gov/frd/cs/profiles/Iran. pdf. on November 10.2010).
- Fishbein, M., & Ajzen, I. (1975). *Belief, attitude, intention and behavior: An introduction to theory and research.* Reading, MA: Addison-Wesley.

- Fishbein, M., & Ajzen, I. (1980). *Understanding attitudes and predicting social behaviour*. Englewood Cliffs, NJ: Prentice-Hall.
- Fishbein, M., & Ajzen, I. (2010). *Predicting and changing behavior: The reasoned acion approach*. New york, NY: Psychologic Press.
- Fisher, H. R. (2001). The needs of parents with chronically sick children: a literature review. *Journal of Advanced Nursing*, *36*(4), 600–607.
- Foster, M. J., Whitehead, L., Maybee, P. & Cullense, V. (2013). The parents', hospitalized child's, and health care providers' perceptions and experiences of Family-Centered Care within a pediatric critical care setting: A metasynthesis of qualitative research. *Journal of Family Nursing*, 19(4), 431-468.
- France, N., Byers, D., Kearney, B., & Myatt, S. (2011). Creating a healing environment: nurse-to-nurse caring in the critical care unit. International *Journal For Human Caring*, 15(1), 44-48.
- Francis, J. J., Eccles, M. P., Johnston, M., Walker, A., Grimshaw, J., Foy, R., & Bonetti, D. (2004). *Constructing questionnaires based on the theory of planned behavior*, Newcastle upon Tyne, UK: Centre for Health Services Research, University of Newcastle. 2-12.
- Friedkin, N. (2010). The attitude-behavior linkage in behavioral cascades. *Social Psychology Quarterly*, 73: 196-213.
- Galvin, E., Boyers, L., Schwartz, P. K., Jones, M. W., Mooney, P., Warwick, J., & Davis, J. (1999). Challenging the precepts of family-centered care: testing a philosophy. *Pediatric Nursing*, 26(6), 625–632.
- Gavaghan, S. R., & Carroll, D. L. (2002). Families of critically ill patients and the effect of nursing interventions. *Dimensions of Critical Care Nursing*, 21 (2), 64–71.
- Ganzalez, S,T., Nipp Lopez., M. C., Marcos, Y.Q., Rodriguez-Marin, J. (2012).

 Development and Validation of the Theory of Planned Behaviour Questionnaire in Physical Activity. *The Spanish Journal os Psychology*. 15(2), 801-816.
- George, D., & Mallery, P. (2003). SPSS for windows step by step: A simple guide and refrence. 11.0 update, 4th edn. Boston, MA: Allyn and Bacon Boston.
- Giles, M., McClenahan, C., Armour, C., Millar, S., Rae, G., Mallett, J., & Stewart-Knox, B. (2014). Evaluation of a theory of planned behaviour–based breastfeeding intervention in Northern Irish Schools using a randomized cluster design. *British Journal of Health Psychology*, 19(1), 16–35.
- Giles, M., Connor, S., McClenahan, C., & Mallett, J. (2010). Attitudes to breastfeeding among adolescents. *Journal of Human Nutrition and Dietetics*, 23, 285–293.

- Godin, G., Bélanger-Gravel, A., Eccles, M., & Grimshaw, J. (2008). Healthcare professionals' intentions and behaviours: A systematic review of studies based on social cognitive theories. *Implementation Science*, *3*(36), 1–12.
- Godin, G., Naccache, H., Morel, S., & Ébacher, M.-F. (2000). Determinants of nurses' adherence to universal precautions for venipunctures. *American Journal of Infection Control*, 28(5), 359–364.
- Gottschall, A. C., West, S. G., & Enders, C. K. (2012). A comparison of item-level and scale-level multiple imputation for questionnaire batteries. *Multivariate Behavioral Research*, 47(1), 1–25.
- Graham, J. W. (2009). Missing data analysis: Making it work in the real world. *Annual Review of Psychology*, 60, 549–576.
- Graham, J. W., Olchowski, A. E., & Gilreath, T. D. (2007). How many imputations are really needed? Some practical clarifications of multiple imputation theory. *Prevention Science*, 8(3), 206–213.
- Grove, S.k., Buknes, N., Gray, J. (2013). The Practice of Nursing Research. Appraisal, Synthesis, and Generation of Evidence, 7thed, Elsevier, Saunders, St. Louis, Missouri 63043.
- Grimshaw, J., Thomas, R., MacLennan, G., Fraser, C., Ramsay, C. R., & Vale, L. (2004). Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technology Assessment*, 8(6), 1–72.
- Grol, R., & Grimshaw, J. (2003). From best evidence to best practice: effective implementation of change in patients' care. *The Lancet*, 362(9391), 1225–1230.
- Guildford, J. P. (1973). *Fundamental statistics in psychology and education*, 5th ed, New York: McGraw-Hill.
- Gulley, T., & Boggs, D. (2013). Timeperspective and Theory of Planned Behaviour:

 Moderate Predictors of Physical Activity Among Central
 AppalachianAdolessents, *Journal of Padiatric Health Care*, 28(5),41-47.
- Guiliano K, et al. (2000). Families first: liberal visitation policies may be in patients' best interests. *Nursing Management Journal*, 31(5), 46-50.
- Hardeman, W., Johnston, M., Johnston, D. W., Bonetti, D., Wareham, N. J. & Kinmonth, A. L. (2002). Application of the theory of planned behaviour in behaviour change interventions: A systematic review. *Psychology and Health*, 17(2), 123–158.
- Hagger, M. S., & Chatzisarantis, N. L. D. (2005). First-and higher-order models of attitudes, normative influence, and perceived behavioural control in the theory of planned behaviour. *British Journal of Social Psychology*, 44(4), 513–535.

- Hair, J. F., Black, W. C., Babin, B. J., Anderson, R. E., & Tatham, R. L. (2010). *Multivariate data analysis*, 7th edn. Englewood Cliffs (NJ): Prentice Hall.
- Hasan Tehrani T, Haghighi M, Bazmamoun H. (2012) Effects of stress on mothers of hospitalized children in a hospital in iran. *Iranian Journal of Child Neurology*, 6(4): 39-45.
- Hanson, S. M. H. (2005). *Family health care nursing: an introduction* (3th edn.). Philadelphia PA: F. A. Davis.
- Haidet, P., Dains, J.E., Paterniti, D.A., Chang, T., Tseng, E., Rogers, J.C. (2001). Medical students' attitudes toward patient-centered care and standardized patients' perceptions of humanism: a link between attitudes and outcomes. *Academic Mededicine Journal*, 76(Suppl 10):S42–S44.
- Harrison, T. M. (2010). Family-centered pediatric nursing care: State of the science. *Journal of Pediatric Nursing*, 25(5), 335–343.
- Harvey, L., Geall, V. and Moon, S. with Aston, J., Bowes, L. and Blackwell, A. (1998). Work Experience: Expanding opportunities for undergraduates. Centre for Research into Quality (CRQ), Birmingham: UK.
- Harvey, L. and Locke, W. with Morey, A., 2002, Enhancing employability, recognising diversity: Making links between higher education and the world of work. London, Universities UK.
- Helmer, S.D., Smith, R.S., Dort, J.M., Shapiro, W.M., Katan, B.S. (2000). Family presence during trauma resuscitation: a survey of AAST and ENA members. American Association for the Surgery of Truma. *Emergency Nursing Association. Journal Trauma*, 26,245-253.
- Henderson, D. P, Knapp, J. F. (2005). Report of the National Consensus Conference on Family Presence During Pediatric Cardiopulmonary Resuscitation and Procedures. *Pediatric Emergency Care Journal*, 21(11), 787-791
- Higman, W., & Shaw, K. (2008). Nurses' understanding about the delivery of family centred care in the neonatal unit. *Journal of Neonatal Nursing*, 14(6), 193–198.
- Hovey R, Cuthbertson K, Birnie K, et al. (2012). The influence of distress on knowledge transfer for men newly diagnosed with prostate cancer. *Journal of Cancer Education*, 27(3),540–545.
- Hockenberry, M. J., & Wilson, D. (2009). Wong's essentials of pediatric nursing. St. Louis: Mosby.
- Hovey R, Craig R. (2011). Understanding the relational aspects of learning with, from, and about the other. *Nursing Philosophy Journal*, 12(4), 262–270.

- Hovey R. (2012). Assessment in health promotion: deconstruction and metaphoric considerations in a nutshell. In: Gilmore GD, editor. *Needs and Capacity Assessment Strategies for Health Education and Health Promotion*, 4th ed. Sudbury, MA: Jones and Bartlett, LLC.
- Hopia, H., Tomlinson, P. S., Paavilainen, E., & \AAstedt-Kurki, P. (2005). Child in hospital: family experiences and expectations of how nurses can promote family health. *Journal of Clinical Nursing*, *14*(2), 212–222.
- Hrisos, S., Eccles, M. P., Francis, J. J., Dickinson, H. O., Kaner, E. F. S., Beyer, F., & Johnston, M. (2009). Are there valid proxy measures of clinical behaviour? a systematic. *Implementation Science*, 4(37), 1–20.
- Huang, J., Jiang, D., Wang, X., Liu, Y., Fennie, K., Burgess, J., & et al. (2002). Changing knowledge, behavior, and practice related to universal precautions among hospital nurses in China. *Journal of Continuing Education in Nursing*, 33, 217–24.
- Hudon, C., Fortin, M., Haggerty, J. L., Lambert, M., & Poitras, M.-E. (2011). Measuring patients' perceptions of patient-centered care: a systematic review of tools for family medicine. *The Annals of Family Medicine*, 9(2), 155–164.
- Hughes, M. (2007). Parents' and nurses' attitudes to family-centered care: an Irish perspective. *Journal of Clinical Nursing*, *16*, 2341–2348.
- Hutchfield, K. (1999). Family-centred care: a concept analysis. *Journal of Advanced Nursing*, 29(5), 1178–1187.
- Institute for Patient & Family Centered Care (IPFCC). (2010). what is patient- and family-centered care? Retrieved from http://www.ipfcc.org/faq.html.
- Iran Demographics Profile. (2014). Demographic of Iran.CIA World Factbook.
- Jaccard, J. J., & Wood, G. (1985). An idiothetic analysis of attitude-behavior models. *Advances in Consumer Research*, 13, 600–605.
- James, G., Witten, D., Hastie, T., & Tibshirani, R. (2013). *An introduction to statistical learning*. New York, NY: Springer.
- Javadi, M., Kadkhodaee, M., Yaghoubi, M., Maroufi, M., & Shams, A. (2013). Applying Theory of Planned Behavior in Predicting of Patient Safety Behaviors of Nurses. *Materia Socio-Medica*, 25(1), 52.
- Jillian, J., Martin, P., Johnston, M., Walker, A., Grimshaw, J., Foy, J., Smith, L., Bonetti, D. (2004). Constructing questionnaires based on the theory of planned behavior: a manual for health services researchers. Newcastle: Centre for Health Services Research, University of Newcastle, United Kingdom.
- Johnson, B., Abraham, M., Conway, J., Simmons, L., Edgman-Levitan, S., Sodomka, P., Ford, D. (2008). Partnering with patients and families to design a patient-

- and family-centered health care system. Bethesda MD: Institute for Family-Centered Care. Retrieved from http://global.factiva.com/sb/default.aspx.
- Kam, L. Y. K., Knott, V. E., Wilson and Suzannek. (2012). Using Theory of Planned Behaviour to understand health professionals' attitudes and intentions to refer cancer patients for psychological support. *Psycho-Oncology*, 21, 316-323.
- Khosravan, S., Mazlom, B., Abdollahzade, N., Jamali, J., & Mansoorian, M. H. (2014). Family Participation in the Nursing Care of the Hospitalized Patients. *Iran Red Crescent Medical Journal*, 16(1):e12686.1-6.
- Kirk, S., Glendinning, C., & Callery, P. (2005). Parent or nurse? The experience of being the parent of a technology-dependent child. *Journal of Advanced Nursing*, 51(5), 456–464.
- Koop, C. E. (1987). Surgeon General's report: Children with special healthcare needs campaign '87. Washington, DC: Department of Health and Human Services.
- Kortteisto, T., Kaila, M., Komulainen, J., Mäntyranta, T., & Rissanen, P. (2010). Health care professionals' intentions to use clinical guidelines: A survey using the theory of planned behaviour. *Implementation Science*, *5*(51), 1–10.
- Kuo, D. Z., Houtrow, A. J., Arango, P., Kuhlthau, K. A., Simmons, J. M., Neff, J. M. (2012). Family-Centered Care: Current Applications and Future Directions in Pediatric Health Care. Matern Child Health J, 16(2),297-305.
- Kuo, D. Z., Frick, K. D., Minkovitz, C. S. (2011). Association of family centered-care with improved anticipatory guidance delivery and reduced unmet needs in child health care. *Journal of Maternal Child Health*, 15, 1228-1237.
- Kumar, U. (2013). Authorized dealer for 3dist....at arihant industrial corporation Itd. Retrieved from www.slideshare.net.
- Kumar, U. (2015). Role of pediatric nurse in child care. Education. Manipal Intl University. Linkedin Corporation@2015.
- Lam, D., Wong, S. N., Hui, H., Lee, W., & So, K. (2007). Attitude of doctors and nurses to family presence during pediatric cardiopulmonary resuscitation. *Journal of Pediatric*, 12, 253–259.
- Lavarakas, P. J. (2008). Encyclopedia of Survey Research Methods. Thousand Oaks, CA: SAGE Publications, Inc. Retrieved from: http://dx.doi.org/10.4135/9781412963947
- Latour, J. M., Hazelzet, J.A., Hugo, J. Duivenvoorden, J. B. van Goudoever. (2010). Perceptions of parents, nurses, and physicians on neonatal intensive care practices. *Journal of Pediatrics*, 157(2), 215-220
- Lanchinger, H.S., Goldenberg, D. (1993). Attitudee of practicing nurses as predictors of intended care behavior with persins who are HIV positive: testing the

- Ajzen-Fishbein theory of reasond action. *Research in Nursing and Health*, 16, 441-450.
- Leech, N. L., Barrett, K. C., & Morgan, G. A. (2008). SPSS for intermediate statistics: Use and interpretation, 3rd ed. New York, NY: Psychology Press.
- Leeman, J., Baernholdt, M., & Sandelowski, M. (2007). Developing a theory-based taxonomy of methods for implementing change in practice. *Journal of Advanced Nursing*, 58(2), 191–200.
- Lee, P. (2004). Family involvement: are we asking too much? *Pediatric Nursing*, 16(10), 37–41.
- Lehto, P. (2004). Shared Being With The Substantive Theory Of Parents' Participation In The Care Of Their Own Child In The Hospita (Academic dissertation). Acta Universitatis Tamperensis, Tampere.
- Lemanske RF, Nayak A, Mcalary M, Everhard F, Flower- Taylor A, & Gupta N. (2011). Omalizumab improves Asthma- Releated Quality of life in children with Allergic Asthma. *Journal of Peadiatric*, 110(5).
- Lemeshow, S. (1990). *Adequacy of sample size in health studies*. New York, NY: Wiley on behalf of the World Health Organization.
- Lopez, V., Molassiotis, A., Chan, W. K., Ng, F., & Wong, E. (2004). An intervention study to evaluate nursing management of peripheral intravascular devices. *Journal of Infusion Nursing*, 27(5), 322–331.
- Lu, H.Y., Hou, H.Y., Dzwo, T.H., Wu, Y.C., Andrews, J. E., Weng, S.T., & Lu, J.Y. (2010). Factors influencing intentions to take precautions to avoid consuming food containing dairy products: Expanding the theory of planned behaviour. *British Food Journal*, 112(9), 919–933.
- MacKay, L. J., & Gregory, D. (2011). Exploring family-centered care among pediatric oncology nurses. *Journal of Pediatric Oncology Nursing*, 28(1), 43–52.
- MacKean, G. L., Thurston, W. E., & Scott, C. M. (2005). Bridging the divide between families and health professionals' perspectives on family-centred care. *Health Expectations*, 8(1), 74–85.
- Macdonald, M. E., Liben, S. & Carnevale, F. A. (2012). An office or a bedroom, challenges for family-centered care in the pediatric intensive care unit. *Journal of Child Health Care*, 16(3), 237-249.
- Maijala, H., & Åstedt-Kurki, P. (2009). From substantive theory towards a family nursing scale, *16*(3), 29–44.
- Matthew, B., & et al. (2007). Applying theory –driven approaches to understanding and modifying clinicians, behavior: What do we know? *Psychiatric Service*, 58, 342–348.

- Matthews, T. L., Hempel, L. M., & Howell, F. M. (2010). Gender and the Transmission of Civic Engagement: Assessing the Influences on Youth Civic Activity. *Sociological Inquiry*, 80(3), 448–474.
- Mann, K., Gordon, J., MacLeod, A. (2009). Reflection and reflective practice in health professions education: a systematic review. *Advances in Health Sciences Education Theory Practice Journal*, 14(4): 595–621.
- McCarthy, M. J., Herbert, R., Brimacombe, M., Hansen, J., Wong, D., & Zelman, M. (2002). Empowering parents through asthma education. *Pediatric Nursing*, 28(5), 465–474.
- McGahey, P. R. (2002). Family presence during pediatric resuscitation: a focus on staff. *Critical Care Nurse*, 22, 29-36.
- McGavin, C. (2013). Colleen's story: Reflection on the Concept of "Patient and Family Centered Care". *Journal of Family Nursing*, 19(4), 418-430.
- McWilliam, C. L. (2007). Continuing education at the cutting edge: promoting transformative knowledge translation. *Journal of Continuing Education in the Health Professions*, 27(2):72–79.
- Merriam, S. B., Caffarella, R.S., Baumgartner, L.M. (2007). *Learning in Adulthood: A Comprehensive Guide*, 3rd ed. San Francisco: John Wiley & Son.
- Mezirow, J. (1991). *Transformative Dimensions of Adult Learning*. San Francisco, CA: Jossey-Bass, Inc, 33.
- Mesibov, G. B., & Shea, V. (2010). Evidence-based practices and autism. *Autism*, 15(1), 114–33.
- Meyers, T. A., Eichhorn, D. J., Guzzetta, C. E., Clark, A. P., Klein, J. D., Taliaferro, E., & Calvin, A. (2000). Family presence during invasive procedures and resuscitation: the experience of family members, nurses, and physicians. *AJN The American Journal of Nursing*, 100(2), 32–43.
- Meyers, T. A., Eichhorn, D. J., Guzzetta, C. E., Clark, A. P., Taliaferro, E. (2004). Family presence during invasive procedures and resuscitation. *Topics in Emergency Medicine*, 26, 61-73.
- Miceli, P. J., & Clark, P. A. (2005). Your patient-My child: Seven priorities for improving pediatric care from the parent's perspective. *Journal of Nursing Care Quality*, 20(1), 43–53.
- Mitchell, M., Chaboyer, W., Burmeister, E., Foster, M. (2009). Positive Effect of a Nursing Intervention on Family-Centered Care in Adult Critical Care. *American Journal of Critical Care*, 18(6), 543-552.

- Molazem, Z., Ahmadi, F., Mohammadi, E., Bolandparvaz, S. (2011). Improvement in the nursing care quality in general surgery wards: Iranian nurses' perceptions. *Scandinavian Journal of Caring Sciences*, 25(2), 350–6.
- Murray, C. A. (1980). "Research: Time of Day Preferences and Achievement." *Human Esources*. N.p., 2004. Web. 6 Feb. 2010. Retrieved from http://learn.humanesources.com/research_24.html.
- Moser, R. P., Green, V., Weber, D., & Doyle, C. (2005). Psychosocial correlates of fruit and vegetable consumption among African American men. *Journal of Nutrition Education and Behavior*, *37*(6), 306–314.
- Nadler & Leonard. (1984). *The Handbook of Human Resource Development*. New York: John Wiley & Sons.
- Nash, R., Edwards, H., & Nebauer, M. (1993). Effect of attitudes, subjective norms and perceived control on nurses' intention to assess patients' pain. *Journal of Advanced Nursing*, 18(6), 941–947.
- Negarandeh, R., Oskouie, F., Ahmadi, F., Nikravesh, M., & Hallberg, I. R. (2006). Patient advocacy: barriers and facilitators. *BMC Nursing*, *5*(1), 3.
- Notani, A. S. (1998). Moderators of perceived behavioral control's productiveness in the theory of planned behavior: a meta-analysis. *Journal of Consumer Psychology*, 7, 247–271.
- Nunnally, J. C. (1978). Psychometric Theory. The University of Michigan McGraw-Hill Book Company.
- O'Boyle, C. A., Henly, S. J., & Duckett, L. J. (2001). Nurses' motivation to wash their hands: a standardized measurement approach. *Applied Nursing Research*, 14(3), 136–145.
- Paliadelis, P., Cruikshank, M., Wainohu, D., Winskill, R., & Stevens, H. (2005). Implementing family-centered care: an exploration of the beliefs and practices of pediatric nurses. *Australian Journal of Advanced Nursing*, 23(1), 31–36.
- Payant, L., Davies, B., Graham, I. D., Peterson, W. E., & Clinch, J. (2008). Nurses' intentions to provide continuous labor support to women. Journal of Obstetrics, *Gynecology and Neonatal Nursing*, 37(4), 405-414.
- Pallant, J. (2005). SPSS survival guide. Maidenhead, England: Open University Press.
- Pallant, J. (2010). Journey in Survey Research, SPSS Survival Manual: A step by Step Guide to Data Analysis Using SPSS for windows, 4th ed. Open University Press; London.
- Pasyar, N., & Gholamzadeh, S. (2009). Effect of Education on Nurses' Knowledge and Performance Regarding AIDS in Emergency Departments of Hospitals

- Affliated to Shiraz University of Medical Scienes (SUMS). *Iranian Journal of Nursing Research*, 4(12-13), 81–90.
- Perkins, M., Jensen, P., Jaccard, J., Gollwitzer, P., Oettingen, G., Pappadopulos, E., & Hoagwood, K. (2007). Applying theory-driven approaches to understanding and modifying clinicians' behavior: What do we know? *Psychiatric Services*, 58(3), 342–348.
- Pegon-Machat, E., Tubert-Jeannin, S., Loignon, C., Landry, A., Bedos, C. (2009). Dentists' experiences with low-income patients benefiting from a public insurance program. *European Journal of Oral Sciences*, 117(4),398–406.
- Peterson, M. F., & et al. (2006). Family-centered care: Do we practice what we preach? Journal of Obstetrics, Gynecology & Neonate Nursing, 33(4), 421–427.
- Pettoello-Mantovani, M., Campanozzi, A., Maiuri, L., & Giardino, I. (2009). Family-oriented and family-centered care in pediatrics. *Italian Journal of Pediatrics*, 35(12), 1–8.
- Phillipi J. (2010). Transformative learning in healthcare. *PAACE Journal of Lifelong Learning*, 19:39–54.
- Polit, D. F., & Beck, C. T. (2004). *Nursing research: Principles and methods*, 7th ed. Philadelphia, New York, Baltimore.: Lippincott Williams & Wilkins.
- Pong, A., & Chow, S.-C. (Eds.). (2010). Handbook of adaptive designs in pharmaceutical and clinical development. CRC Press.
- Potter, P. A., & Perry, A. C. (2009). Fundamental of Nursing, 7th ed. St. Louis, MO: Mosby Co.
- Powers, K. S, Rubenstein, J.S. (1999). Family presence during invasive procedures in the pediatric intensive care unit. *Archives of Pediatric and Adolescent Medicine Journal*, 153, 955-958.
- Puffer, S., & Rashidian, A. (2004). Practice nurses' intentions to use clinical guidelines. *Journal of Advanced Nursing*, 47(5), 500–509.
- Ramezani1, T., Hadian Shirazi1, Z., Sabet Sarvestani2, R., & Moattari, M. (2014). Family-Centered Care in Neonatal Intensive Care Unit: A Concept Analysis. *International Journal of Community Based Nursing and Midwifery*, 2(4), 268-278.
- Redley, B, Hood. K. (1996). Staff attitudes towards family presence during resuscitation. *Accident and Emergency Nursing*, 4, 145-151.
- Rivis, A., & Sheeran, P. (2003). Descriptive norms as an additional predictor in the theory of planned behaviour: A meta-analysis. *Current Psychology*, 22(3), 218–233.

- Robinson, S. M., Mackenzie-Ross, S, Hewson, G. L, Campbell, Egleston, C. V, Prevost, A. T. (1998). Psychological effect of witnessed resuscitation on bereaved relatives. *Lancet*, 352, 614-617.
- Rubin, D. B. (1987). Multiple imputation for nonresponse in surveys. New York: Wiley & Sons.
- Rubin, D. B. (1996). Multiple imputation after 18+ years. *Journal of the American Statistical Association*, *91*(434), 473–489.
- Sacchetti, A, Carraccio, C, Leva, E, Harris, R. H, Lichenstein, R. (2000). Acceptance of family member presence during pediatric resuscitations in the emergency department: effects of personal experience. *Pediatric Emergency Care*, 16, 85-87.
- Saunders, R. P., Abraham, M. R., Crosby, M. J., Thomas, K., & Edwards, W. H. (2003). Evaluation and development of potentially better practices for improving family-centered care in neonatal intensive care units. *Pediatrics*, 111(4), 437–449.
- Shoham, S., & Gonen, A. (2008). Intentions of hospital nurses to work with computers. *CIN: Computers, Informatics, Nursing*, 26(2), 106-115.
- Schafer, J. L. (1997). *Analysis of incomplete multivariate data*. London, England: Chapman & Hall.
- Schafer, J. L. (1999). Multiple imputation: a primer. Statistical Methods in Medical Research, 8(1), 3–15.
- Schafer, J. L., & Graham, J. W. (2002). Missing data: Our view of the state of the art. *Psychological Methods*, 7(2), 147–177.
- Schlomer, G. L., Bauman, S., & Card, N. A. (2010). Best practices for missing data management in counseling psychology. *Journal of Counseling Psychology*, 57(1), 1–10.
- Seccombe, J. A. (2007). Attitudes towards disability in an undergraduate nursing curriculum: the effects of a curriculum change. *Nurse Educ Today*, 27, 445-451.
- Semin-Goossens, A., van der Helm, J. M. J., & Bossuyt, P. M. M. (2003). A Failed Model-Based Attempt to Implement an Evidence-Based Nursing Guideline for Fall Prevention. *Journal of Nursing Care Quality*, *18*(3), 217–225.
- Segal, J., Melinda Smith, M. A., Boose, G., and Jaffe, J. (2014). Guide for Improving your Mental and emotional Health. Nonverbal Communication, Improving Your Nonverbal Skills and Reading Body Language. HELP GUIDE. ORG. E MOTIONAL Intelligence Toolkit.

- Segaric, C.A. & Hall, W.A. (2015). Progressively Engaging: Constructing Nurse, Patient, and Family Relationships in Acute Care Settings. *Journal of Family Nursing*, 21(1) 35-56.
- Shelton, T. (1999). Family-centered care in pediatric practice: When and how? *Development and Behavioral Pediatrics*, 20(2), 117–119.
- Shelton, T., Jepson, E., & Johnson, B. H. (1989). Family-Centered Care for Children with Special Health Care Needs. Washington DC: Association for the Care of Children's Health.
- Sheppard, B. H., Hartwick, J., & Warshaw, P. R. (1988). The theory of reasoned action: A meta-analysis of past research with recommendations for modifications and future research. *Journal of Consumer Research*, 15(3), 325–343.
- Shields, L., & King, S. (2001). Qualitative analysis of the care of children in hospital in four countries—Part 2. *Journal of Pediatric Nursing*, 16(3), 206–213.
- Shields, L., & Nixon, J. (2004). Hospital care of children in four countries. *Journal of Advanced Nursing*, 45(5), 475–486.
- Shields, L., Pratt, J., Davis, L. M., & Hunter, J. (2007). Family-centred care for children in hospital. *Cochrane Database of Systematic Review*, (1): CD004811 (ISSN: 1469-493X). University of Hull, Faculty of Health and Social Care, Cottingham Rd, Hull, UK, HU6 7RX. L. Shields@hull.ac.uk.
- Shields, L., & Tanner, A. (2004). Pilot study of a tool to investigate perceptions of family-centered care in different care settings. *Pediatric Nursing*, 30(3), 189.
- Smith-Idell, C., Grant, M., & Kirk, C. (2007). Alignment of pain reassessment practices and National Comprehensive Cancer Network Guidelines. *Oncology Nursing Forum*, *34*(3), 661–671.
- Smith, L., & et al. (2002). *Family-Centered Care: Concept, Theory and practice*. New York: Basingstoke.
- Soto-Crespo, R. E. (1999). The bounds of hope: Unlearning "Old Eyes" and a pedagogy of renewal. *Thresholds in Education*, 25(23), 42–46.
- Starkweather, J., & Herrington, R. (2012). *Module 6: Multiple Imputation Short Course.* Retrieved from http://www.unt.edu/rss/class/Jon/SPSS_SC/Module6/SPSS_M6_2.htm
- Streiner, D., & Norman, G. (2008). *Biostatistics: The bare essentials*. London, England: McGraw-Hill.
- Summers, J. A., & Turnbull, A. (2010). Characteristics and Trend in Family-Centered Care Conceptualizations. *Journal of Family Social Work*, *13*(3), 269–285.

- Smith, J., Kirksey, K., Becker, H., & Brown, A. (2011). Autonomy and self-efficacy as influencing factors in nurses' behavioral intention to disinfect needleless intravenous systems. *Journal of Infusion Nursing*, 34(3), 193-200.
- Tabachnick, B. G., & Fidell, L. S. (2007). *Using Multivariate Statistics: Pearson Education Inc.* Boston, MA: Allyn and Bacon.
- Taylor. E. W. (2007). An update of transformative learning theory: a critical review of the empirical research (1999–2005). *International Journal of Lifelong Education*, 26(2). 173–191.
- Thompson, D. S., Estabrooks, C. A., Scott-Findlay, S., Moore, K., & Wallin, L. (2007). Interventions aimed at increasing research use in nursing: a systematic review. *Implementation Science*, 2(1), 15.
- Topa, G., & Moriano, J. A. (2010). Theory of planned behavior and smoking: Metaanalysis and SEM model. *Substance Abuse and Rehabilitation*, 1, 23–33.
- Udomratn, P., & Srisurapanont, M. (2002). Impact on Thai psychiatrists of passive dissemination of a clinical practice guideline on prescribing attitudes in treatment-resistant schizophrenia. *Neuropsychobiology*, 45(4), 186–190.
- United Nations World. (2014). Population Data Via United Nations. Aguid to population information on UN system web site, Department of Economic and Social Affairs.
- UNICEF. (2012). Levels and Trends in Child Mortality. New York: United Nations Children's. Educational Fund.
- Yousefi, H., Karami, A., Moeini M., Ganji H. (2012). Effectiveness of nursing interventions based on family needs on family satisfaction in the neurosurgery intensivecare unit. *Iranian Journal of Nursing and Midwifery Research*, 17(4), 296-300.
- Valizadeh, F., & Ghasemi, S. F. (2008). Medical Staff Attitude toward Parents' Participation in the Care of their Hospitalized Children. *Hayat*, *14*(1), 69–76.
- Wakefield, J. G., McLaws, M.-L., Whitby, M., & Patton, L. (2010). Patient safety culture: factors that influence clinician involvement in patient safety behaviours. *Quality and Safety in Health Care*, 19(6), 585–591.
- Waltz, C. F, Bausell, R. B. (1983). *Nursing research: Design, StatisticsandComputer Analysis*. 2nd edn, Philadelphia: FA Davis Company.
- Warshaw, P. R., & Davis, F. D. (1985). Disentangling behavioral intention and behavioral expectation. *Journal of Experimental Social Psychology*, 21(3), 213–228.

- Weiner, B. J., Amick, H., & Shou-Yih, D. L. (2008). Conceptualization and measurement of organizational readiness for change. *Medical Care Research and Review*, 65(4), 379–436.
- Wells, N. (2011). Historical perspective on family-centered care. *Academic Pediatrics*, 11, 100-102.
- Yaghmaie, F. (2003). Content validity and its estimation. *Journal of Medical Education*, 3(1), 25-27.
- Ygge, B. M., Lindholm, C., & Arnetz, J. (2006). Hospital staff perceptions of parental involvement in paediatric hospital care. *Journal of Advanced Nursing*, *53*(5), 534–542.
- Young, J., McCann, D., Watson, K. M., Pitcher, A., Bundy, R., & Greathead, D. (2006). Negotiation of care for a hospitalised child: nursing perspectives. *Journal of Neonatal, Paediatric and Child Health Nursing*, 9(3), 7–14.
- Zarei, S., Taghdisi, M. H., Solhi, M., Tehrani, H., & Shojayizade, D. (2013). The Effect of Educational Intervention Based on BASNEF Model to Improve Interpersonal Communication Skills of Nurses. *Journal of Alborz Health*, *1*(3), 173–178.
- Zamanzadeh, V., Rassouli, M., Abbaszadeh, A., Nikanfar, A., Alavi-Majd, H., & Ghahramanian, A. (2014). Factors Influencing communication between the patients with cancer and their nurses in oncology wards. *Indian Journal of Palliative Care*, 20(1), 12-20.