



UNIVERSITI PUTRA MALAYSIA

***EXTRACTION, CHARACTERISATION, OPTIMISATION AND
MODELING OF PHALERIN FROM PHALERIA MACROCARPA
(SCHEFF.) BOERL LEAVES***

NOR FARIZA ISMAIL

FK 2014 168



**EXTRACTION, CHARACTERISATION, OPTIMISATION AND
MODELING OF PHALERIN FROM *PHALERIA MACROCARPA*
(SCHEFF.) BOERL LEAVES**

By

NOR FARIZA ISMAIL

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy**

November 2014

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



*This thesis is specially dedicated to
my beloved husband Hafidz,
my wonderful kids Haikal, Harith and Haziq Irfan,
my parents
and
my in laws.*

Abstract of thesis presented to the Senate of Universiti of Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**EXTRACTION, CHARACTERISATION, OPTIMISATION AND
MODELING OF PHALERIN FROM *PHALERIA MACROCARPA*
(SCHEFF.) BOERL LEAVES**

By

NOR FARIZA ISMAIL

November 2014

Chairman : Professor Luqman Chuah Abdullah, PhD.
Faculty : Engineering

Phaleria macrocarpa (Scheff.) Boerl is a medicinal plant found in Malaysia and Indonesia. This plant has been used traditionally to treat various diseases. Little studies have been done to identify compound that contributes to medicinal effect. Moreover, there is lack of studies in the method to extract those valuable components from this plant.

This research focuses on the leaves part of the plant. A single compound known as phalerin is targeted and extracted from the leaves. The compound is purified and identified through chemical analysis such as High Performance Liquid Chromatography, Thin Layer Chromatography, Nuclear Magnetic Resonance and Liquid Chromatogram Mass Spectroscopy. Phalerin was tested for the bioactivity study including antioxidant, anti-inflammatory and anticancer. Then the extraction process using solid-liquid extraction method was investigated on the optimum parameters; solvent, extraction temperature, solid-to-solvent ratio and particle size.

Phalerin was proven having antioxidant and anti-inflammatory activity. Water was the most suitable solvent for extraction of phalerin from the leaves compared to methanol, ethanol, ethyl acetate and hexane. Water was not only safe but also extracted most of polar compounds. The optimum temperature was at 70°C as to avoid degradation of compound. The optimum solid to solvent ratio was 1:20 (g/ml) as too much solvent leads to waste of solvent and extra energy to remove excess liquid. The smallest particle size (<250 µm) was the optimum size as smaller particle provide larger surface area for mass transfer.

Extraction kinetics on extraction of phalerin found that the process reach equilibrium at 120 min of extraction duration. The first order kinetics model was applied and successfully described the extraction process. From thermodynamic study, the enthalpy change (ΔH) was found to be positive (72.51 kJ/mol), indicating the extraction process of *Phaleria macrocarpa* leaves is endothermic. The Gibb's free energy was found to be negative with values range from -2.75 to -9.95 kJ/mol, indicated the extraction was spontaneous at the operating temperature. This spontaneity was favored with increasing extraction temperature.

Phalerin is a bioactive compound extracted from *Phaleria macrocarpa* leaves and having antioxidant and anti-inflammatory activities. Standardization of extract and improvement on the extraction process along with clinical study of phalerin may added new information to this study.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENGKESKTRAKAN, PENCIRIAN, PENGOPTIMUMAN DAN
PEMODELAN PHALERIN DARI DAUN *PHALERIA MACROCARPA*
(SCHEFF.) BOERL**

Oleh

NOR FARIZA ISMAIL

November 2014

Pengerusi : Profesor Luqman Chuah Abdullah, PhD.
Fakulti : Kejuruteraan

Phaleria macrocarpa (Scheff.) Boerl adalah tanaman bernilai yang boleh didapati di Malaysia dan Indonesia. Tumbuhan ini telah digunakan secara tradisional untuk merawat pelbagai penyakit. Hanya terdapat sedikit kajian yang telah dijalankan untuk mengenalpasti kandungan bahan yang menyumbang kepada kesan perubatan. Tambahan lagi, masih kurang penyelidikan terhadap kaedah untuk mengekstrak kandungan bahan yang bermanfaat dari tumbuhan ini.

Kajian ini tertumpu kepada bahagian daun pada tumbuhan. Satu kandungan bahan yang dikenali sebagai phalerin telah disasarkan dan diekstrak daripada daun. Kandungan bahan ini telah dikenalpasti melalui analisis kimia seperti *High Performance Liquid Chromatography*, *Thin Layer Chromatography*, *Nuclear Magnetic Resonance* dan *Liquid Chromatogram Mass Spectroscopy*. Phalerin telah dikaji untuk tindakan biologi termasuk antioksidan, anti radang dan anti kanser. Kemudian, proses pengekstrakan menggunakan kaedah pengekstrakan pepejal-cecair telah dikaji terhadap parameter yang optimum iaitu pelarut, suhu pengekstrakan, nisbah pepejal-cecair dan saiz bahan.

Phalerin telah disahkan mengandungi antioksidan dan anti radang efek. Air dikenalpasti sebagai pelarut yang paling sesuai untuk mengekstrak phalerin daripada daun berbanding dengan metanol, etanol, etil asetat dan heksana. Air bukan sahaja selamat tetapi mampu mengekstrak bahan-bahan bersifat polar. Suhu optimum adalah pada 70°C untuk mengelakkan degradasi bahan. Nisbah pepejal kepada cecair yang optimum adalah 1:20 (g/ml) kerana bahan pelarut yang banyak boleh menyebabkan pembaziran dan juga memerlukan tenaga yang lebih untuk diasingkan. Saiz bahan yang paling kecil iaitu <250µm adalah yang optimum kerana saiz yang lebih kecil memberi luas permukaan yang lebih besar untuk process pertukaran bahan.

Pengekstrakan phalerin melalui kajian kinetik pengekstrakan mendapati proses ini mencapai *equilibrium* pada 120 min. Model *first order kinetics* telah diaplikasi dan berjaya menerangkan proses pengekstrakan. Daripada pengkajian termodinamik, perubahan entalpi didapati positif dengan nilai 72.51 kJ/mol dan ini menerangkan proses pengekstrakan daun *Phaleria macrocarpa* adalah endotermik. *Gibb's free*

energy didapati negatif dengan nilai dari -2.75 hingga -9.95 kJ/mol yang menerangkan proses kinetik berlaku secara spontan pada suhu yang dijalankan. Proses yang spontan ini berlaku berkadaran dengan peningkatan suhu.

Phalerin merupakan compound aktif yang diekstrak dari daun *Phaleria macrocarpa* dan mengandungi aktiviti anti oksida dan anti radang. Penyeragaman ekstrak dan menambahbaikkan dalam proses pengekstrakan serta kajian perubatan terhadap phalerin boleh memberi maklumat baru dalam kajian ini.



ACKNOWLEDGEMENT

First of all I would like to thank my advisor and chairman of the supervisory committee, Professor Dr. Luqman Chuah Abdullah, for his advice and guidance in making this research successful. I also would like to thank my supervisory committee members, Dr Pin Kar Yong, Professor Dr. Umi Kalsom Yusuf and Dr Dayang Radiah Awang Biak for giving advice, ideas and comments throughout the project.

I also would like to thank Dr Zunoliza Abdullah, Dr Fadzureena, Hj Rashid, Mr Rashih Li, Mr Farhan Razak, Nur Hazwani and Masitah from Forest Research Institute of Malaysia for helping me in guiding, sharing knowledge and their technical assistance. Special thank you is dedicated to Dr Latifah and Armania from Medical School of UPM for technical guidance and sharing knowledge.

I would like to acknowledge my wonderful family; husband Mohd Hafidz Mohamad Zain for his patient and support, my kids Haikal, Harith and Haziq Irfan who always stay by my side and fill my heart with love, my parents for their prayers and support, my siblings who always be my courage and in laws who steadfast in me pursuing this project.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement of the degree of Doctor of Philosophy. The members of the Supervisor Committee are as follows:

Luqman Chuah Abdullah, PhD

Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Dayang Radiah Awang Biak, PhD

Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Umi Kalsom Yusuf, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Pin Kar Yong, PhD

Forest Research Institute of Malaysia (FRIM)
Selangor
Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: _____

Declaration by Members of Supervisory Committee

That is to confirm that:

- the research conducted and the writing of the this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduated Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of Chairman
of Supervisory
Committee: Prof. Dr. Luqman Chuah Abdullah

Signature: _____
Name of Member of
Supervisory
Committee: Dr. Dayang Radiah Awang Biak

Signature: _____
Name of Member of
Supervisory
Committee: Prof. Dr. Umi Kalsom Yusuf

Signature: _____
Name of Member of
Supervisory
Committee: Dr. Pin Kar Yong

TABLE OF CONTENTS

	ABSTRACT	Page
	ABSTRAK	i
	ACKNOWLEDGEMENT	iii
	APPROVAL	v
	DECLARATION	vi
	LIST OF TABLES	viii
	LIST OF FIGURES	xiii
	LIST OF ABBREVIATIONS	xiv
		xvi
	 CHAPTER	
1	INTRODUCTION	1
	1.1 Background	1
	1.2 <i>Phaleria macrocarpa</i> (Scheff.) Boerl	3
	1.3 Problem statement	3
	1.4 Objectives of the study	4
	1.5 Scope of the study	5
2	LITERATURE REVIEW	6
	2.1 <i>Phaleria macrocarpa</i> (Scheff.) Boerl	6
	2.1.1 Traditional uses of <i>Phaleria macrocarpa</i>	6
	2.1.2 Chemical constituents and structure	7
	2.1.3 Chemical properties/ Bioactivity of <i>Phaleria macrocarpa</i>	9
	2.1.4 Bioactivity	10
	2.2 Extraction	13
	2.2.1 Soxhlet extraction	14
	2.2.2 Reflux	15
	2.2.3 Maceration	15
	2.2.4 Supercritical fluid extraction (SFE)	15
	2.2.5 Subcritical water extraction	15
	2.2.6 Isolation, separation and purification of compound	16
	2.3 Solid-liquid extraction	17
	2.3.1 Parameters of solid-liquid extraction	17
	2.4 Mathematical models for solid liquid extraction	20
	2.4.1 Mass transfer model	21
	2.4.2 Diffusion model	22
	2.5 Chapter summary	25
3	METHODOLOGY	26
	3.1 Introduction	26
	3.2 Plant material	27
	3.3 Extraction, isolation and characterisation	27
	3.3.1 Fractionation and characterisation	27
	3.4 Bioactivity screening	30

3.4.1	<i>In-vitro</i> antioxidant assays	30
3.4.2	<i>In-vitro</i> anti-inflammatory assays	32
3.4.3	<i>In-vitro</i> anticancer assays	34
3.5	Optimisation of solid-liquid extraction	35
3.5.1	Experiments for the effect of extraction parameters on the extract yield	35
3.5.2	Experiments for the effect of extraction parameters on the extract quality	37
3.6	Mathematical modeling	37
3.6.1	Physical properties of sample	37
3.7	Extraction kinetics	38
3.8	Mathematical modeling of solid-liquid extraction	39
3.8.1	Diffusion dependent model	39
3.8.2	First order kinetic model	40
3.8.3	Second order kinetic model	41
3.8.4	Thermodynamics	42
3.9	Statistical analysis	43
4	RESULTS AND DISCUSSION	44
4.1	Extraction and purification of phalerin	44
4.2	Fractionation and identification of the compound	46
4.3	Bioactivity	50
4.3.1	Antioxidant activity	50
4.3.2	Anti-inflammatory activity	51
4.3.3	Anticancer activity	52
4.4	Quantitative and qualitative effect of parameter on solid liquid extraction	55
4.4.1	Effect of solvents on extraction yield and concentration of phalerin	55
4.4.2	Effects of solid to liquid ratio on extraction yield and concentration of phalerin	56
4.4.3	Effects of particle size on extraction yield and concentration of phalerin	58
4.4.4	Effects of temperature on extraction yield and concentration of phalerin	60
4.5	Extraction kinetics	61
4.5.1	First order kinetic model	61
4.5.2	Second order kinetic model	64
4.5.3	Diffusion dependent model	65
4.6	Thermodynamics	67
5	CONCLUSION	69
5.1	Conclusion	69
5.2	Future studies	68
	REFERENCES	71
	APPENDICES	87
A	Physical properties of <i>Phaleria macrocarpa</i> leaves (Result from Malvern Mastersizer)	87
B	Density determined using pycnometer	88

BIODATA OF STUDENT	89
LIST OF PUBLICATIONS	90



LIST OF TABLES

Table		Page
2.1	Classification of <i>Phaleria macrocarpa</i>	6
2.2	Effective diffusion coefficients of food solutes in diverse matrices	24
3.1	Mobile phase used in the operation of HPLC	29
3.2	Properties of solvents	35
3.3	Experiments setup for studying the effect of extraction parameters	36
4.1	Chemical shift of the ^{13}C carbon and ^1H proton of phalerin (dissolved in deuterated methanol, CD_3OD)	47
4.2	DPPH radical and ABTS cation radical scavenging activities of methanol/water fraction and phalerin compound	51
4.3	Inhibitory effects of phalerin from <i>Phaleria macrocarpa</i> extract on anti-inflammatory assays	51
4.4	IC_{50} ($\mu\text{g/ml}$) values obtained from the MTT assays with the tested cancer and normal cell lines	53
4.5	Extraction of phaleria macrocarpa equation parameters for first order kinetic model	63
4.6	Model parameters for second order kinetic model.	64
4.7	Model parameter for the diffusion mode	66
4.8	Summaries of constant that influence the Biot number with respect to the operating temperature.	67
4.9	Thermodynamics activation parameters for phalerin extraction from <i>Phaleria macrocarpa</i> leaves at different temperature	67
4.10	Equilibrium constant the thermodynamics parameters	68

LIST OF FIGURES

Figure	Page
1.1 Herbal processing flow diagram	2
1.2 <i>Phaleria macrocarpa</i>	3
2.1 Chemical structure of phalerin (Adapted from Mae et al., 2005)	8
2.2 Chemical structure of macronone (Adapted from Susilawati et al., 2012)	8
2.3 Chemical structure of mangiferin (Adapted from Kim et al., 2010)	9
2.4 Phytoconstituents isolated from <i>Phaleria macrocarpa</i> with their respective biological activities (Adapted from Altaf et al., 2013).	13
2.5 Soxhlet apparatus	14
2.6 Typical extraction curve of batch solvent extraction of active compounds from plants (Adapted from Chan et al., 2013)	21
3.1 Overview of the methodology	26
3.2 Calibration curve of phalerin	30
3.3 Schematic diagram of a) extraction process b) solid particle	38
4.1 Phalerin structure	44
4.2 Thin layer chromatogram of a) Dichloromethane extract, b) Water/methanol extract, c) Fraction F.6 and d) Fraction F.6.6 (identified as phalerin)	45
4.3 Positive ESI-ms spectrum of phalerin. The ion at m/z 445.25 represents the sodium adduct of that ion $[M+Na]^+$	45
4.4 HPLC Chromatogram of a) phalerin b) crude extract	46
4.5 ^1H -NMR spectra (300 MHz, Deuterated methanol (CD_3OD)) of Phalerin	48
4.6 ^{13}C -NMR spectra (75 MHz, Deuterated methanol (CD_3OD)) of Phalerin	49
4.7 Distortionless Enhancement by Polarization Transfer (DEPT-135°) spectra (75 MHz, Deuterated methanol (CD_3OD)) of Phalerin	49
4.8 MCF7 cell viability after 72h of treatment with <i>Phaleria macrocarpa</i> extract	53
4.9 HT29 cell viability after 72h of treatment with <i>Phaleria macrocarpa</i> extract	54
4.10 3T3 F442A cell viability after 72h of treatment with <i>Phaleria macrocarpa</i> extract	54
4.11 Effect of solvents on a) yield and b) concentration of phalerin at constant temperature of 60 °C and duration of 5 hours	56
4.12 Effect of solid to solvent ratio on a) yield and b) concentration of phalerin at constant temperature of 60 °C, particle size of <250µm and extraction time of 5 hours	57
4.13 Effect of particle size on a) yield and b) concentration of phalerin at constant temperature of 60°C, solid-to-solvent ratio of 1:40 g/ml and extraction time of 5 hours	59

4.14	Effect of temperature on a) yield and b) concentration of phalerin at constant particle size of $<250\text{ }\mu\text{m}$, solid-to-solvent ratio of 1:40 g/ml and extraction time of 5 hours	60
4.15	Comparison between experimental data and first order kinetic model for temperature of 40, 60 and $70\text{ }^{\circ}\text{C}$. Solid lines represent the first order kinetic model	62
4.16	Rate of reaction for temperature of 40, 60 and 70°C	63
4.17	Second order kinetics model and actual experimental data	64
4.18	Extraction kinetics of <i>Phaleria macrocarpa</i> at different temperatures, solid to solvent ratio of 1:20 g/ml and particle size of $<250\text{ }\mu\text{m}$. Points represent average value of three experiments and lines correspond to fitting model of diffusion	65



LIST OF ABBREVIATIONS

A	Arrhenius constant
A_0	Absorbance of negative control
ABTS	2,2-azinobis (3-ethylbenzothiazoline-6-sulfonic acid)
A_c	Absorbance of control
A_{ref}	Absorbance of reference
A_s	Absorbance of test material
A_T	Area of solid liquid interface (m^2)
a	Interfacial surface area (cm^{-1})
C_A	Concentration of solute in the solid phase (kg/m^3)
C_C	Concentration of solute C in the solution (kg/m^3)
CC	Column chromatography
CD_3OD	Deuterated methanol
C_e	Concentration of solute at equilibrium
CE	Capillary electrophoresis
C_e^{d1}	Concentration at equilibrium in diffusion step 1
C_e^{d2}	Concentration at equilibrium in diffusion step 2
C_e^w	Concentration at equilibrium in washing step
CH	Carbon-hydrogen bond
CH_2	Methylene
CH_3CN	Acetonitrile
CNMR	Carbon nuclear magnetic resonance spectroscopy
CO_2	Carbon dioxide
C_s	Concentration of solute at saturation or equilibrium (mgL^{-1})
C_t	Concentration of solute at any time t
D	Diffusivity of the solute in the solvent or inert solid (m^2/s)
D_{eff}	Effective diffusivity
DCM	Dichloromethane
DEPT	Distortionless enhancement by polarization transfer
DMSO	Dimethylsulfoxide
DPPH	2,2-diphenyl-1-picrylhydrazyl
E	Mean relative deviation modulus
h	Initial extraction rate
h'	Plank constant ($6.626 \times 10^{-34} J.s$)
H_2O	Water
HNMR	Hydrogen nuclear magnetic resonance spectroscopy
HPLC	High performance liquid chromatography
HYA	Hyaluronidase
IR	Infrared spectroscopy
IUPAC	International Union of Pure and Applied Chemistry
k	Rate constant
K_1	First order rate constant (min^{-1})
K_2	Second order rate constant ($ml\ g^{-1}min^{-1}$)
K	Equilibrium constant
ka	Volumetric mass transfer coefficient

k_{d1}	Mass transfer coefficient of diffusion step 1 (min^{-1})
k_{d2}	Mass transfer coefficient of diffusion step 2 (min^{-1})
k_w	Mass transfer coefficient of washing step
LOX	Lipoxygenase
M	Mass (g)
MeOH	Methanol
MS	Mass spectroscopy
MTT	3-(4,5-Dimethylthiazol-2-Yl)-2,5-Diphenyltetrazolium Bromide
N	Avogadro number
N_c	Rate of dissolution of solute C in the solution (kg/s)
NDGA	Nordihydroguaiaretic acid
NMR	Nuclear magnetic resonance spectroscopy
OCH_3	Methoxy group
PVDF	Polyvinylidene fluoride
r	Distance from the solid center (cm)
R	Gas constant (8.314 J/mol.K)
R^2	Correlation coefficient
R_f	Retention factor
RMSE	Root mean square error
SLE	Solid liquid extraction
t	Time (min)
TEAC	Trolox equivalent antioxidant capacity
TLC	Thin layer chromatography
UV-Vis	Ultraviolet visible spectroscopy
V	Volume
V_e	Volume of extract used for freeze drying (ml)
V_t	Volume of extract obtained after filtration (ml)
W_d	Weight of dried extract (g)
W_s	Weight of dried solid leaves (g)
X	Distance
XO	Xanthine oxidase
Y_T	Percent phalerin extracted at equilibrium
Y_U	Percent unextracted phalerin at equilibrium
Z	Distance inside the porous of the solid matrix (m)
ΔG	Gibb's free energy (kJ/mol)
ΔG^\ddagger	Activation free energy (kJ/mol)
ΔH	Enthalpy change (kJ/mol)
ΔH^\ddagger	Activation enthalpy (kJ/mol)
ΔS	Entropy change (J/mol.K)
ΔS^\ddagger	Activation entropy (J/mol.K)
ε	Void fraction space or porosity of the solid
λ_n	Eigenvalues
ρ_A	Density of solid (gcm^{-3})
τ	Dimensionless time

CHAPTER 1

INTRODUCTION

1.1 Background

Traditional medicine is the sum of total knowledge, skills, practices based on theory, beliefs and experiences indigenous to different culture, whether explicable or not, used in diagnosis, prevention and elimination of physical, mental or social imbalance (WHO Traditional medicine Programme, See Zhang, 1998). Traditional medicine is being used globally and it is estimated over 64% of world's population rely on plants as a primary health care source. Traditional medicine is relatively healthier, holistic, promote health and life quality and more self-reliance.

World population nowadays is increasingly seeking natural, drugless remedies or solutions to health problems. Moreover, considerable attention has been paid to utilize eco-friendly and natural based product for the prevention and cure of human diseases. Traditional medicine, based on natural product, is highly lucrative in the international medicine where it covers various aspect including photochemical, drugs, extracts, essential oils, nutraceuticals and aromatherapy. With such amount of diverse application of natural product, the demand for it shows increment of 5 to 13 percent annually. The largest market of the natural product in the world is in Europe followed by United State, Asia, Japan and other countries. In America, the spending on the complementary and alternative medicine industry in 2007 was approximately \$34 billion by which \$18 and \$4 billion of this overall expenditure represented dietary supplements and herbal products respectively (National Center for Complementary and Alternative Medicine, 2013). In Malaysia, traditional medicine has penetrate the herbal industry due to its rewarding business by which the per capita consumption of traditional medicine products is more than double the consumption of modern pharmaceuticals.

Among the common herbs being used as traditional medicine in Malaysia are ginger (*Zingiber officinale*), garlic (*Allium sativum*), turmeric (*Curcuma Longa*), piper (*Piper betle* L.), *Centella asiatica* L., *Orthosiphon staminus* and *Eurycoma longifolia*. Most of these common herbs are used traditionally to treat general illness such as flu, fever, allergies and dysentery due to antibacterial, anti-inflammatory, antioxidant and antidiabetic activities (Anand et al., 2010; Bhat & Karim, 2010; Ho et al., 2010; Mesomo et al., 2013; Pin et al., 2010; Singh et al., 2010). Besides aiming for treating certain illnesses, natural herbs have been taken regularly as healthy drink, tea, coffee and salad (Joubert & de Beer, 2011). The old folks' practices are the main sources for the growth of today's products which based on the natural resources and claim for having specific medicinal effects. However, with the increasing demand for natural product, precaution has to be taken seriously to ensure the products available are efficacious, of quality and safe for human use.

Studies on traditional medicine have been conducted to investigate the efficacy of the herbs practices from generation to generations. Normally, the traditional medicine practices on ordinary herbs that can easily be found in backyard. For

example, turmeric or *Curcuma longa* which is a common ingredient in Malaysian cuisine has been identified having antioxidant, antiviral, antifungal, anti-inflammatory and anticancer activities through recent studies. Thus, with supporting information from various discipline regarding natural herbs, an emerging supply and demand for natural product is flocking the market. With that, each party such as medical profession, investigative scientists, regulatory agencies and public have to understand the particular characteristics of natural herbs and derivatives and their potential for success and failure in preventing and confronting diseases (Barnes & Prasain, 2005).

Herbal processing industry is a team to develop natural product which started from the collection of raw product, pretreatment, extraction and finally provide product to the consumer for further usages or applications as shown in Figure 1.1. This process is a part of larger industry incorporating the pharmaceuticals, nutraceuticals, functional foods, herbal remedies and nutritional supplements. In herbal processing industry, extraction oil, bioactive ingredients, aroma or flavor are among the products that are valuable in the market. Thus, a process development in producing these valuable natural products is essential to ensure the herbal product met the quality, safety and efficacy requirements.




Figure 1.1: Herbal processing flow diagram

1.2 *Phaleria macrocarpa* (Scheff.) Boerl

Phaleria macrocarpa (Scheff.) Boerl shown in Figure 1.2 is a plant originally found in Indonesia. It is known by the name of Mahkota dewa (God's crown) in Malaysia and Indonesia. Most parts of this plant have been used to treat various diseases by the old folk Indonesians. All parts including the fruit, leaves, stem and seed have been studied extensively to determine the usages based mainly on the old folk practices. The seed of this plant is toxic to be consumed and thus only be used topically. This is due to the toxicity exhibited by the plant which might cause burning and numb effect if consume without any pretreatment.



Figure 1.2: *Phaleria macrocarpa*

This plant grows in fertile soil at an altitude of 10-1200 meters above sea level. It becomes productive after an age of about 4 months and can be utilized for 10 to 20 years. It can be found in tropical climates and reaches 1-2.5 meters in height in cultivation and up to 6 meters in wild.

Traditionally, this plant has been consumed for allergy, dysentery, skin diseases and energy drink. From the recent studies, *Phaleria macrocarpa* has been identified as a potential plant source which provides biological activities to treat or prevent certain diseases. Among chemical constituents found in the leaves, fruits and the pericarp are alkaloids, saponin, polyphenols, tannins, flavonoids, phenol and sterols (Harmanto, 2003). Moreover, these parts of plant are proven having antioxidant, anticancer, antidiabetic and anti-inflammatory activities.

1.3 Problem statement

Traditional medicine or alternative medicine is widely used in the prevention, diagnosis and treatment of an extensive range of ailments. The facts that the

traditional medicine is easily accessible which commonly found in the backyard, affordable and not require complicated process to consume make it as first remedies rather than the modern medicine to certain people.

Phaleria macrocarpa is one of the medicinal plants that recently penetrate the herbal industry in Malaysia. This plant consists of stem, fruits, leaves, and seed, has multiple medical usages. Through recent studies the stem has potential in treating bone cancer (Hendig & Ermin, 2009), the fruits can treat tumor, the leaves can treat allergy while the seed not only toxic but can be used for skin diseases (Faried et al., 2007). Eventhough the *Phaleria macrocarpa* plant has proven having various medicinal purposes but little findings has done to identify the compounds in the plant that contribute to those effects.

Many studies focused on the constitution of *Phaleria macrocarpa* and the efficacy of the crude extract of *Phaleria macrocarpa* on the certain diseases. However, there is lack of linkage between these two by which on how the compound that contribute to treat certain diseases can be obtained at optimal condition using extraction process. *Phaleria macrocarpa* is reported to contain phalerin which contributes to the anti-inflammatory and cytotoxic effect. Phalerin is not only major but also active compound that can be found only in the leaves of *Phaleria macrocarpa*. Thus, this study focuses on optimum parameters of extracting phalerin from the plant by solid liquid extraction method. Since phalerin standard was not available for the study, the method of extracting phalerin needs to be performed. The characterisation of phalerin obtained is compared and confirmed with previous study done by Mae et al. (2005).

From recent studies, *Phaleria macrocarpa* has been identified as a potential plant source which provided biological activities to treat or prevent certain diseases. However, there is still some area that is needed to study even further. This study focuses on the bioactivity of the extracted compounds that exhibit the biological activity such as antioxidant, anti-inflammatory and anticancer. Upon identifying the active compound, a study to optimize the extraction parameters subject to the active compound is performed. Conventional solid liquid extraction process is commonly used in herbal industry due to it simple process and require less equipment. Therefore, this method is used in this study. Kinetics study of process is then investigated.

1.4 Objectives of the study

The main objective of the study is to identify an active component from the leaves that contribute to the biological effect of the plant and proposed the optimum process parameters that could produce high yield of that compound. Moreover, it provided some useful information on the usages and the related parameters when dealing with the extraction process of this plant.

The study aims for the followings:

1. To identify the characteristics of extracted compound from the *Phaleria macrocarpa* leaves known as phalerin through chemical analysis.

2. To investigate the bioactivity of the extracted compounds through in-vitro study using antioxidant, anti-inflammatory and anticancer assays.
3. To optimize process parameters including type of solvents, ratio of solid to solvent, solid particle size and temperature for solid liquid extraction of *Phaleria macrocarpa* leaves.
4. To investigate extraction kinetics of active compound from *Phaleria macrocarpa* leaves and applied mathematical models for solid liquid extraction of the leaves.

1.5 Scope of the study

This study focuses on the extraction process of *Phaleria macrocarpa* leaves. The leaf has not been studied extensively and thus there is no study yet done on the optimization and kinetics of the extraction process. The findings are supported by the bioactivity related to the single compound extracted.

The extraction process of *Phaleria macrocarpa* leaves initially was focusing in identifying the single component that is major and has medicinal effect. The extraction was carried out through solid liquid extraction. Extraction and fractionation process was done to isolate a single compound from the leaves. High Performance Liquid Chromatography (HPLC), Thin Layer Chromatogram (TLC), Nuclear Magnetic Resonance (NMR) and Liquid Chromatogram Mass Spectroscopy (LCMS) were performed to ensure a single component was extracted and determined the identity of particular compound based on literature. Later, the obtained component was tested for antioxidant, anti-inflammatory and anticancer effect.

The optimization of extraction process was carried out using solid-liquid extraction method. Through this study, type of solvent, particle size, extraction temperature and solid-to-solvent ratio was optimized based on the crude yield and the phalerin concentration. The HPLC was used to determine the concentration of phalerin present in the extract. Moreover, the effect of each extraction parameters on the yield and concentration of phalerin was studied. The solvent effect on the yield was carried out and the reasons for having such solvent was rectified to ensure high quality and quantity of extract was obtained besides the safety requirement for food processing methods. The practical particle size and solid-to-solvent ratio was measured for the optimum yield. Moreover, consideration of waste due to solvent was measure for choice of solid-to-solvent ratio. On the other hand, optimum parameters were evaluated based on yield obtained and the quality of the phalerin present in the extract.

Based on the optimization process, the kinetics study was performed using lab scale solid-liquid extractor. The process was performed until the system reached equilibrium. Mathematical models were proposed to predict the extraction process.

REFERENCES

- Ahmed, Z., Abdeslam-Hassan, M., Ouassila, L., & Danielle, B. (2012). Extraction and Modeling of Algerian Rosemary Essential Oil Using Supercritical CO₂: Effect of Pressure and Temperature. *Energy Procedia*, 18, 1038–1046.
- Ali, R. B., Atangwho, I. J., Kuar, N., Abraika, O. S., Ahmad, M., Mahmud, R., & Asmawi, M. Z. (2012). Bioassay-guided antidiabetic study of *Phaleria macrocarpa* fruit extract. *Molecules (Basel, Switzerland)*, 17, 4986–5002.
- Ali, R. B., Atangwho, I. J., Kuar, N., Ahmad, M., Mahmud, R., & Asmawi, M. Z. (2013). In vitro and in vivo effects of standardized extract and fractions of *Phaleria macrocarpa* fruits pericarp on lead carbohydrate digesting enzymes. *BMC complementary and alternative medicine*, 13(1), 39.
- Ali, R. B., Atangwho, I. J., Kuar, N., Mohamed, E. A. H., Mohamed, A. J., Asmawi, M. Z., & Mahmud, R. (2012). Hypoglycemic and anti-hyperglycemic study of *Phaleria macrocarpa* fruits pericarp. *Journal of Medicinal Plants Research*, 6(10), 1982–1990.
- Altaf, R., Asmawi, M. Z. Bin, Dewa, A., Sadikun, A., & Umar, M. I. (2013). Phytochemistry and medicinal properties of *Phaleria macrocarpa* (Scheff.) Boerl. extracts. *Pharmacognosy reviews*, 7(13), 73–80.
- Anand, T., Mahadeva, N., Phani, K. G., & Farhath, K. (2010). Antioxidant and DNA Damage Preventive Properties of *Centella asiatica* (L) Urb. *Pharmacognosy Journal*, 2(17), 53–58.
- Aripin, A., Arifin, P. F., & Tjandrawinata, R. R. (2011). U.S. Patent No US 2011/0257112 A1. Isolated compounds from *Phaleria macrocarpa* as anti cancer agents. Washington D. C.: U. S Patent and Trademark Office.
- Armenia, E. F., Widya, R. M., Rusdi, D., & Netty, M. (2006). Anti-atherosclerotic effect and liver toxicity of ethanolic extract of *phaleria macrocarpa* (Scheff.) Boerl fruit on Japanese quail. In *Oral Paper ASOMP XII, 13-18 November, Padang, Indonesia* (pp. 1–10).
- Azmir, J., Zaidul, I. S. M., Rahman, M. M., Sharif, K. M., Sahena, F., Jahurul, M. H. A., & Mohamed, A. (2014). Optimization of oil yield of *Phaleria macrocarpa* seed using response surface methodology and its fatty acids constituents. *Industrial Crops and Products*, 52, 405–412.
- Baek, J. Y., Lee, J. M., & Lee, S. C. (2008). Extraction of nutraceutical compounds from licorice roots with subcritical water. *Separation and Purification Technology*, 63(3), 661–664.

- Barnes, S., & Prasain, J. (2005). Current progress in the use of traditional medicines and nutraceuticals. *Current opinion in plant biology*, 8, 324–8.
- Bäumler, E. R., Crapiste, G. H., & Carelli, A. a. (2010). Solvent Extraction: Kinetic Study of Major and Minor Compounds. *Journal of the American Oil Chemists' Society*, 87(12), 1489–1495.
- Bernhoft, A. (2010). A brief review on bioactive compounds in plants. In *Bioactive compounds in plants – benefits and risks for man and animals Oslo: The Norwegian Academy of Science and Letters* (pp. 11–17).
- Bhat, R., & Karim, A. A. (2010). Tongkat Ali (*Eurycoma longifolia* Jack): a review on its ethnobotany and pharmacological importance. *Fitoterapia*, 81, 669–679.
- Bucić-Kojić, A., Sovová, H., Planinić, M., & Tomas, S. (2013). Temperature-dependent kinetics of grape seed phenolic compounds extraction: experiment and model. *Food chemistry*, 136(3-4), 1136–40.
- Cacace, J. E., & Mazza, G. (2003). Mass transfer process during extraction of phenolic compounds from milled berries. *Journal of Food Engineering*, 59(4), 379–389.
- Chan, C. H., Yusoff, R., & Ngoh, G. C. (2014). Modeling and kinetics study of conventional and assisted batch solvent extraction. *Chemical Engineering Research and Design*, 92(6), 1169–1186.
- Chanda, S. V., & Kaneria, M. J. (2011). Optimization of Conditions for the Extraction of Antioxidants from Leaves of *Syzygium cumini* L. Using Different Solvents. *Food Analytical Methods*, 5(3), 332–338.
- Chong, S. C., Dollah, M. A., Chong, P. P., & Maha, A. (2011). *Phaleria macrocarpa* (Scheff.) Boerl fruit aqueous extract enhances LDL receptor and PCSK9 expression in vivo and in vitro. *Journal of ethnopharmacology*, 137(1), 817–27.
- Chua, L. S. (2013). A review on plant-based rutin extraction methods and its pharmacological activities. *Journal of ethnopharmacology*, 150(3), 805–17.
- Ciapetti, G., Cenni, E., Pratelli, L., & Pizzoferrato, A. (1993). In vitro evaluation of cell/biomaterial interaction by MTT assay. *Biomaterials*, 14(5), 359–364.
- De la Rubia, M. A., Fernández-Cegrí, V., Raposo, F., & Borja, R. (2011). Influence of particle size and chemical composition on the performance and kinetics of anaerobic digestion process of sunflower oil cake in batch mode. *Biochemical Engineering Journal*, 58-59, 162–167.

- Diankov, S., Karsheva, M., & Hinkov, I. (2011). Extraction of natural antioxidants from lemon peels. Kinetics and antioxidant capacity. *Journal of the University of Chemical Technology and Metallurgy*, 46(3), 315–319.
- Diantini, A., Achmad, T. H., Julaeha, E., Faried, A., Faried, L. S., Abdullah, R., Wardhani, S. R., Levita, J., Kurnia, D., Subarnas, A., & Supriyatna, S. (2012). Cytotoxicity of fevicordin-A from *Phaleria macrocarpa* (Boerl.) scheff on P388 HeLa, CasKi, TE-2, TE-8 and Prepuce ' s Fibroblast cells. *E3 Journal of Medical Research*, 1(1), 1–5.
- Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y.-H. (2014). Effect of extraction solvent on total phenol content, total flavonoids content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22, 296–302.
- Dorta, E., Lobo, M. G., & Gonzalez, M. (2012). Reutilization of mango byproducts: study of the effect of extraction solvent and temperature on their antioxidant properties. *Journal of food science*, 71(1), C80–8.
- Durling, N., Catchpole, O., Grey, J., Webby, R., Mitchell, K., Foo, L., & Perry, N. (2007). Extraction of phenolics and essential oil from dried sage (*Salvia officinalis*) using ethanol–water mixtures. *Food Chemistry*, 101(4), 1417–1424.
- Ebewele, R. O., Iyayi, A. F., & Hymore, F. K. (2010). Considerations of the extraction process and potential technical applications of Nigerian rubber seed oil. *International Journal of the Physical Sciences*, 5(June), 826–831.
- Effendy, M. A. W., Yosie, A., Sifzizul, T. M. T., & Habsah, M. (2011). Antibacterial, radical-scavenging activities and cytotoxicity properties of *Phaleria macrocarpa* (Scheff.) Boerl. leaves in HepG2 cell lines. *International Journal of Pharmaceutical Sciences and Research*, 2(7), 1700–1706.
- Eikani, M. H., Golmohammad, F., & Rowshanzamir, S. (2007). Subcritical water extraction of essential oils from coriander seeds (*Coriandrum sativum* L.). *Journal of Food Engineering*, 80(2), 735–740.
- Espinoza-Pérez, J. D., Vargas, a., Robles-Olvera, V. J., Rodrí'guez-Jimenes, G. C., & Garcí'a-Alvarado, M. a. (2007). Mathematical modeling of caffeine kinetic during solid–liquid extraction of coffee beans. *Journal of Food Engineering*, 81(1), 72–78.
- Faried, A., Kurnia, D., Faried, L. S., Usman, N., Miyazaki, T., Kato, H., & Kuwano, H. (2007). Anticancer effects of gallic acid isolated from Indonesian herbal medicine, *Phaleria macrocarpa* (Scheff.) Boerl, on human cancer cell lines. *International journal of oncology*, 30(3), 605–613.

- Ferrari, J., Terreaux, C., Sahpaz, S., Msonthi, J. D., Wolfender, L., & Hostettmann, K. (2000). Benzophenone glycosides from *Gnidia involucrata*. *Phytochemistry*, 54, 883–889.
- Gangga, E., Asriani, H., & Novita, L. (2007). Analisis Pendahuluan Metabolit Sekunder dari Kalus Mahkota Dewa (*phaleria macrocarpa* [Scheff .] Boerl .). *Jurnal Ilmu Kefarmasian Indonesia*, 5(I), 17–22.
- Ghoreishi, S. M., & Shahrestani, R. G. (2009). Subcritical water extraction of mannitol from olive leaves. *Journal of Food Engineering*, 93(4), 474–481.
- Ghoreishi, S. M., & Sharifi, S. (2001). Modeling of supercritical extraction of mannitol from plane tree leaf. *Journal of Pharmaceutical and Biomedical Analysis*, 24(5-6), 1037–1048.
- Ghufron, M., Soesatyo, M., Haryana, S. M., & Sismindari. (2008). The effect of mahkota dewa (*Phaleria macrocarpa* (Scheff) Boerl) leaf etanolic extract on splenic NK1 . 1 cells activity. *Berkala Ilmu Kedokteran*, 40(3), 109–118.
- Gironi, F., & Piemonte, V. (2011). Temperature and solvent effects on polyphenol extraction process from chestnut tree wood. *Chemical Engineering Research and Design*, 89, 857–862.
- Gong, X., Zhang, L., Jiang, R., Ye, M., Yin, X., & Wan, J. (2013). Anti-inflammatory effects of mangiferin on sepsis-induced lung injury in mice via up-regulation of heme oxygenase-1. *The Journal of nutritional biochemistry*, 24, 1173–1181.
- Goula, A. M. (2013). Ultrasound-assisted extraction of pomegranate seed oil – Kinetic modeling. *Journal of Food Engineering*, 117(4), 492–498.
- Gülçin, İ., Huyut, Z., Elmastaş, M., & Aboul-Enein, H. Y. (2010). Radical scavenging and antioxidant activity of tannic acid. *Arabian Journal of Chemistry*, 3(1), 43–53.
- Gurib-Fakim, A. (2006). Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*, 27, 1–93.
- Haq, A. U., Malik, A., Anis, I., Khan, S. B., Ahmed, E., Ahmed, Z., Nawaz, S. A., & Choudhary, M. I. (2004). Enzymes Inhibiting Lignans from *Vitex negundo*. *Chem. Pharm. Bull.*, 52(11), 1269–1272.
- Harmanto, N. (2003). Conquering Disease in Unison with Mahkota Dewa, *Phaleria Macrocarpa*. In *First Edition. P. T. Mahkota Dewa Indonesia, Jakarta* (p. 14). Mahkotadewa Indonesia.

- Hassas-Roudsari, M., Chang, P. R., Pegg, R. B., & Tyler, R. T. (2009). Antioxidant capacity of bioactives extracted from canola meal by subcritical water, ethanolic and hot water extraction. *Food Chemistry*, 114(2), 717–726.
- He, L., Zhang, X., Xu, H., Xu, C., Yuan, F., Knez, Ž., Novak, Z., & Gao, Y. (2012). Subcritical water extraction of phenolic compounds from pomegranate (*Punica granatum* L.) seed residues and investigation into their antioxidant activities with HPLC–ABTS+ assay. *Food and Bioproducts Processing*, 90(2), 215–223.
- Hendig, W., & Ermin, K. W. (2009). Benzophenone glucoside isolated from the ethyl acetate extract of the bark of Mahkota dewa [*Phaleria macrocarpa* (Scheff.) Boerl.] and its inhibitory activity on leukemia L1210 cell line. *Indo. J. Chem.*, 9(1), 142–145.
- Hendra, P., Fukushima, Y., & Hashidoko, Y. (2009). Synthesis of Benzophenone Glucopyranosides from *Phaleria macrocarpa* and Related Benzophenone Glucopyranosides. *Bioscience, Biotechnology, and Biochemistry*, 73(10), 2172–2182.
- Hendra, R., Ahmad, S., Sukari, A., Shukor, M. Y., & Oskoueian, E. (2011). Flavonoid Analyses and Antimicrobial Activity of Various Parts of *Phaleria macrocarpa* (Scheff.) Boerl Fruit. *International journal of molecular sciences*, 12, 3422–3431.
- Herodež, Š. S., Hadolin, M., Škerget, M., & Knez, Ž. (2003). Solvent extraction study of antioxidants from Balm (*Melissa officinalis* L.) leaves. *Food Chemistry*, 80(2), 275–282.
- Herzi, N., Bouajila, J., Camy, S., Romdhane, M., & Condoret, J.-S. (2013). Comparison of different methods for extraction from *Tetralinis articulata*: yield, chemical composition and antioxidant activity. *Food chemistry*, 141, 3537–45.
- Ho, C. H., Noryati, I., Sulaiman, S.-F., & Rosma, A. (2010). In vitro antibacterial and antioxidant activities of *Orthosiphon stamineus* Benth. extracts against food-borne bacteria. *Food Chemistry*, 122(4), 1168–1172.
- Hossain, M. B., Barry-Ryan, C., Martin-Diana, A. B., & Brunton, N. P. (2011). Optimisation of accelerated solvent extraction of antioxidant compounds from rosemary (*Rosmarinus officinalis* L.), marjoram (*Origanum majorana* L.) and oregano (*Origanum vulgare* L.) using response surface methodology. *Food Chemistry*, 126, 339–346.
- Hung, S. M., Hsu, B. D., & Lee, S. (2014). Modelling of isothermal chlorophyll extraction from herbaceous plants. *Journal of Food Engineering*, 128, 17–23.

- Irma, H. S., Erni, S., Bayu, F. O., Wulan, T., Sylvia, P., & Yasmina, P. (2010). Phaleria macrocarpa fruit extract as insulinotropic agent in streptozotocin-induced diabetic cynomolgus monkeys (Macaca fascicularis). In *Proceeding of the International Conference on Medicinal Plants-Surabaya Indonesia* (pp. 346–350).
- Jadhav, D., Rekha, B. N., Gogate, P. R., & Rathod, V. K. (2009). Extraction of vanillin from vanilla pods : A comparison study of conventional soxhlet and ultrasound assisted extraction. *Journal of Food Engineering*, 93(4), 421–426.
- Jaganyi, D., & Wheeler, P. J. (2003). Rooibos tea: equilibrium and extraction kinetics of aspalathin. *Food Chemistry*, 83(1), 121–126.
- Jensen, S., & Zanoelo, É. F. (2013). Kinetics of Aqueous Extraction of Mate (Ilex Paraguariensis) Leaves. *Journal of Food Process Engineering*, 36(2), 220–227.
- Jing, P., & Giusti, M. M. (2007). Effects of extraction conditions on improving the yield and quality of an anthocyanin-rich purple corn (Zea mays L.) color extract. *Journal of food science*, 72(7), C363–8.
- Joubert, E., & de Beer, D. (2011). Rooibos (Aspalathus linearis) beyond the farm gate: From herbal tea to potential phytopharmaceutical. *South African Journal of Botany*, 77, 869–886.
- Ju, Z. Y., & Howard, L. R. (2003). Effects of solvent and temperature on pressurized liquid extraction of anthocyanins and total phenolics from dried red grape skin. *Journal of agricultural and food chemistry*, 51(18), 5207–13.
- Karacabey, E., Bayindirli, L., Artik, N., & Mazza, G. (2013). Modeling Solid-Liquid Extraction Kinetics of Trans -Resveratrol and Trans -E-Viniferin From Grape Cane. *Journal of Food Process Engineering*, 36(1), 103–112.
- Katrin, E., Selvie, & Winarno, H. (2011). Chromatogram Profiles and Cytotoxic Activity of Irradiated Mahkota Dewa (Phaleria Macrocarpa (Scheff .) Boerl) Leaves. *Atom Indonesia*, 37(1), 17–23.
- Khan, K. M., Ambreen, N., Mughal, U. R., Jalil, S., Perveen, S., & Choudhary, M. I. (2010). 3-Formylchromones: potential antiinflammatory agents. *European journal of medicinal chemistry*, 45, 4058–64.
- Kim, W. J., Kim, J., Veriansyah, B., Kim, J. D., Lee, Y. W., Oh, S. G., & Tjandrawinata, R. R. (2009). Extraction of bioactive components from Centella asiatica using subcritical water. *The Journal of Supercritical Fluids*, 48(3), 211–216.

- Kim, W. J., Veriansyah, B., Lee, Y. W., Kim, J., & Kim, J. D. (2010). Extraction of mangiferin from Mahkota Dewa (*Phaleria macrocarpa*) using subcritical water. *Journal of Industrial and Engineering Chemistry*, 16(3), 425–430.
- Ko, M.-J., Cheigh, C.-I., Cho, S.-W., & Chung, M.-S. (2011). Subcritical water extraction of flavonol quercetin from onion skin. *Journal of Food Engineering*, 102(4), 327–333.
- Kostić, M. D., Joković, N. M., Stamenković, O. S., Rajković, K. M., Milić, P. S., & Veljković, V. B. (2014). The kinetics and thermodynamics of hempseed oil extraction by n-hexane. *Industrial Crops and Products*, 52, 679–686.
- Kukic, J., Popovic, V., Petrovic, S., Mucaji, P., Ciric, A., Stojkovic, D., & Sokovic, M. (2008). Food Chemistry Antioxidant and antimicrobial activity of *Cynara cardunculus* extracts. *Food Chemistry*, 107, 861–868.
- Kulkarni, V. M., & Rathod, V. K. (2014a). Mapping of an ultrasonic bath for ultrasound assisted extraction of mangiferin from *Mangifera indica* leaves. *Ultrasonics sonochemistry*, 21(2), 606–11.
- Kulkarni, V. M., & Rathod, V. K. (2014b). Extraction of mangiferin from *Mangifera indica* leaves using three phase partitioning coupled with ultrasound. *Industrial Crops and Products*, 52, 292–297.
- Kumar, M. S. Y., Dutta, R., Prasad, D., & Misra, K. (2011). Subcritical water extraction of antioxidant compounds from Seabuckthorn (*Hippophae rhamnoides*) leaves for the comparative evaluation of antioxidant activity. *Food chemistry*, 127(3), 1309–16.
- Latoui, M., Aliakbarian, B., Casazza, A. A., Seffen, M., Converti, A., & Perego, P. (2012). Extraction of phenolic compounds from *Vitex agnus-castus* L. *Food and Bioproducts Processing*, 90(4), 748–754.
- Lee, M. H., & Lin, C. C. (2007). Comparison of techniques for extraction of isoflavones from the root of *Radix Puerariae*: Ultrasonic and pressurized solvent extractions. *Food Chemistry*, 105(1), 223–228.
- Leitão, N. C. M. C. S., Prado, G. H. C., Veggi, P. C., Meireles, M. A. A., & Pereira, C. G. (2013). *Anacardium occidentale* L. leaves extraction via SFE: Global yields, extraction kinetics, mathematical modeling and economic evaluation. *The Journal of Supercritical Fluids*, 78, 114–123.
- Li, F., Wang, F., Yu, F., Fang, Y., Xin, Z., Yang, F., Xu, J., Zhao, L., & Hu, Q. (2008). In vitro antioxidant and anticancer activities of ethanolic extract of selenium-enriched green tea. *Food Chemistry*, 111, 165–170.

- Linares, A. R., Hase, S. L., Vergara, M. L., & Resnik, S. L. (2010). Modeling yerba mate aqueous extraction kinetics: Influence of temperature. *Journal of Food Engineering*, 97(4), 471–477.
- Ling, S., Tanaka, T., & Kouno, I. (2003). Effects of Iridoids on Lipxygenase and Hyaluronidase Activities and Their Activation by β -Glucosidase in the Presence of Amino Acids. *Biol. Pharm. Bull*, 26(3), 352–356.
- Liu, W., Yu, Y., Yang, R., Wan, C., Xu, B., & Cao, S. (2010). Optimization of Total Flavonoid Compound Extraction from *Gynura medica* Leaf Using Response Surface Methodology and Chemical Composition Analysis. *International journal of molecular sciences*, 11, 4750–63.
- Lopresto, C. G., Petrillo, F., Casazza, A. A., Aliakbarian, B., Perego, P., & Calabrò, V. (2014). A non-conventional method to extract D-limonene from waste lemon peels and comparison with traditional Soxhlet extraction. *Separation and Purification Technology*, 137, 13–20.
- Luo, W., Zhao, M., Yang, B., Shen, G., & Rao, G. (2009). Identification of bioactive compounds in *Phyllanthus emblica* L. fruit and their free radical scavenging activities. *Food Chemistry*, 114(2), 499–504.
- Mae Sri Hartati, W., Mubarika, S., Gandjar, I. G., Hamann, M. T., Rao, K. V., & Wahyuono, S. (2005). Phalerin, a new benzophenonic glucoside isolated from the methanolic extract of Mahkota Dewa [*Phaleria macrocarpa* (Scheff.) Boerl] leaves. *Majalah Farmasi Indonesia*, 16(1), 51–57.
- Makare, N., Bodhankar, S., & Rangari, V. (2001). Immunomodulatory activity of alcoholic extract of *Mangifera indica* L. in mice. *Journal of Ethnopharmacology*, 78(2-3), 133–137.
- Maksimovic, S., Ivanovic, J., & Skala, D. (2012). Supercritical Extraction of Essential Oil from *Mentha* and Mathematical Modelling– the Influence of Plant Particle Size. *Procedia Engineering*, 42, 1767–1777.
- Markom, M., Hasan, M., Daud, W. R. W., Singh, H., & Jahim, J. M. (2007). Extraction of hydrolysable tannins from *Phyllanthus niruri* Linn.: Effects of solvents and extraction methods. *Separation and Purification Technology*, 52, 487–496.
- Masibo, M., & He, Q. (2009). In vitro antimicrobial activity and the major polyphenol in leaf extract of *Mangifera*. *Malaysian Journal of Microbiology*, 5(2), 73–80.
- Mesomo, M. C., Corazza, M. L., Ndiaye, P. M., Dalla Santa, O. R., Cardozo, L., & Scheer, A. de P. (2013). Supercritical CO₂ extracts and essential oil of ginger (*Zingiber officinale* R.): Chemical composition and antibacterial activity. *The Journal of Supercritical Fluids*, 80, 44–49.

- Meyer, A. S., & Frankel, E. N. (2001). Antioxidant activity of hydroxycinnamic acids on human low-density lipoprotein oxidation. *Methods in Enzymology*, 335, 256–265.
- Meziane, S., & Kadi, H. (2008). Kinetics and Thermodynamics of Oil Extraction from Olive Cake. *Journal of the American Oil Chemists' Society*, 85(4), 391–396.
- Milojević, S. Ž., Stojanović, T. D., Palić, R., Lazić, M. L., & Veljković, V. B. (2008). Kinetics of distillation of essential oil from comminuted ripe juniper (*Juniperus communis* L.) berries. *Biochemical Engineering Journal*, 39(3), 547–553.
- Miura, T., Ichiki, H., Hashimoto, I., Iwamoto, N., Kato, M., Kubo, M., Ishihara, E., Komatsu, Y., Okada, M., Ishida, T., & Tanigawa, K. (2001). Antidiabetic activity of a xanthone compound, mangiferin. *Phytomedicine*, 8(2), 85–87.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 65, 55–63.
- Muruganandan, S., Srinivasan, K., Gupta, S., Gupta, P. K., & Lal, J. (2005). Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. *Journal of ethnopharmacology*, 97(3), 497–501.
- Nayak, C. A., Chethana, S., Rastogi, N. K. Ā., & Raghavarao, K. S. M. S. (2006). Enhanced mass transfer during solid – liquid extraction of gamma-irradiated red beetroot. *Radiation Physics and Chemistry*, 75, 173–178.
- Noro, T., Oda, Y., Miyase, T., Ueno, A., & Fukushima, S. (1983). Inhibitors of xanthine oxidase from the flowers and buds of *Daphne genkwa*. *Chemical & pharmaceutical bulletin*, 31(11), 3984–3987.
- Nwabanne, J. T. (2012). Kinetics and Thermodynamics Study of Oil Extraction from Fluted Pumpkin Seed. *International Journal of Multidisciplinary Sciences and Engineering*, 3(6), 11–15.
- Ofori-Boateng, C., & Lee, K. T. (2013). Ultrasonic-Assisted Extraction of α -Tocopherol Antioxidants from the Fronds of *Elaeis guineensis* Jacq.: Optimization, Kinetics, and Thermodynamic Studies. *Food Analytical Methods*, 7(2), 257–267.
- Ong, E. S. (2004). Extraction methods and chemical standardization of botanicals and herbal preparations. *Journal of chromatography. B, Analytical technologies in the biomedical and life sciences*, 812(1-2), 23–33.
- Oshimi, S., Zaima, K., Matsuno, Y., Hirasawa, Y., Iizuka, T., Studiawan, H., Indrayanto, G., Zaini, N. C., & Morita, H. (2008). Studies on the constituents

- from the fruits of *Phaleria macrocarpa*. *Journal of natural medicines*, 62(2), 207–210.
- Ozel, M. Z., Gogus, F., & Lewis, A. C. (2003). Subcritical water extraction of essential oils from *Thymbra spicata*. *Food Chemistry*, 82(3), 381–386.
- Ozel, M. Z., & Kaymaz, H. (2004). Superheated water extraction , steam distillation and Soxhlet extraction of essential oils of *Origanum onites*. *Anal Bioanal Chem*, 379, 1127–1133.
- Pacher, P., Nivorozhkin, A., & Szabo, C. (2006). Therapeutic Effects of Xanthine Oxidase Inhibitors : Renaissance Half a Century after the Discovery of Allopurinol. *Pharmacological Reviews*, 58, 87–114.
- Park, B. Y., & Chung, J. (2009). Cadmium increases ferroportin-1 gene expression in J774 macrophage cells via the production of reactive oxygen species. *Nutrition research and practice*, 3(3), 192–9.
- Pavia, D. L., Lampman, G. M., Kriz, G. S., & Vyvyan, J. R. (2009). *Introduction to spectroscopy 4th Edition* (pp. 177–293; Brooks/ Cole, Cengage Learning).
- Peng, T. Y., Don, M. M., & Tahrel, M. A. (2012). Optimisation and Kinetics Studies on the Extraction of Essential Oil from *Zingiber Cassumunar*. *Journal of Physical Science*, 23(1), 65–82.
- Perez, E. E., Carelli, A. A., & Crapiste, G. H. (2011). Temperature-dependent diffusion coefficient of oil from different sunflower seeds during extraction with hexane. *Journal of Food Engineering*, 105, 180–185.
- Pilavtepe, M., & Yesil-Celiktas, O. (2013). Mathematical modeling and mass transfer considerations in supercritical fluid extraction of *Posidonia oceanica* residues. *The Journal of Supercritical Fluids*, 82, 244–250.
- Pin, K. Y., Chuah, A. L., Rashih, A. A., Mazura, M. P., Fadzureena, J., & Vimala, S. (2010). Antioxidant and anti-inflammatory activities of Extracts of betel leaves (*Piper betle*) from solvents with different polarities. *Journal of Tropical Forest Science*, 22(4), 448–455.
- Pin K. Y., A. Luqman Chuah, A. Abdull Rashih, M.A. Rasadah, Law, C. L., & Choong, T. S. Y. (2011). Solid-liquid extraction of betel leaves (*Piper Betle* L.). *Journal of Food Process Engineering*, 34, 549–565.
- Pinelo, M., Rubilar, M., Jerez, M., Sineiro, J., & Núñez, M. J. (2005). Effect of solvent, temperature, and solvent-to-solid ratio on the total phenolic content and antiradical activity of extracts from different components of grape pomace. *Journal of agricultural and food chemistry*, 53(6), 2111–7.

- Pinelo, M., & Sineiro, J. (2006). Mass transfer during continuous solid – liquid extraction of antioxidants from grape byproducts. *Journal of Food Engineering*, 77, 57–63.
- Punín Crespo, M. O., & Lage Yusty, M. A. (2005). Comparison of supercritical fluid extraction and Soxhlet extraction for the determination of PCBs in seaweed samples. *Chemosphere*, 59(10), 1407–13.
- Radha krishnan, K., Sivarajan, M., Babuskin, S., Archana, G., Azhagu Saravana Babu, P., & Sukumar, M. (2013). Kinetic modeling of spice extraction from *S. aromaticum* and *C. cassia*. *Journal of Food Engineering*, 117(3), 326–332.
- Rahimi, E., Prado, J. M., Zahedi, G., & Meireles, M. A. A. (2011). Chamomile extraction with supercritical carbon dioxide: Mathematical modeling and optimization. *The Journal of Supercritical Fluids*, 56(1), 80–88.
- Ramluckan, K., Moodley, K. G., & Bux, F. (2014). An evaluation of the efficacy of using selected solvents for the extraction of lipids from algal biomass by the soxhlet extraction method. *Fuel*, 116, 103–108.
- Rathod, S. S., & Rathod, V. K. (2014). Extraction of piperine from *Piper longum* using ultrasound. *Industrial Crops and Products*, 58, 259–264.
- Razali, N., Mat-Junit, S., Abdul-Muthalib, A. F., Subramaniam, S., & Abdul-Aziz, A. (2012). Effects of various solvents on the extraction of antioxidant phenolics from the leaves, seeds, veins and skins of *Tamarindus indica* L. *Food Chemistry*, 131, 441–448.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., & Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26(9-10), 1231–1237.
- Reis, S. F., Rai, D. K., & Abu-Ghannam, N. (2012). Water at room temperature as a solvent for the extraction of apple pomace phenolic compounds. *Food chemistry*, 135(3), 1991–8.
- Rice, G. R., & Do, D. D. (1994). *Applied mathematics and modelling for chemical engineers* (pp. 405–478; USA: John Wiley & Son, Inc.).
- Rice-evans, C. A., Miller, N. J., Bolwell, P. G., Bramley, P. M., & Pridham, J. B. (1995). The Relative Antioxidant Activities of Plant-Derived Polyphenolic Flavonoids. *Free Radical Research*, 22(4), 375–383.
- Rohyami, Y. (2008). Penentuan Kandungan Flavonoid dari Ekstrak Metanol Daging Buah Mahkota Dewa (*Phaleria macrocarpa* Scheff Boerl). *Jurnal Logika*, 5(1), 1–16.

- Rostami, M., Farzaneh, V., Boujmehrani, A., Mohammadi, M., & Bakhshabadi, H. (2014). Optimizing the extraction process of sesame seed's oil using response surface method on the industrial scale. *Industrial Crops and Products*, 58, 160–165.
- Sagu, S. T., Nso, E. J., Karmakar, S., & De, S. (2014). Optimisation of low temperature extraction of banana juice using commercial pectinase. *Food chemistry*, 151, 182–90.
- Şahin, S., & Samlı, R. (2013). Optimization of olive leaf extract obtained by ultrasound-assisted extraction with response surface methodology. *Ultrasonics sonochemistry*, 20, 595–602.
- Saifudin, A., Dai, M., & Munawaroh, R. (2006). Cytotoxic effect of metanol, ethanol, and chloroform extracts from callus of *Phaleria macrocarpa* (Scheff) Boerl to myeloma cell by MTT assay. *Berkala Ilmiah Biologi*, 5(1), 31–38.
- Sasidharan, I., & Menon, A. N. (2010). A study of antioxidant properties of different extracts of curry leaf (*Murraya Koenigii* L.). *EJEAFChe*, 9(6), 1036–1046.
- Sellamuthu, P. S., Arulselvan, P., Muniappan, B. P., & Kandasamy, M. (2012). Effect of mangiferin isolated from *Salacia chinensis* regulates the kidney carbohydrate metabolism in streptozotocin-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*, 2(3), S1583–S1587.
- Shen, S., Chen, D., Li, X., Li, T., Yuan, M., Zhou, Y., & Ding, C. (2014). Optimization of extraction process and antioxidant activity of polysaccharides from leaves of *Paris polyphylla*. *Carbohydrate Polymers*, 104, 80–86.
- Sheridan, H., Krenn, L., Jiang, R., Sutherland, I., Ignatova, S., Marmann, A., Liang, X., & Sendker, J. (2012). The potential of metabolic fingerprinting as a tool for the modernisation of TCM preparations. *Journal of ethnopharmacology*, 140, 482–91.
- Silva, L. P. S., & Martínez, J. (2014). Mathematical modeling of mass transfer in supercritical fluid extraction of oleoresin from red pepper. *Journal of Food Engineering*, 133, 30–39.
- Simeonov, E., & Koleva, V. (2012). Solid-liquid Extraction of Tannins from *Geranium Sanguineum* L. – Experimental Kinetics and Modelling. *Chem, Biochem, Eng, Q*, 26(3), 249–255.
- Singh, G., Kapoor, I. P. S., Singh, P., de Heluani, C. S., de Lampasona, M. P., & Catalan, C. A. N. (2010). Comparative study of chemical composition and antioxidant activity of fresh and dry rhizomes of turmeric (*Curcuma longa* Linn.). *Food and chemical toxicology*, 48(4), 1026–31.

- So, G. C., & Macdonald, D. G. (1986). Kinetics of oil extraction from canola (rapeseed). *The Canadian Journal of Chemical Engineering*, 64(1), 80–86.
- Soeksmanto, A., Hapsari, Y., & Simanjuntak, P. (2007). Antioxidant content of parts of Mahkota Dewa, *Phaleria macrocarpa* (Scheff) Boerl. (Thymelaceae). *Biodiversitas*, 8(2), 92–95.
- Solana, M., Boschiero, I., Dall'Acqua, S., & Bertucco, A. (2014). Extraction of bioactive enriched fractions from *Eruca sativa* leaves by supercritical CO₂ technology using different co-solvents. *The Journal of Supercritical Fluids*, 94, 245–251.
- Sousa, A., Ferreira, I. C. F. R., Barros, L., Bento, A., & Pereira, J. A. (2008). Effect of solvent and extraction temperatures on the antioxidant potential of traditional stoned table olives “alcaparras.” *LWT - Food Science and Technology*, 41(4), 739–745.
- Spigno, G., Tramelli, L., & De Faveri, D. M. (2007). Effects of extraction time, temperature and solvent on concentration and antioxidant activity of grape marc phenolics. *Journal of Food Engineering*, 81(1), 200–208.
- Su, C. H., Liu, C. S., Yang, P. C., Syu, K. S., & Chiu, C. C. (2014). Solid–liquid extraction of phycocyanin from *Spirulina platensis*: Kinetic modeling of influential factors. *Separation and Purification Technology*, 123, 64–68.
- Sulaiman, S., Abdul Aziz, A. R., & Kheireddine Aroua, M. (2013). Optimization and modeling of extraction of solid coconut waste oil. *Journal of Food Engineering*, 114(2), 228–234.
- Sumarawati, T., Riwanto, I., Hadisaputro, S., & Dharmana, E. (2012). Concentration of Phalerin from *Phaleria macrocarpa* in Vena Porta: An Experimental Study on Adenocarcinoma Mammariae in C3H Mice That Given Adriamycin and Cyclophosphamide. In *International Conference: Research and Application on Traditional Complementary and Alternative Medicine in Health Care (TCAM)* (pp. 18–23).
- Sumastuti. (2002). Efek Antihistamin Ekstrak Daun dan Buah Mahkota Dewa pada Ileum Marmot Terpisah. *PT Mahkotadewa Indonesia*.
- Susilawati, Matsjeh, S., Pranowo, H. D., & Anwar, C. (2011). Antioxidant activity of 2,6,4'-trihydroxy-4-methoxy benzophenone from ethyl acetate extract of leaves of Mahkota Dewa (*Phaleria macrocarpa* (Scheff .) Boerl .). *Indo. J. Chem.*, 11(2), 180–185.
- Susilawati, Matsjeh, S., Pranowo, H. D., & Anwar, C. (2012). Macronone, a novel diepoxylignan from bark of Mahkota Dewa (*Phaleria macrocarpa* (Scheff .) Boerl .) and its antioxidant activity. *Indo. J. Chem.*, 12(1), 62–69.

- Sutiono, A. B., Faried, A., Arifin, M. Z., Wiriadisastra, K., & Ohta, T. (2008). Herbal Medicine Isolation, *Phaleria macrocarpa* for primary glioblastoma multiforme cells. *Annals of Epidemiology*, 18(9), 718.
- Tambunan, R. M., & Simanjuntak, P. (2006). Determination of chemical structure of antioxidant compound benzophenon glycoside from n-butanol extract of the fruits of Mahkota Dewa [*Phaleria macrocarpa* (Scheff) Boerl.]. *Majalah Farmasi Indonesia*, 17(4), 184–189.
- Tan, P. W., Tan, C. P., & Ho, C. W. (2011). Antioxidant properties : Effects of solid-to-solvent ratio on antioxidant compounds and capacities of Pegaga (*Centella asiatica*). *International Food Research Journal*, (18), 557–562.
- Tao, Y., Zhang, Z., & Sun, D. W. (2014). Kinetic modeling of ultrasound-assisted extraction of phenolic compounds from grape marc: influence of acoustic energy density and temperature. *Ultrasonics sonochemistry*, 21(4), 1461–9.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., & Hawkins Byrne, D. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19, 669–675.
- Thoo, Y. Y., Ho, S. K., Liang, J. Y., Ho, C. W., & Tan, C. P. (2010). Effects of binary solvent extraction system, extraction time and extraction temperature on phenolic antioxidants and antioxidant capacity from mengkudu (*Morinda citrifolia*). *Food Chemistry*, 120(1), 290–295.
- Tjandrawinata, R. R., Aripin, A., Arifin, P. F., & Rahmi, D. (2011). Extract of *Phaleria macrocarpa* as an antineoplastics antiinflammatory and antiangiogenic. U.S. Patent No US 2011/091395 A1. Washington D. C.: U. S. Patent and Trademark Office.
- Tjandrawinata, R. R., Nofiarny, D., Susanto, L. W., Hendri, P., & Clarissa, A. (2011). Symptomatic treatment of premenstrual syndrome and/or primary dysmenorrhea with DLBS1442, a bioactive extract of *Phaleria macrocarpa*. *International journal of general medicine*, 4, 465–476.
- Topallar, H., & Gecgel, U. (2000). Kinetics and Thermodynamics of Oil Extraction from Sunflower Seeds in the Presence of Aqueous Acidic. *Turk J Chem*, 24, 247–253.
- Triastuti, A., & Choi, J. W. (2008). Protective effects of ethyl acetate fraction of *Phaleria macrocarpa* (Scheff) Boerl. on oxidative stress associated with alloxan-induced diabetic rats. *Jurnal Ilmiah Farmasi*, 5(1), 9–17.
- Triastuti, A., Park, H., & Choi, J. W. (2009). *Phaleria macrocarpa* Suppress Nephropathy by Increasing Renal Antioxidant Enzyme Activity in Alloxan-Induced Diabetic Rats. *Natural Product Sciences*, 15(3), 167–172.

- Tsui, L., Fong, T., & Wang, I. (2012). YC-1 targeting of hypoxia-inducible factor-1 α reduces RGC-5 cell viability and inhibits cell proliferation. *Molecular Vision*, 18(June), 1594–1603.
- Türker, N., & Erdoğan, F. (2006). Effects of pH and temperature of extraction medium on effective diffusion coefficient of anthocyanin pigments of black carrot (*Daucus carota* var. L.). *Journal of Food Engineering*, 76(4), 579–583.
- Velickovic, D. T., Milenovic, D. M., Ristic, M. S., & Veljkovic, V. B. (2006). Kinetics of ultrasonic extraction of extractive substances from garden (*Salvia officinalis* L .) and glutinous (*Salvia glutinosa* L .) sage. *Ultrasonics sonochemistry*, 13, 150–156.
- Vetal, M. D., Lade, V. G., & Rathod, V. K. (2013). Extraction of ursolic acid from *Ocimum sanctum* by ultrasound: Process intensification and kinetic studies. *Chemical Engineering and Processing: Process Intensification*, 69, 24–30.
- Vuong, Q. V., Golding, J. B., Nguyen, M., & Roach, P. D. (2010). Extraction and isolation of catechins from tea. *Journal of separation science*, 33(21), 3415–28.
- Wahyuningsih, M. S. H., Mubarika, S., & Wahyuono, S. (2008). Effect of phalerin isolated from *Phaleria macrocarpa* (Scheff) Boerl. leaves on EVSA-T p53 protein expression in vitro. *Journal of Traditional Medicines*, 13(44), 83–89.
- Wang, P., Zhang, Q., Wang, Y., Wang, T., Li, X., Ding, L., & Jiang, G. (2010). Evaluation of Soxhlet extraction, accelerated solvent extraction and microwave-assisted extraction for the determination of polychlorinated biphenyls and polybrominated diphenyl ethers in soil and fish samples. *Analytica chimica acta*, 663(1), 43–8.
- Wang, X., Wei, Y., Yuan, S., Liu, G., Zhang, Y. L. J., & Wang, W. (2006). Potential anticancer activity of litchi fruit pericarp extract against hepatocellular carcinoma in vitro and in vivo. *Cancer letters*, 239, 144–50.
- Widyastuti, N., Widiyanti, T., & Listyawati, S. (2006). Terratogenic effect of *Phaleria macrocarpa* (Scheff.) Boerl. fruit extract on rat (*Rattus norvegicus* L.) Galur Winstar. *Bioteknologi*, 3(2), 56–62.
- Wongkittipong, R., Prat, L., Damronglerd, S., & Gourdon, C. (2004). Solid – liquid extraction of andrographolide from plants — experimental study , kinetic reaction and model. *Separation and Purification Technology*, 40, 147–154.
- Xi, J., He, L., & Yan, L. (2015). Kinetic modeling of pressure-assisted solvent extraction of polyphenols from green tea in comparison with the conventional extraction. *Food chemistry*, 166, 287–91.

- Xu, H. N., Huang, W. N., & He, C. H. (2008). Modeling for extraction of isoflavones from stem of *Pueraria lobata* (Willd.) Ohwi using n-butanol/water two-phase solvent system. *Separation and Purification Technology*, 62(3), 590–595.
- Yahya, F., Lu, T., Santos, R. C. D., Fryer, P. J., & Bakalis, S. (2010). Supercritical carbon dioxide and solvent extraction of 2-acetyl-1-pyrroline from Pandan leaf: The effect of pre-treatment. *The Journal of Supercritical Fluids*, 55(1), 200–207.
- Yaich, H., Garna, H., Besbes, S., Paquot, M., Blecker, C., & Attia, H. (2013). Effect of extraction conditions on the yield and purity of ulvan extracted from *Ulva lactuca*. *Food Hydrocolloids*, 31(2), 375–382.
- Yang, Y., Zhang, Z., Li, S., Ye, X., Li, X., & He, K. (2014). Synergy effects of herb extracts: Pharmacokinetics and pharmacodynamic basis. *Fitoterapia*, 92, 133–147.
- Yoshimi, N., Matsunaga, K., Katayama, M., Yamada, Y., Kuno, T., Qiao, Z., Hara, A., Yamahara, J., & Mori, H. (2001). The inhibitory effects of mangiferin, a naturally occurring glucosylxanthone, in bowel carcinogenesis of male F344 rats. *Cancer Letters*, 163, 163–170.
- Zeković, Z., Vidović, S., Vladić, J., Radosavljević, R., Cvejic, A., Elgndi, M. A., & Pavlić, B. (2014). Optimization of subcritical water extraction of antioxidants from *Coriandrum sativum* seeds by response surface methodology. *The Journal of Supercritical Fluids*, 95, 560–566.
- Zhang, S. Y., Zhang, Q. H., Zhao, W., Zhang, X., Zhang, Q., Bi, Y. F., & Zhang, Y. B. (2012). Isolation, characterization and cytotoxic activity of benzophenone glucopyranosides from Mahkota Dewa (*Phaleria macrocarpa* (Scheff.) Boerl). *Bioorganic & medicinal chemistry letters*, 22(22), 6862–6.
- Zhang, Y., Liu, Z., Li, Y., & Chi, R. (2014). Optimization of ionic liquid-based microwave-assisted extraction of isoflavones from *Radix puerariae* by response surface methodology. *Separation and Purification Technology*, 129, 71–79.
- Zhang, Y. B., Xu, X.-J., & Liu, H.-M. (2006). Chemical constituents from Mahkota dewa. *Journal of Asian Natural Products Research*, 8(2), 119–123.
- Zhao, S., & Zhang, D. (2014). Supercritical CO₂ extraction of Eucalyptus leaves oil and comparison with Soxhlet extraction and hydro-distillation methods. *Separation and Purification Technology*, 133, 443–451.