



**UNIVERSITI PUTRA MALAYSIA**

***DETERMINATION OF POLYCYCLIC AROMATIC HYDROCARBONS  
IN GRILLED CHICKEN AND BEEF AND THEIR BIOACCESSIBILITY  
USING IN VITRO HUMAN DIGESTION MODEL***

**ELLIYANA NADIA BT HAMIDI**

**FSTM 2015 33**



**DETERMINATION OF  
POLYCYCLIC AROMATIC HYDROCARBONS  
IN GRILLED CHICKEN AND BEEF AND THEIR BIOACCESSIBILITY USING  
*IN VITRO* HUMAN DIGESTION MODEL**

**By**

**ELLIYANA NADIA BT HAMIDI**

**Thesis Submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfilment of the Requirements for the  
Degree of Master of Science**

**November 2015**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the Degree of Master of Science

**DETERMINATION OF  
POLYCYCLIC AROMATIC HYDROCARBONS  
IN GRILLED CHICKEN AND BEEF AND THEIR BIOACCESSIBILITY USING  
IN VITRO HUMAN DIGESTION MODEL**

By

**ELLIYANA NADIA BT HAMIDI**

**November 2015**

**Chairman : Ahmad Faizal Abdull Razis, PhD**  
**Faculty : Food Science and Technology**

Bioaccessibility of contaminant in food is defined as the fraction of the contaminant that is released from food matrix in the gastrointestinal tract and thus become available for intestinal absorption. As an important human exposure pathway of contaminants, dietary intake of polycyclic aromatic hydrocarbons (PAHs) is of increasing concern for assessing health risk in human body. Therefore, the present study aimed to evaluate the bioaccessibility of PAHs in different parts of digestive system and to determine the effect of degree of doneness and fat content on the bioaccessibility of PAHs in grilled meat. In this study, the bioaccessibility of PAHs was assessed using *in vitro* human digestion model of grilled beef and chicken (satay). The samples were prepared by charcoal grilling and the levels of 15 priority-controlled PAHs in the samples were determined using high performance liquid chromatography (HPLC) equipped with photodiode array detector and fluorescence detector. The level of PAHs in the liquid portion of the digested samples were compared to the level in the undigested samples to determine the percentage of PAHs released i.e. its bioaccessibility. Limit of detection (LOD) of PAHs compounds were ranged from 0.025 to 5.0 ng/g, while limit of quantification (LOQ) were ranged from 0.075 to 15.0 ng/g. Recovery of PAHs was varied in the range 13.68 to 154.6% and 43.37 to 144.79% for beef and chicken samples, respectively. In general, there were significant differences ( $p < 0.05$ ) in PAHs bioaccessibility in different parts of digestive system with higher in stomach part as compared in mouth and small intestine parts of both grilled samples. The bioaccessibility of PAHs were ranged from 3.4% to 96.79% in stomach, 3.18% to 60.44% in mouth and 2.38% to 81.02% in small intestine of grilled meat; while 7.36% to 83.67% in stomach, 0.67% to 51.13% in mouth and 1.26% to 63.55% in small intestine of grilled chicken. There were

also significant differences ( $p < 0.05$ ) in bioaccessibility of PAHs with increased degree of doneness in both grilled beef and chicken. Bioaccessibility of PAHs was higher in well done cooked than in rare and medium cooked meat. Results indicated that the bioaccessibility of PAHs ranged from 2.38% to 43.75% in rare, 5.27 to 79.02% in medium, and 6.51% to 84.83% in well done cooked grilled beef; 0.67% to 8.93% in rare, 1.3% to 21.28% in medium and 2.95% to 21.87% in well done cooked grilled chicken. In addition, there was also a positive correlation ( $R^2 = 0.923$ ) between fat content and bioaccessibility of PAHs in both meat samples. In sum, this study pointed out the importance of food matrix (beef and chicken meat), elemental chemical properties (pH, enzyme) in different part of digestive system, physicochemical (polarity, water solubility) and cooking practices (rare, medium and well done) in the bioaccessibility of PAHs.

Abstraktesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENENTUAN POLISIKLIK AROMATIK HIDROKARBON  
DI DALAM AYAM DAN DAGING PANGGANG  
DAN BIOAKSES MENGGUNAKAN  
MODEL PENGHADAMAN MANUSIA SECARA *IN VITRO***

Oleh

**ELLIYANA NADIA BT HAMIDI**

**November 2015**

**Pengerusi: Ahmad Faizal Abdull Razis, PhD**  
**Fakulti : Sains dan Teknologi Makanan**

Bioakses pada kontaminan dalam makanan ditakrifkan sebagai pecahan kontaminan yang telah dibebaskan daripada matrik makanan di dalam saluran gastro-usus dan berupaya untuk penyerapan usus. Sebagai laluan penting pendedahan manusia kepada kontaminan, pengambilan polisiklik aromatik hidrokarbon (PAHs) dalam makanan adalah masalah yang semakin meruncing dalam menilai risiko kesihatan di dalam badan manusia. Oleh itu, kajian ini bertujuan untuk menilai bioakses PAHs pada bahagian yang berbeza dalam sistem penghadaman dan untuk menentukan kesan tahap kemasakan dan kandungan lemak terhadap bioakses PAHs dalam daging panggang. Dalam kajian ini, bioakses PAHs dinilai menggunakan model penghadaman manusia secara *in vitro* ke atas daging lembu dan ayam panggang (sate). Sampel telah disediakan melalui kaedah memanggang menggunakan arang dan aras 15 PAHs utama yang dikawal dalam sampel ditentukan menggunakan kromatografi cecair berprestasi tinggi (HPLC) dengan bantuan pengesan fotodiod pelbagai dan pengesan pendarfluor. Aras PAHs dalam bentuk cecair pada bahagian sampel yang telah dihadam dibandingkan dengan aras dalam sampel yang tidak dihadamkan untuk menentukan peratusan PAHs yang dibebaskan (pecahan boleh diakses), iaitu bioaksesnya. Had pengesanan (LOD) sebatian PAHs telah dianggarkan dari 0.025 kepada 5.0 ng/g, manakala had pengkuantifan (LOQ) telah dianggarkan dari 0.075 kepada 15.0 ng/g. Pemulihan PAHs masing-masing dianggarkan dalam julat 13.68 hingga 154.6% dan 43.37 hingga 144.79% untuk sampel daging lembu dan ayam. Secara umumnya, terdapat perbezaan yang ketara ( $p < 0.05$ ) pada bioakses PAHs di bahagian yang berbeza dalam sistem pencernaan dengan bioakses didapati tinggi dalam bahagian perut berbanding di bahagian mulut dan usus

ke atas kedua-dua sampel panggang. Bioakses PAHs telah dianggarkan dari 3.4% hingga 96.79% dalam perut, 3.18% hingga 60.44% dalam mulut dan 2.38% hingga 81.02% dalam usus di dalam daging panggang; manakala 7.36% hingga 83.67% dalam perut, 0.67% hingga 51.13% dalam mulut dan 1.26% hingga 63.55% dalam usus di dalam ayam panggang. Terdapat juga perbezaan yang ketara ( $p < 0.05$ ) dalam bioakses PAHs dengan penambahan tahap kadar kemasakan dalam kedua-dua daging lembu dan ayam panggang. Bioakses PAHs didapati meningkat dalam keadaan masak berbanding kurang dan sederhana masak daging panggang. Keputusan menunjukkan bioakses PAHs dianggarkan dari 2.38% hingga 43.75% dalam kurang masak, 5.27 hingga 79.02% dalam sederhana masak dan 6.51% hingga 84.83% dalam daging barbeku yang masak; manakala 0.67% hingga 8.93% dalam kurang masak, 1.3% hingga 21.28% dalam sederhana masak dan 2.95% hingga 21.87% dalam ayam barbeku yang masak. Tambahan pula, terdapat juga perkaitan positif ( $R^2 = 0.923$ ) antara kandungan lemak dan bioakses PAHs dalam kedua-dua sampel daging. Kesimpulannya, kajian ini menunjukkan kepentingan matriks makanan (daging lembu dan daging ayam), sifat-sifat unsur kimia (pH, enzim) di bahagian yang berbeza dalam sistem pencernaan, fizikokimia (kekutuban, keterlarutan air) dan amalan memasak (kurang masak, sederhana masak dan masak) dalam bioakses PAHs.

## ACKNOWLEDGEMENTS

In the Name of Allah, the Most Beneficent, the Most Merciful. All the praises and thanks be to Allah, the Lord of the 'Alamin'. Thanks Allah S.W.T for my ease way in doing and completing this research.

In preparing this thesis, I contacted with many people. They have contributed towards my understanding and thoughts. In particular, I wish to acknowledge my sincere gratitude to my main supervisor, Dr. Ahmad Faizal bin AbdullRazis, for his guidances and advices throughout my studies at Universiti Putra Malaysia. Appreciation is also extended to my co-supervisor, Professor Dr. JinapSelamat for serving on my guidance committee and all the helpful meeting and suggestions for my research. Not to forget, special appreciation to my previous supervisor, DrParvanehHajeb for her invaluable encouragements, guidances and advices in completing this research from the early stages until the end. Without her continued support, this thesis would not have been the same as presented here.

My great appreciation goes to my family for their tremendous supports, encouragements and love throughout my years at Universiti Putra Malaysia. Thanks for consistently be my side in my ups and down.

Sincere thanks are given to Tuan Haji Ismail from Research Instrument (RI) for his assistance with the HPLC analyses. I also would like to thank the lab assistants, for their assistance in supplying the materials for my research and work. Finally, my sincere thanks also extend to all my colleagues and others who have provided assistance at various occasions. Their views and tips are useful indeed. Unfortunately, it is not possible to list all of them in this limited space. I am very gratefull for all the contributions behind the preparation of the thesis.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Ahmad Faizal bin AbdullRazis, PhD**

Senior Lecturer  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**JinapSelamat, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

---

**BUJANG KIM HUAT, PHD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in theUniversiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Elliyana Nadia btHamid ,GS35387

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_

Name of Chairman of

Supervisory Committee:

Dr. Ahmad Faizal bin AbdullRazis

Signature: \_\_\_\_\_

Name of Member of

Supervisory Committee:

Prof.Dr.JinapSelamat

## TABLE OF CONTENTS

		Page
	<b>ABSTRACT</b>	i
	<b>ABSTRAK</b>	iii
	<b>ACKNOWLEDGEMENTS</b>	v
	<b>APPROVAL</b>	vi
	<b>DECLARATION</b>	viii
	<b>LIST OF TABLES</b>	xii
	<b>LIST OF APPENDICES</b>	xiii
	<b>LIST OF FIGURES</b>	xiv
	<b>LIST OF ABBREVIATIONS</b>	xv
<b>CHAPTER</b>		
<b>1</b>	<b>INTRODUCTION</b>	
	1.1 Overview	1
	1.2 Statement of problem	3
<b>2</b>	<b>LITERATURE REVIEW</b>	
	2.1 Polycyclic aromatic hydrocarbons (PAHs)	
	2.1.1 Characteristics of 15 PAHs	4
	2.1.2 Formation of PAHs	7
	2.1.3 Human exposure to PAHs	8
	2.1.4 Carcinogenicity of PAHs	11
	2.1.5 Sources of PAHs as Food Contaminant	12
	2.1.6 PAHs in Processed and Cooked Food	15
	2.1.7 Factors Affecting PAHs Concentration in Grilled Meat	17
	2.2 Bioaccessibility of Food Contaminant	
	2.2.1 Definition	19
	2.2.2 <i>In vitro</i> Digestion Model for Bioaccessibility	20
<b>3</b>	<b>MATERIALS AND METHODS / METHODOLOGY</b>	
	3.1 Materials	
	3.1.1 Chemicals	24
	3.1.2 Equipments	24
	3.1.3 Standard Solutions	24
	3.1.4 Meat Samples	25
	3.2 Methods	
	3.2.1 Sample Preparation of Grilled Beef and Chicken	25
	3.2.2 Marinate Ingredients	25
	3.2.3 Marinade Process	26
	3.2.4 Charcoal Grilling of Satay	26
	3.2.5 Calibration of PAHs standard	27

3.2.6	Linearity Test, Limit of Detection (LOD), Limit of Quantification (LOQ) and Recovery	27
3.2.7	Determination of the Bioaccessibility of PAHs for Grilled Meat	28
3.2.7.1	<i>In vitro</i> Human Digestion Model	28
3.2.7.2	Extraction and clean-up of PAHs from Grilled Meat	31
3.2.7.3	Determination of PAHs by HPLC	33
3.2.8	Fat Measurements of Grilled Meat	33
3.3	Calculation of Bioaccessibility of PAHs	34
3.3.1	Statistical analysis	34
<b>4</b>	<b>RESULTS AND DISCUSSION</b>	
4.1	Methods for PAHs Determination	35
4.1.1	Limit of Detection (LOD) and Limit of Quantification (LOQ)	35
4.1.2	Recovery of PAHs and Precision Study	40
4.2	Bioaccessibility of PAHs in Grilled Beef and Chicken Meat	42
4.2.1	Bioaccessibility of PAHs in Grilled Meat in Different Parts of Digestive System	44
4.2.2	Bioaccessibility of PAHs in Grilled Meat at Different Degree of Doneness	49
4.3	Effect of Fat Content on the Bioaccessibility of PAHs in Grilled Beef and Chicken	53
<b>5</b>	<b>CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	
5.1	Conclusion	56
5.2	Recommendations for Future Research	58
	<b>REFERENCES</b>	60
	<b>APPENDICES</b>	71
	<b>BIODATA OF STUDENT</b>	79
	<b>LIST OF PUBLICATIONS</b>	80

## LIST OF TABLES

Table	Page
2.1 Molecular weights and formula structures of the 16 EPA priority PAHs and European Union(EU) PAHs of concern in food	5
2.2 Groups of PAHs based on their molecular weight	7
2.3 Routes of human exposure to PAHs	9
2.4 Classifications of PAHs according to possibility to be carcinogenic to human	11
2.5 Sources of PAHs contamination in food products	14
2.6 The level of PAHs in different grilled meat dishes among three different cooking methods	16
2.7 Enzymes utilized in <i>in vitro</i> digestion model	21
2.8 Three-step phases of <i>in vitro</i> digestion model (mouth, stomach and small intestine) and its characteristics	22
3.1 Amount of ingredients in the satay marinades (for 1 kg meat)	26
3.2 Cooking parameters for satay prepared by three different degrees of doneness	27
3.3 Composition of digestive juices used per 100 ml of ultrapure water	29
3.4 Digestive phases and processes throughout the <i>in vitro</i> digestive system	31
3.5 Gradient program for HPLC analysis	33
4.1 Linear equations, limit of detection (LOD) and quantification (LOQ) obtained for quantification of PAHs	39
4.2 Precision and recovery of PAHs in Malaysian beef and chicken grilled satay, determined by HPLC	41
4.3 The percentage <sup>a</sup> of fat in grilled beef and chicken samples	53

## LIST OF APPENDICES

Appendix		Page
A.1	Bioaccessibility of PAHs (%) in mouth, stomach and small intestine of rare grilled beef and chicken samples	71
A.2	Bioaccessibility of PAHs (%) in mouth, stomach and small intestine of medium grilled beef and chicken samples	72
A.3	Bioaccessibility of PAHs (%) in mouth, stomach and small intestine of well done grilled beef and chicken samples	73
A.4	PAHs content (ng/g) before <i>in vitro</i> digestion, as well as bioaccessible of PAHs (%) after mouth digestion of grilled beef and chicken samples	74
A.5	PAHs content (ng/g) before <i>in vitro</i> digestion, as well as bioaccessible of PAHs (%) after stomach digestion of grilled beef and chicken samples	75
A.6	PAHs content (ng/g) before <i>in vitro</i> digestion, as well as bioaccessible of PAHs (%) after small intestine digestion of grilled beef and chicken samples	76
B.1	Grilling satay	77
B.2	Extraction of PAHs	77
B.3	HPLC for analysis of PAHs	78

## LIST OF FIGURES

Figure	Page
2.1 Estimated contribution (%) of nine PAHs in various food groups of Swedish diet	13
2.2 Schematic representation of an <i>in vitro</i> digestion model	28
3.1 Schematic diagram of the <i>in vitro</i> digestive procedure	30
3.2 Schematic diagram of the extraction method	32
4.1 HPLC/UV-FLD chromatograms of PAHs mix standard	36
4.2 Typical UV and fluorescence chromatograms of PAHs from the extraction of (a) digested grilled beef sample (b) digested grilled chicken sample	43
4.3 Bioaccessibility of PAHs (%) in different parts of digestive system (mouth, stomach and small intestine) of (a) rare, (b) medium and (c) well done grilled beef samples	45
4.4 Bioaccessibility of PAHs (%) in different parts of digestive system (mouth, stomach and small intestine) of (a) rare, (b) medium and (c) well done grilled chicken samples	46
4.5 Bioaccessibility of PAHs (%) between different degree of doneness (rare, medium and well done) in (a) mouth, (b) stomach and (c) small intestine of grilled beef samples	49
4.6 Bioaccessibility of PAHs (%) between different degree of doneness (rare, medium and well done) in (a) mouth, (b) stomach and (c) small intestine of grilled chicken samples	50
4.7 Relationship between the bioaccessibility of PAHs and the fat content in (a) grilled beef, (b) grilled chicken samples	53

## LIST OF ABBREVIATIONS

µg	Microgram
µm	Micrometer
µL	Microliter
λ <sub>ex</sub>	Excitation wavelength
λ <sub>em</sub>	Emission wavelength
A	Anthracene
Ace	Acenaphthene
ANOVA	Analysis of Variance
AOAC	Association of Official Analytical Chemists
B(a)A	Benzo(a)anthracene
B(a)P	Benzo(a)pyrene
B(b)F	Benzo(b)fluoranthene
B(c)F	Benzo(c)fluorene
B(g)P	Benzo(ghi)perylene
B(j)F	Benzo(j)fluoranthene
B(k)F	Benzo(k)fluoranthene
BSE	Bovine spongiform encephalopathy
Ch	Chrysene
CH <sub>2</sub> Cl <sub>2</sub>	Dichloromethane
D(ah)A	Dibenzo(a,h)anthracene
EFSA	European Food Safety Authority
F	Fluorene
Fl	Fluoranthene
FAO	Food and Agriculture Organization

GI	Gastrointestinal tract
HPLC	High Performance Liquid Chromatography
HPLC/UV-FLD	High Performance Liquid Chromatography/Ultraviolet-Fluorescence detector
HCl	Hydrochloride acid
IARC	International Agency Research on Cancer
i.e	“id est”, that is
IP	Indeno(1,2,3-cd)pyrene
KCl	Potassium chloride
kg	Kilogram
KSCN	Potassium thiocyanate
ml	Mililiter
Na	Naphthalene
NaCl	Sodium chloride
ng	Nanogram
NaOH	Sodium hydroxide
Na <sub>2</sub> SO <sub>4</sub>	Sodium sulfate
NaHCO <sub>3</sub>	Sodium bicarbonate
NaH <sub>2</sub> PO <sub>4</sub>	Sodium dihydrogen phosphate
NH <sub>4</sub> Cl	Ammonium chloride
P	Pyrene
PAH	Polycyclic aromatic hydrocarbon
pbb	Parts per billion
PCB	Polychlorinated biphenyls
Phe	Phenanthrene
PRS SPE	Propyl Sulfonic Acid Solid Phase Extraction

US-EPA

US Environmental Protection Agency

v/v

Volume/volume

WHO

World Health Organization



# CHAPTER 1

## INTRODUCTION

### 1.1 Overview

Diet contains variety of carcinogens which consists of chemicals present naturally, synthetic compounds and compounds produced during cooking (Viegas *et al.*, 2012). Worldwide, health authorities and consumers are increasingly concerned with the exposure to toxic chemical contaminants in the diet, such as heavy metals, persistent organic pollutants (POPs), veterinary drugs, mycotoxins as well as heat-generated contaminants (Marques *et al.*, 2011). Last decades, much attention is being paid to these heat-generated toxicants for examples polycyclic aromatic hydrocarbons (PAHs), heterocyclic amines (HCAs) and acrylamide; concerning their development, occurrence, reduction and effect on human health (Viegas *et al.*, 2012). Avoiding exposure to these toxicants is difficult; however, our eating habits, as well as cooking preparation and composition can limit the exposure to these food contaminants (Krul *et al.*, 2000).

Polycyclic aromatic hydrocarbons (PAHs) have been the subject of much concern as they are ubiquitous environmental contaminants (Phillips, 1999). PAHs are commonly found in meat, fish or other foods which cooked at high temperature. PAHs are formed from the incomplete combustion or pyrolysis of organic matter during industrial processing and various human activities (Onwukeme *et al.*, 2015). Factors responsible for the formation of PAHs include the temperature, time/level of doneness, method of cooking, fat content, type of heat source as well as direct contact with the heat source.

PAHs were believed to make a substantial contribution to the burden of cancers in humans as many of them are carcinogenic in experimental animals (Phillips, 1999). It has been reported that 35% of breast, pancreas, stomach and colon cancer incidences may be related to dietary factors (Krul *et al.*, 2000). Stomach, breast, colorectal, lung and prostate cancer have been demonstrated to have close relationship to exposure of significant levels of dietary and environmental PAHs which generally generated from well-done cooked meat consumption (Sinha *et al.*, 1999). Based on Malaysian Cancer Statistics in 2006, ten frequent cancers among population of Peninsular Malaysia are breast cancer (16.5%), colorectal cancer (13.2%), lung cancer (9.4%), cervical cancer (4.9%), nasopharyngeal cancer (4.5%), thyroid gland cancer (4.1%), liver cancer (3.6%), stomach cancer (3.6%), prostate gland cancer (3.4%) and lymphoma cancer (3.2%).

Nowadays, cooking procedure is one of the aspects not taken into account in food risk assessment of foodstuff, despite it strongly affects the availability of toxic elements in foodstuff. It was well-explained by Afzaninawati *et al.* (2013) who has pointed out that based on Malaysian diet, taste of cooking such as spicy, salty and sour as well as type of cooking for examples frying, grilling and smoking are the main factors contributing to colorectal cancer. According to Jahurul *et al.* (2010), one's daily diet may contain PAHs and HCAs resulting from meat and fish being cooked at high temperature. These heat-induced food toxicants are normally found in grilled and fried meat, fish and poultry (Jahurul *et al.*, 2010).

Marques *et al.* (2011) has also came out with the statistic that approximately 97% of the total daily intake of PAHs in humans derived from their diet. Although grilled food only contributed a small part of PAHs intake, people who regularly consume roasted, barbecued or grilled and even smoked food may have significant intake of PAHs. Viegas *et al.* (2012) has reported the uptake of grilled food can represent a large portion of PAHs up to ~120g or more per meal.

Grilled foods are popular both at home and even in restaurants, where this type of food may be detrimental to human health due to high concentration of carcinogens found in such products compared to food which prepared by alternative cooking methods (Sundararajan *et al.*, 1999). Some Malaysian popular dishes i.e various grilled meats such as satay (grilled beef), ayam bakar (grilled chicken) and ikan bakar (grilled fish) are prepared at high temperature that can produce marked differences in the concentrations of PAHs. A study by Farhadian *et al.* (2010) has found in Malaysian dishes, the maximum concentration of fluoranthene (Fl), one of frequently detected PAHs was reported to be as high as 106 ng/g in beef satay, 33.2 ng/g in chicken satay, 38.4 ng/g in grilled chicken and 11.8 ng/g in grilled fish. Such high values indicate high probability that Malaysians consumed PAHs in the level that may be harmful and can lead to cancer.

Grilled satay is a good example of food prepared at high temperature cooking method which applied grilling and roasting method. Satay has been used in this study since it is a popular dish in Malaysia as well as being well-known grilled dish in many other Southeast Asia countries like Indonesia, Thailand and Singapore and some Western countries. Similar to shish kebab, it consists of chunks or slices of dice-sized boneless meat for examples chicken, mutton, beef, pork and fish on skewers made from the midrib of coconut leaf or bamboo (Safzan, 2009; Jinap *et al.*, 2013). In general, satay is grilled over charcoal fire, and then served with variety of spiced seasoning depending on the satay recipe. In Southeast Asian countries, satay commonly been prepared by open charcoal grilling method (Safzan, 2009; Jinap *et al.*, 2013).

Excessive consumption of grilled meat might place the regular consumers at risk of cancer. Therefore, it is crucial to determine the level of PAHs present in food products and their bioaccessibility after digestion. In this sense, *in vitro* gastrointestinal model which is closely related to the human digestive system was employed. The bioaccessibility of PAHs were quantified in grilled beef and chicken satay digested in different part of digestion system at three degree of doneness. The effect of fat content and the bioaccessibility of PAHs was studied in order to correlate the amount of PAHs released during digestion and fat content of the meat.

## **1.2 Statement of problem**

Dietary style of grilled meat may increase exposure to PAHs and put the consumer at risk of cancer. Diet is one aspect of an individual's lifestyle that may be practically modified. In relation to this, the analysis in some of the most commonly consumed foods in Malaysia such as grilled meat for the occurrence of PAHs is crucial in order to estimate the exposure of the Malaysian population to these toxic compounds by ingestion of food and to identify practices that increase or decrease an individual's risk. Therefore, bioaccessibility of PAHs is needed in order to quantify the release of PAHs naturally formed during cooking and to assess the ingestion of PAHs from grilled meat which becomes available for intestinal absorption after digestion. In addition, previous studies indicated that concentrations of PAHs varies depending on different cooking conditions such as degree of doneness and were significantly correlated with fat content. In that sense, we evaluated to what extent the doneness level and fat content of meat can affects the bioaccessibility of PAHs.

Yet, so far, only few studies addressed the bioaccessibility of PAHs in food. On the other hand, this is the first extensive study on PAHs bioaccessibility in Malaysian food. Thus, the present study was designed to investigate the effect of cooking conditions by varying degree of doneness and different types of meat (chicken and beef) on PAHs formation during cooking and its bioaccessibility to human body. In this concern, together with the role of cooking method in the initiation and promotion of cancer, this present study aimed:

1. To evaluate the bioaccessibility of PAHs from grilled meat in different parts of digestive system.
2. To determine the effect of degree of doneness and fat content on the bioaccessibility of PAHs in grilled meat.

## REFERENCES

- Abou-Arab, A. A. K., Abou-Donia, M. A., El-Dars, F. M. S. E., Ali, O. I. M., Hossam, A. G. (2014). Detection of polycyclic aromatic hydrocarbons levels in Egyptian meat and milk after heat treatment by gas chromatography-mass spectrometry. *International Journal of Current Microbiology and Applied Science* 3(7): 294-305.
- Afzaninawati, S. Y., Zaleha, M. I., Shamsul, A. S. (2013). Perceptions of Malaysian colorectal cancer patients regarding dietary intake: A qualitative exploration. *Asian Pacific Journal of Cancer Prevention* Vol 14(2): 1151-1154.
- Akpambang, V. O. E., Purcaro, G., Lajide, L., Amoo, I. A., Conte, I. S., Moret, S. (2009). Determination of polycyclic aromatic hydrocarbons (PAHs) in commonly consumed Nigerian smoked/grilled fish and meat. *Food Additives & Contaminants: Part A* 26(7): 1096-1103.
- Alomirah, H., Al-Zenki, S., Al-Hooti, S., Zaghloul, S., Sawaya, W., Ahmed, N., Kannan, K. (2011). Concentrations and dietary exposure to polycyclic aromatic hydrocarbons (PAHs) from grilled and smoked foods. *Food Control* 22: 2028-2035.
- Al-Rashdan, A., Helaleh, M. I. H., Nisar, A., Ibtisam, A., Al-Ballam, Z. (2010). Determination of the levels of Polycyclic Aromatic Hydrocarbons Toasted Bread Using Gas Chromatography Mass Spectrometry. *International Journal of Analytical Chemistry*, doi:10.1155/2010/821216.
- Anderson, K. E., Sinha, R., Kulldorff, M., Gross, M., Lang, N. P., Barber, C., Harnack, L., DiMango, E., Bliss, R., Kadlubar, F. F. (2002). Meat intake and cooking techniques: associations with pancreatic cancer. *Mutation Research* 506-507: 225-231.
- Anderson, B. A., Breidenstein, B. B., Kauffman, R. G., Cassens, R. G., Bray, R. W. (1971). Effect of cooking on fatty acid composition of beef lipids. *Journal of Food Technology* 6 : 141-148.

- Chen, B. H., Lin, Y. S. (1997). Formation of polycyclic aromatic hydrocarbons during processing of duck meat. *Journal of Agriculture and Food Chemistry*, 45(4): 1394-1403.
- Chung, S. Y., Yetella R. R., Kim, J. S., Kwon, K., Kim, M. C., Min, D. B. (2011). Effects of grilling and roasting on the levels of polycyclic aromatic hydrocarbons in beef and pork. *Food Chemistry* 129: 1420-1426.
- Coles, L. T., Moughan, P. J., Darragh, A. J. (2005). *In vitro* digestion and fermentation methods, including gas production techniques, as applied to nutritive evaluation of foods in the hindgut of humans and other simple-stomached animals. *Animal Food Science and Technology*, 123-124: 421-444.
- Danyi, S., Brose, F., Brasseur, C., Schneider, Y., Larondelle, Y., Pussemier, L., Robbens, J., Saeger, S. D., Maghuin-Rogister, G., Scippo, M. (2009). Analysis of EU priority polycyclic aromatic hydrocarbons in food supplements using high performance liquid chromatography coupled to an ultraviolet, diode array or fluorescence detector. *Analytica Chimica Acta* 633: 293 – 299.
- Doremire, M. E.; Harmon, G. E.; Pratt, D. E. (1979). 3,4-Benzopyrene in charcoal grilled meats. *Journal of Food Science* 44:622-623.
- El-Badry, N. (2010). Effect of Household Cooking Methods and Some Food Additives on Polycyclic Aromatic Hydrocarbons (PAHs) Formation in Chicken Meat. *World Applied Sciences Journal* 9(9): 963-974.
- Falcó, G., Domingo, J. L., Llobet, J. M., Teixido, A., Casa, C., Müller, L. (2003). Polycyclic aromatic hydrocarbons in foods: Human exposure through the diet in Catalonia, Spain. *Journal of Food Protection*, Vol. 66, No. 12: 2325-2331.
- Farhadian, A., Jinap, S., Faridah, A., Zaidul, I. S. (2010). Determination of polycyclic aromatic hydrocarbon in grilled meat. *Food Control* 21: 606-610.

- Farhadian, A., Jinap, S., Hanifah, H. N., Zaidul, I. S. (2011). Effects of meat preheating and wrapping on the levels of polycyclic aromatic hydrocarbons in charcoal-grilled meat. *Food Chemistry* 124: 141-146.
- Farhadian, A., Jinap, S., Faridah, A., Zaidul, I. S. M. (2012). Effects of marinating on the formation of polycyclic aromatic hydrocarbons (benzo[a]pyrene, benzo[b]fluoranthene and fluoranthene) in grilled beef meat. *Food Control* 28: 420-425.
- Gomes, A., Roseiro, C., Santos, C. (2009). Determination of polycyclic aromatic hydrocarbons profile in Portugese traditional fermented sausage. V. Food Safety European Symposium, Berlim, Germany.
- Gomes, A, Santos, C., Almeida, J., Elias, M., Roseiro, L. C. (2013). Effect of fat content, casing type and smoking procedures on PAHs contents of Portugese traditional dry fermented sausages. *Food and Chemical Toxicology* 58: 369-374.
- Harris, K. L., Banks, L. D., Mantey, J. A., Huderson, A. C., Ramesh, A. (2013). Bioaccessibility of polycyclic aromatic hydrocarbons : relevance to toxicity and carcinogenesis. *Expert Opinion Drug Metabolisme Toxicology* 9(11): 1465-1480.
- Harris, K. B., Harberson, T. J., Savell, J. W., Cross, H. R., Smith, S. B. (1992). Influences of quality grade, external fat level and degree of doneness on beef steak fatty acids. *Journal of Food Composition and Analysis* 5: 84-89.
- Hu, J., Wu, F., Wu, S., Cao, Z., Lin, X., Wng, M. H. (2013). Bioaccessibility, dietary exposure and human risk assessment of heavy metals from market vegetables in Hong Kong revealed with an *in vitro* gastrointestinal model. *Chemosphere* 91: 455-461.
- Hur, S. J., Decker, E. A., McClements, D. J. (2009). Influence of initial emulsifier type on microstructural changes occurring in emulsified lipids during *in vitro* digestion. *Food Chemistry* 114: 253-62.

Hur, S. J., Lim, B. O., Decker, E. A., McClements, D. J. (2011). *In vitro* human digestion models for food applications. Food Chemistry 125: 1-12.

Intawongse, M., Dean, J. R. (2006). *In-vitro* testing for assessing oral bioaccessibility of trace metals in soil and food samples. Trend in Analytical Chemistry, 25: 9.

International Agency research for Cancer (IARC). (1987). Overall evaluations of carcinogenicity. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Lyon, France pp 440, Supplement 7.

International Agency for Research on Cancer (IARC). (2004). Overall evaluations of carcinogenicity: An updating of IARC monographs, Lyon, France Vols. 1-42, Supplement. 7.

International Agency Research for Cancer (IARC). (2010). Monographs on the evaluation of carcinogenic risk to humans. Some on-heterocyclic polycyclic aromatic hydrocarbons and some related exposures. Lyon, France Volume 92: 360-437.

Ishizaki, A., Saito, K., Hanioka, N., Narimatsu, S., Kataoka, H. (2010). Determination of polycyclic aromatic hydrocarbons in food samples by automated on-line in-tube solid-phase microextraction coupled with high-performance liquid chromatography-fluorescence detection. Journal of Chromatography A, 1217: 5555-5563.

Jägerstad, M., Skog, K. (2005). Genotoxicity of heat-processed foods. Mutation Research 574: 156-172.

Jahurul, M. H. A., Jinap, S., Zaidul, I. S. M., Sahena, F., Farhadian, A., Hajeb, P. (2013). Determination of fluoranthene, benzo(b)fluoranthene and benzo(a)pyrene in meat and fish product and their intake by Malaysian. Food Bioscience 1: 73-80.

Janoszka, B., Warzecha, L., Blaszczyk, U., Bodzek, D. (2004). Organic Compound Formed In Thermally Treated High-Protein Food Part I: Polycyclic Aromatic Hydrocarbons. *Acta Chromatographica*, 14: 115-128.

Jarvis, I. W. H., Dreijji, K., Mattson, A.; Jernström, B., Stenius, U. (2014). Interactions between polycyclic aromatic hydrocarbons in complex mixtures and implications for cancer risk assessment. *Toxicology* 321: 27 – 39.

JECFA. Summary and conclusions on the 64<sup>th</sup> meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA), Expert Committee on Food Additives, Food and Agricultural Organization (FAO), Rome/World Health Organization (WHO), Geneva, 2005.

Jinap, S., Mohd-Mokhtar, M. S., Farhadian, A., Hasnol, N. D. S., Jaafar, S. N., Hajeb, P. (2013). Effects of varying degrees of doneness on the formation of heterocyclic aromatic amines in chicken and beef satay. *Meat Science* 94: 202-207.

Jira, W., Ziegenhals, K., Speer, K. (2006). PAH in smoked meat products according to EU standards. *Fleischwirtschaft International* 4: 11-17.

Juan, C., Zinedine, A., Molto, J.C., Idrissi, L., Manes, J. (2008). Aflatoxins level in dried fruits and nuts from Rabat-sale area, Morocco. *Food Control*, 19: 849-853.

Knightes, C. D., Peters, C. A. (2006). Multisubstrate biodegradation kinetics for binary and complex mixtures of polycyclic aromatic hydrocarbons. *Environmental Toxicology and Chemistry*, 25: 1746-1756.

Krul, C., Luiten-Schuite, A., Bann, R., Verhagen, H., Mohn, G., Feron, V., Havenhaar, R. (2000). Application of a dynamic *in vitro* gastrointestinal tract model to study the availability of food mutagens, using heterocyclic aromatic amines as model compounds. *Food and Chemical Toxicology* 38: 783-792.

- Kulp, K. S., Fortson, S. L., Kniza, M. G., Felton, J. S. (2003). An *in vitro* model system to predict the bioaccessibility of heterocyclic amines from a cooked meat matrix. *Food and Chemical Toxicology* 4: 1701- 1710.
- Kumari, R., Chaturvedi, P., Ansari, N. G., Murthy, R. C., Patel, D. K. (2012). Optimization and validation of an extraction method for the analysis of polycyclic aromatic hydrocarbons in chocolate candies. *Journal of Food Science* Volume 71, Nr. 1.
- Ma, J. Xiao, R., Li, J., Yu, J., Zhang, Y., Chen, L. (2010). Determination of 16 polycyclic aromatic hydrocarbons in environmental water samples by solid-phase extraction using multi-walled carbon nanotubes as adsorbent coupled with gas chromatography-mass spectrometry. *Journal of Chromatography A*, 1217 : 5462-5469.
- Marchand, L. L., Hankin, J. H., Pierce, L. M., Sinha, R., Nerurkar, P. V., Franke, A. A., Wilkens, L. R., Kolonel, L. N., Donlon, T., Seifried, A., Custer, L. J., Lum-Jones, A., Chang, W. (2002). Well-done red meat, metabolic phenotypes and colorectal cancer in Hawaii. *Mutation Research* 506-507: 205-214.
- Marques, A., Lourenco, H. M., Nunes, M. L., Roseiro, C., Santos, C., Barranco, A., Rainieri, S., Langerholc, T., Cencic, A. (2011). New tools to assess toxicity, bioaccessibility and uptake of chemical contaminants in meat and seafood. *Food Research International* 44: 510 – 522.
- Martí-Cid, R.; Llobet, J. M.; Castell, V.; Domingo, J. L. (2008). Evolution of the dietary exposure to polycyclic aromatic hydrocarbons in Catalonia, Spain. *Food and Chemical Toxicology* 46: 3163-3171.
- Martorell, I., Perelló, G., Martí-Cid, R., Castell, V., Llobet, J. M., Domingo, J. L. (2010). Polycyclic aromatic hydrocarbons (PAHs) in foods and estimated PAH intake by the population of Catalonia, Spain, temporal trend. *Environment Introduction*. 36: 424-432.

Maulvault, A. L., Raquel, M., Cláudia, A., Helena M. L., Maria, L. N., Coelho, I., Langerholc, T., Marques, A. (2011). Bioaccessibility of Hg, Cd and As in cooked black scabbard fish and edible crab. Food and Chemical Toxicology 49: 2808-2815.

McClements, D. J., Decker, E. A., Park, Y. (2009). Controlling lipid bioavailability through physicochemical and structural approaches. Critical Reviews in Food Science and Nutrition 49: 48–67.

Mello'D, J. P. F. (2003). Food safety and contaminants and toxins (1<sup>st</sup> ed.). CABI Publishing.

National Cancer Registry..(2006)..Malaysia Cancer Statistics – Data and Figure Peninsular Malaysia. Ministry of Health, Malaysia.

Nielsen, S.S. (2003). Food analysis (3<sup>rd</sup> ed.). New York: Kluwer Academic/Plenum Publishers.

NIOSH Manual of Analytical Methods (NMAM), Fourth Edition. (1998). Polycyclic aromatic hydrocarbons by HPLC.

Omar, N. A., Praveena, S. M., Aris, A. Z., Hashim, Z. (2013). Bioavailability of heavy metals in rice using *in vitro* digestion model. International Food Research Journal 20: 2979 – 2985.

Omwukeme, V. I., Obijiofor, O. C., Asomugha, R .N., Okafor, F. A. (2015). Impact of Cooking Methods on the Levels pf Polycyclic Aromatic Hydrocarbons (PAHs) in Chicken Meat. IOSR Journal of Environment Science, Toxicology and Food Technology 9: 21-27.

Oomen, A. G., Rempelberg, C. J. M., Bruil, M. A., Dobbe, C. J. G., Pereboom, D. P. K. H., Sips, A. J. A. M. (2003). Development of an *in vitro* digestion model to estimating the bioaccessibility of soil contaminants. Archives of Environmental Contamination and Toxicology 44: 281-287.

- Palm, L. M. N., Derick, C. Yeboah, P. O., Quasie, W. J., Gorleku, M. A., Darko, A. (2011). Characterization of polycyclic aromatic hydrocarbon (PAHs) present in smoked fish from Ghana. *Advance Journal of Food Science and Technology* 3(5): 332-338.
- Pan, H., Cao, Y. (2010). Optimization of pretreatment procedures for analysis of polycyclic aromatic hydrocarbons in charcoal-grilled pork. *Analytical Letters* 43: 97-109.
- Parrish, F. C. JR., Olson, D. G., Miner, B. E., Rust, R. E. (1973). Effect of degree of marbling and internal temperature of doneness on beef rib steaks. *Journal of Animal Science* 37 : 430-434.
- Perelló, G., Martí-Cid, Castell, V., Llobet, J. M., Domingo, J. L. (2009). Concentrations of polybrominated diphenyl ethers, hexachlorobenzene and polycyclic aromatic hydrocarbons in various foodstuffs before and after cooking. *Food and Chemical Toxicology* 47: 709-715.
- Phillips, D. H. (1999). Polycyclic aromatic hydrocarbons in the diet. *Mutation Research* 443: 139-147.
- Plaza-Bolanós, P., Frenich, A. G., Vidal, J. L. M. (2010). Polycyclic aromatic hydrocarbons in food and beverages. *Analytical methods and trends. Journal of Chromatography A* 1217: 6303 – 6326.
- Public Health England. (2008). Polycyclic aromatic hydrocarbons (Benzo[a]pyrene): Toxicological Overview.
- Purcaro, G., Moret, S., S.Conte, L. (2013). Overview on polycyclic aromatic hydrocarbons : Occurrence, legislation and innovative determination in foods. *Talanta* 105: 292 – 305.
- Ramesh, A., Walker, S. A., Hood, D. B., Guillen, M. D., Shneider, K., Weyand, E. H. (2004). Bioavailability and risk assessment of orally ingested polycyclic aromatic hydrocarbons. *International Journal of Toxicology* 23: 301-333.

- Reinik, M., Tamme, T., Roasto, M., Juhkam, A., Tenno, T., Kiis, A. (2007). Polycyclic aromatic hydrocarbons (PAHs) in meat products and estimated PAH intake by children and the general population in Estonia. *Food Additives & Contaminants* 24: 429-437.
- Roseiro, L. C., Gomes, A., Santos, C. (2008). Polycyclic aromatic hydrocarbons profile in a Portuguese traditional meat product. IAFP's Fourth European Symposium on Food Safety. *Advancements in Food Safety*, 19-21 November, Lisbon, Portugal. P47.
- Safzan, M. (2009). Effect of cooking methods and conditions on heterocyclic amines content in satay and roasted marinated chicken. Master Thesis. Universiti Putra Malaysia.
- Silva, B. O., Adetunde O. T., Oluseyi T. O., Olayinka, K. O., Alo, B. I. (2011). Effects of the methods of smoking on the levels of PAH in some locally consumed fishes in Nigeria. *African Journal of Food Science* 5: 284-391.
- Šimko, P. (2002). Determination of polycyclic aromatic hydrocarbons in smoked meat products and smoke flavouring food additives. *Journal of Chromatography B*, 770: 3-18.
- Sinha, R., Rothman, N., Salmon, C. P., Knize, M. G., Brown, E. D., Swanson, C. A., Rhodes, D., Rossi, S., Felton, J. S., Levander, O. A. (1998). Heterocyclic amine content in beef cooked by different methods to varying degrees of doneness and gravy made from meat drippings. *Food and Chemical Toxicology* 36: 279-287.
- Sinha, R., Chow, W.H., Kulldorff, M., Denobile, J., Butler, J., Garcia-Closas, M., Weil, R., Hoover, R. N., Rothman, N. (1999). Well-done grilled red meat increases the risk of colorectal adenomas. *Cancer Research*. 59: 4320 - 4324.
- Sinha, R., Rothman, N. (1999). Role of well-done, grilled red meat, heterocyclic amines (HCAs) in the etiology of human cancer. *Cancer Letters*. 143: 189-194.

- Sundararajan, N., Sundararajan Nadife, M., Basel, R., Green, S. (1999). Comparison of sensory properties of hamburgers cooked by conventional and carcinogen reducing safe grill equipment. *Meat Science* 51: 289-295.
- Tang, X., Tang, L., Zhu, Y., Xing, B., Duan, J., Zheng, M. (2006). Assessment of the bioaccessibility of polycyclic aromatic hydrocarbons in soils from Beijing using an *in vitro* test. *Environmental Pollution* 140: 279-285.
- Versantvoort, C., Van de Kamp, E., Rempelberg, C. (2004). Development and applicability of an *in vitro* digestion model in assessing the bioaccessibility of contaminants from food. Report no. 320102002. National Institute for Public Health and the Environment, Bilthoven, The Netherlands.
- Viegas, O., Novo, P., Pinto, E., Pinho, O., Ferreira, I. M. P. L. V. O. (2012). Effect of charcoal types and grilling conditions on formation of heterocyclic aromatic amines (HAs) and polycyclic aromatic hydrocarbons (PAHs) in grilled muscle foods. *Food and Chemical Toxicology* 50: 2128 – 2134.
- Wang, H., Zhao, Y., Man, Y., Wong, C. K. C., Wong, M. (2011). Oral bioaccessibility and human risk assessment of organochlorine pesticides (OCPs) via fish consumption, using an *in vitro* gastrointestinal model. *Food Chemistry* 127: 1673 – 1679.
- Yang, L. S., Zhang, X. W., Li, Y. H., Li, H. R., Wang, Y., Wang, W. Y. (2012). Bioaccessibility and risk assessment of cadmium from uncooked rice using an *in vitro* digestion model. *Biological Trace Element Research* 145 (1): 81-86.
- Yu, Y., Li, J., Zhang, X., Yu, Z., Wiele, T. V., Han, S., Wu, M., Sheng, G., Fu, J. (2010). An assessment of the bioaccessibility of polybrominated diphenyl ethers in foods and the correlations of the bioaccessibility with nutrient contents. *Journal of Agricultural and Food Chemistry*. 58: 301 – 308.

Yu, Y., Huang, N., Zhang, X., Li, J., Yu, Z., Han, S., Lu, M., Wiele, T. V., Wu, M., Sheng, G., Fu, J. (2011). Polybrominated diphenyl ethers in food and associated human daily intake assessment considering bioaccessibility measured by simulated gastrointestinal digestion. *Chemosphere* 83: 152 – 160.

Yu, Y., Chen, L., Yang, D., Pang, Y., Zhang, S., Zhang, X., Yu, Z., Wu, M., Fu, J. (2012). Polycyclic aromatic hydrocarbons in animal-based foods from Shanghai : bioaccessibility and dietary exposure. *Food Additives & Contaminants A*, 29: 1465-1474.

Yusty, M. A. L., Daviña, J. L. C. (2005). Supercritical fluid extraction and high-performance liquid chromatography-fluorescence detection method for polycyclic aromatic hydrocarbons investigation in vegetable oil. *Food Control* 16: 59-64.