



UNIVERSITI PUTRA MALAYSIA

***SIMULTANEOUS DETERMINATION OF AFLATOXINS AND
OCHRATOXIN A IN SELECTED SPICES USING
HIGH PERFORMANCE LIQUID CHROMATOGRAPHY
WITH FLUORESCENCE DETECTOR***

WAN AINIZA BINTI WAN MUSTAPHA

FSTM 2015 38



**SIMULTANEOUS DETERMINATION OF AFLATOXINS AND
OCHRATOXIN A IN SELECTED SPICES USING
HIGH PERFORMANCE LIQUID CHROMATOGRAPHY
WITH FLUORESCENCE DETECTOR**

By

WAN AINIZA BINTI WAN MUSTAPHA

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfillment of the Requirement for the Degree of Master of Science**

October 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATION

This work is especially dedicated to my loving husband (Jailani Talib), who always given me the encouragement, support, patience, understanding and love throughout my research period.

Lots of love to my beautiful daughters (Syakirah Jaida, allahyarhamah Adriana Jaida and Karlissa Jaida). They are my guiding lights.

Also never forget my parents, a bundle of love and hugs to my mother, Raimah Baheran, also with respect and love to my late father, allahyarham Wan Mustapha Wan Taib.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfillment of the requirement for the degree of Master of Science

**SIMULTANEOUS DETERMINATION OF AFLATOXINS AND
OCHRATOXIN A IN SELECTED SPICES USING
HIGH PERFORMANCE LIQUID CHROMATOGRAPHY
WITH FLUORESCENCE DETECTOR**

By

WAN AINIZA BINTI WAN MUSTAPHA

October 2015

Chairman : Professor Jinap binti Selamat, PhD
Faculty : Food Science and Technology

Mycotoxins are found in a wide range of food making it possible for aflatoxins (AFs) and ochratoxin A (OTA) to occur simultaneously in spices and other commodities. Previously, immunoaffinity columns were designed to be used for single mycotoxin clean up and separate extractions was required if many mycotoxins were to be determined in a single food sample. Now, due to advances in technology, multi-mycotoxins immunoaffinity column were developed, making simultaneous determination possible. The objectives of this study were to develop a simultaneous determination method of AFs and OTA in selected spices, to validate the extraction procedure and to verify the method performance using curry mixture samples which were collected from retail markets. The simultaneous determination developed was a modification of a published method using multi-toxin immunoaffinity column and reverse-phased HPLC-FLD with photochemical derivatization system. Recoveries from spiked samples for AFB₁, AFB₂, AFG₁ and AFG₂ were ranged from 77.2% to 105.7% for coriander, chilli, cumin, curry mixture and soup mixtures; and from 63.6% to 90.1% for fennel, turmeric and kurma mixture. Recoveries for OTA ranged from 70.0% to 109.3% in coriander, chilli, cumin, fennel, turmeric and all spice mixtures. Acceptable recoveries are ranged from 70% to 110%. Relative standard deviation (RSD) for AFB₁, AFB₂, AFG₁, AFG₂ and OTA in all spices ranged from 3.18% to 10.81%. The method was validated using curry powder samples for selectivity, limit of detection (LOD), limit of quantification (LOQ), linearity and working range, accuracy (taken as recovery) and precision. The LOQ was found to be at 0.4 µg/kg for AFB₁/AFG₁, 0.2 µg/kg for AFB₂/AFG₂ and 0.5 µg/kg for OTA respectively. The correlation coefficients of >0.99 have been obtained for all AFs and OTA. The accuracy and precision for this method were acceptable. The recoveries were between 77.7% to 88.1% for all analytes, meanwhile the precision were < 7.3% for repeatability and < 8.7% for reproducibility. This method was verified when 30 retail samples were analysed and 63.3% were found to contain AFs >LOQ, and 70.0% contained OTA >LOQ. The current results indicated potential of the extraction procedure

to be used as a simultaneous determination for official food safety program and also for research purposes.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENENTUAN SERENTAK AFLATOKSIN DAN
OKRATOKSIN A DALAM REMPAH TERPILIH
MENGUNAKAN KROMATOGRAFI CECAIR BERPRESTASI
TINGGI DENGAN PENGESAN PNDARFLUOR**

Oleh

Wan Ainiza binti Wan Mustapha

Oktober 2015

Pengerusi : Profesor Jinap binti Selamat, PhD
Fakulti : Sains dan Teknologi Makanan

Mikotoksin boleh didapati dalam pelbagai jenis makanan, ini membolehkan aflatoksin (AFs) dan okratoksin A (OTA) untuk hadir serentak dalam rempah ratus dan komoditi-komoditi lain. Sebelum ini, turus *immunoaffinity* telah dibangunkan untuk pembersihan mikotoksin secara tunggal dan pengekstrakan berasingan perlu dilakukan sekiranya penentuan pelbagai jenis mikotoksin hendak dilakukan dalam satu jenis sampel makanan. Pada masa terkini, dengan kemajuan dalam teknologi, lajur *immunoaffinity* untuk pembersihan pelbagai mikotoksin telah dibangunkan, oleh itu penentuan serentak boleh dijalankan. Objektif kajian ini adalah untuk membangunkan satu kaedah penentuan serentak AFs dan OTA dalam rempah terpilih, untuk mengesahkan prosedur pengekstrakan dan untuk mengesahkan prestasi kaedah tersebut ke atas sampel campuran serbuk kari yang dikumpulkan dari pasaran runcit. Penentuan serentak yang telah dibangunkan adalah suatu pengubahsuaian dari kaedah yang diterbitkan menggunakan lajur *immunoaffinity* pelbagai toksin dan HPLC-FLD secara fasa terbalik, dengan sistem penerbitan secara fotokimia. Perolehan semula ke atas sampel yang telah ditambah AFB₁, AFB₂, AFG₁ dan AFG₂ adalah dari 77.2% hingga 105.7% untuk ketumbar, cili, jintan putih, campuran kari dan campuran sup; dan dari 63.6% hingga 90.1% untuk jintan manis, kunyit dan campuran kurma. Perolehan semula untuk OTA adalah dari 70.0% hingga 109.3% dalam ketumbar, cili, jintan putih, jintan manis, kunyit dan semua campuran rempah. Perolehan semula yang diterima adalah antara 70% hingga 110%. Sisihi piawai relative (RSD) untuk AFB₁, AFB₂, AFG₁, AFG₂ dan OTA dalam semua rempah telah adalah antara 3.18% hingga 10.81%. Kaedah ini telah divalidasi menggunakan sampel serbuk kari untuk kepilihan, had pengesanan (LOD), had kuantifikasi (LOQ), kelinearan, ketepatan (diambil sebagai perolehan semula) dan kejituan. Nilai LOQ diperolehi pada 0.4 µg/kg untuk AFB₁/AFG₁, 0.2 µg/kg untuk AFB₂/AFG₂ dan 0.5 µg/kg untuk OTA masing-masing. Pekali korelasi >0.990 telah diperolehi untuk semua AFs and OTA. Ketepatan dan kejituan bagi kaedah ini adalah diterima. Perolehan semula adalah antara 77.7% hingga 88.1% untuk semua

analit, manakala kejituan adalah $< 7.3\%$ for keterulangan dan $< 8.7\%$ untuk kebolehulangan semula. Kaedah ini telah diverifikasi dengan menganalisa 30 sampel pasaran 63.3% dijumpai mengandungi AFs $>LOQ$, dan 70.0% mengandungi OTA $>LOQ$. Keputusan kajian telah menunjukkan bahawa kaedah pengekstrakan ini mempunyai potensi untuk digunakan sebagai penentuan serentak untuk program rasmi keselamatan makanan dan juga untuk tujuan-tujuan penyelidikan.



ACKNOWLEDGEMENT

First and foremost, I would like to expressed greatest appreciation and gratitude to my supervisor Professor Jinap Selamat for her guidance, patience, encouragement and support throughout the research from start until completion. Also not to forget, I would also like to give my appreciation to my co-supervisor Dr. Maimunnah Sanny for her attention and support all along the course of work.

I am grateful and indebted to Ministry of Health (MOH), Malaysia for honoring me scholarship during my study. I am also thankful to the Food Safety and Quality Division and National Public Health Laboratory, MOH for funding of research materials and providing the facilities to conduct the research.

My deepest appreciation and love goes to my husband Jelani and children (Kirah, Anna and Lissa) for the endless support and understanding. Greatest thanks and appreciation to all staffs of Food Division, National Public Health Laboratory especially to Mycotoxin Unit (Sharmili, Shafie, Zuliana, Roshanida, Nur Azlina and Junizah) and HPLC team for assisting me in many aspects during the research. Last but not least, personal thanks to all my colleagues for your support and advise especially Rafidah, Norazlina, Jamilah, Zawiyah, Ghanthimathi, Zailina, Tuan Zainazor, Tosiah and Mazliza.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of the Master of Science. The members of the Supervisory Committee were as follows:

Jinap binti Selamat, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Maimunnah Sanny, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PHD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- This thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature : _____ Date: _____

Name and Matric No.: Wan Ainiza Binti Wan Mustapha, GS29237 _____

TABLES OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENT	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xv
 CHAPTER	
1 INTRODUCTION	1
 2 LITERATURE REVIEW	3
2.1 Aflatoxin	3
2.1.1 Occurrence	5
2.2 Ochratoxin A	8
2.2.1 Occurrence	9
2.3 Health effects of aflatoxins and ochratoxin A	10
2.4 Spices	11
2.4.1 Challenges in determination of aflatoxins and ochratoxin A in spices	12
2.4.2 Legislation on spices	13
2.5 Determination of aflatoxins and ochratoxin A	14
2.5.1 Extraction solvents	14
2.5.2 Purification using immunoaffinity column clean-up	15
2.5.3 Other purification techniques	16
2.6 Quantification of aflatoxins and ochratoxin A	19
2.6.1 High performance liquid chromatography	19
2.6.2 Others	20
2.7 Method Validation	20
2.7.1 Specificity	21
2.7.2 Limit of detection and quantification	21
2.7.3 Linearity and working range	21
2.7.4 Accuracy and trueness	22
2.7.5 Precision	22
 3 SELECTION OF EXTRACTION METHOD FOR SIMULTANEOUS DETERMINATION OF AFLATOXINS AND OCHRATOXINS IN SPICES	
3.1 Introduction	23
3.2 Materials and methods	24
3.2.1 Chemical and solvents	24
3.2.2 Preparation of stock solutions	24
3.2.3 HPLC condition	25
3.2.4 Preparation of samples	27

3.2.5	Selection of extraction methods	27
3.2.6	Data Analysis	32
3.3	Results and discussion	33
3.3.1	Selection of the extraction methods	33
3.3.2	Performance of extraction method C on single spice samples.	33
3.3.3	Performance of extraction method C on mixed spices samples	37
3.4	Conclusion	41
4	VALIDATION OF THE EXTRACTION METHOD FOR SIMULTANEOUS DETERMINATION OF AFLATOXINS AND OCHRATOXINS IN CURRY POWDER	
4.1	Introduction	42
4.2	Materials and methods	43
4.2.1	Chemicals and solvents	43
4.2.2	Preparation of stock solutions	43
4.2.3	Preparation of samples for validation	43
4.2.4	Collection of samples from retail market for verification	44
4.2.5	Extraction method	44
4.2.6	HPLC condition	44
4.2.7	Validation procedure	44
4.3	Results and discussion	46
4.3.1	Specificity and selectivity	46
4.3.2	Limit of detection and quantification	48
4.3.3	Linearity and working range	50
4.3.4	Accuracy and trueness	52
4.3.5	Precision	53
4.3.6	Verification on real samples	54
4.4	Conclusion	56
5	GENERAL CONCLUSION AND RECOMMENDATIONS	
5.1	General conclusions	57
5.2	Recommendation for future research	58
	REFERENCES	59
	APPENDICES	68
	BIODATA OF STUDENT	71
	LIST OF PUBLICATIONS	72

LIST OF TABLES

Table	Page
2.1 The physical and chemical properties of aflatoxins	3
2.2 Occurrence of aflatoxins in various spices	6
2.3 Occurrence of ochratoxin A in spices	9
3.1 Concentration of mixed working standard solutions for calibration	25
3.2 The concentration of spiking for recoveries study	25
3.3 Summary of HPLC condition	26
3.4 List of spices involved in curry powder production and its percentage	27
3.5 Recoveries of AFs and OTA in spiked coriander powder using two different extraction procedures.	33
3.6 Recoveries of AFs and OTA in spiked coriander powder using method C	34
3.7 Recoveries of AFs and OTA in spiked chili powder using method C	35
3.8 Recoveries of AFs and OTA in spiked cumin powder using method C	36
3.9 Recoveries of AFs and OTA in spiked fennel powder using method C	36
3.10 Recoveries of AFs and OTA in spiked turmeric powder using method C	37
3.11 Recoveries of simultaneous determination AFs and OTA in different spices using method C	38
3.12 Recoveries of AFs and OTA in curry mixture using method C	39
3.13 Recoveries of AFs and OTA in kurma mixture using method C	40
3.14 Recoveries of AFs and OTA in soup mixture using method C	40
3.15 Recoveries (%), standard deviation (SD) and relative standard deviation (RSD) in mixed spices using method C (n=6)	41
4.1 Formulation of curry powder mixture	43

4.2	Concentration of mixed working standard solutions for linearity study	45
4.3	The concentration of spiking of the mycotoxins for accuracy and precision study	46
4.4	LOD and LOQ of validated method	49
4.5	Comparison of limit of quantification (LOQ) in $\mu\text{g/kg}$	49
4.6	Linear equation, coefficient correlation (R^2) and working range	52
4.7	Mean recoveries and standard deviation for recovery study	53
4.8	The relative standard deviation (RSD) for precision study	54
4.9	Results of verification on the commercial samples	55

LIST OF FIGURES

Figure	Page
2.1 Chemical structures of AFB ₁ , AFB ₂ , AFG ₁ and AFG ₂ .	4
2.2 Chemical structures of ochratoxin A	8
2.3 Diagram of immunoaffinity column (IAC)	16
2.4 Steps of using SPE cartridges	17
2.5 Diagram of Multifunctional column (MFC)	18
2.6 Basic outline of an SFE system	18
3.1 Research design for preliminary testing	28
3.2 Design of evaluation of method performance on selected spices	29
3.3 Flow diagram of method A and method B	31
3.4 Flow diagram for method C	32
3.5 Chromatogram of (a) blank coriander sample and (b) spiked coriander sample at concentration of AFB ₁ and AFG ₁ at 40.0 µg/kg, AFB ₂ and AFG ₂ at 20 µg/kg, OTA at 50.0 µg/kg	35
3.6 Chromatogram of (a) blank curry powder sample and (b) spiked curry powder sample at concentration of AFB ₁ and AFG ₁ at 40.0 µg/kg, AFB ₂ and AFG ₂ at 20 µg/kg, OTA at 50.0 µg/kg.	39
4.1 Chromatogram of blank curry powder (a) and spiked curry powder (b) (spiked level at 40 µg/kg for AFB ₁ and AFG ₂ , 20 µg/kg for AFB ₂ and AFG ₂ , 50 µg/kg for OTA)	47
4.2 Chromatogram of blank standard solution (a) and mixed standard solution (b) (Concentration at 40 µg/kg for AFB ₁ and AFG ₂ , 20 µg/kg for AFB ₂ and AFG ₂ , 40 µg/kg for OTA)	48
4.3 The calibration curve for AFB ₁	50
4.4 The calibration curve for AFB ₂	50
4.5 The calibration curve for AFG ₁	51
4.6 The calibration curve for AFG ₂	51
4.7 The calibration curve for OTA	51

LIST OF ABBREVIATIONS

µg	microgram
AFs	Aflatoxin
AFB ₁	Aflatoxin B ₁
AFB ₂	Aflatoxin B ₂
AFG ₁	Aflatoxin G ₁
AFG ₂	Aflatoxin G ₂
EC	European Commission
EU	European Union
FAO	Food and Agricultural Organisation
FLD	Fluorescent detector
FTIR	Fourier transforms infrared spectroscopy
GC	Gas Chromatography
HPLC	High Performance Liquid Chromatography
IAC	Immunoaffinity column
IARC	International Agency of Research on Cancer
ISO	International Standard Organisation
kg	kilogram
L	litre
LC MS/MS	Liquid Chromatography Tandem Mass Spectrometry
LLC	Liquid-liquid extraction
LOD	Limit of Detection
LOQ	Limit of Quantification
OTA	Ochratoxin A
mL	milliliter

MFC	Multifunctional column
MPP	Maximum permitted proportion
MS-FID	MS-flame ionization detector
ng	nanogram
PVDF	Polyvinylidene fluoride
SFE	Supercritical fluid extraction
SPE	Solid phase extraction
TFA	Trifluoroacetic acid
QuEChERS	Quick, Easy, Cheap, Effective, Rugged and Safe
WHO	World Health Organization

CHAPTER 1

INTRODUCTION

Mycotoxins are found in a wide range of food, therefore aflatoxins (AFs) and ochratoxin A (OTA) could occur simultaneously in spices or any other commodities. Spices are widely used in our daily cooking making it possible for the mycotoxin to accumulate in the body if the spices are contaminated. As mycotoxins are highly stable compounds, cooking at normal temperature will not cause any or cause only little degradation. AFs and OTA are classified as Group 1 and Group 2 carcinogen respectively by International Agency of Research on Cancer (IARC). AFs are known to be potent toxic, carcinogenic, mutagenic, immunosuppressive agents, produced as secondary metabolites by the fungus *Aspergillus flavus* and *Aspergillus parasiticus* on variety of food products. Major members of AFs are aflatoxin B₁ (AFB₁), aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁) and aflatoxin G₂ (AFG₂). Food products contaminated with AFs include cereal (maize, sorghum, pearl millet, rice, wheat), oilseeds (groundnut, soybean, sunflower, cotton), spices (chillies, black pepper, coriander, turmeric, ginger), tree nuts (almonds, pistachio, walnuts, coconut) and milk. OTA is regarded as carcinogenic and nephrotoxic. It is mainly produced by *Aspergillus ochraceus* and *Penicillium verrucosum*. Food that is susceptible to OTA contamination are cereals, spices, coffee beans, cocoa beans, wine and beer.

Spices are commonly used in Malaysia to give flavours into food and are also well known for their medicinal value. Spices are mostly produced in tropical countries and often dried up in open air that could lead to fungus growth and cause mycotoxin contaminations. Currently there is no complete baseline data on mycotoxin contamination in such commodities in Malaysia. There are reports on the occurrence of AFs and OTA in European countries and in the Middle East. Spices that will be studied are in single form (individual spice) and also in mixed form (mixture of spices). These spices are chosen because they are the most commonly used in making curry, kurma and soup mixture.

Currently, there are very limited research conducted on the simultaneous determination of AFs and OTA in spices in Malaysia and also in other countries. Most of the references obtained provide studies on single mycotoxin analysis, either on AFs or OTA only. There are some studies on simultaneous occurrence of AFs and OTA but the determination or extraction were done separately on the same samples. Most of the researches conducted in the recent years very much concentrated on black pepper, red paprika, chilli powder or ginger. There are many more types of spices to be analysed because these spices are susceptible to mycotoxin contamination. Most mycotoxin extraction methods are matrix dependant meaning that one method may only suit one type of spices. Therefore, there will be a lot of extraction procedures to conduct and it will incur time and cost.

The complexity of spices due to high pigmentation, high lipid, essential oil compounds and its inhibitory effect on mycotoxin formation will interfere during the extractions of mycotoxin. In the past, immunoaffinity columns are designed for single mycotoxin clean up and this will require separate extractions, clean-up and quantification if many mycotoxin are to be analysed in a single sample. However due to advances in technology, multi-mycotoxin immunoaffinity column are developed making simultaneous determination possible.

Due to these limitations, there is a need to develop a simultaneous determination of AFs and OTA in spices using immunoaffinity column clean-up and quantification using high performance liquid chromatography. The Malaysian Food Regulation 1985 specified that the maximum level of AFs in processed groundnuts and other food shall not be greater than 10 ug/kg, while groundnut for further processing not greater than 15 ug/kg. Currently there is no regulatory limit set for OTA in spice in Malaysia. Legal limits should therefore be set to as low as reasonably achievable as both aflatoxins and ochratoxins A are carcinogen.

The main objective of this study is to select and validate the most suitable method of extraction that could simultaneously determine AFs and OTA in a variety of spices using one single extraction procedure. To accomplish this, the study will address the following specific objectives:

- i) To select the extraction methods for the simultaneous determination of AFs and OTA in spices.
- ii) To validate the extraction method for the simultaneous determination of AFs and OTA in curry powder mixture, and then to verify the method performance on real samples from the retail markets.

Hypothesis or the expected outcomes are as below:

- i) At least one of the extraction methods tested is able to simultaneously extract both AFs and OTA from variety of spices mixture.
- ii) To be able to use this extraction method as the standard method for simultaneous determination of AFs and OTA in spices.

REFERENCES

- Abdulkadar, A. H. W., Al-Ali, A. A., Al-Kildi, A. M., & Al-Jedah, J. H. (2004). Mycotoxins in food products available in Qatar. *Food Control*, 15(7), 543-548.
- Aflatoxins: Naturally Occurring Aflatoxins (Group1) Aflatoxin M₁ (Group 2B) (1993). *International Agency for Research on Cancer (IARC) - Summary and Evaluation, Volume 56*, 245.
- Afsah-Hejri, L., Jinap, S., & Mirhosseini, H. (2012). Ochratoxin A quantification: newly developed HPLC conditions. *Food Control*, 23(1), 113-119.
- Afsah-Hejri, L., Jinap, S., Arzandeh, S., & Mirhosseini, H. (2011). Optimization of HPLC conditions for quantitative analysis of aflatoxins in contaminated peanut. *Food Control*, 22(1), 381-388.
- Ahn, J., Kim, D., Jang, H., Kim, Y., Shim, W., & Chung, D. (2010). Occurrence of ochratoxin A in Korean red paprika and factors to be considered in prevention strategy. *Mycotoxin Research*, 26, 279-286.
- Ali, N., Hashim, N.H., Saad, B., Safan, K., Nakajima, M., & Yoshizawa, T. (2005). Evaluation of a method to determine the natural occurrence of aflatoxins in commercial traditional herbal medicines from Malaysia and Indonesia. *Food and Chemical Toxicology*, 43, 1763-1772.
- Al-juraini, A.A. (2011). Natural occurrence of fungi and aflatoxins of cinnamon in the Saudi Arabia. *African Journal of Food Science*, 5(8), 460-465.
- Almela, L., Rabe, V., Sanchez, B., Torella, F., Lopez-Perez, J.P., Gabaldon, J.A., & Guardiola, L. (2007). Ochratoxin A in red paprika: Relationship with the origin of the raw material. *Food Microbiology*, 24, 319-327.
- Anklam, E., Stroka, J., & Boenke, A. (2002). Acceptance of analytical methods for implementation of EU legislation with a focus on mycotoxins. *Food Control*, 13, 173-183.
- Ardic, M., Karakaya, Y., Atasevar, M., & Durmaz, H. (2008). Determination of aflatoxin B₁ levels in deep-red ground pepper (isot) using immunoaffinity column combined with ELISA. *Food and Chemical Toxicology*, 46, 1596-1599.
- Aydin, A., Emin Erkan, M., Başkaya, R., & Ciftcioglu, G. (2007). Determination of aflatoxin B₁ levels in powdered red pepper. *Food Control*, 18(9), 1015-1018.
- Binder, E.M., Tan, L.M., Chin, L.J., Handl, J., & Richard, J. (2007). Worldwide occurrence of mycotoxins in commodities, feed and feed ingredients. *Animal Feed Science and Technology*, 137, 265-282.

- Bircan, Cavit. (2005). The determination of aflatoxins in spices by immunoaffinity column extraction using HPLC. *International Journal of Food Science and Technology*, 40, 929-934.
- Bircan, C., Barringer, S.A., Ulken, U., & Pehlivan, R. (2008). Aflatoxin levels in dried figs, nuts and paprika for export from Turkey. *International Journal of Food Science and Technology*, 43, 1492-1498.
- Boonzaaijer, G., van Osenbruggen, W.A., Kleinnijenhuis, A.J., & van Dongen, W.D. (2008). An exploratory investigation of several mycotoxins and their natural occurrence in flavour ingredients and spices, using a multi-mycotoxin LC-MS/MS method. *World Mycotoxin Journal*, 1(20), 167-174.
- Brera, C., Debegnach, F., De Santis, B., Pannunzi, E., Berdini, C., Prantera, E., Miraglia, M. (2011). Simultaneous determination of aflatoxins and ochratoxin A in baby foods and paprika by HPLC with fluorescence detection: A single-laboratory validation study. *Talanta*, 83(5), 1442-1446.
- Bryden, W.L. (2007). Mycotoxins in the food chain: human health implications. *Asia Pacific Journal of Clinical Nutrition*, 16(1), 95-101.
- Buttinger, Gerhard. (2010). Aflatoxin measurements: how HPLC methods have evolved over the last 20 years. *Food Additives and Contaminants*, 27(9), 1266-1272.
- Chan, D., MacDonald, S.J., Boughtflower, V., & Brereton, P. (2004). Simultaneous determination of aflatoxins and ochratoxin A in food using a fully automated immunoaffinity column clean-up and liquid chromatography-fluorescence detection. *Journal of Chromatography A*, 1059, 13-16.
- Chao, T.C. (1991). Forensic medicine: the changing roles for the challenges ahead. *Malaysian Journal of Pathology*, 13(1), 1-4.
- Cho, Sung-Hye., Lee, Cyhang-Hee., Jang, Mi-Ran., Son, Young-Wook., Lee, Sang-Mok., Choi, In-Sun., Kim, So-Hee., & Kim, Dai-Byung. (2008). Aflatoxins contamination in spices and processed spice products commercialized in Korea. *Food Chemistry*, 107, 1283-1288.
- Commission Regulation (EC) No 401/2006 of 23 February 2006 laying down the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs.
- Commission Regulation (EU) No 105/2010 of 5 February 2010 amending Regulation EC No 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards ochratoxin A.

- Creppy, E.E. (2002). Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicology Letters*, 127, 19-28.
- Curticapean, A., Toma, F., Tarcea, M., Curticapean, M., Samarghitan, V., Pop, I.A., & Gulea, A. (2011). HPLC method validation for simultaneous determination of three mycotoxins from corn seeds. *Croatia Chemical Acta*, 84(3), 413-417.
- D'Ovidio, K., Trucksess, M., Weaver, C., Horn, E., McIntosh, M., & Bean, G. (2006). Aflatoxins in ginseng roots. 174-180.
- Duarte, S.C., Pena, A., & Lino, C.M. (2009). Ochratoxin A non-conventional exposure sources – a review. *Microchemical Journal*, 3, 115-120.
- Ediage, E.N., Di Mavungu, J.D., Monbaliu, S., Van Peteghem, C., & De Saeger, S. (2011). A validated multianalyte LC-MS/MS method for quantification of 25 mycotoxins in cassava flour, peanut cake and maize samples. *Journal of Agricultural and Food Chemistry*, 59, 5173-5180.
- Erdogan, Ahmet. (2004). The aflatoxin contamination of some pepper types sold in Turkey. *Chemosphere*, 56, 321-325.
- El tawila, M.M., Neamatallah, A., & Serdar, S.A. (2013). Incidence of aflatoxins in commercial nuts in the holy city of Mekkah. *Food Control*, 29, 121-124.
- Eurachem Guide: The Fitness for Purpose of Analytical Methods: A Laboratory Guide to Method Validation and Related Topics (1998).
- Fazekas, B., Tar, A., & Kovacs, M. (2005). Aflatoxin and ochratoxin A content of spices in Hungary. *Food Additives and Contaminants*, 22(90), 856-863.
- Garcia-Villanova, F.J., Cordon, C., Paramas, A.M.G., Aparicio, P., & Rosales, M.E.G. (2004). Simultaneously immunoaffinity column cleanup and HPLC analysis of aflatoxins and ochratoxin A in Spanish bee pollen. *Journal of Agricultural and Food Chemistry*, 52, 7235-7239.
- Golge, O., Hepsag, F., & Kabak, B. (2013). Incidence and level of aflatoxin contamination in chili commercialized in Turkey. *Food Control*, 33, 514-520.
- Gonzalez, A.G., & Herrador, M.A. (2007). A practical guide to analytical method validation, including measurement uncertainty and accuracy profiles. *Trends in Analytical Chemistry*, 26(3), 227-238.
- Goryacheva, I.Y., De Saeger, S., Lobeau, M., Eremin, S.A., Barna-Vetro, I., & Van Peteghem, C. (2006). Approach of ochratoxin A fast screening in spices using clean-up tandem immunoassay columns with confirmation by high performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS). *Analytica Chimica Acta*, 577, 38-45.

- Hierro, J.M.H., Garcia-Villanova, R.J., Torrero, P.R., & Fonseca, I.M.T. (2008). Aflatoxins and ochratoxin A in red paprika for retail sale in Spain: Occurrence and evaluation of a simultaneous analytical method. *Journal of Agricultural and Food Chemistry*, 56, 751-756.
- Hitokoto, H., Morozumi, S., Wauke, T., Sakai, S., & Kurata, H. (1980). Inhibitory effects of spices on growth and toxin production of toxigenic Fungi. *Applied and Environmental Microbiology*, 39(4), 818-822.
- Horwitz, W., & Albert, R. (2006). The Horwitz Ration (HorRat): a useful index of method performance with respect to precision. *Journal of AOAC International*, 89(4), 1095-1109.
- How to Meet ISO 17025 Requirements for Method Verification (2007) ALACC Guide, AOAC International.
- Hussein, H.S., & Brasel, J.M. (2001). Toxicity, metabolism, and impact of mycotoxins on human and animals. *Toxicology*, 167, 101-134.
- Ibanez-Vea, M., Corcuera, L.A., Remiro, R., Murillo-Arbizu, M.T., Gonzales-Penaz, E., & Lizarraga, E. (2011). Validation of a UHPLC-FLD method for the simultaneous quantification of aflatoxin, ochratoxin A and zearalenone in barley. *Food Chemistry*, 137, 351-358.
- Iha, M.H., & Trucksess, M.W. (2010). Aflatoxins and ochratoxin A in tea prepared from naturally contaminated powdered ginger. *Food Additives and Contaminants*, 27(8), 1142-1147.
- Imperato, R., Campone, I., Piccinelli, A.L., Veneziano, A., & Rastrelli, L. (2011). Survey of aflatoxins and ochratoxin A contamination in food products imported in Italy. *Food Control*, 22, 1905-1910.
- Iqbal, S.Z., Asi, M.R., Zuber, M., & Akhtar, J. (2013). Natural occurrence of aflatoxins and ochratoxin A in commercial chilli and chilli sauce samples. *Food Control*, 30(2), 621-625.
- Iqbal, S.Z., Paterson, R.R.M., Bhatti, I.A., & Asi, M.R. (2011). Aflatoxin concentration in chillies vary depending on variety. *Mycoscience*, 52(2), 296-299.
- Jalili, M., & Jinap, S. (2012). Natural occurrence of aflatoxins and ochratoxin A in commercial dried chili. *Food Control*, 24, 160-164.
- Kabak, Bulent. (2012). Determination of aflatoxins and ochratoxin A in retail cereal products from Turkey by high performance liquid chromatography with fluorescence detection. *Food Control*, 28, 1-6.
- Koppen, R., Koch, M., Siegel, D., Merkel, S., Maul, R., & Nehls, I. (2010). Determination of mycotoxins in foods: current state of analytical methods and limitations. *Applied Microbiology and Biotechnology*, 86, 1595-1612.

- Kursun, O., & Mutlu, A.G. (2010). Aflatoxin in spices marketed in the West Meditternenian region of Turkey. *Journal of Animal and Veterinary Advances*, 9(23), 2979-2981.
- Llewellyn, G.C., Mooney, R.L., Cheatle, T.F., & Flannigan, B. (1992). Mycotoxin contamination of spices – An update. *International Biodetermination & Biodegradation*, 29, 111-121.
- Marasas, W.F.O., Gelderblom, W.C.A., Shepard, G.S., & Vismer, H.F. (2008). Mycotoxin : A Global Problem. In Leslie, J.F., Bandyopadhyay, R., & Visconti, A. (Eds.), *Mycotoxins: Detection Methods, Management, Public Health and Agricultural Trade* (pp.171-183). Oxfordshire : CAB International
- Monaci, L., & Palmisano, F. (2004). Determination of ochratoxin A in foods: state-of-the-art and analytical challenges. *Anal Bioanal Chem*, 378, 96-103.
- Monbaliu, S., Wu, A., Zhang, D., Van Peteghem, C., & De Saeger, S. (2010). Multimycotoxin UPLC-MS/MS for tea, herbal infusions and the derived drinkable products. *Journal of Agricultural and Food Chemistry*, 58, 12664-12671.
- Monbaliu, S., Van Poucke, C., Van Peteghem, C., Van Poucke, K., Heugens, K., & De Saeger, S. (2009). Development of multi-mycotoxin liquid chromatography/tandem mass spectrometry method for sweet pepper analysis. *Rapid Communications in Mass Spectrometry*, 23, 3-11.
- Ochratoxin A (Group 2B) (1993). *International Agency for Research on Cancer (IARC) - Summary and Evaluation, Volume 56*, 489.
- O'Riordan, M.J., & Wilkinson, M.G. (2008). A survey of the incidence and level of contamination in a range of imported spice preparations on the Irish retail market. *Food Chemistry*, 107, 1429-1435.
- O'Riordan, M.J., & Wilkinson, M.G. (2009). Comparison of analytical methods for aflatoxin determination in commercial chilli spice preparation and subsequent development of an improved method. *Food Control*, 20, 700-705.
- Ozbey, F., & Kabak, B. (2012). Natural co-occurrence of aflatoxins and ochratoxin A in spices. *Food Control*, 28, 354-361.
- Pascale, M., & Visconti, A. (2008). Overview of detection methods for Mycotoxins. In Leslie, J.F., Bandyopadhyay, R., & Visconti, A. (Eds.), *Mycotoxins: Detection Methods, Management, Public Health and Agricultural Trade* (pp.171-183). Oxfordshire : CAB International.
- Paterson, R. Russel M. (2007). Aflatoxins contamination in chilli samples from Pakistan. *Food Control*, 18, 817-820.

- Peraica, M., Radic, B., Lucic, A., & Pavlovic, M. (1999). Toxic effects of mycotoxins in humans. *Bulletin of the World Health Organization*, 77, 754-766.
- Petzinger, E., & Weidenbach, A. (2002). Mycotoxins in the food chain: the role of ochratoxins. *Livestock Production Science*, 76, 245-250.
- Pietri, A., Rastelli, S.R., Mulazzi, A., & Bertuzzi, T. (2012). Aflatoxins and ochratoxin A in dried chestnuts and chestnut flour produced in Italy. *Food Control*, 25, 601-606.
- Rahman, A., and Jinap, S. (2010). Validation of the procedure for the simultaneous determination of aflatoxins ochratoxin A and zearalenone in cereals using HPLC-FLD. *Food Additives and Contaminants*, 27(12), 1683-1693.
- Rahmani, A., Jinap, S., Khatib, A., & Tan, C.P. (2013). Simultaneous determination of aflatoxins, ochratoxin A, and zearalenone in cereals using a validated RP-HPLC Method and PHRED derivatization system. *Journal of Liquid Chromatography & Related Technologies*, 36:5, 600-617.
- Rahmani, A., Selamat, J., Soleimany, F. (2011). Optimization and validation of a HPLC method for simultaneous determination of aflatoxin B₁, B₂, G₁, G₂, ochratoxin A and zearalenone using an experimental design. *Food Additives and Contaminant*, 28(7), 902-912.
- Reddy, K.R.N., Farhana, N.I., & Salleh, B. (2011). Occurrence of *Aspergillus* spp. and aflatoxin B₁ in Malaysian foods used for human consumption. *Journal of Food Science*, 00, T1-T6.
- Reif, K., & Wolfgang, M. (1995). Determination of aflatoxins in medicinal herbs and plant extract. (1995). *Journal of Chromatography A*, 692, 131-136.
- Richard, J.L. (2007). Some major mycotoxins and their mycotoxicoses – An overview. *International Journal of Food Microbiology*, 119, 3-10.
- Riwei Wei, Feng Qiu, Weijun Kong, Jianhe Wei, Meihua Yang, Zuliang Luo, Jiepan Qin & Xiujing Ma. (2013). Co-occurrence of aflatoxins B₁, B₂, G₁, G₂ and ochratoxin A in *Glycyrrhiza uralensis* analyzed by HPLC-MS/MS. *Food Control*, 32, 216-221.
- Romagnoli, B., Menna, V., Gruppioni, N., & Bergamini, C. (2007). Aflatoxins in spices, aromatic herbs, herb-teas and medicinal plants marketed in Italy. *Food Control*, 18, 697-701.
- Salem, N.M., & Ahmad, R. (2010). Mycotoxins in food from Jordan: preliminary survey. *Food Control*, 21, 1099-1103.

- Santos, L., Marin, S., Mateo, E.M., Gil-Serna, J., Valle-Algarra, F.M., Patino, B., & Ramos, A.J. (2011). Mycobiota and co-occurrence of mycotoxins in Capsicum powder. *International Journal of Food Microbiology*, 151, 270-276.
- Santos, L., Marin, S., Sanchis, V., & Ramos, A.J. (2010). Co-occurrence of aflatoxins, ochratoxin A and zearalenone on the Spanish market. *Food Chemistry*, 122, 826-830.
- Saxena, J., & Mehrotra, B.S. (1989). Screening of spices commonly marketed in India for natural occurrence of Mycotoxins. *Journal of Food Composition and Analysis*, 2, 286-292.
- Schweiggert, U., Carle, R., & Schieber, A. (2007). Conventional and alternative processes for spice production – a review. (2007). *Trend in Food Science & Technology*, 18, 260-268.
- Sedmikova, M., Reisnerova, H., Dufkova, Z., Barta, I., & Jilek, F. (2001). Potential Hazard of simultaneous occurrence of aflatoxin B1 and ochratoxin A. *Veterinary Medical – Czech*, 46(6), 169-174.
- Senyuva, H., Gilbert, J., ozcan, S., & Gurel, N. (2008). Rapid LC and LC/MS routine analysis for mycotoxins in foods. *World Mycotoxin Journal*. 1(3), 229-235.
- Set, E., & Erkmen, O. (2010). The aflatoxin contamination of ground pepper and pistachio nuts sold in Turkey. *Food and Chemical Toxicology*, 48, 2532-2537.
- Shephard, G.S., Berthiller, F., Burdaspal, P.A., Crews, C., Jonker, M.A., Krska, R., MacDonald, S., Malone, R.J., Maragos, C., Sabino, M., Solfrizzo, M., Van Egmond, H.P., & Whitaker, T.B. (2012). *Developments in mycotoxin analysis: an update for 2010-2011*. *World Mycotoxin Journal*, 5(1), 3-30.
- Shundo, L., Almeida, A.P., Alaburda, J., Lamardo, L.C.A., Navas, S.A., Ruvieri, V., & Sabino, M. (2009). Aflatoxins and ochratoxin A in Brazilian paprika. *Food Control*, 20, 1099-1102.
- Siu-Po, Ip., & Chun-Tao, Che. (2006). Determination of aflatoxins in Chinese medicinal herbs by high-performance liquid chromatography using immunoaffinity column cleanup improvement of recovery. *Journal of Chromatography A*, 1135, 241-244.
- Škrbić, B., Koprivic, S., & Godula, M. (2013). Validation of a method for determination of mycotoxins subjected to the EU regulations in spices: The UHPLC-HESI-MS/MS analysis of the crude extracts. *Food Control*, 31, 461-466.
- Smith, J.S. (2010). Food Analysis. *Evaluation of Analytical Data* (pp.53-67). Springer Science.

Steyn, P.S., (1995). Mycotoxin, general view, chemistry and structure. *Toxicology Letters*, 82/83, 843-851.

Tajkarimi, M.M., Ibrahim, S.A., & Cliver, D.O. (2010). Antimicrobial herb and spice compound in food. *Food Control*. 21, 1189-1218.

Thirumala-Devi, K., Mayo, M.A., Reddy, G., Emmanuel, K.E., & Reddy, D.V.R. (2001). Occurrence of ochratoxin A in black pepper, coriander, ginger and turmeric in India. *Food Additives and Contaminants*, 18(9), 830-835.

Thompson, M., Ellison, S.L.R., & Wood, R. (2002). Harmonized guidelines for single-laboratory validation of methods of analysis. *International Union of Pure and Applied Chemistry*. Vol. 74, No. 5, 835-855.

Trucksess, M.W., Weaver, C.M., Oles, C.J., Fry Jr, F.S., & Noonan, G.O. (2008). Determination of aflatoxin B₁, B₂, G₁, and G₂ and ochratoxin A in Ginseng and Ginger by multitoxin immunoaffinity column cleanup and liquid chromatography quantification: Collaborative study. *Journal of AOAC International*, 91(3), 511-523.

Trucksess, M.W., & Scott, P.M. (2008). Mycotoxins in botanicals and dried fruits: a review. *Food Additives and Contaminants*, 25(2), 181-192.

Turner, N. W., Subrahmanyam, S., & Piletsky, S. A. (2009). Analytical methods for determination of mycotoxins: A review. *Analytica Chimica Acta*, 632(2), 168-180.

Waltking, A.E., & Wilson, D. (2006). Liquid chromatographic analysis of aflatoxin using post-column photochemical derivatization: collaborative study. *Journal of AOAC international*, 89(3), 678-692.

Whitaker, T.B. (2003). Standardisation of mycotoxin sampling procedure: an urgent necessity. *Food Control*, 14, 233-237.

Wei, R., Qiu, F., Kong, W., Wei, J., Yang, M., Luo, Z., Qin, J., & Ma, X. (2013). Co-occurrence of aflatoxin B₁, B₂, G₁, G₂ and ochratoxin A in *Glycyrrhiza uralensis* analyzed by HPLC-MS/MS. *Food Control*, 32, 216-221.

Yogendrarajah, P., Van Poucke, C., De Meulenaer, B., & De Saeger, S. (2013). Development and validation of a QuEChERS based liquid chromatography tandem mass spectrometry method for the determination of multiple mycotoxin in spices. *Journal of Chromatography A*, 1297, 1-11.

Zain, M. E. (2011). Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society*, 15(2), 129-144.

Zhang, J.M., Wu, Y.L., & Lu, Y.B. (2013). Simultaneous determination of carbamate insecticides and mycotoxins in cereals by reversed phase liquid chromatography tandem mass spectrometry using a quick, easy, cheap, effective, rugged and safe extraction procedure. *Journal of Chromatography B*, 915-916, 13-20.

Zinedine, A., & Manes, J. (2009). Occurrence and legislation of mycotoxins in food and feed from Morocco. *Food Control*, 20, 334-344.

Zinedine, A, Brera, C., Elakhdari, S., Catano, C., Debegnach, F., Angelini, S., De Santis, B., Faid, M., Benlemlih, M., Minardi, V., & Miraglia, M. (2006). Natural occurrence of mycotoxins in cereals and spices commercialized in Morocco. *Food Control*, 17, 868-874.

