



**UNIVERSITI PUTRA MALAYSIA**

***ASSESSMENT OF HAZARDOUS SUBSTANCES IN THE SNAIL, *Nerita lineata* GMELIN, AND SEDIMENTS FROM SELECTED MANGROVE AREAS OF PENINSULAR MALAYSIA***

***CHENG WAN HEE***

**FS 2015 38**



**ASSESSMENT OF HAZARDOUS SUBSTANCES IN THE SNAIL, *Nerita lineata*  
GMELIN, AND SEDIMENTS FROM SELECTED MANGROVE AREAS OF  
PENINSULAR MALAYSIA**

By

**CHENG WAN HEE**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
fulfillment of the Requirement for the degree of Doctor of Philosophy**

**December 2015**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Doctor of Philosophy

**ASSESSMENT OF HAZARDOUS SUBSTANCES IN THE SNAIL, *Nerita lineata*  
GMELIN, AND SEDIMENTS FROM SELECTED MANGROVE AREAS OF  
PENINSULAR MALAYSIA**

By

**CHENG WAN HEE**

**December 2015**

**Chair : Associate Professor Yap Chee Kong, PhD**  
**Faculty : Science**

Rapid growth of the land and marine based industries has generated various contaminations to the coastal areas of Malaysia. Thus, continuous assessments and monitoring of pollutants especially trace metals and polycyclic aromatic hydrocarbons (PAHs) are crucial. The aim of this study is to determine the baseline data, ecological risk and health risk assessments of trace metals and PAHs in mangrove snails, *Nerita lineata* and surface sediments of at least 9 sampling sites in Peninsular Malaysia between 2010 – 2012. Metal analysis was done by using the Atomic Absorption Spectrophotometer (AAS) and Inductively Coupled Plasma-mass Spectrometer (ICPMS). Analysis of PAHs was carried out by using gas chromatography-mass spectrophotometry (GC-MS) while Inter-Simple Sequence Repeat (ISSR) markers were applied for genetic polymorphism study in the snails. All 18 metals in the snails ranged from BDL (below detection limit) – 3235 µg/g dw, with higher accumulation of essential metals in the soft tissues and non-essential metals in the shells and opercula. For sediments, all 18 metals ranged from 0.003 – 48916 µg/g dw. The ∑16 PAHs concentrations (ng/g dw) ranged from 871 – 3066 and 273 – 18030 for soft tissues and sediments, respectively. To determine the ecological risks of habitat on surface sediments, the sediment quality guidelines (SQGs), geochemical indices, potential ecological risk index (PERI) and mean ERM quotient (m-ERM-q) were used. Based on the SQGs, all the metals investigated were unlikely to cause any adverse ecological effects. The ecological risk indices revealed that the pollution and ecological risks of all metals and PAHs were minimal, except for PAHs in JPKetam being ‘moderately to highly polluted’. Potential health risks were assessed from calculations of estimated daily intakes (EDI), target hazard quotients (THQ) and total THQ (TTHQ) of the snails’ soft tissues for trace metals, while excess cancer risks (ECR) were applied for PAHs assessments. Non-carcinogenic risks estimation (THQ and TTHQ) for trace metals showed that average snails consumers were exposed to low potential health risk whereas the high consumers were at high risks (TTHQ>1). The carcinogenic risks approach (ECR) showed that the snail consumption from most sites were in a range of moderate to high carcinogenic risks exposure ( $1 \times 10^{-6} < \text{ECR} < 1 \times 10^{-4}$ ) to PAHs, implying moderation (<17.86 g/day) in snail consumption. The ISSR markers were found to have positive relationships from comparison of patterns of genetic clustering and trace metal levels of the snail muscle tissues but not PAHs. This implied that the

polymorphic loci could be used as potential biomarkers for metals pollution. Generally this study suggested that all metals and PAHs from the sampling sites were not likely to cause hazardous effects to the environment. Consumption of the snails should be of moderation especially to high consumers of the snails as the data showed possible health risks for both trace metals and PAHs. This study has produced important data of the hazardous pollutants in the snails and sediments which can be used for environmental management in future.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENILAIAN BAHAN PENCEMAR DALAM SIPUT, *Nerita lineata* GMELIN,  
DAN SEDIMEN DARI KAWASAN BAKAU DI SEMENANJUNG MALAYSIA**

Oleh

**CHENG WAN HEE**

**Disember 2015**

**Pengerusi : Profesor Madya Yap Chee Kong, PhD**  
**Fakulti : Sains**

Pertumbuhan pesat dalam industri (daratan dan marin) telah menjana pelbagai pencemaran ke atas kawasan pantai di Malaysia. Oleh itu, penilaian dan pemantauan pencemaran yang berterusan terutamanya logam surih dan hidrokarbon polisiklik aromatik (PAHs) adalah penting. Tujuan kajian ini dijalankan adalah untuk menentukan data asas, risiko ekologi dan risiko kesihatan bagi logam surih dan PAHs dalam siput bakau, *Nerita lineata* dan permukaan sedimen dalam sekurang-kurangnya 9 tapak persampelan di Semenanjung Malaysia dari tahun 2010 - 2012. Analisis logam dilakukan dengan menggunakan 'Atomic Absorption Spectrophotometer' (AAS) dan 'Inductively Coupled Plasma-mass Spectrometer' (ICPMS). Analisis PAH dilakukan dengan menggunakan gas chromatography-mass spectrophotometry (GC-MS) manakala 'Inter-Specific Sequence Repeat' (ISSR) telah digunakan untuk mengkaji polimorfisme genetik dalam siput. Julat kandungan semua 18 logam dalam siput adalah BDL – 3235 µg/g berat kering, dengan akumulasi logam esensial yang lebih tinggi dalam tisu lembut dan logam tidak esensial dalam cengkerang dan operkula. Dalam sedimen, julat semua 18 logam adalah 0.003 – 48916 µg/g berat kering. Kepekatan Σ16 PAH (ng/g berat kering) adalah antara 871 – 3066 and 273 – 18030 untuk tisu-tisu lembut siput dan sedimen, masing-masing. Untuk menentukan risiko ekologi habitat pada permukaan sedimen, garis panduan kualiti sedimen (SQGs), indeks geokimia, indeks risiko potensi ekologi (PERI) dan hasil bahagi min ERM (m-ERM-q) telah digunakan. Berdasarkan kepada SQGs, semua logam dalam kajian ini tidak berkemungkinan untuk menyebabkan kesan ekologi yang buruk. Indeks risiko ekologi menunjukkan bahawa pencemaran dan risiko ekologi untuk semua logam dan PAHs adalah minimum, kecuali PAHs di JPKetam yang dikategorikan sebagai 'sederhana hingga sangat tercemar'. Risiko potensi ke atas kesihatan dinilai melalui anggaran pemakanan harian (EDI), 'total hazard quotient' (THQ) dan jumlah THQ (TTHQ) di dalam tisu lembut siput bagi logam surih, manakala risiko kanser (ECR) telah digunakan untuk penilaian risiko PAHs. Anggaran risiko bukan karsinogen (THQ dan TTHQ) telah menunjukkan bahawa risiko potensi kesihatan yang disebabkan oleh pendedahan kepada logam surih bagi konsumen siput sederhana adalah rendah manakala konsumen siput tinggi adalah berisiko tinggi (TTHQ>1). Anggaran risiko karsinogenik (ECR) pula menunjukkan bahawa risiko yang disebabkan oleh pendedahan kepada PAHs bagi konsumen siput adalah tinggi ( $1 \times 10^{-6} < \text{ECR} < 1 \times 10^{-4}$ ) dan pemakanan siput secara sederhana (<17.86 g/day) adalah digalakkan. Analisis

kluster (melalui petanda ISSR) telah mendapati hubungan positif antara corak pengelompokan genetik dan kandungan logam surih di dalam tisu otot siput tetapi tidak untuk PAHs. Ini telah membuktikan bahawa lokus polimorfik berpotensi sebagai penanda biologi untuk pencemaran logam. Secara keseluruhannya, kajian ini telah mencadangkan bahawa semua logam dan PAH dari semua sampel tidak akan membawa kemudaran terhadap persekitaran. Pengambilan siput secara kesederhanaan dari segi pemakanan terutamanya kepada konsumen tinggi siput. Kajian ini telah menghasilkan data bahan pencemar merbahaya yang penting dalam siput dan sedimen untuk pengurusan alam sekitar pada masa akan datang.



## ACKNOWLEDGEMENTS

It has been a big challenge throughout my studies and I would like to take this opportunity to express my gratitude towards a few groups of individuals.

First and foremost, I would also like to thank my supervisory committee members. They are Assoc. Prof. Dr. Yap Chee Kong, Prof. Mohamad Pauzi Zakaria, Prof. Tan Soon Guan and Assoc. Prof. Dr. Ahmad Zaharin Aris for their advice and guidance. It is because of their undying support that had helped me through the completion my studies.

I would also like to thank to all my friends and faculty members. Without your help and support, I would have had numerous problems during my laboratory work and analysis. It has been a great pleasure working with them.

Last but not least, to my family and siblings, I am truly blessed to have them around. They had supported me with patience through many ways, be it financial and morale. Thank you so much and may god blessed you all.



I certify that a Thesis Examination Committee has met on 1 December 2015 to conduct the final examination of Cheng Wan Hee on his thesis entitled "Assessment of Hazardous Substances in the Snail, *Nerita lineata* Gmelin, and Sediments from Selected Mangrove Areas of Peninsular Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Rosimah binti Nulit, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Faridah binti Qamaruz Zaman, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Hishamuddin bin Omar, PhD**

Senior Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Szefer Piotr, PhD**

Professor  
Medical University of Gdansk  
Poland  
(External Examiner)



---

**ZULKARNAIN ZAINAL, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 12 January 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy in Science. The members of the Supervisory Committee were as follows:

**Yap Chee Kong, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Mohamad Pauzi Zakaria, PhD**

Professor  
Faculty of Environmental Sciences  
Universiti Putra Malaysia  
(Member)

**Tan Soon Guan, PhD**

Professor  
Faculty of Biotechnology and Molecular Sciences  
Universiti Putra Malaysia  
(Member)

**Ahmad Zaharin Aris, PhD**

Associate Professor  
Faculty of Environmental Science  
Universiti Putra Malaysia  
(Member)

---

**BUJANG KIM HUAT, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Cheng Wan Hee GS24422

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: \_\_\_\_\_  
Name of Chairman of  
Supervisory  
Committee: Yap Chee Kong, PhD

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: Mohamad Pauzi Zakaria, PhD

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: Tan Soon Guan, PhD

Signature: \_\_\_\_\_  
Name of Member of  
Supervisory  
Committee: Ahmad Zaharin Aris, PhD

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>APPROVAL</b>	vi
<b>DECLARATION</b>	viii
<b>LIST OF TABLES</b>	xiii
<b>LIST OF APPENDICES</b>	xvi
<b>LIST OF FIGURES</b>	xvii
<b>LIST OF ABBREVIATIONS</b>	xix
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>3</b>
2.1 Monitoring and baseline studies of hazardous pollutants	3
2.2 Pollutants in marine ecosystem	4
2.3 Trace metals	4
2.3.1 Fate of trace metals in the environment	5
2.3.2 Uptake, bioaccumulation and toxicity of trace metals gastropods	6
2.4 Polycyclic aromatic hydrocarbons (PAHs)	7
2.4.1 Fate of PAHs in the environment	8
2.4.2 Toxicity, uptake and bioaccumulation of PAHs in	9
2.5 Analytical detection of pollutants	10
2.5.1 Detection of trace metals	10
2.5.2 Detection of PAHs	11
2.6 Risk assessments of trace metals and PAHs in sediments	13
2.6.1 Biomonitoring assessment of hazardous substances in marine environment	13
2.6.2 Sediment Quality Guidelines (SQGs)	14
2.6.3 Enrichment factor (EF) and index of geoaccumulation (Igeo)	14
2.6.4 Ecological Risk Assessment Indices	15
2.6.5 Molecular markers for sources determination of PAHs	15
2.7 Risks assessments of trace metals and PAHs in snails	18
2.7.1 Non-cancerous risks assessments (trace metals)	18
2.7.2 Cancer risk assessments (PAHs)	18
2.8 Genetic assessment based on genetic markers	20
2.9 Common DNA markers used for genetic variability	21
2.9.1 Restriction fragment length polymorphism (RFLP)	21
2.9.2 Random amplified polymorphic DNA (RAPD)	21
2.9.3 Simple sequence repeat (SSR)	21
2.9.4 Inter-simple sequence repeat (ISSR)	22
2.10 Statistical analysis in the interpretation of environmental data	23

2.11	<i>Nerita lineata</i>	25
2.12	<i>Nerita lineata</i> in ecotoxicological study	26
<b>3</b>	<b>MATERIALS AND METHODS</b>	27
3.1	Sampling collections	27
3.2	Trace metal analysis by using AAS	27
3.2.1	Samples preparation and digestion	27
3.2.2	Metal determination	31
3.3	Trace metal analysis by using ICPMS	32
3.3.1	Preparation of sediment and snail samples for microwave digestion	32
3.3.2	Metal determination	32
3.3.3	Quality control	32
3.3.4	Statistical analysis	33
3.3.5	Trace metal data treatment	33
3.4	PAHs analysis	37
3.4.1	Preparation of chemicals and materials	37
3.4.2	Samples preparation	38
3.4.3	Determination of PAHs in soft tissues of snails	38
3.4.4	Determination of PAHs in sediments	39
3.4.5	Determination of dry weight	40
3.4.6	Determination of lipid content in soft tissues	40
3.4.7	Analysis by using GC-MS	40
3.4.8	Quality control and quality assurance	41
3.4.9	PAHs data treatment	41
3.5	Inter-specific sequence repeat (ISSR)	42
3.5.1	DNA isolation	42
3.5.2	PCR amplification	43
3.5.3	Electrophoresis	43
3.5.4	Data analysis	44
<b>4</b>	<b>RESULTS</b>	45
4.1	Trace metals	45
4.1.1	Background of trace metal levels in <i>N. lineata</i>	45
4.1.2	BSAF of trace metals in different parts of the snails	48
4.1.3	Background of trace metal levels in sediments	49
4.1.4	Enrichment factor and geoaccumulation index of sediments	50
4.1.5	Ecological risk assessment	53
4.1.6	Estimated dietary intake (EDI) of metals, target hazard quotient (THQ) and total target hazard quotient (TTHQ)	53
4.2	Polycyclic aromatic hydrocarbon (PAHs)	59
4.2.1	PAH profile in sediments and snails	59
4.2.2	Spatial distribution and molecular indices of PAHs in sediments and snails	59
4.2.3	Dietary intake of PAHs from consumption of snails	62
4.2.4	Carcinogenic risks from consumption of snails	62
4.2.5	Potential risk assessments of PAH in surface sediments	62
4.3	Inter-simple sequence repeat (ISSR)	67

	4.3.1	Profile of ISSR banding and genetic polymorphism	67
	4.3.2	Genetic distance	67
<b>5</b>	<b>DISCUSSION</b>		<b>74</b>
	5.1	Background profile of trace metals and PAHs in <i>N. lineata</i> and surface sediments collected from the southern and western part of Peninsular Malaysia	74
	5.1.1	Profile of trace metals in snails	74
	5.1.2	Profiles of trace metals in sediments	75
	5.1.3	Relationship between trace metal concentrations in <i>Nerita lineata</i> and sediments	77
	5.1.4	Profiles of PAHs in snails	83
	5.1.5	Profiles of PAHs in sediments	84
	5.1.6	Sources identification of PAHs in snails and sediments	84
	5.1.7	Multivariate analysis of PAH concentrations in snails and sediments	86
	5.1.8	Relationship between PAH concentrations in <i>Nerita lineata</i> and sediments based on molecular indices and multivariate analysis	90
	5.2	Ecological risk assessment of heavy metals and PAHs on the surface sediments at the mangrove snail habitats	91
	5.2.1	Ecotoxicological significance of trace metal concentrations in sediments	92
	5.2.2	Risk assessments by using contamination indices and risk categories	97
	5.2.3	Comparison of PAH concentrations with SQGs	98
	5.2.4	Risk assessment of total mixture of PAHs in surface sediments	98
	5.3	Assessment of the levels of human health risk based on the estimated daily intake and risk indices of the snail consumption	100
	5.3.1	Comparisons of trace metal concentrations with food safety guidelines	100
	5.3.2	Potential health risk assessment of trace metals	103
	5.3.3	Dietary risk assessment of PAHs	105
	5.3.4	Cancer risk assessment	105
	5.4	Relationship between the trace metal levels, PAH and the genetic polymorphisms of <i>N. lineata</i> collected from selected sites in the coastal areas of Peninsular Malaysia	106
<b>6</b>	<b>CONCLUSIONS</b>		<b>110</b>
	<b>REFERENCES</b>		<b>111</b>
	<b>APPENDICES</b>		<b>134</b>
	<b>BIODATA OF STUDENT</b>		<b>147</b>
	<b>LIST OF PUBLICATIONS</b>		<b>148</b>

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
2.1 List of essential and non-essential metals	5
2.2 List of PAH compounds focused in the present study	8
2.3 Terminologies for the values of the potential risk of individual metal ( $Er$ ) and the sum of potential risk of individual metal ( $RI$ ) (Håkanson, 1980)	16
2.4 Toxic equivalent factors (TEFs) for 16 PAHs in this study	19
2.5 Common statistical analysis for interpretation of environmental data	24
3.1 Site description and shell length (mm), shell width (mm) and shell height (mm) of <i>Nerita lineata</i> , collected from the mangrove areas of Peninsular Malaysia	30
3.2 Comparisons of metal concentrations ( $\mu\text{g}/\text{kg}$ dry weight) between certified and measured values of certified reference materials (CRMs). NIST SRM 1646 Estuarine Sediment (SED); NIST No. 2976 Mussel soft tissue (Mussel); Dogfish liver DOLT-3 (Dogfish) (NA= not available)	34
3.3 Primers and sequences used for ISSR in this study	43
4.1 Biota-Sediment Accumulation Factor (BSAF) of the three parts of <i>Nerita lineata</i> based on mean metal concentrations of 10 sampling sites	49
4.2 Enrichment factor (EF) and geoaccumulation index ( $I_{geo}$ ) of sediments collected from 10 sites of Peninsular Malaysia	51
4.3 The values of the contamination factor ( $C_f$ ), potential risk of individual metal ( $Er^i$ ) and the sum of potential risk of individual metal ( $RI$ ) of all sampling sites of surface sediments from Peninsular Malaysia	54
4.4 Estimated daily intake ( $\mu\text{g kg}^{-1} \text{day}^{-1}$ ) of <i>Nerita lineata</i> for average (ALC) and high (HLC) consumers	55
4.5 Target hazard quotients (THQs) for individual metals and total hazard quotient (TTHQ) for average (AC) and high (HC) consumers	57
4.6 Total 16 PAH concentrations (ng/g) and molecular indices for PAHs in sediments and snails (including lipid content (%), dry weight (dw), wet weight (ww) and lipid weight (lw)) from mangrove areas of Peninsular Malaysia	61



4.7	Estimated daily intake (ng TEQ/day ww) for individual PAHs from consumption of <i>Nerita lineata</i> for average (ALC) and high (HLC) consumers from 10 sampling sites	63
4.8	The concentrations of BaP (ng/g ww), total benzo(a)pyrene toxic equivalent quotient (TEQ) (ng TEQ/g ww), estimated daily intake (ALC/HLC) (ng TEQ/day ww) and excess cancer risks (ECR for ALC and HLC) of <i>Nerita lineata</i> collected from 10 sampling sites	65
4.9	Summary of ISSR bands generated from the muscle tissues of six populations of <i>Nerita lineata</i>	68
4.10	Genetic distance of 6 populations of <i>Nerita lineata</i> based on Dice similarity coefficients	73
5.1	Reference values ( $\mu\text{g/g}$ ), and values of toxic-response factors (unitless) employed in this study	76
5.2	Factor loadings (Varimax normalized), eigenvalue, percentage total variances and cumulative total variances for factors derived by factor analysis based on $\log_{10}(\text{mean} + 1)$ for 18 metals in soft tissues of <i>Nerita lineata</i> collected from 10 sampling sites (N= 10). Extraction: Principal components (Bolded figures indicate loadings > 0.70)	80
5.3	Factor loadings (Varimax normalized), eigenvalue, percentage total variances and cumulative total variances for factors derived by factor analysis based on $\log_{10}(\text{mean} + 1)$ for 18 metals in surface sediments collected from 10 sampling sites (N= 10). Extraction: Principal components (Bolded figures indicate loadings > 0.70)	81
5.4	Comparison of PAH concentrations (ng/g dw) in sediments of current study and previous reported studies	85
5.5	Factor loadings (Varimax normalized), eigenvalue, percentage total variances and cumulative total variances for factors derived by factor analysis based on $\log_{10}(\text{mean} + 1)$ for 16 PAHs in soft tissues of <i>Nerita lineata</i> collected from 8 sampling sites (N= 8). Extraction: Principal components (Bolded figures indicate loadings > 0.50)	87
5.6	Factor loadings (Varimax normalized), eigenvalue, percentage total variances and cumulative total variances for factors derived by factor analysis based on $\log_{10}(\text{mean} + 1)$ for 16 PAHs in surface sediments collected from 9 sampling sites (N= 9). Extraction: Principal components (Bolded figures indicate loadings > 0.50)	88
5.7	Sediment quality guidelines employed in this study	93
5.8	Comparison of trace metal concentrations ( $\mu\text{g/g dw}$ ) in sediments of current study and previous reported studies	95

5.9	Individual PAH concentrations (ng/g dw) of sediments (n=9) in comparison with SQGs (ERL, ERM, TEL and PEL values)	99
5.10	Metal concentrations ( $\mu\text{g/g}$ ww) in the wet weight soft tissues (ST) of snails collected from 11 sampling sites in mangrove area of Peninsular Malaysia	101



## LIST OF APPENDICES

Appendix		Page
1	Mean concentrations ( $\mu\text{g/g dw}$ ) of 18 trace metals in the shells of the <i>Nerita lineata</i> collected from 11 sites of Peninsular Malaysia	134
2	Mean concentrations ( $\mu\text{g/g dw}$ ) of 18 trace metals in the opercula of the <i>Nerita lineata</i> collected from 11 sites of Peninsular Malaysia	136
3	Mean concentrations ( $\mu\text{g/g dw}$ ) of 18 trace metals in the soft tissues of the <i>Nerita lineata</i> collected from 11 sites of Peninsular Malaysia	138
4	Periodic Table (source: Ivanov, 2006)	140
5	Mean concentrations ( $\mu\text{g/g dw}$ ) of 18 trace metals in the sediments collected from 10 sites of Peninsular Malaysia	141
6	Relationship of trace metal concentrations [ $\log_{10}(\text{mean}+1)$ ] in shells, opercula and soft tissues of <i>Nerita lineata</i> and sediments based on Pearson's correlations collected from 10 sampling sites of Peninsular Malaysia	143
7	Relationship of PAH concentrations [ $\log_{10}(\text{mean}+1)$ ] in soft tissues of <i>Nerita lineata</i> and sediments based on Pearson's correlations collected from 9 sampling sites of Peninsular Malaysia	144
8	Benzo(a)pyrene toxic equivalent quotient (TEQ) ( $\text{ng/TEQ/g ww}$ ) of individual PAHs in the soft tissues of <i>Nerita lineata</i> collected from 10 sampling sites from Peninsular Malaysia	145

## LIST OF FIGURES

Figure	Page
2.1 Distribution of trace metals in different compartments	7
2.2 <i>Nerita lineata</i> on rocks	25
3.1 Summary of this study on trace metals and PAHs in <i>Nerita lineata</i> and sediments	28
3.2 Sampling Map (Source: Google map)	29
3.3 The (a) shells, (b) opercula, (c) soft tissues and (d) intact figure of the <i>Nerita lineata</i>	31
4.1 Concentrations (ng/g dw) of individual PAHs in sediments collected from 9 sites of mangrove areas in Peninsular Malaysia	60
4.2 Concentrations (ng/g dw) of individual PAHs in soft tissues of <i>Nerita lineata</i> collected from 10 sites of mangrove areas in Peninsular Malaysia	60
4.3 Spatial distribution of mean ERM quotient (m-ERM-q) values of $\sum 16$ PAHs in surface sediments collected from 9 sites of Peninsular Malaysia	66
4.4 ISSR profile of <i>Nerita lineata</i> population generated by 8 primers collected from 6 sampling sites	72
5.1 Cluster analysis based on Single Linkage Euclidean distances on 18 metals concentrations [ $\log_{10}(\text{mean}+1)$ ] in the soft tissues of <i>Nerita lineata</i> collected from 11 sampling sites from Peninsular Malaysia	78
5.2 Cluster analysis based on Single Linkage Euclidean distances on 18 metals concentrations [ $\log_{10}(\text{mean}+1)$ ] in the surface sediments collected near the habitat of <i>Nerita lineata</i> collected from 10 sites from Peninsular Malaysia	79
5.3 Cluster analysis based on Single Linkage Euclidean distances on 16 PAHs concentrations [ $\log_{10}(\text{mean}+1)$ ] in the soft tissues of <i>Nerita lineata</i> collected from 10 sites from Peninsular Malaysia	89
5.4 Cluster analysis based on Single Linkage Euclidean distances on 16 PAHs concentrations [ $\log_{10}(\text{mean}+1)$ ] in the surface sediments collected near the habitat of <i>Nerita lineata</i> collected from 9 sites from Peninsular Malaysia	90
5.5 UPGMA dendrogram of genetic relationships of 6 populations of <i>Nerita lineata</i> based on Dice similarity coefficients derived from 113 ISSR bands	107

5.6	Cluster analysis based on Single Linkage Euclidean distances on $\Sigma 18$ metals concentrations in the soft tissues of <i>Nerita lineata</i> collected from 6 selected sampling sites	108
5.7	Cluster analysis based on Single Linkage Euclidean distances on $\Sigma 18$ metals concentrations in the sediments collected from 6 selected sampling sites	108
5.8	Cluster analysis based on Single Linkage Euclidean distances on $\Sigma 16$ PAHs concentrations in the soft tissues of <i>Nerita lineata</i> collected from 6 selected sampling sites	109
5.9	Cluster analysis based on Single Linkage Euclidean distances on $\Sigma 16$ PAHs concentrations in the sediments collected from 6 selected sampling sites	109

## LIST OF ABBREVIATIONS

AC	Average consumers
Ace	Acenaphthene
Acy	Acenaphthylene
AHR	Aryl hydrocarbon receptor
Ant	Anthracene
BaA	Benzo(a)anthracene
BaP	Benzo(a)pyrene
BbF	Benzo(b)fluoranthene
BghiP	Benzo(g,h,i)perylene
BkF	Benzo(k)fluoranthene
BSAF	Biota-sediment accumulation factors
<i>C<sub>f</sub></i>	Contamination factor
CF	Conversion factor
Chry	Chrysene
CYP1A	Cytochrome P4501A
DahA	Dibenzo(a,h)anthracene
DCM	Dichloromethane
DDW	Double distilled water
dw	dry weight
EADI	Estimated average daily intake
ECD	Electron capture detector
ECR	Excess cancer risks
ECRA	Excess cancer risks for average consumers
ECRH	Excess cancer risks for high consumers

EDI	Estimated daily intake
EDI	Estimated daily intakes
EF	Enrichment factor
EHDI	Estimated high daily intake
$E_r^i$	Potential risk of individual metal
ERL	Effect range low
ERM	Effect range median
FID	Flame ionization detector
Flu	Fluorene
Fth	Fluoranthene
HC	High consumers
HMW	High molecular weight polycyclic hydrocarbons
IcdP	Indeno[1,2,3-cd]pyrene
IIS	Internal injection Standard
ISQV-high	Interim sediment quality value-high
ISQV-low	Interim sediment quality value-low
LMW	High molecular weight polycyclic hydrocarbons
lw	lipid weight
MS	Mass spectrophotometry
m-ERM-q	Mean effect range median quotient
Nap	Naphthalene
PEL	Probable effect level
Phe	Phenanthrene
Pyr	Pyrene
$RI$	sum of potential risk of individual metal
SD	Standard deviation

SIS	Surrogate internal standard mixture
SQGs	Sediment quality guidelines
ST	Soft tissues
TCD	Thermal conductivity detector
TEF	Toxic equivalent factor
TEL	Threshold effect level
TEQ/B(a)P <sub>cq</sub>	Benzo(a)pyrene toxic equivalent quotient
TFM	Tetraflouroethylene Modified
THQ	Target hazard quotients
tPAH	Total polycyclic aromatic hydrocarbons
TTHQ/HI	Total target hazard quotient/hazard index
ww	wet weight



## CHAPTER 1

### INTRODUCTION

In the field of ecotoxicology studies, marine coastal pollution has always been a matter of concern. With the rapid growth of the industries in Malaysia, various organic and inorganic contaminations were generated to the coastal areas of Malaysia. Inorganic contaminants such as heavy metals occur naturally in the environment. Human activities which cause elevated levels of metals input to the environment will cause detrimental effects on the marine coastal ecosystem. Organic pollutants such as polycyclic aromatic hydrocarbons (PAHs) on the other hand, is an important class of persistent pollutants known to have carcinogenic and mutagenic properties and they are introduced to the environment by two sources of anthropogenic activities namely, pyrogenic and petrogenic (Wang et al., 2014). Such pollution will not only threaten the ecology of marine system but also pose danger to humans, as most marine organisms are consumed by humans. Therefore, a continuous study on monitoring of heavy metal pollution is crucial.

In this study the analyses of pollution levels of trace metals and PAHs were conducted through the assessment biomonitors, sediments and snails, from mangrove areas of Klang (Selangor), Negeri Sembilan and Johore between the period of 2010 – 2011. These sites are chosen due to their location on important shipping lane across the coastal line of Peninsular Malaysia and susceptible to various marine base pollution sources. Various land base industries were observed in these sites. On top of that, mangrove areas occupying majority of the coastal area in Peninsular Malaysia, is an important ecosystem and exchange point of numerous organic and inorganic substances from various sources, i.e. marine, terrestrial or atmospheric (Alongi et al., 2004).

Biomonitoring is a widely procedure that has been practically utilized as a monitoring and assessment tool for heavy metal pollutions in order to remedy the aggravation of the environment. While sediment has a reputation as an ultimate biomonitor for environmental pollution, application of local or endemic biological species are confronted with challenges of the wide range of environmental conditions that may limit the availability of the species (Kang et al., 2000). Apart from their massive species number and ubiquitous distribution in marine coastal ecosystems, the biological data from these species are directly related to the ecological condition and or 'ecological health' of the marine aquatic ecosystem (Campbell, 2002).

The species collected for the current study were the mangrove snails, *Nerita lineata*. The *N. lineata* is a common snail found mainly in the intertidal area of Malaysia, Indonesia and Singapore. This species is studied for it fulfils some basic criteria of a good biomonitor such as easily collected and weighed, abundant, sedentary and of relatively high longevity. It is also observed that many people in this region are taking

*N. lineata* as source of food. Therefore it is essential to provide the heavy metal data of the snails as a guideline for safety consumption.

Assessment of risks of pollutants, such as metal and PAHs, in sediments were done through three approaches namely, i) established Sediment Quality Guidelines (SQGs), ii) ecological risks indices, and iii) geochemical pollution indexes. Screening-level of risk assessment in the *N. lineata* was conducted through consideration of internationally accepted dietary guidelines and the calculation of estimated daily intakes (EDI), target hazard quotients (THQ) and excess cancer risks (ECR). These guidelines were developed by researchers or organizations (USEPA) and they are widely used as a reliable risk assessment tool.

However, ecotoxicological studies should not be restricted to the context of metal levels as the impact of the pollution is also revealed in the biological response of the organisms. Marine organisms exhibit genetic polymorphisms due to adaptation and response towards environmental contaminants (Basu et al., 2014). Thus, inter-simple sequence repeat (ISSR) markers were used for the analysis of genetic polymorphisms in the *N. lineata*.

The objectives of this study were:

1. To determine the baseline levels of trace metals and PAHs in different parts of the snails, *N. lineata* and sediments from selected mangrove areas of Peninsular Malaysia,
2. To determine the ecological risk assessment of trace metals and PAH on the surface sediments at the mangrove snail's habitats,
3. To determine the human health assessments based on the estimated daily intake and risk indices of trace metals and PAHs in the *N. lineata* collected from selected mangrove areas of Peninsular Malaysia,
4. To investigate the relationships between genetic variation and hazardous substances (trace metals and PAHs) in *N. lineata* collected from selected mangrove areas of Peninsular Malaysia.

## REFERENCES

- Alongi, D. M., Sasekumar, A., Chong, V. C., Pfitzner, J., Trott, L. A., Tirendi, F., Dixon, P., & Brunskill, G. J. (2004). Sediment accumulation and organic material flux in a managed mangrove ecosystem: estimates of land–ocean–atmosphere exchange in peninsular Malaysia. *Marine Geology*, 208, 383-402.
- Al-Qurainy, F. (2010). Application of inter simple sequence repeat (ISSR marker) to detect genotoxic effect of heavy metals on *Eruca sativa* (L.). *African Journal of Biotechnology*, 9(4), 467-474.
- Amiard, J. C., Amiard-Triquet, C., Barka, S., Pellerin, J., & Rainbow, P. S. (2006). Metallothioneins in aquatic invertebrates: Their role in metal detoxification and their use as biomarkers. *Aquatic Toxicology*, 76, 160-202.
- Amin, B., Ismail, A., Arshad, A., Yap, C. K. & Kamarudin, M. S. (2006). A comparative study of heavy metal concentrations *Nerita lineata* from the intertidal zone between Dumai Indonesia and Johor Malaysia. *Journal of Coastal Development*, 10, 19-32.
- Amin, B., Ismail, A. & Yap, C. K. (2008). Heavy metal concentrations in sediment and intertidal gastropod *Nerita lineata* from two opposing sites in the Straits of Malacca. *Wetland Science*, 6(3), 411-421.
- Amin, B., Ismail, A., Arshad, A., Yap, C. K. and Kamarudin, M. S. (2009). Gastropod assemblages as indicators of sediment metal contamination in mangroves of Dumai, Sumatra, Indonesia. *Water, Air, & Soil Pollution*, 201, 9-18.
- ANZFA, (2011). Australian and New Zealand Food Standards Code, Standard 1.4.1-contaminants and Natural Toxicants (F2011C 00542).
- Bai, J., Xiao, R., Zhang, K., & Gao, H. (2012). Arsenic and heavy metal pollution in wetland soils from tidal freshwater and salt marshes before and after the flow-sediment regulation regime in the Yellow River Delta, China. *Journal of Hydrology*, 450-451, 244-253.
- Bakhtiari, A. R., Zakaria, M. P., Yaziz, M. I., Lajis, M. N. H., Bi, X., & Rahim, M. C. A. (2009). Vertical distribution and source identification of polycyclic aromatic hydrocarbons in anoxic sediment cores of Chini Lake, Malaysia: Perylene as indicator of land plant-derived hydrocarbons. *Applied Geochemistry*, 24, 1777-1787.
- Bandowe, B. A. M., Bigalke, M., Boamah, L., Nyarko, E., Saalia, F. K., & Wilcke, W. (2014). Polycyclic aromatic compounds (PAHs and oxygenated PAHs) and trace metals in fish species from Ghana (West Africa): Bioaccumulation and health risk assessment. *Environment International*, 65, 135-146.
- Barone, G., Storelli, A., Garofalo, R., Busco, V. P., Quaqlia, N. C., Centrone, G., & Storelli, M. M. (2015). Assessment of mercury and cadmium via seafood

consumption in Italy: Estimated dietary intake (EWI) and target hazard quotient. *Food Additives & Contaminants: Part A*, 13, 1-10.

- Batista, M. J., Abreu, M. M., Locutura, J., De Oliveira, D., Matos, J. X., Silva, C., Bel-Lan, A., & Martins, L. (2012). Evaluation of trace elements mobility from soils to sediments between the Iberian Pyrite Belt and the Atlantic Ocean. *Journal of Geochemical Exploration*, 123, 61-68.
- Barker, C. S., Medrano-Gonzalez, L., Calambokidis, J., Perry, A., Pichler, F., Rosenbaum, H., Straley, J. M., Urrban-Ramirez, J., Yamaguchi, M., & von Ziegeler, O. (1998). Population structure of nuclear and mitochondrial DNA variation among humpback whales in the North Pacific. *Molecular Ecology*, 7, 695-707.
- Basu, N., Goodrich, J. M., & Head, J. (2014). Ecogenetics of mercury: From genetic polymorphisms and epigenetics to risk assessment and decision-making. *Environmental Toxicology and Chemistry*, 33(6), 1248-1258.
- Baumard, P., Budzinski, H., Garrigues, P., Sorbe, J. C., Burgeot, T., & Bellocq, J. (1998). Concentrations of PAHs (polycyclic aromatic hydrocarbons) in various marine organisms in relation to those in sediments and to trophic level. *Marine Pollution Bulletin*, 36, 951-960.
- Beach, D. G., Quilliam, M. A., & Hellou, J. (2009). Analysis of pyrene metabolites in marine snails by liquid chromatography using fluorescence and mass spectrometry detection. *Journal of Chromatography B*, 877, 2142-2152.
- Brakstad, F. (1992). A comprehensive pollution survey of polychlorinated dibenzo-p-dioxins and dibenzofurans by means of principal component analysis and partial least square regression. *Chemosphere*, 25, 1611-1629.
- Bodin, N., N'Gom-Kâ, R., Kâ, S., Thiaw, O. T., Tito de Morais, L., Le Loc'h, F., Rozuel-Chartier, E., Auger, D. and Chiffolleau, J. F. (2013). Assessment of trace metal contamination in mangrove ecosystems from Senegal, West Africa. *Chemosphere*, 90, 150-157.
- Botello, A. V., Soto, L. A., Ponce-Vélez, G., & Villanueva, S. F. (2015). Baseline for PAHs and metals in NW Gulf of Mexico related to the Deepwater Horizon oil spill. *Estuarine, Coastal and Shelf Science*, 156, 124-133.
- Boyle, W. R., & Robinson, H. A. (1988). Nickel in the natural environment. In H. Sigel & A. Sigel (Eds.), *Metal ions in biological systems: Nickel and its role in biology* (pp. 1-29). New York and Basel: Marcel Dekker, Inc.
- Buat-Menerd, P., & Chesselt, R. (1979). Variable influence of the atmosphere flux on the trace metal chemistry of oceanic suspended matter. *Earth and Planetary Science Letter*, 42, 398-411.
- Cai, X. W., Shao, Y., & Lin, Z. M. (2014). Genetic differentiation caused by chromium treatment in *Leersia hexandra* swartz revealed by RAPD analysis. *Open Journal of Soil Science*, 4, 366-373.

- Cai, L., Xu, Z., Bao, P., He, M., Dou, L., Chen, L., Zhou, Y., & Zhu, Y. (2015). Multivariate and geostatistical analyses of the spatial distribution and source of arsenic and heavy metals in the agricultural soils in Shunde, Southeast China. *Journal of Geochemical Exploration*, 148, 189-195.
- Campbell, I. C. (2002). Biological monitoring and assessment using invertebrates. In R.B. Frank, I. McKelvie, U. Förstner, & A. Guenther (Eds.), *Environmental monitoring handbook* (pp. 5.1-5.16). New York: Mc Grawhill.
- Carla de Aranzamendi, M., Gardenal, C. N., Martin, J. P., & Bastida, R. (2009). Limpets of the genus *Nacella* (patellogastropoda) from the southwestern Atlantic: Species identification based on molecular data. *Journal of Molluscan Studies*, 75, 241-251.
- Carré, M., Bentaleb, I., Bruguier, O., Ordinola, E., Barrett, N. T., & Fontugne, M. (2006). Calcification rate influence on trace element concentrations in aragonitic bivalve shells: Evidences and mechanisms. *Geochimica et Cosmochimica Acta*, 70, 4906-4920.
- Casu, M., Casu, D., Lai, T., Cossu, P., & Curini-Galletti, M. (2006). Inter-simple sequence repeat markers reveal strong genetic differentiation among populations of the endangered mollusk *Patella ferruginea* (Gastropoda: Patellidae) from two Sardinian marine protected areas. *Marine Biology*, 149, 1163-1174.
- Casu, M., Rivera-Ingraham, G. A., Cossu, P., Lai, T., Sanna, D., Dedola, G. L., Sussarellu, R., Sella, G., Cristo, B., Curini-Galletti, M., Garcí'a-Go'mez, J. C., & Espinosa, F. (2011). Patterns of spatial genetic structuring in the endangered limpet *Patella ferruginea*: implications for the conservation of a Mediterranean endemic. *Genetica*, 139, 1293-1308.
- CCME (Canadian Council of Ministers of the Environment) (2002). Canadian Sediment Quality Guidelines for the Protection of Aquatic Life. [Accessed on May 2015]. [http://www.ccme.ca/assets/pdf/sedqg\\_summary\\_table.pdf](http://www.ccme.ca/assets/pdf/sedqg_summary_table.pdf)
- CEC (The Commission of the European Communities) (2006). Regulation (EC) No 1881/2006 of the European parliament and the council of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. Official Journal of the European Communities, L364/5-24. [Accessed on November 2014]. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2006:364:0005:0024:EN:PDF>
- Chandru, K., Zakaria, M. P., Anita, S., Shahbazi, A., Sakari, M., Bahry, P. S., & Mohamed, C. A. R. (2008). Characterization of alkanes, hopanes, and polycyclic aromatic hydrocarbons (PAHs) in tar-balls collected from the East Coast of Peninsular Malaysia. *Marine Pollution Bulletin*, 56, 950-962.
- Chapman, P. M., Patrick, J. A., & Vigers, G. A. (1999). Development of sediment quality values for Hong Kong special administrative region: A possible model for other jurisdictions. *Marine Pollution Bulletin*, 38, 161-169.

- Cheng, W. H. (2008). *Distribution and concentrations of several heavy metals in snails (Nerita lineata) from the intertidal areas of Peninsular Malaysia*, Unpublished Master Thesis, Universiti Putra Malaysia.
- Christensen, E. R., & Bzdusek, P. A. (2005). PAHs in sediments of Black River and the Ashtabula River, Ohio: Source apportionment by factor analysis. *Water Research*, 39(4), 511-524.
- Christensen, E. R., & Arora, S. (2007). Source apportionment of PAHs in sediments using factor analysis by time records: Application to Lake Michigan, USA. *Water Research*, 41, 168-176.
- Clark, J. R. (1995). *Coastal zone management handbook*. United States of America: CRR Press.
- Conti, M. E., Cucina, D., & Mecozzi, M. (2008). Regression Analysis Model Applied to Biomonitoring Studies. *Environmental Modeling & Assessment*, 13, 553-565.
- Cravo, A., Bebianno, M. J., & Foster, P. (2004). Partitioning of trace metals between soft tissues and shells of *Patella aspera*. *Environment International*, 30, 87-98.
- Cuong, D. T., Bayen, S., Wurl, O., Subramanian, K., Wong, K. K. S., Sivasothi, N., & Obbard, J. (2005). Heavy metal contamination in mangrove habitats of Singapore. *Marine Pollution Bulletin*, 50, 1713-1744.
- Dallinger, R. (1993). Strategies of metal detoxification in terrestrial invertebrates. In R. Dallinger, & P.S. Rainbow (Eds), *Ecotoxicology of Metals in Invertebrates* (pp. 246–332). Boca Raton, FL: Lewis Publisher.
- DeCoster, J. (1998). *Overview of Factor Analysis*. [Accessed on September 2011]. <http://www.stat-help.com/notes.html>
- Deng, G., Yang, W., Zhou, G., Li, Y., & Liu, S. (2014). Heavy metals and polycyclic aromatic hydrocarbons in sediments from the Shenzhen River, South China. *Environmental Science and Pollution Research*, 21, 10594-10600.
- Ding, C., Ni, H. and Zeng, H. (2012). Parent and halogenated polycyclic aromatic hydrocarbons in rice and implications for human health in China. *Environmental Pollution*, 168, 80-86.
- Dong, C. D., Chen, C. F., & Chen, C. W. (2014). Vertical profile, sources, and equivalent toxicity of polycyclic aromatic hydrocarbons in sediment cores from the river mouths of Kaohsiung Harbor, Taiwan. *Marine Pollution Bulletin*, 85, 665-671.
- Duan, Y., Guttman, S. I., Oris, J. T., & Bailer, A. J. (2001). Differential survivorship among allozyme genotypes of *Hyaella azteca* exposed to cadmium, zinc or low pH. *Aquatic Toxicology*, 54, 15-28.

- Eisler, R. (1987). *Mercury hazards to Fish, wildlife, and invertebrates: a synoptic review*. US Fish and Wildlife Service Biological Report 85 (1.10).
- Ergin, M., Saydam, C., Baştürk, Ö., Erdem E., & Yörük, R. (1991). Heavy metal concentrations in surface sediments from the two coastal inlets (Golden Horn Estuary and Izmit Bay) of the northeastern Sea of Marmara. *Chemical Geology*, *91*, 269-285.
- eSilva, C. A. R., Smith, B. D., & Rainbow, P. S. (2006). Comparative biomonitors of coastal trace metal contamination in tropical South America (N. Brazil). *Marine Environmental Research*, *61*, 439-455.
- FAO (Food Administration Organization) (1983). Compilation of legal limits for hazardous substances in fish and fishery products. FAO Fisheries Circular. 764. FAO, Rome. [Accessed on November 2014]. [http://www.fao.org/fi/oldsite/eims\\_search/1\\_dett.asp?calling=simple\\_s\\_result&lang=fr&pub\\_id=65155](http://www.fao.org/fi/oldsite/eims_search/1_dett.asp?calling=simple_s_result&lang=fr&pub_id=65155)
- Feldstein, T., Kashman, Y., Abelson, A., Fishelson, L., Mokady, O., B., Vladimir, B., & Erel, Y. (2003). Marine molluscs in environmental monitoring. *Helgoland Marine Research*, *57*, 212-219.
- Fent, K. (2004). Ecotoxicological effects at contaminated sites. *Toxicology*, *205*, 223-240.
- Fernández-Tajes, J., Flórez, F., Pereira, S., Rábade, T., Laffon, B., & Méndez, J. (2011). Use of three bivalve species for biomonitoring a polluted estuarine environment. *Environmental Monitoring and Assessment*, *177*, 289-300.
- Finley, B. L., Monnot, A. D., Paustenbach, D. J., & Gaffney, S. H. (2012). Derivation of a chronic oral reference dose for cobalt. *Regulatory Toxicology and Pharmacology*, *64*, 491-503.
- Förstner, U. (2002). *Introduction to soils and sediments*. New York: The McGraw-Hill Companies, Inc.
- Foster, P., & Chacko, J. (1995). Minor and trace elements in the shell of *Patella vulgata* (L.). *Marine Environmental Research*, *40*(1), 55-76.
- Foster, P., & Cravo, A. (2003). Minor elements and trace metals in the shells of marine gastropods from a shore in tropical East Africa. *Water, Air and Soil Pollution*, *145*, 53-65.
- Fowles, I.A. (1995). *Gas Chromatography: Analytical chemistry by open learning*. England: John Wiley and Sons.
- Fritsch, C., Coeurdassier, M., Gimbert, F., Crini, N., Scheifler, R., & de Vaufleury, A. (2011). Investigations of responses of metal pollution in land snail populations (*Cantareus aspersus* and *Cepaea nemoralis*) from a smelter-impacted area. *Ecotoxicology*, *20*(4), 739-59.

- Fritz, M., & Berger, P. D. (2015a). Chapter 3 – Comparing two designs (or anything else!) using paired sample T-tests. *Improving the User Experience Through Practical Data Analytic*, 2015, 71-89.
- Fritz, M., & Berger, P. D. (2015b). Chapter 10 – Can you relate in multiple ways? Multiple linear regression and stepwise regression. *Improving the User Experience Through Practical Data Analytic*, 2015, 239-269.
- Gay, D., & Maher, W. (2003). Natural variation of copper, zinc, cadmium and selenium concentrations in *Bembicium nanum* and their potential use as a biomonitor of trace metals. *Water Research*, 37, 2173-2185.
- Gewurtz, S. B., Lazar, R., & Haffner, G. D. (2003). Biomonitoring of bioavailable PAH and PCB water concentrations in the Detroit River using the freshwater mussel, *Elliptio complanata*. *Journal of Great Lakes Research*, 29(2), 242-255.
- Gharibreza, M., Raj, K. R., Yusoff, I., Othman, Z, Tahir, W. Z. W. M., & Ashraf, M. A. (2013). Historical variations of Bera Lake (Malaysia) sediments geochemistry using radioisotopes and sediment quality indices. *Journal of Radioanal Nuclear Chemistry*, 295, 1715-1730.
- Gomes, F., Oliveira, M., Ramalhosa, M. J., Delerue-Matos, C., & Morais, S. (2013). Polycyclic aromatic hydrocarbons in commercial squids from different geographical origins: Levels and risks for human consumption. *Food and Chemical Toxicology*, 59, 46-54.
- Gong, Q., Deng, J., Xiang Y., Wang, Q., & Yang L. (2008). Calculating Pollution Indices by Heavy Metals in Ecological Geochemistry Assessment and a Case Study in Parks of Beijing. *Journal of China University of Geosciences*, 19(3), 230-241.
- Gorell, J. M., Johnson, C. C., Rybicki, B. A., Peterson, E. L., Kortsha, G. X., Brown, G. G., & Richardson, R. J. (1997). Occupational exposures to metals as risk factors for Parkinson's disease. *Neurology*, 48, 650-658.
- Gralla, F., Abson, D. J., Møller, A. P., Lang, D. J., and von Wehrden, H. (2014). The impact of nuclear accidents on provisioning ecosystem services. *Ecological Indicators*, 41, 1-14.
- Grande, J. A., Borrego, J., & Morales, J. A. (2000). A study of heavy metal pollution in the Tinto-Odiel Estuary in Southern Spain by using factor analysis. *Environmentl Geology*, 39(10), 1095-1101.
- Guérin, T., Chekri, R., Vastel, C., Sirot, V., Volatier, J., Leblanc, J., & Noël, L. (2011). Determination of 20 trace elements in fish and other seafood from the French market. *Food Chemistry*, 127, 934-942.
- Gumpu, M. B., Sethuraman, S., Krishnan, U. M., & Rayappan, J. B. B. (2015). A review on detection of heavy metal ions in water – An electrochemical approach. *Sensors and Actuators B*, 213, 515-533.



- Guo, W., Liu, X., Liu, Z., & Li, G. (2010). Pollution and Potential Ecological Risk Evaluation of Heavy Metals in the Sediments around Dongjiang Harbor, Tianjin. *Procedia Environmental Sciences*, 2, 729-736.
- Håkanson, L. (1980). An ecological risk index for aquatic pollution control. A sedimentary approach. *Water Research*, 14, 975-1001.
- Hamed, M. A., & Emara, A. M. (2006). Marine molluscs as biomonitors for heavy metal levels in the Gulf of Suez, Red Sea. *Journal of Marine Systems*, 60(3-4), 220-234.
- Hang, X. S., Wang, H. Y., Zhou, J. M., Ma, C. L., Du, C. W., & Chen, X. Q. (2009). Risk assessment of potentially toxic element pollution in soils and rice (*Oryza sativa*) in a typical area of the Yangtze River Delta. *Environmental Pollution*, 157, 2542-2549.
- Haris, H., & Aris, A. Z. (2013). The geoaccumulation index and enrichment factor of mercury in mangrove sediment of Port Klang, Selangor, Malaysia. *Arabian Journal of Geosciences*, 6, 4119-4128.
- HKDJ (Hong Kong Department of Justice) (2000). The Judiciary of the Hong Kong Special Administrative Region (HKSAR). Food Adulteration (metallic contamination) Regulations.
- Höllriegel, V. & München, H. Z. (2011). Strontium in the environment and possible human health effects. *Reference Module in earth System and Environmental Sciences, Encyclopedia of Human Health*, 2011, 268-275.
- Hseu, Z. (2004). Evaluating heavy metal contents in nine composts using four digestion methods. *Bioresource Technology*, 95, 53-59.
- IARC (International Agency for Research on Cancer) (2004). IARC monographs on the evaluation of carcinogenic risks to humans. Volume 88. Lyon, France.
- Ikem, A., & Egiebor, N. O. (2005). Assessment of trace elements in canned fishes (mackerel, tuna, salmon, sardines and herrings) marketed in Georgia and Alabama (United States of America). *Journal of Food Composition and Analysis*, 18, 771-787.
- Iqbal, J., & Shah, M. H. (2014). Study of seasonal variations and health risk assessment of heavy metals in *Cyprinus carpio* from Rawal Lake, Pakistan. *Environmental Monitoring and Assessment*, 186, 2025-2037.
- Ireland, M. P. (1986). Studies on the effects of dietary beryllium at two different calcium concentrations in *Achatina fulica* (pulmonata). *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology*, 83(2), 435-438.
- Ireland, M. P. (1988). A comparative study of the uptake and distribution of silver in a slug *Arion ater* and a snail *Achatina fulica*. *Comparative Biochemistry and Physiology Part C: Comparative Pharmacology*, 90(1), 189-194.

- IRIS (2014). Integrated risk information system. US Environmental Protection Agency. [Accessed on November 2014]. <http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showSubstanceList>
- Irwin, R. J., VanMouwerik, M., Stevens, L., Seese, M. D., & Basham, W. (1997). Environmental Contaminants Encyclopedia. National Park Service, Water Resources Division, Fort Collins, Colorado. Distributed within the Federal Government as an Electronic Document. [Accessed on December 2015]. <http://www.nature.nps.gov/hazardssafety/toxic/strontiu.pdf>
- Islam, M. S., & Tanaka, M. (2004). Impacts of pollution on coastal and marine ecosystems including coastal and marine fisheries and approach for management: A review and synthesis. *Marine Pollution Bulletin*, 48, 624-649.
- Ismail, A., & Ramli, R. (1997). Trace metals in sediments and molluscs from an estuary receiving pig farms effluents. *Environmental Technology*, 18, 509-515.
- Isobe, K. O., Tarao, M., Zakaria, M. P., Chiem, N. H., Minh, L. Y., & Takada, H. (2002). Quantitative application of fecal sterols using gas chromatography-mass spectrometry to investigate fecal pollution in tropical waters: Western Malaysia and Mekong Delta, Vietnam. *Environmental Science & Technology*, 36, 4497-4507.
- Isobe, T., Takada, H., Kanai, M., Tsutsumi, S., Isobe, K. O., Boonyatumanond, R., & Zakaria, M. P. (2007). Distribution of Polycyclic Aromatic Hydrocarbons (PAHs) and phenolic endocrine disrupting chemicals in South and Southeast Asian mussels. *Environmental Monitoring and Assessment*, 135, 423-440.
- Ivanov, S. (2006). *Theoretical and Quantum Mechanics Fundamentals for Chemists*. The Netherlands: Springer.
- Jabbarzadeh, Z., Khosh-khui, M., Salehi, H., & Saberivand, A. (2010). Inter simple sequence repeat (ISSR) markers as reproducible and specific tools for genetic diversity analysis of rose species. *African Journal of Biotechnology*, 9(37), 6091-6095.
- Javed, M., & Usmani, N. (2014). Assessment of heavy metals (Cu, Ni, Fe, Co, Mn, Cr, Zn) in rivulet water, their accumulations and alterations in hematology of fish *Channa punctatus*. *African Journal of Biotechnology*, 13(3), 492-501.
- Jilani, S. (2014). Status of Metal Pollution in the River and Coastal Areas of Karachi. *Middle-East Journal of Scientific Research*, 22(9), 1288-1293.
- Joshi, S. P., Gupta, V. S., Aggarwal, R. K., Ranjekar, P. K., & Brar, D. S. (2000). Genetic diversity and phylogenetic relationship as revealed by inter-simple sequence repeat (ISSR) polymorphism in the genus *Oryza*. *Theoretical and Applied Genetics*, 100, 1311-1320.
- Jović, M., & Stanković, S. (2014). Human exposure to trace metals and possible public health risks via consumption of mussels *Mytilus galloprovincialis* from the Adriatic coastal area. *Food and Chemical Toxicology*, 70, 241-251.

- Kalender, L., & Uçar, S. Ç. (2013). Assessment of metal contamination in sediments in the tributaries of the Euphrates River, using pollution indices and the determination of the pollution source, Turkey. *Journal of Geochemical Exploration*, 134, 73-84.
- Kanakaraju, D., & Anuar, A. (2009). Accumulation and Depuration of Lead and Chromium Using *Nerita lineata*. *World Applied Sciences Journal*, 6(9), 1205-1208.
- Kang, S. G., Wright, D. A., & Koh, C. H. (2000). Baseline metal concentration in the Asian periwinkle *Littorina brevicula* employed as a biomonitor to assess metal pollution in Korean coastal water. *The Science of the Total Environment*, 263, 143-153.
- Karditsa, A., Poulos, S. E., Botsou, F., Alexakis, D., & Stamatakis, M. (2014). Investigation of major and trace element distribution patterns and pollution status of the surficial sediments of a microtidal inner shelf influenced by a transboundary river. The case of the Alexandroupolis Gulf (northeastern Aegean Sea, Greece). *Journal of Geochemical Exploration*, 146, 105-118.
- Karickhoff, S. W., Brown, D. S., & Scott, T. A. (1978). Sorption of hydrophobic pollutants of natural sediments. *Water Research*, 13, 241-248.
- Khalili, N. R., Scheff, P. A. and Holsen, T. M. (1995). PAH source fingerprinting for coke ovens, diesel and gasoline engines, highway tunnels, and wood combustion emissions. *Atmospheric Environment*, 29, 533-542.
- Khillare, P. S., Jyethi, D. S., & Sarkar, S. (2012). Health risk assessment of polycyclic aromatic hydrocarbons and heavy metals via dietary intake of vegetables grown in the vicinity of thermal power plants. *Food and Chemical Toxicology*, 50, 1642-1652.
- Kiss, T., & Osipenko, O. N. (1994). Toxic effects of heavy metals on ionic channels. *Pharmacological Reviews*, 46(3), 245-263.
- Külahci, F., & Şen, Z. (2008). Multivariate statistical analyses of artificial radionuclides and heavy metals contaminations in deep mud of Keban Dam Lake, Turkey. *Applied Radiation and Isotopes*, 66, 236-246.
- Kumar, K. S., Sajwan, K. S., Richardson, J. P., & Kannan, K. (2008). Contamination profiles of heavy metals, organochlorine pesticides, polycyclic aromatic hydrocarbons and alkylphenols in sediment and oyster collected from marsh/estuarine Savannah GA, USA. *Marine Pollution Bulletin*, 56, 136-162.
- Kumar, K. S., Dahms, H. U., Won, E. J., Lee, J. S., & Shin, K. H. (2015). Microalgae – A promising tool for heavy metal remediation. *Ecotoxicology and Environmental Safety*, 113, 329-352.
- Lam, P. K. S. (2009). Use of biomarkers in environmental monitoring. *Ocean & Coastal Management*, 52(7), 348-354.

- Larsen, R. K., & Baker, J. E. (2003). Source apportionment of polycyclic aromatic hydrocarbons in the urban atmosphere: a comparison of three methods. *Environmental Science & Technology*, 37, 1873-1881.
- Lavie, B., & Nevo, E. (1986). Genetic selection of homozygote allozyme genotypes in marine gastropods exposed to cadmium pollution. *The Science of the Total Environment*, 57, 91-98.
- Li, J., Huang, Z. Y., Hu, Y., & Yang, H., (2013). Potential risk assessment of heavy metals by consuming shellfish collected from Xiamen, China. *Environmental Science and Pollution Research*, 20, 2937-2947.
- Li, C., Huo, S., Yu, Z., Xi, B., Zeng, X., & Wu, F. (2014). Spatial distribution, potential risk assessment, and source apportionment of polycyclic aromatic hydrocarbons (PAHs) in sediments of Lake Chaohu, China. *Environmental Science and Pollution Research*, 21, 12028-12039.
- Li, J., Dong, H., Zhang, D., Han, B., Zhu, C., Liu, S., Liu, X., Ma, Q., & Li, X. (2015). Sources and ecological risk assessment of PAHs in surface sediments from Bohai Sea and northern part of the Yellow Sea, China. *Marine Pollution Bulletin*, 96, 485-490.
- Lidén, C., Bruze, M., Thyssen, J. P., & Menné, T. (2006). Metals. In: Contact Dermatitis ed. J.D. Johansen et al. 643-679. Springer-Verlag Berlin Heidelberg.
- Lokeshappa, B., Shivpuri, K., Tripathi, V., & Dikshit, A. K. (2012). Assessment of Toxic Metals in Agricultural Produce. *Food and Public Health*, 2(1), 24-29.
- Long, E. R., MacDonald, D. D., Smith, S. L., & Calder, F. D. (1995). Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management* 19, 81-97.
- Long, E. R. and MacDonald, D. D., (1998). Recommended uses of empirically derived, sediment quality guidelines for marine and estuarine ecosystems. *Human and Ecological Risk Assessment*, 4, 1019-1039.
- Long, E. R., MacDonald, D. D., Severn, C. G., & Hong, C. B. (2000). Classifying probabilities of acute toxicity in marine sediments with empirically derived sediment quality guidelines. *Environmental Toxicology and Chemistry*, 19(10), 2598-2601.
- MacDonald, D. D., Carr, R. S., Calder, F. D., Long, E. R., & Ingersoll, C. G. (1996). Development and evaluation of sediment quality guidelines for Florida coastal waters. *Ecotoxicology*, 5, 253-278.
- MAFF (Ministry of Agriculture, Fisheries and Food) (2000). Monitoring and surveillance of non-radioactive contaminants in the aquatic environment and activities regulating the disposal of wastes at sea, 1997. Aquatic environment monitoring report number 52. Lowestoft: MAFF.

- Maie, N., Pisani, O., & Jaffe, R. (2008). Mangrove tannins in aquatic ecosystems: Their fate and possible influence on dissolved organic carbon and nitrogen cycling. *Limnology and Oceanography*, 53(1), 160-171.
- Maiz, I., Arambarri, I., Garcia, R., & Millán, E. (2000). Evaluation of heavy metal availability in polluted soils by two sequential extraction procedures using factor analysis. *Environmental Pollution*, 110, 3-9.
- Markert, B. A., Breure, A. M., & Zechmeister, H. G. (2003). Definitions, strategies and principles for bioindication/biomonitoring of the environment. *Bioindicators and biomonitors*, 6, 3-39.
- Mehdinia, A., Aghadadashi, V., & Fumani, N. S. (2015). Origin, distribution and toxicological potential of polycyclic aromatic hydrocarbons in surface sediments from the Bushehr coast, The Persian Gulf. *Marine Pollution Bulletin*, 90, 334-338.
- Mertz, W. (1981). The essential trace elements. *Science*, 213, 1332-1338.
- Meyer, W., Seiler, T. B., Christ, A., Redelstein, R., Püttmann, W., Hollert, H., & Achten, C. (2014). Mutagenicity, dioxin-like activity and bioaccumulation of alkylated picene and chrysene derivatives in a German lignite. *Science of the Total Environment*, 497-498, 634-641.
- MFR (Malaysian Food Regulation) (1985). *Malaysian law on food and drugs*. Kuala Lumpur: Malaysian Law Publishers.
- Mirsadeghi, S. A. (2010). *Bioaccumulation and hazard assessment of polycyclic aromatic hydrocarbons in cockle (Anadara granosa) from selected intertidal mudflats of Peninsular Malaysia*. PhD Thesis, Universiti Putra Malaysia.
- Mirsadeghi, S. A., Zakaria, M. P., Yap, C. K., & Shahbazi, A. (2011). Risk assessment for the daily intake of polycyclic aromatic hydrocarbons from the ingestion of cockle (*Anadara granosa*) and exposure to contaminated water and sediments along the west coast of Peninsular Malaysia. *Journal of Environmental Sciences*, 23(2), 336-345.
- Mirsadeghi, S. A., Zakaria, M. P., Yap, C. K., & Gobas, F. (2013). Evaluation of the potential bioaccumulation ability of the blood cockle (*Anadara granosa* L.) for assessment of environmental matrices of mudflats. *Science of the Total Environment*, 454-455, 584-597.
- Mourier, B., Fritsch, C., Dhivert, E., Gimbert, F., Coeurdassier, M., Pauget, B., de Vaufleury, A., & Scheifler, R. (2011). Chemical extractions and predicted free ion activities fail to estimate metal transfer from soil to field land snails. *Chemosphere*, 85, 1057-1065.
- Muel, B., & Saguem, S. (1985). Determination of 23 polycyclic aromatic hydrocarbons in atmospheric particulate matter of the Paris area and photolysis by sunlight. *International Journal of Environmental Analytical Chemistry*, 19(2), 111-131.
- Muller, G. (1969). Index of geoaccumulation in sediments of the Rhine River. *GeoJournal*, 2, 108-118.

- Munoz, O., Devesa, V., Suner, M. A., Velez, D., Montoro, R., Urieta, I., Macho, M. L., & Jalon, M. (2000). Total and inorganic arsenic in fresh and processed fish products. *Journal of Agricultural and Food Chemistry*, 48, 4369-4376.
- Næs, K., Oug, E., & Knutzen, J. (1998). Source and species-dependent accumulation of polycyclic aromatic hydrocarbons (PAHs) in littoral indicator organisms from Norwegian smelter-affected marine waters. *Marine Environmental Research*, 45, 193-207.
- Naser, H. A. (2013). Assessment and management of heavy metal pollution in the marine environment of the Arabian Gulf: A review. *Marine Pollution Bulletin*, 72, 6-13.
- Nasher, E., Lee, Y. H., Zakaria, Z., & Surif, S. (2013). Assessing the Ecological Risk of Polycyclic Aromatic Hydrocarbons in Sediments at Langkawi Island, Malaysia. *The Scientific World Journal*, 2013, 1-13.
- Neff, J. M. (1979). *Polycyclic aromatic hydrocarbons in the aquatic environment: Source, fates and biological effect*. London: Applied Science Publisher LTD.
- Nei, M., & Li, W. H. (1979) Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences*, 76, 5269-5273.
- Neşer, G., Kontas, A., Ünsalan, D., Altay, O., Darılmaz, E., Uluturhan, E., Küçüksezgin, F., Tekoğul, N., & Yercan, F. (2012). Polycyclic aromatic and aliphatic hydrocarbons pollution at the coast of Aliağa (Turkey) ship recycling zone. *Marine Pollution Bulletin*, 64, 1055-1059.
- Ng, W. L., & Tan, S. G. (2015). Inter-simple sequence repeat (ISSR) markers: Are we doing it right? *ASM Science Journal*, 9(1), 48-57.
- Ng, J. C. (2005). Environmental contamination of arsenic and its toxicological impact on humans. *Environmental Chemistry*, 2, 146-160.
- Nie, J., Shi, J., Duan, X., Wang, B., Huang, N., & Zhao, X. (2014). Health risk assessment of dietary exposure to polycyclic aromatic hydrocarbons in Taiyuan, China. *Journal of Environmental Sciences*, 26, 432-439.
- Nisbet, I. C. T., & Lagoy, P. K. (1992). Toxic equivalency factors (TEFs) for polycyclic aromatic hydrocarbons (PAHs). *Regulatory Toxicology and Pharmacology*, 16, 290-300.
- Niu, H., Deng, W., Wu, Q., & Chen, X. (2009). Potential toxic risk of heavy metals from sediment of the Pearl River in South China. *Journal of Environmental Sciences*, 21, 1053-1058.
- Nkongolo, K. K., Gervais, S., Michael, P., & Zhou, Y. (2014). Comparative analysis of inter simple sequence repeats and simple sequence repeats markers: Genetic analysis of *Deschampsia cespitosa* populations growing in metal contaminated

- regions in Canada. *American Journal of Biochemistry and Biotechnology*, 10(1), 69-80.
- Nordberg, G. F., Fowler, B. A., Nordberg, M., & Friberg, L. T. (2007). *Handbook on the toxicology of metals* (3rd ed.). Academic Press is an imprint of Elsevier.
- Oehlmann, J., & Schulte-Oehlmann, U. (2003). Molluscs as bioindicators. In B. A. Markert, A. M. Breure, & H. G. Zechmeister (Eds.), *Bioindicators and biomonitors* (pp. 557-635). Elsevier Science Ltd.
- Oen, A. M. P., Schaanning, M., Ruus, A., Cornelissen, G. Källqvist, T., & Breedveld, G. D. (2006). Predicting low biota to sediment accumulation factors of PAHs by using infinite-sink and equilibrium extraction methods as well as BC-inclusive modeling. *Chemosphere*, 64, 1412-1420.
- Ogata, Y., Takada, H., K., Mizukawa, Hirai, H., Iwasa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M. P., Dung, L. Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagioti, H. K., Weerts, S., McClurg, T., Burres, E., Smith, W., Van Velkenburg, M., Lang, J. S., Lang, R. C., Laursen, D., Danner, B., Stewardson, N., & Thompson, R. C. (2009). International Pellet Watch: Global monitoring of persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs. *Marine Pollution Bulletin*, 58(10), 1437-1446.
- Ong, G. H. (2012). *Effects of heavy metals accumulation on biological activities and genetic variation of Centella asiatica (L.) urban*. Unpublished PhD Thesis, Universiti Putra Malaysia.
- Otachi, E. O., Körner, W., Avenant-Oldewage, A., Fellner-Frank, C., & Jirsa, F. (2014). Trace elements in sediments, blue spotted tilapia *Oreochromis leucostictus* (Trewavas, 1933) and its parasite *Contracaecum multipapillatum* from Lake Naivasha, Kenya, including a comprehensive health risk analysis. *Environmental Science and Pollution Research*, 21, 7339-7349.
- Otitoloju, A. A. (2002). Evaluation of the joint-action toxicity of binary mixtures of heavy metals against the mangrove periwinkle *Tympanotonus fuscatus* var. *radula* (L.). *Ecotoxicology and Environmental Safety*, 53, 404-415.
- Pan, K., & Wang, W. X. (2011). Trace metal contamination in estuarine and coastal environments in China. *Science of the Total Environment*, 421-422, 3-16.
- Paramasivam, K. Ramasamy, V., & Suresh, G. (2015). Impact of sediment characteristics on the heavy metal concentration and their ecological risk level of surface sediments of Vaigai River, Tamilnadu, India. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 137, 397-407.
- Parengam, M., Judprasong, K., Srianujata, S., Jittinandana, S., Laoharajanaphand, S., & Busamongko, A. (2010). Study of nutrients and toxic minerals in rice and legumes by instrumental neutron activation analysis and graphite furnace

atomic absorption spectrophotometry. *Journal of Food Composition and Analysis*, 23, 340-345.

- Park, J., & Presley, B. J. (1997). Trace metal contamination of sediments and organisms from the Swan Lake area of Galveston Bay. *Environmental Pollution*, 98(2), 209-221.
- Pearce, N. J. G., & Mann, V. L. (2006). Trace metal variations in the shells of *Ensis siliqua* record pollution and environmental conditions in the sea to the west of mainland Britain. *Marine Pollution Bulletin*, 52, 739-755.
- Perkins, J., & Gadd, G. M. (1993). Caesium toxicity, accumulation and intracellular localization in yeasts. *Mycological Research*, 97(6), 717-724.
- Peters, C. A., Knightes, C. D., & Brown, D. G. (1999). Long-term composition dynamics of PAH-containing NAPLs and implications for risk assessment," *Environmental Science and Technology*, 33(24), 4499-4507.
- Petry, T., Schmid, P., & Schlatter, C. (1996). The use of toxic equivalency factors in assessing occupational and environmental health risk associated with exposure to airborne mixtures of polycyclic aromatic hydrocarbons (PAHs). *Chemosphere*, 32(4), 639-648.
- Prasad, A. S., (1983). The role of zinc in gastrointestinal and liver disease. *Clinics in Gastroenterology*, 12, 713-741.
- Rachdawong, P., Christensen, E. R., & Karls, J. F. (1998). Historical PAH fluxes to Lake Michigan sediments determined by factor analysis. *Water Research*, 32(8), 2422-2430.
- Radwan, M. A., El-Gendy, K. S., & Gad, A. F. (2010). Biomarkers of oxidative stress in the land snail, *Theba pisana* for assessing ecotoxicological effects of urban metal pollution. *Chemosphere*, 79, 40-46.
- Rainbow, P. S., Phillips, D. J. H., & Depledge, M. H. (1990). The significance of trace metal concentrations in marine invertebrates: A need for laboratory investigation of accumulation strategies. *Marine Pollution Bulletin*, 21(7), 321-324.
- Rainbow, P. S. (1995). Biomonitoring of heavy metal availability in the marine environment. *Marine Pollution Bulletin*, 31, 183-192.
- Rainbow, P. S. (1997). Trace metal accumulation in marine invertebrates: marine biology or marine chemistry? *Journal Marine Biology Association UK*, 77, 195-210.
- Rainbow, P. S. (2002). Trace metal concentrations in aquatic invertebrates: why and so what? *Environmental Pollution*, 120, 497-507.
- Rainbow, P. S. (2007). Trace metal bioaccumulation: Models, metabolic availability and toxicity. *Environment International*, 33, 576-582.



- Real, C., Barreiro, R., & Carballeira, A. (1994). The application of microwave heating in sequential extractions of heavy metals in estuarine sediments. *The Science of the Total Environment*, 152, 135-142.
- Retnam, A. (2013). *Distribution, characterization and risk assessment of polycyclic aromatic hydrocarbons in selected aquaculture farms in Peninsular Malaysia: A forensic and chemometric approach*. Unpublished PhD Thesis, Universiti Putra Malaysia.
- Retnam, A., Zakaria, M. P., Juahir, H., Aris, A. Z., Zali, M. A., & Kasim, M. F. (2013). Chemometric techniques in distribution, characterization and source apportionment of polycyclic aromatic hydrocarbons (PAHS) in aquaculture sediments in Malaysia. *Marine Pollution Bulletin*, 69, 55-66.
- Rezaee, K. H., Abdi, M. R., Saion, E. B., Naghavi, K., & Shafaei, M. A. (2011). Distribution of trace elements in the marine sediments along the South China Sea, Malaysia. *Journal of Radioanalytical and Nuclear Chemistry*, 287, 733-740.
- Rittmann, B. E., & McCarty, P. L. (2001). *Environmental Biotechnology: Principles and applications*. New York: McGraw-Hill Companies Inc.
- Rizo, O. D., Reumont, S. O., Fuente, J. V., Arado, O. D., Pino, N. L., Rodríguez, K. D., López, J. O. A., Rudnikas, A. G., & Carballo, G. A. (2010). Copper, Zinc and Lead Bioaccumulation in Marine Snail, *Strombus gigas*, from Guacanayabo Gulf, Cuba. *Bulletin of Environmental Contamination & Toxicology*, 85, 330-333.
- Rudnick, R. L., & Gao, S., (2003). Composition of the continental crust. In R.L. Rudnick (Ed.), *The Crust* (pp. 1-64). Elsevier Science Ltd.
- Ruus, A., Schaanning, M., Øxnevad, S., & Hylland, K. (2005). Experimental results on bioaccumulation of metals and organic contaminants from marine sediments. *Aquatic Toxicology*, 72, 273-292.
- Rossmann, T. G. (2003). Mechanism of arsenic carcinogenesis: An integrated approach. *Mutation Research*, 533, 37-65.
- Saha, M., Togo, A., Mizukawa, K., Murakami, M., Takada, H., Zakaria, M. P., Chiem, N. H., Tuyen, B. C., Prudente, M., Boonyatumanond, R., Sarkar, S. K., & Bhattacharya, B. (2009). Sources of sedimentary PAHs in tropical Asian waters: Differentiation between pyrogenic and petrogenic sources by alkyl homolog abundance. *Marine Pollution Bulletin*, 58, 189-200.
- Sarkar, A., Ray, D., Shrivastava, A. N., & Sarker, S. (2006). Molecular biomarkers: Their significance and application in marine pollution monitoring. *Ecotoxicology*, 15, 333-340.
- Sarkar, A., Dipak, C. S. G., Amit, S., Vashisthaa, D., Classy, D., & Dalala, S. G. (2008). Evaluation of impairment of DNA integrity in marine gastropods (*Cronia contracta*) as a biomarker of genotoxic contaminants in coastal water

around Goa, West coast of India. *Ecotoxicology and Environmental Safety*, *71*, 473-482.

- Sakan, S., Dević, G., Relić, D., Anđelković, I., Sakan, N., & Đorđević, D. (2014). Risk assessment of trace element contamination in river sediments in Serbia using pollution indices and statistical methods: a pilot study. *Environmental Earth Sciences*, *73*(10), 6625-6638.
- Sany, S. B. T., Salleh, A., Sulaiman, A. H., Sasekumar, A., Tehrani, G., & Rezayi, M. (2012). Distribution characteristics and ecological risk of heavy metals in surface sediments of West Port, Malaysia. *Environment Engineering Protection*, *38*(4), 139-155.
- Sawasdee, B., Köhler, H. R., & Triebkorn, R. (2011). Histopathological effects of copper and lithium in the ramshorn snail, *Marisa cornuarietis* (Gastropoda, Prosobranchia). *Chemosphere*, *85*, 1033-1039.
- Schi, K. C., & Weisberg, S. B. (1999). Iron as a reference element for determining trace metal enrichment in Southern California coast shelf sediments. *Marine Environmental Research*, *48*, 161-176.
- Schrauzer, G. N. (2002). Lithium: Occurrence, dietary intakes, nutritional essentiality. *Journal of the American College of Nutrition*, *21*(1), 14-21.
- Shahbazi, A. (2009). *Distribution and sources of petroleum hydrocarbons in green-lipped mussels (Perna viridis) from selected coastal waters of Peninsular Malaysia*. Unpublished PhD Thesis, Universiti Putra Malaysia.
- Shahbazi, A., Zakaria, M. P., Yap, C. K., Tan, S. G., Surif, S., Mohamed, C. A. R., Sakari, M., Bakhtiari, A. R., Bahry, P. S., Chandru, K., & Mirsadeghi, S. A. (2010a). Use of Different Tissues of *Perna Viridis* as Biomonitor of polycyclic aromatic hydrocarbons (PAHs) in the coastal waters of Peninsular Malaysia. *Environmental Forensics*, *11*, 248-263.
- Shahbazi, A., Zakaria, M. P., Yap, C. K., Surif, S., Bakhtiari, A. R., Chandru, K., Bahry, P. S., & Sakari, M. (2010b). Spatial distribution and sources of polycyclic aromatic hydrocarbons (PAHs) in green mussels (*Perna viridis*) from coastal areas of Peninsular Malaysia: implications for source identification of perylene. *International Journal of Environmental Analytical Chemistry*, *90*(1), 14-30.
- Shafie, N. A., Aris, A. Z., Zakaria, M. P., Haris, H., Lim, W. Y., and Isa, N. M. (2013). Application of geoaccumulation index and enrichment factors on the assessment of heavy metal pollution in the sediments. *Journal of Environmental Science and Health, Part A*, *48*, 182-190.
- Shaw, D. M., Dostal, J., & Keays, R. R. (1976). Additional estimates of continental surface Precambrian shield composition in Canada. *Geochimica et Cosmochimica Acta*, *40*, 73-84.

- Shehata, A. I., Al-Ghethar, H. A., & Al-Homaidan, A. A. (2009). Application of simple sequence repeat (SSR) markers for molecular diversity and heterozygosity analysis in maize inbred lines. *Saudi Journal of Biological Sciences*, 16, 57-62.
- Showler, A. T. James, W. D. Armstrong, J. S., & Westbrook, J. K. (2006). An experiment using neutron activation analysis and a rare earth element to mark cotton plants and two insects that feed on them. *Applied Radiation and Isotopes*, 64, 875-880.
- Shukla, S. P., & Sharma, M. (2010). Neutralization of rainwater acidity at Kanpur, India. *Tellus*, 62B, 172-180.
- Siddiquee, S., Tan, S. G., Yusuf, U. K., Fatihah, N. H. N., & Hasan, M. M. (2012). Characterization of Malaysian *Trichoderma* isolates using random amplified microsatellites (RAMS). *Molecular Biology Reports*, 39, 715-722.
- Sivaperumal, P., Sankar, T. V., & Nair, P. G. V. (2007). Heavy metal concentrations in fish, shellfish and fish products from internal markets of India vis-a-vis international standards. *Food Chemistry*, 102, 612-620.
- Słomka, A., Sutkowska, A., Szczepaniak, M., Malec, P., Mitka, J., & Kuta, E. (2011). Increased genetic diversity of *Viola tricolor* L. (Violaceae) in metal-polluted environments. *Chemosphere*, 83, 435-442.
- Sparks, D. L., (2005). Toxic metals in the environment: The role of surfaces. *Elements*, 1, 193-197.
- Spencer, K. L., MacLeod, C. L., Tuckett, A., & Johnson, S. M. (2006). Source and distribution of trace metals in the Medway and Swale estuaries, Kent, UK. *Marine Pollution Bulletin*, 52, 214-238.
- Sutherland, R. A. (2000). Bed sediment-associated trace metals in an urban stream, Oahu, Hawaii. *Environmental Geology*, 39(6), 611-627.
- Srogi, K. (2007). Monitoring of environmental exposure to polycyclic aromatic hydrocarbons: a review. *Environmental Chemistry Letters*, 5, 169-195.
- Stafne, E. T., Clark, J. R., Weber, C. A., Graham, J., & Lewers, K. S. (2005). Simple Sequence Repeat (SSR) Markers for Genetic Mapping of Raspberry and Blackberry. *Journal of the American Society for Horticultural Science*, 130(5), 722-728.
- Storelli, M. M. (2008). Potential human health risks from metals (Hg, Cd, and Pb) and polychlorinated biphenyls (PCBs) via seafood consumption: Estimation of target hazard quotients (THQs) and toxic equivalents (TEQs). *Food and Chemical Toxicology*, 46, 2782-2788.
- Sukumaran, S., & Grant, A. (2013). Effects of genotoxicity and its consequences at the population level in sexual and asexual *Artemia* assessed by analysis of inter-simple sequence repeats (ISSR). *Mutation Research*, 757, 8-14.

- Sychra, J., Čelechovská, O., Svobodová, Z., & Sychra, O. (2011). Lead, mercury and cadmium content in bottom sediments, reed (*Phragmites australis*) beds and great pond snails (*Lymnaea stagnalis*) in fishponds and the role of littoral zones in their accumulation. *Acta Veterinaria Brno*, 80, 313-321.
- Szefer, P., Ali, A. A., Ba-Haroon, A. A., Rajeh, A. A., Geldon, J., & Nabrzyski, M. (1999). Distribution and relationships of selected trace metals in molluscs and associated sediment from the Gulf of Aden, Yemen. *Environmental Pollution*, 106, 299-314.
- Tanguy, A., Boutet, I., Bonhomme, F., Boudry, P., & Moraga, D. (2002). Polymorphism of metallothionein genes in the Pacific oyster *Crassostrea gigas* as a biomarker of response to metal exposure. *Biomarkers*, 7(6), 439-450.
- Taylor, S. R., & McLennan, S. M. (1995). The geochemical evolution of the continental crust. *Reviews of Geophysics*, 33, 241-265.
- Tessier, A., & Campbell, P. G. C. (1987). Partitioning of trace metals in sediments: relationships with bioavailability. *Hydrobiology*, 149, 43-52.
- Thompson, P. A., Kurias, J., & Mihok, S. (2005). Derivation and use of sediment quality guidelines for ecological risk assessment of metals and radionuclides released to the environment from uranium mining and milling activities in Canada. *Environmental Monitoring and Assessment*, 110, 71-85.
- Topcuoğlu, S., Kırbaşoğlu, Ç., & Güngör, N. (2002). Heavy metals in organisms and sediments from Turkish Coast of the Black Sea, 1997–1998. *Environment International*, 27, 521-526.
- Tranvik, L., Sjogren, M., & Bengtsson, G. (1994). Allozyme polymorphism and protein profile in *Orchesella bifasciata* (Collembola)" Indicative of Extended metal pollution? *Biochemical Systematics and Ecology*, 22(1), 13-23.
- UNESCO (1976). A comprehensive plan for global investigation of pollution in the marine environment and baseline study guidelines. [Accessed on June 2015]. [http://www.jodc.go.jp/info/ioc\\_doc/Technical/021095eo.pdf](http://www.jodc.go.jp/info/ioc_doc/Technical/021095eo.pdf)
- Ungherese, G., Mengoni, A., Somigli, S., Baroni, D., Focardi, S., & Ugolini, A. (2010). Relationship between heavy metals pollution and genetic diversity in Mediterranean populations of the sandhopper *Talitrus saltator* (Montagu) (Crustacea, Amphipoda). *Environmental Pollution*, 158, 1638-1643.
- USEPA (US Environmental Protection Agency) (1983). Methods for chemical analysis of water and waste. EPA Report 600/4-79-020. Cincinnati: Office of Water, USEPA.
- USEPA (US Environmental Protection Agency) (1989). Guidance manual for assessing human health risks from chemically contaminated, fish and shellfish. EPA-503/8-89-002. USEPA, Washington DC.

- USEPA (US Environmental Protection Agency) (2000). Risk-Based Concentration Table. Philadelphia PA: USEPA, Washington, DC.
- USEPA (US Environmental Protection Agency), (2002). Polycyclic organic matter. Washington: Environmental Protection Agency. [Accessed on March 2015]. <http://www.epa.gov/ttn/atw/hlthef/polycycl.html>
- USEPA (US Environmental Protection Agency) (2008). USEPA, Integrated Risk Information System. CRC. [Accessed on January 2015]. <http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.showSubstanceList>
- USEPA (US Environmental Protection Agency) (2009). Silver and compound summary document. [Accessed June 2015]. [http://epa.gov/oppsrrd1/registration\\_review/silver/index.html](http://epa.gov/oppsrrd1/registration_review/silver/index.html).
- USFDA (US Food and Drug Administration) (1993a). Food and drug administration, Guidance document for nickel in shell fish. DHHS/PHS/ FDA/CFSAN/office of seafood, Washington D.C.
- USFDA (US Food and Drug Administration) (1993b). Guidance Documents for Trace Elements in Seafood. Center for Food Safety and Applied Nutrition, Washington, DC, USA.
- USFDA (US Food and Drug Administration) (2011). Fish and fisheries products hazards and controls guidance, third ed. Center for food safety and applied nutrition. US Food and Drug Administration.
- Usman, A. R. A. (2008). The relative adsorption selectivities of Pb, Cu, Zn, Cd and Ni by soils developed on shale in New Valley, Egypt. *Geoderma*, 144, 334-343.
- Valadez-Vega, C., Zúñiga-Pérez, C., Quintanar-Gómez, S., Morales-González, J. A., Madrigal-Santillán, E., Villagómez-Ibarra, J. R., Sumaya-Martínez, M. T., & García-Paredes, J. D. (2011). Lead, cadmium and cobalt (Pb, Cd, and Co) leaching of glass-clay containers by pH effect of food. *International Journal of Molecular Sciences*, 12, 2336-2350.
- van Straalen, N. M. (2008). Contaminant concentrations in organisms as indicators of bioavailability: a review of kinetic theory and the use of target species in biomonitoring. *Developments in Soil Science*, 32, 449-477.
- Verhaert, V., Covaci, A., Bouillon, S., Abrantes, K., Musibono, D., Bervoets, L., Verheyen, E., & Blust, R. (2013). Baseline levels and trophic transfer of persistent organic pollutants in sediments and biota from the Congo River Basin (DR Congo). *Environment International*, 59, 290-302.
- Vijver, M., Vangestel, C. M., Lanno, R. P., Vanstraelen, N., & Peijnenburg, W. G. M. (2004). Internal metal sequestration and its ecotoxicological relevance: A review. *Environmental Science & Technology*, 38(18), 4705-4712.
- Walker, C. H., Hopkins, S. P., Sibly, R. M., & Peakall, D. B. (2001). *Principles of ecotoxicology*. London: Taylor and Francis.

- Wang, Y., Liang, L., Shi, J., & Jiang, G. (2005). Study on the contamination of heavy metals and their correlations in mollusks collected from coastal sites along the Chinese Bohai Sea. *Environment International*, 31, 1103-1113.
- Wang, Z., Yang, C., Parrott, J. L., Frank, R. A., Yang, Z., Brown, C. E., Hollebone, B. P., Landriault, M., Fieldhouse, B., Liu, Y., Zhang, G., & Hewitt, L. M. (2014). Forensic source differentiation of petrogenic, pyrogenic, and biogenic hydrocarbons in Canadian oil sands environmental samples. *Journal of Hazardous Materials*, 271, 166-177.
- Wang, Y., Yang, L., Kong, L., Liu, E., Wang, L., & Zhu, J. (2015). Spatial distribution, ecological risk assessment and source identification for heavy metals in surface sediments from Dongping Lake, Shandong, East China. *Catena*, 125, 200-205.
- Wedepohl, K. H. (1995). The composition of the continental crust. *Geochimica et Cosmochimica Acta*, 59(7), 1217-1232.
- Williams, C. (1996). Combating marine pollution from land-based activities: Australian initiatives. *Ocean and Coastal Management*, 33(1-3), 87-112.
- WHO (World Health Organization) (1989). Heavy metals-environmental aspects. Environment Health Criteria No. 85, Geneva.
- WHO (World Health Organization) (1990). Methylmercury. Environmental Health Criteria 101, 1-80.
- WHO (World Health Organization) (1996). Health criteria other supporting information. In Guidelines for drinking water quality, 2nd ed. (pp. 31-388). Geneva: WHO.
- WHO (World Health Organization) (2001). Beryllium and beryllium compounds. Concise international chemical assessment document: 32. Published under the joint sponsorship of the United Nations Environment Program, the International Labour Organization, and the WHO, and produced within the framework of the Inter-Organization Program for the Sound Management of Chemicals. Stuttgart.
- Yap, C. K. (2012). Application of factor analysis in geochemical fractions of heavy metals in the surface sediments of the offshore and intertidal areas of Peninsular Malaysia. *Sains Malaysiana*, 41(4), 389-394.
- Yap, C. K., Ismail, A., Tan, S. G., & Omar, H. (2002a). Concentration of Cu and Pb in the offshore and intertidal sediments of the west coast of Peninsular Malaysia. *Environment International* 28, 467-479.
- Yap, C. K., Ismail, A., & Omar, H. (2002b). Correlation between speciation of Cd, Cu, Pb and Zn in sediment and their concentrations in total soft tissue of green-lipped mussel (*Linnaeus*) from the west coast of Peninsular Malaysia. *Environment International*, 28, 117-126.

- Yap, C. K., Tan, S. G., Ismail, A., & Omar, H. (2004a). Allozyme polymorphisms and heavy metal levels in the green-lipped mussel *Perna viridis* (Linnaeus) collected from contaminated and uncontaminated sites in Malaysia. *Environment International*, 30, 39-46.
- Yap, C. K., & Cheng, W. H. (2008). Heavy metal concentrations in *Nerita lineata*: the potential as a biomonitor for heavy metal bioavailability and contamination in the tropical intertidal area. *Marine Biodiversity Records* 2, Doi: 10.1017/S1755267209000505
- Yap, C. K., Cheng, W. H., Ismail, A., Ismail, A. R., & Tan, S. G. (2009a). Biomonitoring of heavy metal (Cd, Cu, Pb, and Zn) concentrations in the west intertidal area of Peninsular Malaysia by using *Nerita lineata*. *Toxicological & Environmental Chemistry*, 91(1), 29-41.
- Yap, C. K., Edward, F. B., & Tan, S. G. (2009b). Similarities and differences of metal distributions in the tissues of molluscs by using multivariate analyses. *Environment Monitoring and Assessment*, 165(1-4), 39-53.
- Yap, C.K., Noorhaidah, A., Azlan, A., Nor Azwady, A. A., Ismail, A., Ismail, A. R., Siraj, S. S., & Tan, S. G. (2009c). *Telescopium telescopium* as potential biomonitors of Cu, Zn, and Pb for the tropical intertidal area. *Ecotoxicology and Environmental Safety*, 72, 496-506.
- Yap, C. K., Mohd Ruszaidi, S., Cheng, W. H., & Tan, S. G. (2010). Heavy-metal concentrations in the mangrove snail, *Nerita lineata* and surface sediments collected from Klang river estuary, Selangor, Malaysia. *Journal of Sustainability Science and Management*, 5(1), 1-12.
- Yap, C. K., & Cheng, W. H. (2010). Depuration of gut contents in the intertidal snail *Nerita lineata* is not necessary for the study of heavy metal contamination and bioavailability: A laboratory study. *Pertanika Journal of Tropical Agricultural Science*, 33(2), 167-170.
- Yap, C. K., Chong, C. M., & Tan, S. G. (2011a). Allozyme polymorphisms in horseshoe crabs, *Carcinoscorpius rotundicauda*, collected from polluted and unpolluted intertidal areas in Peninsular Malaysia. *Environmental Monitoring Assessment*, 174, 389-400.
- Yap, C. K., Azmizan, A. R., & Hanif, M. S. (2011b). Biomonitoring of trace metals (Fe, Cu, and Ni) in the mangrove area of Peninsular Malaysia using different soft tissues of flat tree oyster *Isognomon alatus*. *Water, Air, & Soil Pollution*, 218, 19-36.
- Yap, C. K., Shahbazi, A., & Zakaria, M. P. (2012). Concentrations of heavy metals (Cu, Cd, Zn and Ni) and PAHs in *Perna viridis* collected from seaport and non-seaport waters in the Straits of Johore. *Bulletin of Environmental Contamination and Toxicology*, 89, 1205-1210.

- Ylmaz, A. B., Sangun, M. K., Yag, D., & Turan, C., (2010). Metals (major, essential to non-essential) composition of the different tissues of three demersal fish species from Iskenderun Bay, Turkey. *Food Chemistry*, 123, 410-415.
- Yoon, E., Park, K., Lee, H., Yang, J., & Lee, C. (2007). Estimation of excess cancer risk on time-weighted lifetime average daily intake of PAHs from food ingestion. *Human and Ecological Risk Assessment*, 13, 669-680.
- Yu, K., Tsai, L., Chen, S., & Ho, S. (2001). Correlation analyses on binding behavior of heavy metals with sediment matrices. *Water Research*, 35(10), 2417-2428.
- Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D., & Sylvestre, S., (2002). PAHs in the Fraser river basin: a critical appraisal of PAHs ratios as indicator of PAHs sources and composition. *Organic Geochemistry*, 33, 489-515.
- Zakaria, M. P., Takada, H., Keiohno, S., Erikokouno, J., & Kumata, H. (2002). Distribution of Polycyclic Aromatic Hydrocarbons (PAHs) in Rivers and Estuaries in Malaysia: A Widespread Input of Petrogenic PAHs. *Environmental Science & Technology*, 36, 1907-1918.
- Zakaria, M. P., & Mahat, A. A. (2006). Distribution of polycyclic aromatic hydrocarbon (PAHs) in sediment in the Langat Estuary. *Coastal Marine Science*, 30(1), 387-395.
- Zar, J. H. (1996). *Biostatistical Analysis*. New Jersey: Prentice Hall.
- Zhang, Y., Yu, H., Xi, B., Hou, H., Gao, R., Liang, Q., Zhang, H., & Xu, F. (2014). Levels, sources, and potential ecological risks of polycyclic aromatic hydrocarbons (PAHs) in a typical effluent-receiving river (Wangyang River), North China. *Arabian Journal of Geosciences*, Doi: 10.1007/s12517-014-1695-1.
- Zheng, N., Wang, Q., Zhang, X., Zheng, D., Zhang, Z., & Zhang, S. (2007). Population health risk due to dietary intake of heavy metals in the industrial area of Huludao city, China. *Science of the Total Environment*, 387, 96-104.
- Zhao, G., Li, J., Zou, F., Mo, X., Yuan, Z., Lin, R., Weng, Y., & Zhu, X. (2009). ISSR, an effective molecular approach for studying genetic variability among *Schistosoma japonicum* isolates from different provinces in mainland China. *Infection, Genetics and Evolution*, 9, 903-907.
- Zhao, S., Feng, C., Yang, Y., Niu, J., & Shen, Z., (2012a). Risk assessment of sedimentary metals in the Yangtze Estuary: New evidence of the relationships between two typical index methods. *Journal of Hazardous Materials*, 241-242, 164-172.
- Zhao, G., Li, J. Blair, D., Li, X., Elsheikha, H. M., Lin, R., Zou, F., & Zhu, X. (2012b). Biotechnological advances in the diagnosis, species differentiation and



phylogenetic analysis of *Schistosoma* spp. *Biotechnology Advances*, 30, 1381-1389.

Zhao, L., Yang, F., & Yan, X. (2013). Concentrations of selected trace elements in marine bivalves and the dietary risk to residents of Dalian City, Northern China. *Human and Ecological Risk Assessment: An International Journal*, 19(1), 145-150.

Zhao, Z., Zhang, L., Cai, Y., & Chen, Y. (2014) Distribution of polycyclic aromatic hydrocarbon (PAH) residues in several tissues of edible fishes from the largest freshwater lake in China, Poyang Lake, and associated human health risk assessment. *Ecotoxicology and Environmental Safety*, 104, 323-331.

Zhou, Q., Zhang, J., Fu, J., Shi, J., & Jiang, G. (2008). Biomonitoring: An appealing tool for assessment of metal pollution in the aquatic ecosystem. *Analytica Chimica Acta*, 606, 135-150.

Zhu, H., Yuan, X., Zeng, G., Jiang, M., Liang, J., Zhang, C., Yin, J., Huang, H., Liu, Z., & Jiang, H. (2012). Ecological risk assessment of heavy metals in sediments of Xiawan Port based on modified potential ecological risk index. *Transaction Nonferrous Metals Society of China*, 2, 21470-1477.

Zietkiewicz, E., Rafalski, A., & Labuda, D. (1994). Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics*, 20, 176-183.