

UNIVERSITI PUTRA MALAYSIA

MODIFIED SPLINE FUNCTIONS AND CHEBYSHEV POLYNOMIALS FOR THE SOLUTION OF HYPERSINGULAR INTEGRALS PROBLEMS

LAWAN SIRAJO BICHI

FS 2015 40



MODIFIED SPLINE FUNCTIONS AND CHEBYSHEV POLYNOMIALS FOR THE SOLUTION OF HYPERSINGULAR INTEGRALS PROBLEMS



By

LAWAN SIRAJO BICHI

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of Philosophy

December 2015



COPYRIGHT

All materials contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.



DEDICATIONS

I dedicate this work of mine to my parents Alhaji Lawan Mai-goro and late Khadijah as well as my family. May the mercy of Allah be upon us all. Your support and courage is what made me who I am today. Thank you.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

MODIFIED SPLINE FUNCTIONS AND CHEBYSHEV POLYNOMIALS FOR THE SOLUTION OF HYPERSINGULAR INTEGRALS PROBLEMS

By

LAWAN SIRAJO BICHI

December 2015

Chair: Prof. Madya. Zainidin Eshkuvatov, PhD Faculty: Science

The research work studied the singular integration problems of the form

$$H_k(x,y) = \begin{cases} \iint\limits_{\Omega} \frac{h(x,y)}{|\bar{x} - \bar{x}_0^*|^{2-\gamma}} dA, 0 \le \gamma \le 1, \quad k = 1, \\ \iint\limits_{\Omega} h(x,y) \log |\bar{x} - \bar{x}_0^*| dA, \quad k = 2, \end{cases}$$

where $\Omega = [a_1, a_2] \times [b_1, b_2]$, $\bar{x} = (x, y) \in \Omega$ and fixed point $\bar{x}_0^* = (x_0^*, y_0^*) \in \Omega$. The density function h(x, y) is given, continuous and smooth on the rectangle Ω and belong to the class of functions $C^{2,\gamma}(\Omega)$.

Cubature formulas for double integrals with algebraic and logarithmic singularities on a rectangle Ω are constructed using the modified spline function $S_{\Lambda}(P)$ of type (0,2). Exactness of the cubature formulas for the two cases $k \in \{1,2\}$ together with tested examples are shown each for linear and quadratic functions. Highly accurate numerical results for the cubature formulas are given for both tested density function h(x, y) as linear and quadratic functions. The results are in line with the theoretical findings. Further more, Hadamard type hypersingular integral (HSI) of the form

$$H_i(h,x) = \frac{w_i(x)}{\pi} \int_{-1}^{1} \frac{h(t)}{w_i(t)(t-x)^2} dt, x \in (-1,1), i \in \{1,2,3,4\},$$

where
$$w_1(t) = \sqrt{1-t^2}$$
, $w_2(t) = \frac{1}{\sqrt{1-t^2}}$, $w_3(t) = \sqrt{\frac{1-t}{1+t}}$ and $w_4(t) = \sqrt{\frac{1+t}{1-t}}$

are the weights and h(t) is a smooth function, are considered. Automatic quadrature schemes (AQSs) in each case for $i \in \{1,2,3,4\}$ are constructed via approximating the density function h(t) by the first, second, third and fourth kind truncated series of Chebyshev polynomials, respectively. Error estimations in the cases $i \in \{1,2,3,4\}$ are obtained via approximating the density function by truncated series of Chebyshev polynomials of the first, second, third and fourth kind, respectively, in the class of function $C^{N,\alpha}[-1,1]$. Exactness of the methods each for $i \in \{1,2,3,4\}$ are shown for the degree 3 polynomial functions and the results of tested examples are presented and discussed. Numerical results of the obtained quadrature schemes revealed that the proposed methods are highly accurate for the tested density function h(t) as polynomial and rational functions. Comparisons made with other known methods showed that the automatic quadrature schemes (AQSs) constructed in this research has better results than others. The results are in line with the theoretical findings. Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

FUNGSI SPLIN TERUBAHSUAI DAN POLINOMIAL CHEBYSHEV BAGI PENYELESAIAN MASALAH KAMIRAN HIPERSINGULAR

Oleh

LAWAN SIRAJO BICHI

Disember 2015

Pengerusi: Prof. Madya. Zainidin Eshkuvatov, PhD Fakulti: Sains

Penyelidikan ini mengkaji masalah pengamiran singular dalam bentuk

$$H_k(x,y) = \begin{cases} \iint \frac{h(x,y)}{|\bar{x} - \bar{x}_0^*|^{2-\gamma}} dA, 0 \le \gamma \le 1, \quad k = 1, \\ \iint \int h(x,y) \log |\bar{x} - \bar{x}_0^*| dA, \quad k = 2, \end{cases}$$

dengan $\Omega = [a_1, a_2] \times [b_1, b_2], \ \bar{x} = (x, y) \in \Omega$ dan titik tetap $\bar{x}_0^* = (x_0^*, y_0^*) \in \Omega$. Fungsi ketumpatan h(x, y) diberi, selanjar dan mulus di atas segi empat Ω dan tergolong dalam kelas fungsi $C^{2,\gamma}(\Omega)$.

Formula pengkiuban untuk kamiran ganda dua dengan singulariti berbentuk aljabar dan logaritma di atas segiempat Ω dibina menggunakan fungsi splin terubahsuai $S_{\Lambda}(P)$ jenis (0,2). Ketepatan formula pengkiuban untuk keduadua kes $k \in \{1,2\}$ bersama-sama dengan contoh teruji bagi fungsi linear dan kuadratik ditunjukkan. Keputusan berangka yang sangat tepat bagi formula pengkiuban disertakan bagi kedua-dua fungsi ketumpatan teruji h(x,y) sebagai fungsi linear dan kuadratik. Keputusan yang diperolehi adalah selari dengan penemuan secara teori. Selanjutnya, kamiran hipersingular jenis Hadamard (KHH) dalam bentuk

$$H_i(h,x) = \frac{w_i(x)}{\pi} \int_{-1}^{1} \frac{h(t)}{w_i(t)(t-x)^2} dt, x \in (-1,1), i \in \{1,2,3,4\},$$

dengan
$$w_1(t) = \sqrt{1-t^2}, w_2(t) = \frac{1}{\sqrt{1-t^2}}, w_3(t) = \sqrt{\frac{1-t}{1+t}} \, dan \, w_4(t) = \sqrt{\frac{1+t}{1-t}}$$

adalah pemberat dan h(t) adalah fungsi mulus. Skim kuadratur automatik (SKA) bagi setiap kes $i \in \{1,2,3,4\}$ dibina dengan menghampirkan fungsi ketumpatan h(t) dengan siri polinomial Chebyshev terpangkas jenis pertama, kedua, ketiga dan keempat. Anggaran ralat bagi kes $i \in \{1,2,3,4\}$ diperolehi dengan menghampirkan fungsi ketumpatan dengan siri polinomial chebyshev terpangkas jenis pertama, kedua, ketiga dan keempat, dalam fungsi kelas $C^{N,\alpha}[-1,1]$. Ketepatan kaedah bagi kes $i \in \{1,2,3,4\}$ dipamerkan untuk fungsi polinomial berdarjah 3 dan keputusan bagi contoh teruji dibentangkan dan dibincangkan. Keputusan berangka bagi skim kuadratur yang diperolehi mendedahkan bahawa kaedah yang dicadangkan adalah sangat tepat untuk fungsi ketumpatan teruji h(t) sebagai fungsi polinomial dan fungsi nisbah. Pembandingan yang telah dibuat dengan kaedah lain yang diketahui menunjukkan bahawa skim kuadratur automatik (SKA) yang telah dibina dalam penyelidikan ini mempunyai keputusan yang lebih baik berbanding dengan yang lain. Keputusan yang diperolehi adalah setara dengan penemuan secara teori.

ACKNOWLEDGEMENTS

For most I am highly indebted to my creator Allah Subhanahu Wataalah for His guidance, strength and health He rendered to me. My father and late mother (May "jannatul firdaus" be her final destination) are my courage and happiness. Words are inadequate to express the love I have for you. My efforts in this project could not have been achieved and made possible without the help and support from many individuals and organizations. First of all I would like to extend my sincere appreciation to all of them.

My appreciation goes to the Department of Mathematics, Universiti Putra Malaysia (UPM) and Bayero University Kano (BUK) for the provision of necessary and enabling environment regarding the project work and also for their support in completing the project. There is no amount of words that can adequately express the debt I owe to my Supervisor and guide, Assoc. Prof. Dr. Zainidin Eshkuvatov, Mathematics Department, Faculty of Science, UPM (currently in Faculty of Science and Technology, Universiti Sains Islam, Malaysia (USIM)) for his unrelenting and continuous words of encouragement and educative discussion during the course of the present work. I am very much grateful for introducing me to the field of Integration problems, appreciating my ideas and giving me the freedom to take on the tasks independently, enriching me with amassed knowledge in the area and helping to explore the things myself. My special appreciation and thanks also goes to the members of my supervisory committee; Prof. Madya Dr. Nik Mohd Asri Nik Long (Currently the Head of Mathematics Department, UPM), Prof Fudziah Ismail (Former Head of Mathematics Department, UPM), Assistant Prof. Dr. Abdurrahim Okhunov (From Internatinal Islamic Universiti Malaysia (IIUM)) for providing me with need support and encouragement academically; giving me attention and time. I would also want to take the opportunity to express my deep appreciation to the entire staff members of the Department of Mathematics and INSPEM, UPM for helping me in my work, and always willing to engage in thoughtful discussion. From Hameed Husam Hameed, Fatima Samihah Zulkarnain a research group members, I appreciate all the attention and time.

Other Nigerians; Dr. Abbas J. Badakaya, Dr. M.Y Waziri, Dr. Sani Jibril, Kabiru Umar, Ibrahim Arzuka to mention but a few, I appreciate your support. Other non Nigrians such as Abubakar Somali and Ghadeer Omar Algabishi, I appreciate your encouragement.

Specially, I would like to thank Prof. Abubakar Adamu Rasheed (the former Vice Chancellor of Bayero University), the Vice Chancellor of Bayero University in person of Prof. Muhammad Yahuza Bello, Prof. Abba Gumel (University of Manitoba), Prof Aisha Halliru (former Head of Mathematics Department, Bayero University Kano (BUK)), Assoc. Prof Dr Bashir Ali (Head of Mathematics Department, Bayero University Kano (BUK)), Prof Bala Sidi Aliyu (former Dean Faculty of Science, Bayero University Kano (BUK)), to mention but a few .

Finally, yet very important, my heartfelt thanks to my beloved wife for always being with me and supporting me, the families of Ilyasu (Great grand father), Alhaji Baba, Alhaji Baba yaro OS, Alhaji Rabiu dan baffa, Alhaji Aliyu A. Sidi for their blessings, my friends/classmates for their help and success wishes.



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy.

The members of the Supervisory Committee were as follows:

Zainidin K. Eshkuvatov, PhD

Associate Professor Faculty of Science Universiti Putra Malaysia (Chairperson)

Nik Mohd Asri Nik Long, PhD

Associate Professor Faculty of Science Universiti Putra Malaysia (Member)

Fudziah Ismail, PhD

Professor Faculty of Science Universiti Putra Malaysia (Member)

Abdurahim, Okhunov, PhD

Assistant Professor Kulliyyah of Engineering International Islamic Univeristi Malaysia (Member)

BUJANG KIM HUAT, PhD Professor and Dean School of Graduate Studies Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by University Putra Malaysia, as according to the University Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature –

Date:

Name and Matric No.: Lawan Sirajo Bichi (GS35080)

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
	vvi
LIST OF APPPEVIATIONS	AVI VViii
LIST OF ADDREVIATIONS	XVIII
CHAPTER	
1 INTRODUCTION	1
1.1 Background	1
1.2 Problem statement	4
1.3 Objectives of the thesis	4
1.4 Scope of the study	5
1.5 Limitations of the study	6
1.6 Outline of the thesis	6
1.7 Basic concepts and definitions	7
1.7.1 Standard Quadrature Formula of Gauss-type	7
1.7.2 Change of interval	8
1.7.3 Gaussian rules	8
1.7.4 Extended Gaussian Rules	12
1.7.5 Gauss-Kronrod rules	21
1.7.6 Other Forms of Gauss Quadrature	22
1.7.7 Fundamental theorem	23
1.9. Summers	24
1.8 Summary	24
	26
2 LITERATURE REVIEW	26
2.1 Quadrature formulas in Numerical Integration	$\frac{26}{27}$
2.2 Cubature formulas approaches to integrations with singular	The $2/$
2.5 Automatic quadrature formulas approaches to hypersingula	11 IIIte- 20
	<u> </u>
2.4 Summary	32

3	CUBATURE FORMULA FOR DOUBLE INTEGH	RALS WITH ALGEBRAIC
	AND LOGARITHMIC SINGULARITY	33
	3.1 Introduction	33
	3.2 Concept of double integration by spline poly	nomial 33
	3.3 Construction of cubature formulas for doub	le integration with al-
	gebraic and logarithmic singularities	36
	3.4 Exactness of cubature formulas for Quadrati	c functions 44
	3.5 Numerical results and discussion	54
	3.6 Summary	61
4	BOUNDED AND UNBOUNDED HADAMARD	TYPE HYPERSINGU-
	LAR INTEGRALS	62
	4.1 Introduction	62
	4.2 Mathematical concepts of the first and sec	cond kinds Chebyshev
	polynomials and singular integrals	62
	4.3 Automatic quadrature schemes for bounded	and unbounded Hyper-
	singular integrals	66
	4.4 Error estimation for the bounded and unbou	nded hypersingular in-
	tegrals	67
	4.5 Exactness of bounded and unbounded HSI:	s for polynomial func-
	tions of degree 3	75
	4.6 Numerical results and discussion	77
	4.7 Summary	84
5	SEMI BOUNDED HADAMARD TYPE HYPERS	INGULAR INTEGRALS 85
0	5.1 Introduction	85
	5.2 Mathematical concepts of third and fourth k	ind Chebyshev polyno-
	mials and singular integrals	85
	5.3 Automatic guadrature schemes for semi bou	nded HSIs 87
	5.4 Error estimation and exactness for the semi h	ounded hypersingular
	integrals	89
	5.4.1 Error estimates for semi bounded HSI	s 89
	5.4.2 Exactness of semi bounded HSIs for n	olynomial functions of
	degree 3	96
	5.5 Numerical Results and Discussion	97
	5.6 Summary	106
	o.o Summary	100
6	SUMMARY, CONCLUSIONS AND RECOMME	NDATIONS 107
	6.1 Summary	107
	6.2 Conclusion	107
	6.3 Recommendations	108
	FEDENCES	110
R	EFEKENUES	110
Α	PPENDICES	116

xii

BIODATA OF STUDENT LIST OF PUBLICATIONS 176 177



LIST OF TABLES

Tab	le	Page
1.1	Details are given in Abramowitz and Stegun (1972)	23
3.1 3.2	Solutions of Example (3.5.1) obtained by <i>CF</i> (3.14) for $k = 1$ when $h(x, y)$ is a linear function and $\Omega = [-1, 1] \times [-1, 1]$. The solution of Example (3.5.1)by <i>CF</i> (3.14) for $k = 1$ when $h(x, y)$ is a linear function and $\Omega = [0, 1] \times [0, 1]$	54
3.3	Solution of Example 3.5.2 <i>CF</i> (3.14) when $h(x, y)$ is a quadratic function.	55
3.4	Solution of Example 3.5.3 by <i>CF</i> (3.14) for $k = 1$ when $h(x, y)$ is linear function.	56
3.5	Solutions of Example (3.5.4) by <i>CF</i> (3.14) for $k = 2$ when $h(x, y)$ is a linear function.	57
3.6	Solutions of Example (3.5.5) by <i>CF</i> (3.14) for $k = 2$, when $h(x, y)$ is linear function Besults obtained for the test of identity (3.29) for exactness of <i>CF</i>	58
3.8	(3.14) with $k \in \{1,2\}$ when $h(x,y)$ is any linear function. Values chosen $1-8$ to check exactness of cubature formula (3.14)	59
3.9	for Example (3.5.7) when $h(x, y)$ is any quadratic function. Results obtained to confirm the identities (3.29) for exactness of substance formula (3.14) when $h(x, y)$ is any quadratic function	60
3.10	Continuation of results in Table 3.9 obtained for the remaining identities (3.30) - (3.35) for exactness of cubature formula (3.14) when $h(x, y)$ is any quadratic function.	61
4.1	Results of HSI for $w_1(x) = \sqrt{1-x^2}$	78
4.2	Results of HSI for $w_2(x) = \frac{1}{\sqrt{1-x^2}}$	79
4.3	Results of HSI for $w_1(x) = \sqrt{1-x^2}$	80
4.4 4.5	Results of HSI for $w_2(x) = \frac{1}{\sqrt{1-x^2}}$ Numerical results compared with Obavis et al. (2013), Hui and Shia	81
4.6	(1999) and Kutt (1975) for the function (4.65) Results of AOS (4.13) for $i=2$ compared with Obavis et al. (2013)	82
4.7	and Hui and Shia (1999) for the function (4.66) Results of AOS (4.13) for $i=2$ compared with Hui and Shia (1999)	82
4.8	and Kutt (1975) for the function (4.66) Results of exactness of AQS (4.13) for degree 3 polynomials for the	83
4.9	case $i = 1$ Exactness of AQS (4.13) for degree 3 polynomials for the case $i = 2$	83 84
5.1	Results of HSIs for $w_3(x)$	98

- 5.2 Results of HSIs for $w_4(x)$ 99
- 5.3 Results of HSIs for $w_3(x)$ 100 103
- 5.4 Results of HSIs for $w_4(x)$
- 5.5 Results of exactness of AQS (5.15) with i = 3 for degree 3 polyno-105 mials

106

5.6 Results of exactness of AQS (5.15) with i = 4 for degree 3 polynomials



LIST OF FIGURES

Figu	ure	Page
5.1	Exact and AQS solutions (a) and Absolute error (b) for HSIs with $w_3(t)$ for $h(t)$ guadratic	98
5.2	Exact and AQS solutions (a) and Absolute error (b) for HSIs with	
	$w_4(t)$ for $h(t)$ quadratic	99
5.3	Exact and AQS solutions (a) and Absolute error (b) for HSIs with	
- 4	$w_3(t)$ for $h(t)$ rational and $N = 10$	101
5.4	Exact and AQS solutions (a) and Absolute error (b) for HSIs with $w_{i}(t)$ for $h(t)$ rational and $N = 20$	101
55	$w_3(t)$ for $h(t)$ rational and $N = 20$ Exact and AOS solutions (a) and Absolute error (b) for HSIs with	101
0.0	$w_3(t)$ for $h(t)$ rational and $N = 40$	102
5.6	Exact and AQS solutions (a) and Absolute error (b) for HSIs with	
	$w_4(t)$ for $h(t)$ rational and $N = 10$	104
5.7	Exact and AQS solutions (a) and Absolute error (b) for HSIs with	
	$w_4(t)$ for $h(t)$ rational and $N = 20$	104
5.8	Exact and AQS solutions (a) and Absolute error (b) for HSIs with (t) for $h(t)$ mating land h	105
	$w_4(t)$ for $n(t)$ rational and $N = 40$	105
A.1	Maple code for Computing Exact Values of coefficients $A'_{i,i}(\bar{x}^*_0)$ for	
	k = 1	117
A.2	Maple code for Computing Exact Values of coefficients $B'_{i,j}(\bar{x}_0^*)$ for	
	k = 1	119
A.3	Maple code for Computing Exact Values of coefficients $C'_{i,j}(\bar{x}^*_0)$ for	
	k=1	122
A.4	Maple code for Computing Exact values of coefficients $A_{i,j}(x_0)$ for $l = 1$	104
Δ 5	$\kappa = 1$ Maple code for Computing Exact Values of coefficients $B'(\bar{x}^*)$ for	124
A.J	Maple code for computing Exact values of coefficients $B_{i,j}(x_0)$ for $k-1$	126
A.6	Maple code for Computing Exact Values of coefficients $C'_{\cdot}(\bar{x}^*)$ for	120
11.0	k=2	128
B.1	Programme for Computing Exact Values of Double Integrals with	
	Algebraic Singularity for Linear Functions	129
B.2	Programme for Computing cubature formula for Double integrals	
	with algebraic singularity for Linear functions	135
C.1	Programme for Computing Exact Values of Double Integrals with	107
C^{2}	Algebraic Singularity for Quadratic Functions	13/
0.2	with Algebraic Singularity for Quadratic Functions	147
D.1	Programme for Computing values of cubature formula for Double	± ± /
. –	integrals with algebraic singularity for absolute value functions	156

C

ធ 1	Programme for Computing Exact values of Double integrals with	
Ľ.1	la negitierite ein melegite for lin een fer etiene	1 - 7
	logarithmic singularity for linear functions	157
E.2	Programme for Computing values of cubature formula for Double	
	integrals with logarithmic singularity for linear functions	163
F.1	Programme for Computing Exact values of Double integrals with	
	logarithmic singularity for absolute value functions	164
F.2	Programme for Computing values of cubature formula for Double	
	integrals with logarithmic singularity for absolute value functions	170
G.1	Programme for Computing HSIs of $w_1(x)$	171
H.1	Programme for Computing HSIs of $w_2(x)$	173
I.1	Programme for Computing HSIs of $w_3(x)$	174
J.1	Programme for Computing HSIs of $w_4(x)$	175



G

LIST OF ABBREVIATIONS

BCs	Boundary Conditions
HSI	Hypersingular integrals
CPV	Cauchy principal-value integrals
AQS	Automatic quadrature scheme
В	Closed region in d-dimensional Euclidean space
BS	B-Spline
D	Closed bounded connected set in R ²
K(x,y)	Regular or singular kernel
c_1, a_1 and b_1	Lower limits of integrals
c_2 , a_2 and b_2	Upper limits of integrals
$S_{\Lambda}(P)$	Modified spline function of type $(0,2)$
QT	Quadrature type
GLQ	Gauss-Legendre quadrature
GJQ	Gauss Jacobi quadrature
GChQ	Gauss Chebyshev quadrature
GLgQ	Gauss Laguerre quadrature
GHQ	Gauss Hermite quadrature
$C^{N,\alpha}[-1,1].$	Class of function which is N times continuous
	differentiable and belong to Hölder continuous
	functions
$C^{2,\gamma}(\Omega),$	Class of function which is 2 times continuous
	partially differentiable and satify Hölder condition
CC	Clenshaw-Curtis Chebyshev quadrature methods
LP	Legendre Polynomial
ChP	Chebyshev Polynomials
QF	Quadrature formula
AS	Algebraic singularity
LS	Logarithmic singularity
CF	Cubature formula

C

CHAPTER 1

INTRODUCTION

1.1 Background

Splines can be considered as a mathematical model that associate a continuous representation of a curve or surface with a discrete set of points in a given space (Weston, 2002). The general form of cubic spline is given by

$$S(x) = \begin{cases} s_0(x), & \text{if } x_0 \le x \le x_1, \\ s_1(x), & \text{if } x_1 \le x \le x_2, \\ \vdots & \vdots \\ s_{n-1}(x), & \text{if } x_{n-1} \le x \le x_n, \end{cases}$$
(1.1)

where each $s_i(x)$ is cubic polynomial for n data points say $(x_0, y_0), (x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$.

A cubic spline interpolant S, for given function h satisfies

- 1. S is a cubic polynomial, $s_i(x)$ on $[x_i, x_{i+1}]$, for $i = 0, 1, \dots, n-1$
- 2. $S(x_i) = h(x_i)$, for $i = 0, 1, 2, \dots, n-1, n$
- 3. $S_{i+1}(x_{i+1}) = S_i(x_{i+1})$, for $i = 0, 1, 2, \dots, n-2$
- 4. $S'_{i+1}(x_{i+1}) = S'_i(x_{i+1})$, for $i = 0, 1, 2, \dots, n-2$
- 5. $S_{i+1}''(x_{i+1}) = S_i''(x_{i+1})$, for $i = 0, 1, 2, \dots, n-2$
- 6. One of the following boundary conditions is satisfied
 - $S''(x_0) = S''(x_n) = 0$ (free or natural BCs)
 - $S'(x_0) = h'(x_0)$ and $S'(x_n) = h'(x_n)$ (clamped BCs)

The subject of singular integrals is highly significant tool in the pure Mathematics as well as the applied mathematics. Significant contribution had been made by many researchers in both singular and hypersingular integrals. The evaluation of HSI has been a case of interest to many researchers tackling many unsolved problems. To achieve this, there is the need to implement various techniques via approximate methods for evaluating HSI. In some cases, however, require the transformation of HSIs into singular or weakly singular integrals by using various techniques proposed in Clenshaw and Curtis (1960); Monegato (1960); Hasegawa and Torii (1991); Martin and Rizzo (1996); Eskhuvatov et al. (2011); Eskhuvatov and Nik Long (2011); Tadeu and Antonio (2012). Others, may require the numerical computation of finite part integrals by various quadrature or cubature formulas as in the research work done in Kutt (1975); Hui and Shia (1999); Colm and Rokhlin (2001); Yang (2012). Some ideas used the quadrature formula as in the work of Clenshaw and Curtis (1960); Hasegawa and Torii (1991); Eskhuvatov et al. (2011); Obayis et al. (2013), which take advantage of the collocation points to solve the Cauchy principalvalue and hypersingular integrals problems.

Consider the Hadamard finite-part integral (or hypersingular integral) of the form

$$H_p(a,b;s,h) = \int_a^b \frac{h(t)}{(t-s)^{p+1}}, s \in (a,b), p = \{1,2\}$$
(1.2)

with p + 1-order singularity and *s* is the singular point.

Definition 1.1.1 (Boykov et al. (2009) and Zhang et al. (2009)) HSI (1.2) must be understood in the Hadamard finite-part sense and is defined as

$$\int_{a}^{b} \frac{h(t)}{(t-s)^{p+1}} dt = \lim_{\epsilon \to 0} \left\{ \int_{a}^{s-\epsilon} \frac{h(t)}{(t-s)^{p+1}} dt + \int_{s+\epsilon}^{b} \frac{h(t)}{(t-s)^{p+1}} dt - \frac{2h^{(p-1)}(s)}{\epsilon} \right\}, (1.3)$$

h(t) is said to be finite-part integrable (or integrable in the Hadamard sense) with the kernel $(t-s)^{-p-1}$ if the limit on the right-hand side of (1.3) exist.

An important and widely used definition of HSI in the society of engineers is that if h(t), $a \le t \le b$, satisfies a Hölder continuous first-derivative condition

$$|h(t) - h(x) - (t - x)h'(x)| \le A|t - x|^{\alpha + 1},$$
(1.4)

where *A* is a positive constant and $\alpha \in (0, 1]$.

The HSI is defined as

$$H(h,x) = \int_{-1}^{1} \frac{h(t)}{(t-x)^2} dt = \lim_{\epsilon \to 0} \left[\left(\int_{-1}^{x-\epsilon} + \int_{x+\epsilon}^{1} \right) \frac{h(t)}{(t-x)^2} dt - \frac{2h(x)}{\epsilon} \right], \quad (1.5)$$

where this limits of integration exists and bounded. The high order accuracy was obtained by Hui and Shia (1999) for hypersingular integrals with secondorder singularities of the form

$$H(h,x) = \int_{a}^{b} \frac{w(t)h(t)}{(t-x)^2} dt,$$
 (1.6)

in which the Gaussian quadrature rule have been used based on the Legendre and Chebyshev series expansion.

Differentiation of Cauchy principle-value integral (CPVI)

$$C(h,x) = \int_{-1}^{1} \frac{w(t)h(t)}{t-x} dt$$
 (1.7)

with respect to the singular point x, gives

$$H(h,x) = \int_{-1}^{1} \frac{w(t)h(t)}{(t-x)^2} dt = \frac{d}{dx} \int_{-1}^{1} \frac{w(t)h(t)}{t-x} dt, \ x \in (-1,1)$$
(1.8)

which is also another definition of HSI according to Hui and Mukherjee (1997). Thus HSI represent a natural extension of singular integrals in the Cauchy principal-value (CPV). i.e

$$H(h,x) = \frac{d}{dx}C(h,x).$$
(1.9)

In order to approximate integration in two or more dimensions, Let *B* designate a closed region in *d*-dimensional Euclidean space and let dV designate the *d*dimensional volume element in Davis and Rabinowitz (1984). Find fixed points P_1, P_2, \dots, P_n (preferably in B) and fixed weights w_1, w_2, \dots, w_n such that

$$\int_{B} w(P)h(P)dV \approx \sum_{k=1}^{n} w_k h(P_k).$$
(1.10)

This is a useful approximation to the integral on the left for a reasonably large class of functions of d variables defined on B.

The problem of approximating double integral of the form

$$H(h) = \int_{D} \int K(x, y)h(x, y)dA, \qquad (1.11)$$

where K(x, y) is a regular or singular kernel and D is closed bounded connected set in R^2 was studied by Eshkuvatov et al. (2013). In particular, the problem of evaluations of double integrals with the function of algebraic singularity

$$H(h) = \int_{D} h(\bar{x}) |\bar{x} - \bar{x_0}|^{\gamma - 2} dA, \ 0 < \gamma \le 1,$$
(1.12)

where D is a closed bounded simple connected region in \mathscr{R}^2 and $\bar{x} = (x, y) \in D$ are variables and $\bar{x_0} = (x_0, y_0)$ is a fixed point in D was studied. Double integral in (1.12) has algebraic singularity at a point of the plane D where $|\bar{x} - \bar{x_0}| \to 0$. Therefore the numerical calculation of (1.12) requires a care about subtle point. Various approximations of an integral

$$H(h) = \int_{\Omega} W(\mathbf{x}) h(\mathbf{x}) d\mathbf{x},$$

where $\Omega \in \mathfrak{R}^n$, and $W(\mathbf{x}) \ge 0$, $\forall \mathbf{x} \in \mathfrak{R}^n$, $n \ge 2$, by

$$H(h) = Q(h) + R(h),$$

where Q(h) is the cubature formula (CF) and R(h) is the remainder are presented by Cools and Rabinowitz (1993) and Cools (2003). To solve multivariate singular integrals problems there is need to find an efficient approximate formulas as in Stroud (1973); Davis and P. (1984); Evans (1993); Krommer and Ueberhuber (1998).

1.2 Problem statement

Many problems of multivariate integration problems arise in applications but the analytic solutions does not exist. Hence it become necessary to use the numerical techniques to arrive at the solutions. One such approaches to numerical solutions of double integrations problems is by using the spline interpolation. Although the modified spline function $S_{\Lambda}(P)$ of type (0,2) was established theoretically to have interpolate the density functions in the double integration problems, the justifications in the accuracy has not been established numerically.

In the evaluation of HSIs problems, there the need to develop accurate AQSs via suitable approximations approach. Some researches have been conducted previously, for the constructions of AQSs particularly with the weight functions $w_0 = 1$, $w_1 = \sqrt{1-x^2}$ and $w_2 = \frac{1}{\sqrt{1-x^2}}$ for some HSIs problems. Additionally, error estimation was obtained for w_0 case only. Hence the need to investigate constructions of AQSs for similar and other weight functions via different approximations approaches and also obtain the error estimations for those weight functions that have not been considered. More investigations are needed for the exactness of the AQSs.

1.3 Objectives of the thesis

The problems of double integrals of the form

$$H_{k}(x,y) = \begin{cases} \iint_{\Omega} \frac{h(x,y)}{|\bar{x}-\bar{x}_{0}|^{2-\gamma}} dA, & 0 \le \gamma \le 1, \quad k = 1, \\ \iint_{\Omega} h(x,y) \log |\bar{x}-\bar{x}_{0}^{*}| dA, \quad k = 2, \end{cases}$$
(1.13)

where $\Omega = [a_1, a_2] \times [b_1, b_2]$, $\bar{x} = (x, y) \in \Omega$ and fixed point $\bar{x}_0^* = (x_0^*, y_0^*) \in \Omega$, the density function $h(\bar{x})$ is assumed continuous and smooth enough on the rectangle Ω and belong to the class of function $C^{2,\gamma}(\Omega)$, i.e. $h(\bar{x})$, $h_x(\bar{x})$, $h_y(\bar{x})$, $h_{xx}(\bar{x})$, $h_{yy}(\bar{x})$ and $h_{xy}(\bar{x}) = h_{yx}(\bar{x})$ in Ω are all continuous and all second partial derivatives satify Hölder condition are considered. The aims of the thesis here are:

- To construct cubature formulas and develop Maple codes for the constructed cubature formulas in the two cases $i \in \{1,2\}$ for double integrals with algebraic and logarithmic singularities respectively, on a rectangle Ω using the modified spline function $S_{\Lambda}(P)$ of type (0,2).
- To obtain the exactness and provide numerical results of the cubature formulas for the two cases $i \in \{1,2\}$ together with tested examples each for linear and quadratic functions.

Furthermore, the problem of Hadamard type hypersingular integral (HSI) of the form

$$H_{i}(h,x) = \frac{w_{i}(x)}{\pi} \int_{-1}^{1} \frac{h(t)}{w_{i}(t)(t-x)^{2}} dt, x \in (-1,1), i \in \{1,2,3,4\},$$
(1.14)

where $w_1(t) = \sqrt{1-t^2}$, $w_2(t) = \frac{1}{\sqrt{1-t^2}}$, $w_3(t) = \sqrt{\frac{1-t}{1+t}}$ and $w_4(t) = \sqrt{\frac{1+t}{1-t}}$ are the weights and h(t) is a smooth function are considered. The aims of the thesis here are:

- To construct an automatic quadrature schemes (AQSs) together with their Maple codes and obtain error estimations in each case for $i \in \{1,2,3,4\}$ via approximating the density function h(t) by the first, second, third and fourth kind truncated series of Chebyshev polynomials, respectively.
- To obtain exactness for the degree 3 polynomial functions and obtain numerical results of the AQSs each for $i \in \{1,2,3,4\}$ for the tested density function h(t) as polynomial and rational functions. However, compare the efficiency and accuracy of the AQSs constructed with other known methods in the available literature.

1.4 Scope of the study

The research work was based on the following scopes;

- The solution of double integration problems in the rectangular domain using the modified spline function $S_{\Lambda}(P)$ of type (0,2).
- Use of Gauss chebyshev quadrature formula approaches to solve the hypersingular integrations problems by approximating the density functions with first, second, third and fourth kind truncated series of chebyshev polynomials.

• Use of Maple 14 to develop the codes in both cubature formulas and automatic quadrature formulas to solve the double and hypersingular integrations problems.

1.5 Limitations of the study

Our investigation in this research work is limited the use of test function as linear and quadratic function on the constructed cubature formulas (3.14) with $k \in \{1,2\}$ for double integration with algebraic and logarithmic singularities, $H_k(x, y)$ for k = 2 in (3.1). In an attempt to test the methods with the rational function due to the large computations involved in the method and the number of iterations needed to arrive at the solutions, it was found impossible to manage this with smaller processing unit computers. A powerful and higher processing computing machine (HPC) which understand the Maple computing is needed to handle the job.

1.6 Outline of the thesis

In Chapter 1, motivation and reason of the research is given. Problem statement, objectives, scope of the study, limitations, outline of the thesis and basic concepts and definitions related to research work are discussed.

Chapter 2 is devoted to earlier research on cubature and quadrature in numerical integration as well as their approach to the double integration and hypersingular integration problems, respectively.

Chapter 3 presents the concept of double integration by spline polynomial. Demonstrated the construction of cubature formulas for double integration problems with algebraic and logarithmic singularities. Exactness of the obtained cubature formulas each for linear and quadratic functions and finally numerical results and discussion are provided and discussed.

Chapter 4 provides the mathematical concepts of the first and second kind Chebyshev polynomials and hypersingular integrals. The detail construction of automatic quadrature schemes for bounded and unbounded hypersingular integrals and rate of convergence of the suggested methods are discussed. Additionally, the exactness of the AQSs constructed each for polynomials of degree 3 are Shown and finally numerical results are given and discussed.

Chapter 5 provides the mathematical concepts of the third and fourth kind Chebyshev polynomials and hypersingular integrals. The construction of automatic quadrature schemes for semi bounded solutions of hypersingular integrals and the rate of convergence of the suggested methods are presented. Additionally, the exactness of the AQSs each for polynomials of degree 3 are shown and finally numerical results are given and discussed. Finally, Chapter 6 gives conclusion of the research work. Some future works are also suggested on Cubature and Quadrature formulas.

1.7 Basic concepts and definitions

To facilitate understanding in this thesis write up, the following concepts and definitions are useful. Beginning with the continuous orthogonality which can be seen from the following

Definition 1.7.1 (Mason and Handscomb (2003)) Two functions $h_1(t)$ and $h_2(t)$ in $L_2[c_1,c_2]$ are said to be orthogonal on the interval $[c_1,c_2]$ with respect to weight function w(t) if

$$\langle h_1, h_2 \rangle = \int_{c_1}^{c_2} w(t) h_1(t) h_2(t) dt = 0.$$
 (1.15)

Infact, it is possible to convert continuous orthogonality relationship (1.15) into a discrete orthogonality relationship by replacing the integral with a summation and the result here is in general, only approximately true. Particularly, in case of discrete orthogonality for trigonometric functions or Chebyshev polynomials, there are many cases which the formulas have been shown to hold exactly by Mason and Handscomb (2003).

1.7.1 Standard Quadrature Formula of Gauss-type

A quadrature rule is an approximation of the definite integral of a function, usually stated as a weighted sum of function values at specified points within the domain of integration (Teubner and Leipzig, 1993). An n-point Gaussian quadrature rule, named after Carl Friedrich Gauss, is a quadrature rule constructed to yield an exact result for polynomials of degree 2n-1 or less by a suitable choice of the points x_i and weights w_i for $i = 1, \dots, n$. The domain of integration for such a rule is conventionally taken as [-1, 1], so the rule is stated as

$$\int_{-1}^{1} h(x)dx \approx \sum_{i=1}^{n} w_i h(x_i).$$
(1.16)

Gaussian quadrature as above will only produce accurate results if the function h(x) is well approximated by a polynomial function within the range [-1, 1]. The method is not, for example, suitable for functions with singularities. However, if the integrated function can be written as $h(x) = \omega(x)g(x)$, where g(x) is approximately polynomial and $\omega(x)$ is known weight function, then alternative weights w'_i and points x'_i that depend on the weighting function $\omega(x)$ may give

better results, where

$$\int_{-1}^{1} h(x)dx = \int_{-1}^{1} \omega(x)g(x)dx \approx \sum_{i=1}^{n} w'_{i}h(x'_{i}).$$
(1.17)

Common weighting functions include $\omega(x) = 1/\sqrt{1-x^2}$ (Gauss Chebyshev) and $\omega(x) = e^{-x^2}$ (Gauss Hermite).

1.7.2 Change of interval

If integral over $[c_1, c_2]$ need to be changed into an integral over [-1, 1] then this change of interval can be done in the following way:

$$\int_{c_1}^{c_2} w(x)h(x)dx = \frac{c_2 - c_1}{2} \int_{-1}^{1} w_1(z)h(\frac{c_2 - c_1}{2}z + \frac{c_2 + c_1}{2})dz \quad (1.18)$$

$$\approx \sum_{i=1}^{n} w_i h(\frac{c_2 - c_1}{2} z_i + \frac{c_2 + c_1}{2}).$$
(1.19)

1.7.3 Gaussian rules

Let $v_n(x)$ denote a sequence of polynomials, where $v_n(x) \in \mathscr{P}$ and the polynomials $v_n(x)$ are orthogonal with respect to a weight function w(x) over the interval $[c_1, c_2]$. We can write $v_n(x) = b_n p_n(x) + b_{n-1} p_{n-1}(x) + \dots + b_0$, where $b_n \neq 0$ is the coefficient of x^n in $v_n(x)$, and $p_n(x)$ is an orthogonal polynomial. Then

$$\int_{c_1}^{c_2} w(x)v_i(x)v_j(x)dx = 0, \text{ for } i \neq j.$$
 (1.20)

with the Chrstoffel-Darbourx identity

$$\sum_{k=0}^{n} \frac{\upsilon_k(x)\upsilon_k(y)}{\gamma_k} = \frac{\upsilon_{n+1}(x)\upsilon_n(y) - \upsilon_n(x)\upsilon_{n+1}(y)}{\alpha_n\gamma_n(x-y)},$$
(1.21)

where

$$\alpha_n = \frac{b_{k+1}}{b_k}, \ \gamma_n = \int_{c_1}^{c_2} w(x) v_k^2(x) dx.$$
(1.22)

Now, let $y = x_i$, where x_i are the zeros of $v_n(x)$, then

$$\sum_{k=0}^{n-1} \frac{\upsilon_k(x)\upsilon_k(x_i)}{\gamma_k} = \frac{\upsilon_{n+1}(x)\upsilon_n(x_i) - \upsilon_n(x)\upsilon_{n+1}(x_i)}{\alpha_n\gamma_n(x - x_i)}.$$
 (1.23)

Multiply both sides of (1.23) by $w(x)v_0(x)$ and integrate over $[c_1, c_2]$, then applying (1.20) we have

$$\upsilon_0(x_i) = -\frac{\upsilon_{n+1}(x_i)}{\alpha_n \gamma_n} \int_{c_1}^{c_2} w(x) \frac{\upsilon_0(x)\upsilon_n(x)}{x - x_i} dx.$$
 (1.24)

Recalling the definition of the Lagrangian interpolation polynomial

$$l_i(x) = \frac{\pi_n(x)}{(x - x_i)\pi'_n(x_i)} = \frac{\upsilon_n(x)}{(x - x_i)\upsilon'_n(x_i)},$$
(1.25)

and since for some constant $v_0(x) = c$, we find from (1.24) and (1.25) that

C-

$$1 = -\frac{\upsilon_{n+1}(x_i)}{\alpha_n \gamma_n} \int_{c_1}^{c_2} w(x) \frac{\upsilon_n(x)}{x - x_i} dx$$
$$= -\frac{\upsilon_{n+1}(x_i)\upsilon_n'(x_i)}{\alpha_n \gamma_n} \int_{c_1}^{c_2} w(x)l_i(x) dx$$
$$= -\frac{\upsilon_{n+1}(x_i)\upsilon_n'(x_i)}{\alpha_n \gamma_n} w_i$$
(1.26)

where

$$w_i = -\frac{b_{n+1}\gamma_n}{b_n v_{n+1}(x_i)v'(x_i)} \text{ for } i = 1, \cdots, n.$$
(1.27)

The error is given by

$$E_n = \frac{\gamma_n}{b_n^2(2n)!} h^{(2n)}(\xi), \ c_1 < \xi < c_2.$$
(1.28)

Note that for the Legendre polynomials $P_n(x)$ these formulae give

$$\gamma_{n} = \int_{-1}^{1} P_{n}^{2}(x) dx = \frac{2}{2n+1},$$

$$b_{n} = \frac{(2n)!}{2^{n}(n!)^{2}},$$

$$w_{i} = -\frac{2}{(n+1)P_{n+1}(x_{i})P_{n}'(x_{i})},$$

$$E_{n} = \frac{2^{2n+1}(n!)^{4}}{(2n+1)[(2n)!]^{3}}h^{(2n)}(\xi).$$
(1.29)
(1.30)

Consider quadrature rules of the form

$$I_{c_1}^{c_2}(h) = \int_{c_1}^{c_2} h(x) dx = \sum_{i=0}^m w_i h(x_i), \qquad (1.31)$$

where x_i are not equally spaced. Replacing h by an interpolating polynomial $p_n(x)$ constructed at the points x_i , $i = 0, 1, \dots, n$, then the polynomial will have Lagrangian form

$$p_n(x) = l_0(x)h(x_0) + \dots + l_n(x)h(x_n),$$
(1.32)

where $l_i(x)$, $i = 0, 1, \dots, n$, are defined by

$$l_i(x) = \prod_{j=0, j \neq i}^n \frac{x - x_j}{x_i - x_j}.$$
 (1.33)

By integrating (1.32) over $c_1 \le x \le c_2$, we get the right hand side of (1.31), in which case the weights w_i are given by

$$w_i = \int_{c_1}^{c_2} l_i(x) dx, \ 0 \le i \le n,$$
(1.34)

For an equally-spaced nodes x_i this procedure leads to a different method of deriving the Newtons-cotes quadrature formulas. The error term

$$E(x) = (x - x_0)(x - x_1) \cdots (x - x_n) \frac{h^{n+1}(\xi)}{(n+1)!}, \ c_1 < \xi < c_2$$
(1.35)

 $h^{n+1}(\xi)$ is continuous and ξ depends on x on integration over the interval $[c_1, c_2]$ gives

$$I_{c_1}^{c_2}h - \sum_{i=0}^{n} w_i h(x_i) = \frac{1}{(n+1)!} \int_{c_1}^{c_2} \pi_{n+1}(x) h^{n+1}(\xi), \qquad (1.36)$$

where $\pi_{n+1}(x) = (x-x_0)(x-x_1)\cdots(x-x_n)$. Thus, the quadrature rule (1.31) is exact if $h \in \mathcal{P}_n$ because since $h^{n+1}(x) \equiv 0$ the right side of (1.36) vanishes. The rule (1.31) is exact for any linear combination of functions h and g if it is exact for the functions h and g separately. Thus, if α and β are any arbitrary real numbers, we have

$$\int_{c_1}^{c_2} (\alpha h(x) + \beta g(x)) dx = I_{c_1}^{c_2} h + I_{c_1}^{c_2} g = \sum_{i=0}^n w_i [\alpha h(x_i) + \beta g(x_i)].$$
(1.37)

This shows that the rule (1.31) is exact for $h \in \mathcal{P}_n$ if it is exact for the monomials $1, x, x^2, \dots, x^n$. In fact, the rule (1.31) is exact for x^j if

$$\int_{c_1}^{c_2} x^j dx = \sum_{i=0}^n w_i x_i^j.$$
 (1.38)

Since the left side of (1.38) is known, we take $j = 0, 1, \dots, 2n+1$ and obtain a system (2n+2) equations to solve 2n+2 unknowns w_i and for $i = 0, 1, \dots, n$. If this system has a solution, the resulting quadrature rule will be exact for $h \in \mathscr{P}_n$ for j, k < n, so that

$$\sum_{i=0}^{n} w_i p_j(x_i) p_k(x_i) = \int_{c_1}^{c_2} w(x) p_j(x) p_k(x) dx = \delta_{jk}.$$
 (1.39)

Thus

$$\langle h, g \rangle = \sum_{i=0}^{n} w_i h(x_i) g(x_i).$$
 (1.40)

We will develop the error (1.36) in terms of the divided differences. Since

$$h(x) - p_n(x) = \pi_{n+1}(x)h[x, x_0, \cdots, x_n], \qquad (1.41)$$

we have

$$I_{c_1}^{c_2}h - \sum_{i=0}^{n} w_i h(x_i) = \int_{c_1}^{c_2} \pi_{n+1}(x) h[x, x_0, \cdots, x_n] dx.$$
(1.42)

If $h[x, x_0, \dots, x_n] \in \mathscr{P}_m$, m > 0, then $h[x, x_0, \dots, x_{k+1}] \in \mathscr{P}_{m-1}$. This result follows from

$$h[x, x_0, \cdots, x_n] = \frac{h[x_0, \cdots, x_n] - h[x, x_0, \cdots, x_{n-1}]}{x_n - x_0},$$
(1.43)

called a divided difference, which gives

$$h[x, x_0, \cdots, x_{k+1}] = \frac{h[x_0, \cdots, x_{k+1}] - h[x, x_0, \cdots, x_k]}{x_{k+1} - x},$$
(1.44)

and since the numerator on the right side is a polynomial of degree m and has $(x-x_{k+1})$ as one of the factors, then $h \in \mathscr{P}_{m-1}$. Hence, if $h \in \mathscr{P}_{2n+1}$, we can show by induction that $h[x, x_0, \cdots, x_n] \in \mathscr{P}_n$. Now, let $\{p_0, p_1, \cdots, p_{n+1}\}$ be a set of orthogonal polynomials $[c_1, c_2]$, and let $p_r \in \mathscr{P}_r$, $r = 0, 1, \cdots, n+1$. Then for some set of real numbers $\alpha_0, \alpha_1, \cdots, \alpha_n$, we have $h[x, x_0, \cdots, x_n] = \alpha_0 p_0 + \alpha_1 p_1 + \cdots + \alpha_n p_n$. Then, because of the orthogonality relations

$$\int_{c_1}^{c_2} p_r(x) p_s(x) dx \begin{cases} 0, & \text{for } r \neq s, \\ ||p_s||^2, & \text{for } r = s, \end{cases}$$
(1.45)

the right side of (1.42) is zero if $\pi_{n+1}(x) = \alpha p_{n+1}(x)$ for some real $\alpha \neq 0$.

The Gaussian quadrature rules cannot in general be used for functions that are defined only at discrete points since such rules require that the functions be evaluated at specific points ξ_t which are the zeros of the related polynomials. However, in some experimental cases where there is complete freedom of choosing the data of functional values at the specific nodes ξ , the corresponding Gaussian rules can be used to evaluate $I_{c_1}^{c_2}(h)$. The error bounds are virtually impossible to obtain in such cases. But if the data is mostly error free, then a very high order interpolating polynomial may be effectively used to obtain a significantly accurate result. There is no practical advantage in using a Gaussian quadrature if the data is noisy and some smoothing is needed prior to the application of a specific Gaussian quadrature rule.

It must also be noted that the Gaussian rules of all orders are also Riemann sums. These rules integrate exactly polynomials of any degree by a formula of the form (1.31).

1.7.4 Extended Gaussian Rules

The extended (repeated or compund) Gaussian rules are similar to the repeated trapezoidal, midpoint and Simpson's rules.

Theorem 1.7.1 (Abramowitz and Stegun (1972)) If $p_{2n+1}(y) \in \mathscr{P}_{2n+1}$, then

the formula

$$\int_{c_1}^{c_2} w(x) p_{2n+1}(y) dy = \sum_{j=0}^m \gamma_j p_{2n+1}(y_j)$$

is exact if the points y_j , $j = 0, 1, \cdots, m$, are the zeros of the orthogonal polynomial v_{m+1} , and γ_j are defined by

$$\gamma_j = \int_{c_1}^{c_2} w(x) l_j(x) dx$$

Theorem 1.7.2 (Johnson and Riess (1977)) Gaussian quadrature formula has precision 2m + 1 only if the points x_j , $j = 0, 1, \dots, m$, are the zeros of v_{m+1} .

Theorem 1.7.3 (Abramowitz and Stegun (1972)) Let $h(y) \in C^{2m+2}[c_1, c_2]$, then the error in the Gaussian quadrature is

$$I_{c_1}^{c_2} - I_m = \frac{h^{2m+2}(\xi)}{(2m+2)!} \int_{c_1}^{c_2} w(y) T_{m+1}^2(y) dy$$

where T_m is the Chebyshev polynomial of the first kind.

1.7.4.1 Gauss-Jacobi Rule

The Jacobi polynomials $P^{(\gamma,\sigma)}$ are the orthogonal with weight function $w(y) = (1-y)^{\gamma}(1+y)^{\sigma}$, $\gamma > -1$, $\sigma > -1$. The Gauss-Jacobi rule (also known as the Mehler quadrature formula) is defined by

$$\int_{-1}^{1} (1-y)^{\gamma} (1+y)^{\sigma} h(y) dy = \sum_{j=1}^{m} w_j h(y_j) + E, \qquad (1.46)$$

where

$$w_{j} = -\frac{2m+\gamma+\sigma+2}{2m+\gamma+\sigma+1} \frac{\Gamma(m+\sigma+1)\Gamma(m+\gamma+\sigma+1)}{(m+1)!\Gamma(m+\gamma+\sigma+1)} \times \frac{2^{\gamma+\sigma}}{P_{m}^{\prime(\gamma+\sigma)}(y)P_{m+1}^{(\gamma+\sigma)}(y)} (1.47)$$

and the series form of Jacobi polynomials $P^{(\gamma,\sigma)}$ (Prem and Michael, 2005) is

$$P^{(\gamma,\sigma)}(y) = \frac{1}{2^m} \sum_{k=0}^{[m/2]} {m+\gamma \choose k} {m+\sigma \choose m-k} (y-1)^{m-k} (y+1)^k, \qquad (1.48)$$

and the error term E_m in the m-point rule is

$$E_{m} = \frac{\Gamma(m+\gamma+1)\Gamma(m+\sigma+1)\Gamma(m+\gamma+\sigma+1)}{(2m+\gamma+\sigma+1)[\Gamma(2m+\gamma+\sigma+1)]^{2}} \times \frac{m!2^{2m+\gamma+\sigma+1}}{(2m)!} h^{2m}(\xi), \ \xi \in (-1,1).$$
(1.49)

The Gauss-Legendre rule is a special case of formula (1.47) with $\gamma = \sigma = 0$. The Gauss-Chebyshev rule is another special case with $\gamma = \sigma = -\frac{1}{2}$ For integrands with the Jacobi weight function $w(y) = (1-y)^{\gamma}(1+y)^{\sigma}$ in (Piessens and Branders, 1973) use the formulas

$$\int_{-1}^{1} (1-y)^{\gamma} (1+y)^{\sigma} g(y) \approx \sum_{k=0}^{M} b_k G_k(\gamma,\sigma) + E'_M, \quad (1.50)$$

$$\int_{-1}^{1} (1-y)^{\gamma} (1+y)^{\sigma} \ln \frac{1+y}{2} g(y) dy \approx \sum_{k=0}^{M} b_k I_k(\gamma, \sigma) + E_M^2, \qquad (1.51)$$

where g(y) is assume to have a rapidly convergent Chebyshev series expansion

$$g(y) = \sum_{k=0}^{\infty} a_k T_k(y),$$
 (1.52)

and

$$b_k = \frac{2}{M} \sum_{k=0}^{\infty} {}'' g(y_i) T_k(y_i), \ y_i = \cos \frac{i\pi}{M},$$
(1.53)

$$G_m(\gamma,\sigma) = 2^{\gamma+\sigma+1} \frac{\Gamma(\gamma+1)\Gamma(\sigma+1)}{\Gamma(\gamma+\sigma+2)} 3F \begin{bmatrix} m,-m,\gamma+1\\ \frac{1}{2},\gamma+\sigma+2 \end{bmatrix}; 1 \end{bmatrix},$$
(1.54)

$$E_k^1 \approx a_{k+1}(G_{k+1}(\gamma,\sigma) - I_{k-1}(\gamma,\sigma)),$$
 (1.55)

$$E_k^2 \approx a_{k+1}(I_{k+1}(\gamma,\sigma) - I_{k-1}(\gamma,\sigma)),$$
 (1.56)

and $I_m(\gamma, \sigma)$ are obtained from the recurrence relation

$$(\gamma + \sigma + m + 2)I_{m+1}(\gamma, \sigma) + 2(\gamma - \sigma)I_m(\gamma, \sigma) + (\gamma + \sigma - m + 2)I_{m-1}(\gamma, \sigma)$$

= 2G_m(\gamma, \sigma) - G_{m-1}(\gamma, \sigma) - G_{m+1}(\gamma, \sigma). (1.57)

1.7.4.2 Gauss-Legendre Rule

This rule is a special case of the Gauss-Jacobi formula (1.47) with $\gamma = \sigma = 0$. The weight function w(y) = 1 and the orthogonal polynomials $v_j(y)$ are the Legendre polynomials $P_j(y)$, $j = 0, \cdots$, with $p_j(1) = 1$. This rule on the interval [-1,1] is given by (1.31), where the nodes y_j are the j-th zero of P_m ; the weight are given by

$$w_j = \frac{2}{(1 - y_j^2)[P'_m(y_j)]^2};$$
(1.58)

and the remainder is

$$R_m = \frac{2^{2m+1}(m!)^4}{(2m+1)[(2m)!]^3} h^{2m}(\xi), -1 < \xi < 1.$$
(1.59)

The Gauss-Legendre rule fore arbitrary interval $[c_1, c_2]$ is

$$\int_{c_1}^{c_2} h(y) dy = \frac{c_2 - c_1}{2} \sum_{j=1}^m w_j h(y_j) + R_m$$
(1.60)

where the nodes are

$$y_i = \left(\frac{c_2 - c_1}{2}x_j + \frac{c_2 + c_1}{2}\right); \tag{1.61}$$

the related nodes x_i and the weights w_i are defined above; and

$$R_m = \frac{(c_2 - c_1)^{2m+1} 2^{2m+1} (m!)^4}{(2m+1)[(2m)!]^3} h^{2m}(\xi), \ c_1 < \xi < c_2.$$
(1.62)

The rule (1.31) are exact for $h \in \mathscr{P}_{2m+1}$.

Using the first legendre polynomial

$$P_0(y) = 1, P_1(y) = y, P_2(y) = (3y^2 - 1)/2, P_3(y) = (2y^3 - 3y)/2,$$

we get:

- 1. For m = 2 in (1.31) we have the Gauss-Legendre rule with one node at y = 0 which is the zero of $P_1(y)$: $I_{-1}^1 h = 2h(0)$, which is exact for $h \in \mathscr{P}_1$. This is comparable to the midpoint rule.
- 2. For m = 1 in (1.31) we have the Gauss-Legendre rule with two nodes at $y = \pm 1/\sqrt{3}$ which are the zeros of $P_2(y)$: $I_{-1}^1 h = h(-1/\sqrt{3}) + h(1/\sqrt{3})$, which is exact for $h \in \mathcal{P}_3$. This is comparable to the Simpson's rule.
- 3. For m = 2 in (1.31) we have the Gauss-Legendre rule with three nodes at $y = 0, \pm \sqrt{3/5}$ which is the zero of $P_3(y)$: $I_{-1}^1 h = 2[5h(-\sqrt{3/5}) + 8h(0) + 5h(0) + 5h(\sqrt{3/5})]/9$, which is exact for $h \in \mathcal{P}_5$.

For the Gauss-Legendre rule, an interval $[c_1, c_2]$ can be transformed into [-1, 1] by using the transformation

$$c_1 = \frac{c_2 + c_1}{2} + \frac{c_2 - c_1}{2}\xi, \ \xi \in [-1, 1].$$
(1.63)

Then the Gauss-Legendre quadrature rule becomes

$$Hh = \frac{c_2 - c_1}{2} \sum_{k=1}^{m} w_k h(\xi_k).$$
(1.64)

where the nodes ξ_k are the *m* zeros of the n-th degree Legendre polynomials.

1.7.4.3 Gauss-Laguerre Rule.

A Gaussian quadrature over the interval $[0, \infty)$ with weight function $w(x) = e^{-x}$ (Davis and Rabinowitz, 1984) is Gauss-Laguerre. The abscissae for quadrature order *n* are given by the roots of the Laguerre polynomials $L_n(x)$. The weights are

$$w_{i} = -\frac{A_{n+1}\gamma_{n}}{A_{n}L_{n}'(x_{i})L_{n+1}(x_{i})} = \frac{A_{n}}{A_{n-1}}\frac{\gamma_{n-1}}{L_{n-1}(x_{i})L_{n}'(x_{i})},$$
(1.65)

where A_n is the coefficient of x^n in $L_n(x)$, given by $A_n = \frac{(-1)^n}{n!}$. Thus, $\frac{A_{n+1}}{A_n} = -\frac{1}{n+1}$, and $\frac{A_n}{A_{n-1}} = -\frac{1}{n!}$. Also, $\gamma_n = \int_0^\infty w(x) [L_n(x)]^2 dx = 1$, which gives

$$w_{i} = \frac{1}{(n+1)L'_{n}(x_{i})L_{n+1}(x_{i})} = \frac{1}{nL_{n-1}(x_{i})L'_{n}(x_{i})},$$

$$= \frac{1}{x_{i}[L'_{n}(x_{i})]^{2}} = \frac{x_{i}}{(n+1)^{2}[L_{n+1}(x_{i})]^{2}}$$
(1.66)

The error term is given by

$$E_n = \frac{(n!)^2}{(2n)!} h^{(2n)(\xi)}.$$
(1.67)

1.7.4.4 Gauss-Hermite Rule.

This is a Gauss quadrature over the interval $(-\infty,\infty)$ with weight function $w(x) = e^{x^2}$ (Davis and Rabinowitz, 1984). The nodes for quadrature of order n are given by the roots x_i of the Hermite polynomials H_n , which occur symmetrically about 0. Th weights are

$$w_{i} = -\frac{A_{n+1}\gamma_{n}}{A_{n}H_{n}'(x_{i})H_{n+1}(x_{i})} = \frac{A_{n}}{A_{n-1}}\frac{\gamma_{n-1}}{H_{n-1}(x_{i})H_{n}'(x_{i})},$$
(1.68)

where A_n is the coefficient of x^n in $H_n(x)$. For Hermite polynomials, $A_n = 2^n$, so $\frac{A_{n+1}}{A_n} = 2$. Additionally, $\gamma_n = \sqrt{\pi} 2^n n!$, so (see Abramowitz and Stegun 1968, p.

890).

$$w_{i} = -\frac{2^{n+1}n!\sqrt{\pi}}{H_{n+1}(x_{i})H_{n}'(x_{i})} = \frac{2^{n}(n-1)!\sqrt{\pi}}{H_{n-1}(x_{i})H_{n}'(x_{i})}$$
$$= \frac{2^{n+1}n!\sqrt{\pi}}{\left[H_{n}'(x_{i})\right]^{2}} = \frac{2^{n+1}n!\sqrt{\pi}}{\left[H_{n+1}(x_{i})\right]^{2}} = \frac{2^{n-1}n!\sqrt{\pi}}{n^{2}\left[H_{n-1}(x_{i})\right]^{2}}, \quad (1.69)$$

where the following recurrence relations is used:

$$H'_{n}(x) = 2nH_{n-1} = 2xH_{n}(x) - H_{n+1}(x).$$

The error term is

$$E_n = \frac{n!\sqrt{\pi}}{2^n(2n)!} h^{(2n)(\xi)}.$$
(1.70)

The nodes and weights can be computed analytically for small n as:

$$n = 2: \quad x_i = \pm \sqrt{2}, \quad w_i = \frac{1}{2}\sqrt{\pi}, \ i \in \{1, 2\}$$

$$n = 3: \quad x_0 = 0, \quad w_i = \frac{2}{3}\sqrt{\pi},$$

$$x_i = \pm \frac{1}{2}\sqrt{6}, \quad w_i = \frac{\sqrt{\pi}}{6}, \ i \in \{1, 2\}$$

$$n = 4: \quad x_i = \pm \sqrt{\frac{3 - \sqrt{6}}{2}}, \quad w_i = \frac{\sqrt{\pi}}{4(3 - \sqrt{6})}, \ i \in \{1, 2\}$$

$$x_i = \pm \sqrt{\frac{3 + \sqrt{6}}{2}}, \quad w_i = \frac{\sqrt{\pi}}{4(3 - \sqrt{6})}, \ i \in \{3, 4\}.$$

1.7.4.5 Gauss-Radau Rule.

A Gaussian quadrature-like formula for numerical estimation of integrals. It requires m+1 points and fits all polynomials to degree 2m, so it effectively fits exactly all polynomials of degree 2m-1. It uses a weight function w(x) = 1 in which the endpoint -1 in the interval [-1,1] is included in a total of n abscissae, r-n-1 free abscissae. The general formula is

$$\int_{-1}^{1} h(x)dx = \frac{2}{n^2}h(-1) + \sum_{i=2}^{n} w_i h(x_i).$$
(1.71)

The free nodes for $i = 2, \dots, n$ are the roots of the polynomial $\frac{P_{n-1}(x)+P_n(x)}{1+x}$ where $P_n(x)$ is a Legendre polynomial. The weights of the free nodes are

$$w_i = \frac{1 - x_i}{n^2 [P_{n-1}(x_i)]^2} = \frac{1}{(1 - x_i) [P'_{n-1}(x_i)]^2}, \ i = 1, \cdots, n.$$
(1.72)

The error term is

$$E_n = \frac{2^{2n-1}n[(n-1)!]^4}{[(2n-1)!]^3} h^{(2n-1)}(\xi), -1 < \xi < 1.$$
(1.73)

1.7.4.6 Gauss-Lobatto Rule.

Also known as Lobatto quadrature (Abramowitz and Stegun, 1972), named after Dutch mathematician Rehuel Lobatto. It is similar to Gaussian quadrature with the following differences:

- 1. The integration points include the end points of the integration interval.
- 2. It is accurate for polynomials up to degree 2n-3, where *n* is the number of integration points (Quarteroni et al., 2000).

Lobatto quadrature of function h(x) on interval [-1, 1]: When the Gaussian formula (1.31) is used with the polynomials $P'_{n-1}(x)$, where $P_n(x)$ are the Legendre polynomials, we get Lobatto's rule:

$$\int_{-1}^{1} h(x)dx = \frac{2}{n(n-1)} [h(1) + h(-1)] + \sum_{i=1}^{n-1} w_i h(x_i) + R_n, \quad (1.74)$$

where the quadrature points x_i is the (i-1)-st zero of $P'_{n-1}(x)$; this formula is exact if $h \in \mathcal{P}_{2n-1}$, and the weights $w_i = \frac{2}{n(n-1)[P_{n-1}(x_i)]^2}$ for $x_i \neq 1$; and

$$R_n = \frac{-n(n-1)^3 2^{2n-1} [(2n-1)!]^4}{(2n-1)[(2n-2)!]^3} h^{2n-2}(\xi), \ -1 < \xi < 1.$$
(1.75)

The 3-point Lobbato rule is Simpson's rule.

1.7.4.7 Gauss-Chebyshev Rule.

In the quadrature rule

$$\int_{c_1}^{c_2} w(x)h(x)dx = \sum_{i=0}^n w_i h(x_i),$$
(1.76)

the weight w_i and the nodes x_i can be found for different orthogonal polynomials approximations of h(x) such that formula (1.76) is exact for all $h \in \mathscr{P}_{2n+1}$.

Thus, in particular, if $[c_1, c_2]$ is taken as [-1,1] and $w(x) = (1-x^2)^{-1/2}$, the orthogonal polynomial are the Chebyshev polynomials $T_N(x)$ of the first kind, and the resulting formulas are known as the Gauss-Chebyshev rule. Thus, by taking the interpolating polynomials as

$$p_n(x) = \sum_{i=0}^n l_i(x)h(x_i),$$
(1.77)

where x_i are the zeros of $T_n(x)$. Then, in view of (1.47) the weights w_i are given by

$$w_i = -\frac{\pi}{T_{n+1}(x_i)T'_n(x_i)}.$$
(1.78)

Putting $x = \cos \theta$, we have $T_n(x) = \cos n\theta$, and $T'_n(x) = \frac{n \sin n\theta}{\sin \theta}$. Now we write $x_i = \cos \theta_i$. Then

$$T_{n+1}(x_i) = \cos(n+1)\theta_i = \cos n\theta_i \cos \theta_i - \sin n\theta_i \sin \theta$$
$$= \pm \sqrt{1-x_i},$$

since $\cos n\theta = 0$,

$$T'_n(x_i) = \frac{n \sin n\theta_i}{\sin \theta_i} = \frac{\pm n}{\sqrt{1 - x_i^2}}$$

which gives

$$w_i = \frac{\pi}{n} \tag{1.79}$$

This shows that for all Chebyshev rules the weights are all equal. The error in the n-point rule is given by

$$E_n = \frac{2\pi}{2^{2n}(2n)!} h^{(2n)}(\xi), \ -1 < \xi < 1.$$
(1.80)

If we choose $[c_1, c_2] = [-1, 1]$, $w(x) = (1-x^2)^{-1/2}$, and the polynomials $q_i(x)$ as the normalized Chebyshev polynomials

$$q_i(x) = \begin{cases} \sqrt{\frac{2}{\pi}} T_i(x) & \text{for } i = 1, 2, \cdots, n-1 \\ \sqrt{\frac{1}{\pi}} T_0(x) & \text{for } i = 0. \end{cases}$$

The nodes ξ_i for the *n*-point quadrature are zero of $T_n(x)$, defined by

$$\xi_i = \cos\frac{(2i-1)\pi}{2n}, \ i = 1, 2, \cdots, n.$$
 (1.81)

The rule with ξ_i given by (1.82) and the weight (1.78) is of degree 2n-1, and is known as the *open Gauss-Chebyshev rule*, since the endpoints are not included as the nodes. The word 'open' is generally omitted since all Gaussian rules with positive weight function are of the open type. This rule is also known as Fejér's first integration formula

$$\int_{-1}^{1} h(x)dx = \sum_{i=1}^{n} w_i h\left(\cos\frac{(2i-1)\pi}{2n}\right), \ h \in \mathcal{P}_{n-1}$$
(1.82)

where the weights w_i are given by

$$w_i = \frac{2}{\pi} \left[1 - 2 \sum_{k=1}^{\lfloor n/2 \rfloor} \frac{\cos(2k\theta_k)}{4k^2 - 1} \right].$$
(1.83)

The (open) Gauss-Chebyshev rule is related to the midpoint rule with w(x) = 1, which is defined by

$$H_{c_1}^{c_2}h \approx h \sum_{i=1}^n h\left(c_1 + \frac{(2i-1)h}{2}\right), \ h = \frac{c_2 - c_1}{2}.$$
 (1.84)

This is easily seen by setting $[c_1, c_2] = [0, \pi]$ and $x = \cos \theta$; then the above midpoint rule changes to

$$\int_{-1}^{1} \frac{h(\cos^{-1}x)}{\sqrt{1-x^2}} dx \approx \frac{\pi}{n} \sum_{i=1}^{n} h\left(\frac{(2i-1)\pi}{2n}\right).$$
(1.85)

We can, however, constructed the 'closed' Gauss-Chebyshev rule by including the endpoints $\xi_0 = 1$ and $\xi_n = -1$ as additional nodes. Then we get a rule with nodes $\xi_i = \cos(i\pi/n), i = 0, 1 \cdots$, *n*, and weights $w_i = \pi/n$ for $i = 1, 2, \cdots, N$, and $w_0 = w_n = \pi/(2n)$. Thus, the 'closed' Gauss-Chebyshev rule is

$$\int_{-1}^{1} \frac{h(x)}{\sqrt{1-x^2}} dx \approx \frac{\pi}{n} \sum_{j=0}^{n} h\left(\cos\frac{i\pi}{n}\right).$$
(1.86)

This rule can be obtained from the repeated trapezoidal rule by the same method as used in showing the relationship of the open Gauss-Chebyshev rule to the midpoint rule, although the relation does not clarify why the rule is of high degree. In practice we do not generally use the closed rule because sometimes it is not possible to include the endpoint nodes. However, the closed rule has the advantage in a situation where closed rule the *n* points ξ_i are the subject of the 2n points ξ_i , and the previous function evaluations can be used again.

There are two Gauss-Chebyshev Rules on an arbitrary interval $[c_1, c_2]$:

1. Using the Gauss-Chebyshev polynomials of the first kind, $T_n(x)$, where $T_n(1) = 1/2^{n-1}$, we have

$$\int_{c_1}^{c_2} \frac{h(y)}{\sqrt{(y-c_1)(c_2-y)}} dy = \frac{c_2-c_1}{2} \sum_{i=1}^n w_i h(y_i) + R_n, \quad (1.87)$$

where the nodes $y_i = \frac{c_2+c_1}{2} + \frac{c_2-c_1}{2}x_i$, with $x_i = \cos\frac{(2i-1)\pi}{2n}$; the weights are the same as above.

2. Using the Chebyshev polynomials of the second kind

$$U_n(x) = \frac{\sin[(n+1)\arccos x]}{\sin(\arccos x)},$$
(1.88)

we get

$$\int_{c_1}^{c_2} h(y)\sqrt{(y-c_1)(c_2-y)}dy = \left(\frac{c_2-c_1}{2}\right)^2 \sum_{i=1}^n w_i h(y_i) + R_n, \quad (1.89)$$

where the nodes are y_i and the weights are the same as in 1 above.

1.7.5 Gauss-Kronrod rules

A (2n+1)-point Gauss-Kronrod rule by DIRK (1997) for the integral

$$Ih = \int_{c_1}^{c_2} h(x) ds(x),$$
 (1.90)

where s is a nonnegative measure on the interval $[c_1, c_2]$, is a formula of the form

$$K^{(2m+1)}h = \sum_{i=1}^{2m+1} w_i h(x_i)$$
(1.91)

with the properties:

- *m* nodes of $K^{(2m+1)}$ coincide with those of the *m*-point Gaussian quadrature rule $G^{(n)}$ for the same measure;
- $K^{(2m+1)}h = Ih$ whenever *h* is polynomial of degree less than or equal to 3n+1.

To approximate definite integrals of the form

$$\int_{c_1}^{c_2} h(x) dx \tag{1.92}$$

by using *n*-point Gaussian quadrature

$$\int_{c_1}^{c_2} h(x) dx \approx \sum_{i=1}^n w_i h(x_i),$$
(1.93)

where w_i , x_i are the weights and points at which to evaluate the function h(x). If the interval $[c_1, c_2]$ is subdivided, the Gauss evaluation points of the new subintervals never coincide with the previous evaluation points (except at the midpoint for odd numbers of evaluation points), and thus the integrand must be evaluated at every point. Gauss Kronrod formulas are extensions of the Gauss quadrature formulas generated by adding n+1 points to an n-point rule in such a way that the resulting rule is of order 3n+1. These extra points are the zeros of Stieltjes polynomials. This allows for computing higher-order estimates while reusing the function values of a lower-order estimate. The difference between a Gauss quadrature rule and its Kronrod extension are often used as an estimate of the approximation error.

Patterson (1968) showed how to find further extensions of this type.

In numerical mathematics, the Gauss-Kronrod quadrature formula is a method for numerical integration (calculating approximate values of integrals). Gauss Kronrod quadrature is a variant of Gaussian quadrature, in which the evaluation points are chosen so that an accurate approximation can be computed by re-using the information produced by the computation of a less accurate approximation. It is an example of what is called a nested quadrature rule: for the same set of function evaluation points, it has two quadrature rules, one higher order and one lower order (the latter called an embedded rule). The difference between these two approximations is used to estimate the calculational error of the integration.

These formulas are named after Alexander Kronrod, who invented them in the 1960s, and Carl Friedrich Gauss. Gaussian quadrature is used in the QUAD-PACK library, the GNU Scientific Library, the NAG Numerical Libraries and R (Hazewinkel and Michiel, 2001).

1.7.6 Other Forms of Gauss Quadrature

The integration problem can be expressed in a slightly more general way by introducing a positive weight function $\omega(x)$ into the integrand, and allowing an interval other than [-1,1] say $[c_1,c_2]$. That is, the problem is to calculate

$$\int_{c_1}^{c_2} \omega(x)h(x)dx \tag{1.94}$$

for some choices of c_1 , c_2 , and $\omega(x)$. For $c_1 = -1$, $c_2 = 1$, and $\omega(x) = 1$, the problem is the same as that considered above (section 1.1). Other choices lead to other integration rules mentioned in the table below.

Interval	$\omega(x)$	Orthogonal polynomials	QT
[-1,1]	1	Legendre polynomials	GLQ
(-1,1)	$(1-x)^{\alpha}(1+x)^{\beta}$, $\alpha, \beta > -1$	Jacobi polynomials ($\beta = 0$)	GJQ
(-1,1)	$\frac{1}{\sqrt{1-x^2}}$	Chebyshev polynomials (first kind)	GChQ
[-1,1]	$\sqrt{1-x^2}$	Chebyshev polynomials (second kind)	GChQ
$[0,\infty)$	e^{-x}	Laguerre polynomials	GLgQ
$[0,\infty)$	$x^{\alpha}e^{-x}$, $\alpha > -1$	Generalized Laguerre polynomials	GLgQ
$(-\infty,\infty)$	e^{-x^2}	Hermite polynomials	GHQ

 Table 1.1: Details are given in Abramowitz and Stegun (1972)

1.7.7 Fundamental theorem

Let p_n be a nontrivial polynomial of degree n such that

$$\int_{c_1}^{c_2} \omega(x) x^k p_n(x) dx = 0, \text{ for all } k = 0, 1, 2, \cdots, n-1$$
(1.95)

If we pick the *n* nodes x_i to be the zeros of p_n , then there exist *n* weights w_i which make the Gauss-quadrature computed integral exact for all polynomials h(x) of degree 2n-1 or less. Furthermore, all these nodes x_i will lie in the open interval (c_1, c_2) (Stoer and Bulirsch, 2002).

The polynomial p_n is said to be an orthogonal polynomial of degree n associated to the weight function $\omega(x)$. It is unique up to a constant normalization factor.

The idea underlying the proof is that, because of its sufficiently low degree, h(x) can be divided by $p_n(x)$ to produce a quotient q(x) of degree strictly lower than n, and a remainder r(x) of still lower degree, so that both will be orthogonal to $p_n(x)$, by the defining property of $p_n(x)$. Thus

$$\int_{c_1}^{c_2} \omega(x)h(x)dx = \int_{c_1}^{c_2} \omega(x)r(x)dx$$
(1.96)

Because of the choice of nodes x_i , the corresponding relation

$$\sum_{i=1}^{n} \omega_i h(x_i) = \sum_{i=1}^{n} \omega_i r(x_i)$$
(1.97)

holds also. The exactness of the computed integral for h(x) then follows from corresponding exactness for polynomials of degree only n or less (as is r(x)).

1.7.8 Errors of Gauss-Quadrature formula

The error of a Gaussian quadrature rule can be stated as follows (Stoer and Bulirsch, 2002). For an integrand which has 2n continuous derivatives,

$$\int_{c_1}^{c_2} w(x)h(x)dx - \sum_{i=1}^n w_i h(x_i) = \frac{h^{2n}(\zeta)}{(2n)!} \langle p_n, p_n \rangle$$
(1.98)

for some $\zeta \in (c_1, c_2)$, where p_n is the monic (i.e. the leading coefficient is 1) orthogonal polynomial of degree n and where

$$\langle h,g\rangle = \int_{c_1}^{c_2} w(x)h(x)g(x)dx \tag{1.99}$$

In the important special case of w(x) = 1, we have the error estimate (Kahaner et al., 1989)

$$\frac{(c_2 - c_1)^{2n+1} (n!)^4}{(2n+1)[(2n)!]^3} h^{2n}(\zeta), c_1 < \zeta < c_2$$
(1.100)

In the research work of Stoer and Bulirsch (2002) remark that this error estimate is inconvenient in practice, since it may be difficult to estimate the order 2n derivative, and furthermore the actual error may be much less than a bound established by the derivative. Another approach is to use two Gaussian quadrature rules of different orders, and to estimate the error as the difference between the two results.

Important consequence of the above equation is that Gaussian quadrature of order *n* is accurate for all polynomials up to degree 2n-1.

1.8 Summary

In this chapter we discussed on background of the research study, provided the problem statement, presented the objectives of the research work, gives the scope of the study, presented the limitations of the study, briefly gives the outline of the thesis and provided the basic concepts and definitions useful in understanding the study area of the research.



REFERENCES

- Abramowitz, M. and Stegun, I. A. 1972. *Hand book of mathematical functions with formulas, graphs and tables*. Dover, ISBN 978-0-486-61272-0.
- Allouch, C., Sablonniére, P., Sbibih, D. and Tahrichi, M. 2010. Product Integration Methods Based on Discrete Spline Quasi-Interpolants and Application to Weakly Singular Integral Equations. *Journal of Computational and Applied Mathematics* 233: 2855–2866.
- Auricchio, F., Calabróc, F., Hughes, T. J. R., Reali, A. and Sangalli, G. 2012. A Simple Algorithm for Obtaining Nearly Optimal Quadrature Rules for Nurbs-Based Isogeometric Analysis. *Journal of Comput. Methods Appl. Mech. Engrg.* 249-252: 15–27.
- Bakhvalov, N. S. and Vasil'eva, L. G. 1968. Evaluation of the integrals of oscillating functions by interpolation at nodes of Gaussian quadratures (Russian) USSR. *Computational Mathematics and Mathematical Physics* 8: 241–249.
- Bonnet, M. and Guiggiani, M. 2003. Direct Evaluation of Double Singular Integrals and New Free Terms in 2D (Symmetric) Galerkin BEM. *Comput. Methods Appl. Mech. Engrg.* 192: 2565–2596.
- Boykov, I. V. 2001. Numerical methods of computation of singular and hypersingular integrals. *Int. J. Math. Sci.* 28(3): 127–79.
- Boykov, I. V., Ventsel, E. S. and Boykova, A. I. 2009. Accuracy optimal methods for evaluating hypersingular integrals. *J. Appl. Numer. Maths.* 59: 1366–1385.
- Chan, Y., Fannjiang, A. C. and Paulino, G. H. 2003. Integral equations with hypersingular kernels-theory and applications to fracture mechanics. *Int. J. Eng. Science* 41: 683–720.
- Chan, Y. S., Fannjiang, A. C., Paulino, G. H. and Feng, B. F. 2007. *Finite part integrals and hypersingular kernels*., vol. 14. DCDIS a supplement, advances in dynamical systems: DCDIS a supplement, advances in dynamical systems.
- Chen, Y. Z. 1993. A numerical solution technique of hypersingular integral equation for curved cracks. *Eng. Frac. Mech.* 46: 215–283.
- Clenshaw, C. W. and Curtis, A. R. 1960. A Method for Numerical Integration on An Automatic Computer. *Numer. Math.* 2: 197.
- Colm, P. and Rokhlin, V. 2001. Numerical Quadratures for Singular and Highpersingular Integrals. *Computers and Mathematics for Application* 41: 327– 352.
- Cools, R. 2003. An Encyclopaedia of Cubature Formulas. *Journal of Complexity* 19: 445–453.

- Cools, R. and Rabinowitz, P. 1993. Monomial cubature rules since ï£iStroudï£i: a compilation. *Journal of Computational and Applied Mathematics* 48: 309– 326.
- Davis, P. and Rabinowitz, P. 1984. *Methods of Numerical Integration , Second Addition*. New York London: Academic Press (INC).
- Davis, P. J. and P., R. 1984. Methods of numerical integration, in: Computer Science and Applied Mathematics. New York: Academic Press.
- Demichelis, V. and Sablonniére, P. 2010. Cubature Rule Associated With a Discrete Blending Sum of Quadratic Spline Quasi-Interpolants. *Journal of Computational and Applied Mathematics* 235: 174–185.
- DIRK, P. L. 1997. Calculation of Gauss-Kronrod Quadrature Rules. *Mathematics* of Computation 66(219): 1133–1145.
- Elliott, D. 1965. Truncation errors in two chebyshev series approximations. *Math. Comp.* 19: 234–248.
- Eshkuvatov, Z. K., Nik Long, N. M. A., Midi, H. and Khaldjigitov, A. 2013. Polynomial Spline Approach for Double Integrals with Algebraic Singularity. *IOP Publishing, Journal of Physics: Conference Series* **43**5 doi:10.1088/1742-6596/435/1/012009: 1–7.
- Eskhuvatov, Z.K.and Ahmedov, A., Nik Long, N. M. A. and Amalina, N. J. 2011. Approximating Cauchytype Singular Integral by an Automatic Quadrature Scheme. J. Comp. Appl. Math. 235: 4675–4686.
- Eskhuvatov, Z. K. and Nik Long, N. M. A. 2011. Approximating the Singular Integrals of Cauchy Type With Weight Function on the Interval. *J. Comp. Appl. Math.* 235: 4742–4753.
- Evans, G. A. 1993. *Practical Numerical Integration*. Chichester: John Wiley & Sons.
- Filon, L. N. G. 1928, On a quadrature formula trigonometric integrals, Tech. rep., Proc. Roy. Soc.
- Gao, X. 2010. An Effective Method for Numerical Evaluation of General 2D and 3D High Order Singular Boundary Integrals. *Comput. Methods Appl. Mech. Engrg.* 199: 2856–2864.
- Gao, X. W. 2002. The Radial Integration Method for Evaluation of Domain Integrals With Boundary-Only Discretization. *Engrg. Anal. Boundary Elem.* 26: 905–916.
- Hadamard, J. 1932. Le Probleme de Cauchy et les Equations aux Derivees Particelles Lineaires Hyperboliques. Paris: Hermann.

- Hadamard, J. 1952. Lectures on Cauchy's problem in linear partial differential equations. New York: Dover Publications.
- Hasegawa, T. and Torii, T. 1991. An Automatic Quadrature for Cauchy Principle Value Integrals. *Math. Comp.* 56: 741–754.
- Hazewinkel and Michiel. 2001. *Gaussï£iKronrod quadrature formula*. Springer, ISBN 978-1-55608-010-4: Encyclopedia of Mathematics.
- Hui, C. Y. and Mukherjee, S. 1997. Evaluation of Hypersingular Integrals in the Boundary Element Method by Complex Variable Techniques. Int. J. Solids Struct. 34: 203–221.
- Hui, C. Y. and Shia, D. 1999. Evaluation of hypersingular integrals using Gaussian Quadrature. *Int. J. Numer. Meth. Engng.* 44: 205–214.
- Iserles, A. and Norsett, S. P. 2005. Efficient quadrature of highly oscillatory integrals using derivatives. *Uzbek Mathematical Journal (ISSN 2010-7269), Proc. R. Soc. A* 461: 1383–1399.
- Israilov, M. I. 1986. Estimate of error of quadrature formula to singular integrals of special type, in: differential equations and inverse problems. *Press FAN, Tashkent* 1.
- Israilov, M. I. 1990. Semi-analytic solution of singular integral equations of first kind by quadrature formula. *Inst. Maths. under Nat. Univ. Uzbek.* 1.
- Johnson, L. W. and Riess, R. D. 1977. *Numerical Analysis*. London: Addison-Wesley.
- Kahaner, D., Moler, C. and Nash, S. 1989. *Numerical Methods and Software*. Prentice-Hall, ISBN 978-0-13-627258-8.
- Kim, P. 2001. A Chebyshev quadrature rule for One sided finite parts integrals. JAT 111: 196–219.
- Kowalewski. 1984. Interpolation und Genährte. Academic press, INC: Computer Science and Applied Mathematics.

Krommer, A. R. and Ueberhuber, C. W. 1998. *Computational Integration*. Philadelphia: SIAM.

- Kutt, H. R. 1975, Quadrature formulae for finite part integrals, Tech. Rep. Report WISK 178, The National Research Institute for Mathematical Sciences, Pretoria.
- Li, C. and Dagnino, C. 2010. An Adaptive Numerical Integration Algorithm for Polygons. *Applied Numerical Mathematics* 60: 165–175.

- Li, C., Lamberti, P. and Dagnino, C. 2009. Numerical Integration Over Polygons Using an Eight-Node Quadrilateral Spline Finite Element. *Journal of Computational and Applied Mathematics* 233: 279–292.
- Li, C. J. and Wang, R. H. 2006. A new 8-node quadrilateral spline finite element. J. Comp. Appl.Math. 195: 54–65.
- Lifanov, I. K., P. L. N. V. G. M. 2004. *Hypersingular integral equations and their applications*. London: Chapman & Hall/CRC, A CRC Press Company.
- Lutz, E., Gray, L. and Ingraffea, A. 1991. An overview of integration methods for hypersingular integrals. *J. Bound. Ele, Southampton: CMP.* 8: 913–25.
- Martin, P. A. and Rizzo, F. J. 1996. Hypersingular Integrals: How Smooth Must The Density Be? Int. J. Num. Meth. Engng 39: 687–704.
- Mason, J. C. and Handscomb, D. C. 2003. *Chebyshev polynomials*. LLC: CRC Press.
- Mazzia, F. and Sestini, A. 2012. Quadrature Formulas Descending from BS Hermite Spline Quasi-Interpolation. *Journal of Computational and Applied Mathematics* 236: 4105–4118.
- Mennouni, A. and Guedjiba, S. 2010. A note on Solving integro-differential equation with Cauchy kernel. *Mathematical and Computer Modelling* 52: 1634– 1638.
- Milovanovic, G. V. 1998. Numerical calculation of integrals involving oscillatory and singular kernels and some applications of quadratures. *Computers Math. Applic.* 36, no. 8: 19–39.
- Milovanovic, G. V. and Stanic, M. P. 2014. Numerical integration of highly oscillating functions II Analytic Number Theory, Approximation Theory, and Special Functions. *Springer* 1: 613–649.
- Monegato, G. 1960. Definitions, Properties and Applications of Finite-Part Integrals. *Journal of Computational and Applied Mathematics* 2009: 425–439.
- Nik Long, N. M. A. and Eskhuvatov, Z. K. 2009. Hypersingular integral equation for multiple curved cracks problem in plane elasticity. *Int. J. Solids Struct.* 46: 2611–2617.
- Novak, E., Ullrich, M. and Wozniakowski, H. 2015. Complexity of oscillatory integration for univariate Sobolev space. *Journal of Complexity* 31: 15–41.
- Obayis, S. J., Eshkuvatov, Z. K. and Nik Long, N. M. A. 2013. On error estimation of automatic quadrature scheme for the evaluation Of hadamard integral of second order singularity. *U.P.B. Sci. Bull., Series A* 75: 85–98.

- Olver, S. 2008. Numerical approximation of highly oscillatory integrals. *PhD* dissertation, University of Cambridge 1: 613–649.
- Patterson, T. N. L. 1968. The Optimum Addition of Points to Quadrature Formulae. Math. Comput. (American Mathematical Society) and C1-C11, doi:10.2307/2004583 22 (104): 847–856.
- Piessens, R. and Branders, M. 1973. The Evaluation and Application of some Modified Moments. BIT, 443-450.
- Prem, K. K. and Michael, R. S. 2005. Handbook of Computational Methods for Integration. New York Washington, D.C.: A CRC Press Company Boca Raton London.
- Quarteroni, A., Sacco, R. and Saleri, F. 2000. Numerical Mathematics. New York: Springer-Verlag, ISBN 0-387-98959-5 22 (104): 422–425.
- Rohman, N. A. A. 1996. Approximation of functions of two variables with modified spline functions of type (0,2). Annales Univ. Sci. Budapest., Sect. Comp. 16: 303–318.
- Samko, S. G. 2002. *Hypersingular integrals and their applications*. London: Taylor & Francis.
- Shadimetov Kh, M. 1999. Weight optimal cubature formulas in sobolev's periodic space (Russian). Siberian J. Numer. Math. -Novosibirsk, 2, no. 2: 185–196.
- Stoer, J. and Bulirsch, R. 2002. Introduction to Numerical Analysis (3rd ed.). Springer, ISBN 978-0-387-95452-3.
- Stroud, A. H. 1973. Approximate Calculation of Multiple Integrals. *W. G. Mathematics of Computation* 27: 437–440.
- Stroud, A. H. and Secrest, D. 1966. *Gaussian Quadrature Formulas*. Englewood Cliffs: N.J.:Prentice-Hall.
- Tadeu, A. and Antonio, J. 2012. 3D Acoustic Wave Simulation Using Bem Formulations: Closed Form Integration of Singular and Hypersingular Integrals. *Engineering Analysis with Boundary Elements* 36: 1389–1396.

Teubner and Leipzig. 1993, Tech. rep.

- Weston, S. 2002, An introduction to the mathematics and construction of splines, Tech. rep., Addix Software Consultancy Limited.
- Yaghobifar, M., Nik Long, N. M. A. and Eshkuvatov, Z. K. 2011. Analytical-Approximate Solution of Abel Integral Equations. *International Mathematical Forum* 6: 211–221.

- Yang, C. 2012. A Unified Approach With Spectral Convergence for the Evaluation of Hypersingular and Supersingular Integrals With a Periodic Kernel. J. Comp. Appl. Math., DOI 10.1016/j.cam.2012.08.028.
- Zhang, X., Wu, J. and Yu, D. 2009. Superconvergence of the composite Simpsons rule for a certain finite-part integral and its applications. *J. Appl. Numer. Maths.* 223: 598–613.

