



UNIVERSITI PUTRA MALAYSIA

***GOLD POTENTIAL MAPPING USING BIVARIATE
AND MULTIVARIATE STATISTICAL MODELS IN GIS
AT GUA MUSANG, MALAYSIA***

SUHAIMIZI BIN YUSOFF

FK 2015 173



**GOLD POTENTIAL MAPPING USING BIVARIATE
AND MULTIVARIATE STATISTICAL MODELS IN GIS
AT GUA MUSANG, MALAYSIA**

By

SUHAIMIZI BIN YUSOFF

**Thesis Submitted to the School of Graduates Studies,
Universiti Putra Malaysia, in Fulfilment of the
Requirements for the Degree of Master of Science**

JANUARY 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Master of Science

GOLD POTENTIAL MAPPING USING BIVARIATE AND MULTIVARIATE STATISTICAL MODELS IN GIS AT GUA MUSANG, MALAYSIA

By

SUHAIMIZI BIN YUSOFF

January 2015

Chairman: Biswajeet Pradhan, PhD

Faculty: Engineering

Throughout the history of humanity, gold remains as the most desired metals. Due to its small occurrences in the earth's crust, this precious metal acquisition is very difficult. In Malaysia, gold potential map have been generated using conventional methods and are inadequate and lacking the ability to assess the accuracy. Develop new techniques has overcome this weakness and currently used around the world. The first time to be applied in Malaysia, this study aim to assess the capability of data-driven Geographical Information System (GIS) modelling technique for mapping gold potential areas of Kelantan, Malaysia. The study area is located at the south of Kelantan state, Malaysia borders Pahang state. The study area covers about 593 km² and is situated approximately 186 km from Kota Bharu, Kelantan. In this study, six gold deposit controlling factors that influence the gold deposit occurrences were extracted from available maps and spatial databases. These controlling factors are: lithology, fault, geochemical data of Copper (Cu), Lead (Pb) and Tungsten (W) and geophysical data of Potassium (K). In the GIS environment, all six controlling factors were integrated and modelled based on data-driven technique. The generation of the gold potential map was carried out using two different types of GIS modelling techniques. The models applied are evidential belief functions (EBF) and logistic regression (LR). The spatial relationship between gold deposit and its controlling parameters was assessed. The predicted gold potential map was classified into four distinct zones based on the classification scheme from the literatures. The analysis and comparison of these results indicate that: (1) The gold potential map generated by EBF model is considered as the best results with prediction accuracy of 81%, (2) The gold potential map generated using LR model has low prediction accuracy of 62.67% and (3) The most influential controlling factors for gold deposit occurrences is lithology, followed by Cu, W, fault, K and Pb. The predicted gold potential map of the study area generated using EBF technique indicated that about 4.8% or 28.49 km² are in the very high potential zone, with about 10.9% or 65.22 km² in high potential zone, with about 38.7% or 229.6 km² fall in the moderate potential zone, and about 45.6% or 269.69 km² constituting the low potential zone. The results indicate that it can be used for future planning of gold exploration by providing a rapid reproduction approach with reduce

time and cost. The results also demonstrate that this modelling technique may also apply to other area with similar parameters.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Sarjana Sains

PEMETAAN POTENSI EMAS MENGGUNAKAN MODEL STATISTIK BIVARIAT DAN MULTIVARIAT DALAM GIS DI GUA MUSANG, MALAYSIA

Oleh

SUHAIMIZI BIN YUSOFF

Januari 2015

Pengerusi: Biswajeet Pradhan, PhD

Fakulti: Kejuruteraan

Sepanjang sejarah manusia, emas kekal sebagai logam yang paling dikehendaki. Disebabkan kewujudan kecil dalam kerak bumi, pemerolehan logam berharga ini adalah sangat sukar. Di Malaysia, peta potensi emas telah dihasilkan menggunakan kaedah konvensional yang selalunya tidak memadai dan tidak mempunyai keupayaan untuk dinilai ketepatannya. Penghasilan teknik-teknik baru telah mengatasi kelemahan ini dan pada masa ini sedang digunakan di seluruh dunia. Kali pertama untuk digunakan di Malaysia, kajian ini bertujuan untuk menilai keupayaan teknik pemodelan panduan data Sistem Maklumat Geografi (GIS) untuk pemetaan kawasan potensi emas di Kelantan. Kawasan kajian terletak di selatan negeri Kelantan, Malaysia bersempadan dengan negeri Pahang. Kawasan kajian meliputi kira-kira 593 km² dan terletak kira-kira 186 km dari Kota Bharu, Kelantan. Dalam kajian ini, enam faktor yang mempengaruhi kewujudan deposit emas diekstrak daripada peta-peta dan pangkalan data spatial yang ada. Faktor-faktor yang mempengaruhi adalah: litologi, sesar, data geokimia tembaga (Cu), Plumbum (Pb) dan Tungsten (W) dan data geofizik kalium (K). Kesemua enam faktor pengawal disepadukan dan dimodelkan berdasarkan teknik panduan data di dalam persekitaran GIS. Penghasilan peta potensi emas dilakukan dengan menggunakan dua teknik pemodelan GIS yang berbeza. Model-model yang digunakan adalah *evidential belief functions (EBF)* dan *logistic regression (LR)*. Hubungan spatial antara deposit emas dengan parameter-parameter yang mengawalinya dinilai. Peta ramalan potensi emas diklasifikasikan kepada empat zon berbeza berdasarkan skema klasifikasi dari literasi. Analisis dan perbandingan keputusan menunjukkan bahawa: (1) Peta potensi emas yang dihasilkan menggunakan model *EBF* dianggap sebagai hasil yang terbaik dengan ketepatan ramalan sebanyak 81%, (2) Peta potensi emas yang dijana menggunakan model *LR* mempunyai ketepatan ramalan yang rendah iaitu 62.67% dan (3) Faktor yang paling berpengaruh untuk kehadiran deposit emas adalah litologi, diikuti dengan Cu, W, sesar, K dan Pb. Peta ramalan potensi emas kawasan kajian menggunakan teknik *EBF* menunjukkan kira-kira 4.8% atau 28.49 km² berada di dalam zon potensi sangat tinggi, kira-kira 10.9% atau 65.22 km² di zon berpotensi tinggi, kira-kira 38.7% atau berjumlah 229.6 km² di zon potensi sederhana, dan kira-kira 45.6% atau 269.69 km² membentuk zon potensi rendah. Keputusan menunjukkan bahawa ia boleh digunakan untuk perancangan masa depan penerokaan

emas dengan menyediakan pendekatan penghasilan semula peta ramalan yang pantas dengan pengurangan masa dan kos. Keputusan ini juga menunjukkan bahawa teknik pemodelan ini juga boleh digunakan di kawasan lain dengan parameter yang sama.



ACKNOWLEDGEMENTS

First of all praise to Allah the Almighty for His blessings that gives me strength which enable me to complete this MSc thesis. Honestly, I face many obstacles to complete this study within the time period given of two years by the financial sponsor. So I would like to take the opportunity to thank my supervisor, Associate Prof. Habil Dr. Biswajeet Pradhan for checking and editing the draft of my thesis. He also gives many valuable suggestions as well as continuous encouragement during the preparation of this thesis.

I also want to thank other members of supervisor committee Associate Prof. Dr. Helmi Zulhaidi Mohd. Shafri who have helped and give me a lot of invaluable advices. Thousands of thanks to Prof Dr. Wan Fuad Wan Hassan, Dr. Mohamad Abdul Manap, Dr. Nuraini Surip, Wan Mohd Razi Idris, Mohd Zukeri Ab Ghani, Mustafa Hamzah, Abdul Halim Hamzah, Abdul Hadi Abdul Rahman, Zulkiflee Che Soh, John Joseph Jinap, Mohd Anuar Ishak, Mohamed Hizam Abdul Kadir, Mohamed Asri Omar, and Zaki Alias who give cooperation and suggestion.

Besides that, I would like to express my gratitude to my current employer the Department of Minerals and Geoscience (JMG) and Department of National Mapping and Survey (JUPEM) for providing the necessary data and information to undertake during this study.

I also want to express my appreciation to JMG for selecting me to pursue my study at the MSc level. I would like to express my gratitude to the Government of Malaysia through the Public Services Department (JPA) for providing the financial support.

To my colleagues and friends particularly in the JMG Terengganu, JMG headquarters in Kuala Lumpur, JMG state office, Universiti Putra Malaysia (UPM) and Universiti Kebangsaan Malaysia (UKM), many thanks to all of you for your supports and assistances.

Last but not least, my warmest thanks to my family members especially my beloved wife Seri Hayati Annuar, my three charming prince and princesses namely Nur Husna, Mohamad Haris and Nur Hazirah, my parent Hajjah Karimah Abdullah and Haji Yusoff Ali, my wife parent Che Mahani Tengku Ibrahim and Annuar Yahya, my siblings as well as my wife siblings who have been very supportive and faithfully praying all the time for my success.

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Biswajeet Pradhan, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Helmi Zulhaidi Mohd. Shafri, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: _____

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
CHAPTER	
1 INTRODUCTION	
1.1 Background of the research	1
1.2 Problem statement	2
1.3 Goal of study	2
1.4 Objectives	2
1.5 Research questions	2
1.6 Scope and limitation of the study	3
1.7 The structure of the thesis	4
2 LITERATURE REVIEW	
2.1 Introduction	5
2.2 Geographical information system (GIS)	5
2.3 Remote sensing (RS)	5
2.4 Gold status in Malaysia	5
2.5 Previous works of using RS and GIS on gold potential mapping	8
2.6 Controlling factors used in gold potential mapping	9
2.7 GIS modelling techniques for gold potential mapping	10
2.7.1 Knowledge-driven method	10
2.7.1.1 Fuzzy logic	10
2.7.1.2 Evidential belief functions (EBF)	12
2.7.2 Data-driven method	12
2.7.2.1 Frequency ratio (FR)	12
2.7.2.2 Weights-of-evidence (WoE)	13
2.7.2.3 Logistic regression (LR)	13
2.7.2.4 Artificial neural network (ANN)	13
2.8 Model validation	14
2.9 Summary	14

3	METHODOLOGY	
3.1	Introduction	15
3.2	General information of the study area	15
3.2.1	Geography and study area selection	15
3.2.2	Climate and vegetation	15
3.2.3	Topography	15
3.2.4	General geology and gold mineralization	16
3.3	Data sources	18
3.4	Methodology	19
3.4.1	Extraction of gold deposit location	20
3.4.2	Extraction of gold deposit controlling factors	21
3.4.2.1	Lithology	21
3.4.2.2	Structural controlling features	21
3.4.2.3	Geochemical	21
3.4.2.4	Geophysical	27
3.4.3	GIS modelling	27
3.4.3.1	Evidential belief functions	27
3.4.3.2	Logistic regression	31
3.4.3.3	Model validation	32
3.5	Summary	32
4	RESULTS AND DISCUSSION	
4.1	Introduction	33
4.2	Results for evidential belief functions model	33
4.2.1	Predicted gold potential map	35
4.2.2	Model validation	35
4.3	Results for logistic regression model	40
4.3.1	Predicted gold potential map	42
4.3.2	Model validation	43
4.4	Comparison of gold potential map	45
4.4.1	Comparisons of gold potential maps obtained by evidential belief functions and logistic regression model	45
4.4.2	Comparisons of maps obtained by evidential belief functions model with JMG existing gold potential map	47
4.4.3	The final gold potential map	49
4.5	Summary	50
5	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	
5.1	Summary and main conclusions	51
5.2	Recommendations	52
5.3	Future research	52
	REFERENCES	53
	APPENDICES	65
	BIODATA OF STUDENT	67
	LIST OF PUBLICATIONS	68

LIST OF TABLES

Table		Page
2.1	Controlling factors used by several researchers for mineral potential mapping using GIS (from year of 2007 to 2013)	11
3.1	Description of every data set applied in the current study	19
3.2	Statistics description of the geochemical elements and skewness of the log-transformed data	27
4.1	The evidential belief functions value of gold controlling factors	34
4.2	The logistic regression coefficient value of gold controlling factors	40
4.3	Comparison of evidential belief functions and logistic regression model for gold potential mapping	45
4.4	Controlling factors spatial correlation using area under the curve approach	49

LIST OF FIGURES

Figure	Page
3.1	16
3.2	17
3.3	20
3.4	22
3.5	23
3.6	24
3.7	25
3.8	26
3.9	28
4.1	36
4.2	38
4.3	39
4.4	43
4.5	44
4.6	46
4.7	48
4.8	49

LIST OF ABBREVIATIONS

ANN	Artificial Neural Network
ASTER	Advanced Spaceborne Thermal Emission and Reflection Radiometer
cps	Counts per Second
EBF	Evidential Belief Functions
ETM+	Enhanced Thematic Mapper Plus
FR	Frequency Ratio
GDP	Gross Domestic Product
GIS	Geographical Information Systems
JMG	Minerals and Geoscience Department
LR	Logistic Regression
MINGEOSIS	Minerals and Geoscience Information System
PALSAR	Phased Array type L-band Synthetic Aperture Radar
SPOT	Satellite Pour l'Observation de la Terre
UKM	National University of Malaysia
UM	University of Malaya
WoE	Weights-of-Evidence
WGS84	World Geodetic System 1984

CHAPTER 1

INTRODUCTION

1.1 Background to the research

For centuries mankind has been captivated by the absolute beauty of gold and its many unique properties. In recent years, the price of gold has risen sharply due to an ever increasing demand for gold that is always exceeding supply. World gold supply in 2013 mainly comes from China, Australia, Russia, United States of America, Peru, South Africa and other countries (Australian Gold, 2013). Throughout the history, foreign traders have acknowledged the Malay Peninsula as a gold producing country or Aurum Khersonese as written by Claudius Ptolemy in the second century AD (Wheatley, 1961).

Gold abundance in the earth's crust is very rare, making the metal acquisition very difficult. The abundances of gold in igneous rock and sediment earth's crust are very small between 4-6ppb (Turekian and Wedepohl, 1961), like 4 peas in 1,000 tonnes of sand. In Malaysia, the figure is also quite similar with granite rocks having 1.0 ppb (Hassan et al., 1997). This small abundance contributes to the low content of gold in earth. As a reference, a total of 174,100 tonnes of gold have been extracted until 2012 (World Gold Council, 2013a). This is roughly equivalent to about 9,261 m³.

Approximately 50% of the world's gold has been used as jewelry, 40% in finance, and 10% in manufacturing. The price of gold as of 2013 was RM159 per gram and nearly 60% of the gold's price goes to the extraction process (World Gold Council, 2013b).

Currently, gold occurrences in Peninsular Malaysia are predominantly found in the Central Belt, stretching from Batu Melintang in Kelantan in the North and southwards through Sokor, Pulai, Selinsing, Raub, Chenderas to Gunung Ledang in Johor (Chu, 2000). Although seemingly random, gold distribution is actually controlled by geological processes, mostly by the Permian-Triassic volcanic arc of Peninsular Malaysia. Types of gold mineralization in Peninsular are quartz vein, massive sulphide, skarn type and intrusion-related gold.

According to MacDonald (1967), gold production in Kelantan started from Pulai through Galas and Pergau until the border of Thailand. At that time, the most important area for gold production was Pulai (Middlebrook, 1933). Kelantan produced a great deal of gold in early 19th century that Dodge (1977) assumed Kelantan was the main gold producer in Peninsular Malaysia. In Kelantan, gold exploration by the Europeans was pioneered by Duff Company in 1903 using dredge for placer gold in Sungai Galas and adit for primer gold in Sokor. However, due to lack of profit, the works stopped in 1907 (Low, 1921).

1.2 Problem statement

In Malaysia, less effort is made on gold potential mapping using Geographic Information System (GIS) and Remote Sensing (RS). Only Surip et al., (2007) presented some case studies in Penjom-Merapoh, Pahang using knowledge- driven modeling techniques. This present study is the first attempt to utilize data-driven techniques in GIS environment. GIS is used in the integration and analyzing of a selection of Geoscience information, namely geological, geochemical and geophysical factors. The spatial relationships between geochemical, geological and geophysical factors with gold deposits occurrences are unknown. No gold potential map has been generated from multiple controlling factors such as geochemical, geological and geophysical data. The existing published gold potential map by JMG in the year of 1987 is based only on the conventional approach of overlaying factors without qualitative analysis with the gold deposits and this can be further improved using new techniques.

1.3 Goal of study

The goal of this study is to evaluate the capability of data-driven GIS modelling techniques for mapping gold potential areas at Gua Musang, Kelantan, Malaysia.

1.4 Objectives

There are four main objectives of this study as follows:

- i. To identify the spatial relationship between geochemical, geological and geophysical factors with the occurrence of gold;
- ii. To apply bivariate and multivariate prediction models (evidential belief functions and logistic regression) for finding the best prediction approach for gold potential mapping in Malaysia;
- iii. To generate gold potential maps using a quantitative gold potential index;
- iv. To compare the predicted gold potential map with the existing published gold potential map by JMG;

1.5 Research questions

This study will try to answer four research questions, which are:

- i. What are the factors that control the occurrences of gold in the study area?
- ii. Which controlling factors have significant association with occurrences of gold in the study area?

- iii. Which data-driven GIS modeling technique is the most suitable for delineation of the gold potential map in the study area?
- iv. Are the generated gold potential maps from GIS performing better than the currently published gold potential map?

1.6 Scope and limitation of the study

The list below contains the scope of the study:

- i. Gold deposit.

There are eight known gold deposits in the study area. This research is to assess and understand the spatial correlation with features controlling its occurrences.

- ii. Gold deposit occurrences controlling factors.

Six gold deposit controlling factors were used in this study i.e. lithology, fault, geochemical data of Copper (Cu), Lead (Pb) and Tungsten (W) and geophysical data of Potassium (K).

- iii. Generation of predicted gold potential map by using two different types of data-driven GIS modeling techniques.

The models are evidential belief functions (EBF) and logistic regression (LR).

- iv. Model validation using area under the curve (AUC) and the existing gold deposit.

An available gold deposits was used for model validation.

- v. Map comparison for the best data-driven GIS modelling technique.

Comparison between predicted gold potential maps generated from evidential belief functions and logistic regression.

This study is limited by two factors, namely:

- i. The type of gold deposit.

Due to lack of information, it is unknown whether gold deposits used are primary or secondary. Knowing the type will provide a more focused study.

- ii. Limitation of gold potential map produced.

Gold potential in this study is rather estimated. The economic value is not considered as the size and grade of deposit cannot be determined.

1.7 The structure of the thesis

This thesis is split into five chapters. The summary of each chapter is:

- i. **CHAPTER 1: INTRODUCTION.** This chapter briefly discusses the problem statement of the study, its goal, objectives and the scope of the study. Also included in this chapter are research questions and the overall structure of the thesis.
- ii. **CHAPTER 2: LITERATURE REVIEW.** This chapter provides an overview of the gold status and previous works on gold potential mapping based on GIS and RS in Malaysia. It also presents the gold controlling factors that influence the gold occurrences and a discussion describing the GIS modeling techniques applied to delineate the gold potential areas. Lastly, the validation approaches were used to evaluate the accuracy of the prediction maps.
- iii. **CHAPTER 3: METHODOLOGY.** This chapter briefly describes the characteristic of the study area. It then focuses on methodology, the sources of all the data, the GIS model used and their validation approach for gold potential mapping of the study area.
- iv. **CHAPTER 4: RESULTS AND DISCUSSION.** This chapter shows the findings of the two different types of data-driven GIS modelling techniques supported by diagrams and tables, followed by a comparison between the two data-driven GIS modelling technique in gold potential mapping and a comparison of the best predicted gold potential map with the existing gold prospective map of JMG. Finally, a selection of the best prediction gold potential maps and the significant relationship between gold controlling factors with gold occurrences in the study area will be discussed.
- v. **CHAPTER 5: SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH.** This chapter offers the overall conclusion from this study, recommendation and further research for the study area.

REFERENCES

- Agterberg, F.P. (1971). A probability index for detecting favourable geological environments. *Canadian Institute of Mining and Metallurgy* 10: 82–91.
- Agterberg, F.P. (1973). Probabilistic models to evaluate regional mineral potential. *Mathematical methods in Geoscience, Symposium held at Pribram, Czechoslovakia*: 3–38.
- Agterberg, F.P. (1974). Automatic contouring of geological maps to detect target areas for mineral exploration. *Mathematical Geology* 6: 373–395.
- Agterberg, F.P. (1988). Application of recent developments of regression analysis in regional mineral resource evaluation. In: C.F. Chung, A.G. Fabbri, R. Sinding-Larsen (Eds.), *Quantitative Analysis of Mineral and Energy Resources*, D. Reidel Publishing Company, Dordrecht: 1-28.
- Agterberg, F.P., Bonham-Carter, G. F., Cheng, Q. and Wright, D. F. (1993). Weights of evidence modeling and weighted logistic regression for mineral potential mapping. In: J.C. Davis & U.C. Herzfeld (eds.) *Computers in Geology-25 Years of Progress*. Oxford University Press, International Association of Mathematical Geology. *Studies in Mathematical Geology* 5:13-32.
- Airo, M.-L. (2000). Aeromagnetic and aeroradiometric response to hydrothermal alteration. *Surveys in Geophysics*, 23 (4): 273-302.
- Althuwaynee, O.F., Pradhan, B. and Ahmad, N. (2014). Landslide susceptibility mapping using decision-tree based CHI-squared automatic interaction detection (CHAID) and Logistic regression (LR) integration. *IOP Conference Series: Earth and Environmental Science*, 20 (1): 012032.
- Althuwaynee, O.F., Pradhan, B. and Lee, S. (2012). Application of an evidential belief function model in landslide susceptibility mapping. *Computers & Geosciences* 44: 120-135.
- Althuwaynee, O.F., Pradhan, B., Park, H.J. and Lee, J.H. (2014). A novel ensemble bivariate statistical evidential belief function with knowledge-based analytical hierarchy process and multivariate statistical logistic regression for landslide susceptibility mapping. *Catena* 114: 21-36.
- An, P., Moon, W.M. and Rencz, A.N. (1991). Application of fuzzy set theory to integrated mineral exploration. *Canadian Journal of Exploration Geophysics* 27:1-11.
- An, P., Moon, W.M. and Bonham-Carter, G.F. (1992). On knowledge-based approach on integrating remote sensing, geophysical and geological information. *Proceedings of International Geoscience and Remote Sensing Symposium (IGARSS)*: 34-38.

- An, P., Moon, W.M. and Bonham-Carter, G.F. (1994). Uncertainty management in integration of exploration data using the belief function. *Nonrenewable Resources* 3(1): 60-71.
- Arianne, F. and Craig, J.R.H. (2013). Mineral potential mapping in frontier regions: A Mongolian case study. *Ore Geol. Rev.* 51:15-26.
- Ariffin, K.S. (2012). Mesothermal lode gold deposit central belt Peninsular Malaysia, Earth Sciences, Dr. Imran Ahmad Dar (Ed.), ISBN: 978-953-307-861-8, InTech.
- Austin, J.R. and Blenkinsop, T.G. (2009). Local to regional scale structural controls on mineralisation and the importance of a major lineament in the eastern Mount Isa Inlier, Australia: Review and analysis with autocorrelation and weights of evidence. *Ore Geol. Rev.* 35:298-316.
- Australian Gold. (2013). Gold production (tonnes) by country. <http://www.australian-gold.com>. Retrieved on 16 September 2013.
- Aw, P.C. (1975). Sampling and distribution of alluvial gold in the Sungai Chendawan-Sungai Lalat area, Southeast Kelantan. Geological Survey Malaysia Annual Report 1975.
- Bateman, A.M. (1951). *Economic Mineral deposits*. Wiley and Sons, New York.
- Beane, R.E. (1982). Hydrothermal alteration in silicate rocks, southwestern North America; In: *Advances in Geology of the Porphyry Copper Deposits, Southwestern North America: Tucson*(ed.) Titley S.R., Univ. Ariz. Press, Chapter 6.
- Bonham-Carter, G.F., Agterberg, F.P. and Wright, D.F. (1988). Integration of geological datasets for gold exploration in Nova Scotia. *Photogrammetric Engineering and Remote Sensing* 54:1585-1592.
- Bonham-Carter, G.F., Agterberg, F.P. and Wright, D.F. (1989). Weights of evidence modeling: A new approach to mapping mineral potential, In: *Statistical Applications in the Earth Sciences*, Agterberg, F.P., Bonham-Carter, G.F., (Ed.): 171-183, Geological Survey of Canada 98, Canadian Government Publishing Centre.
- Bonham-Carter, G.F. (1994). *Geographic Information Systems for Geoscientists: Modeling with GIS*. New York, Pergamon/Elsevier.
- Bui, D.T., Pradhan, B., Lofman, O., Revhaug, I. and Dick, O.B. (2012). Spatial prediction of landslide hazards in Hoa Binh province (Vietnam): a comparative assessment of the efficacy of evidential belief functions and fuzzy logic models. *Catena* 96: 28-40.
- Carranza, E.J.M. (2002). Geological-constrained mineral potential mapping (examples from the Philippines). International institute for Aerospace Survey and Earth Science. PhD Thesis.

- Carranza, E.J.M. (2004). Weights of evidence modeling of mineral potential: a case study using small number of prospects, Abra, Philippines. *Nat. Resour. Res.* 13:173-187.
- Carranza, E.J.M. (2009a). Geochemical anomaly and mineral prospectivity mapping in GIS. *Handbook of Exploration and Environmental Geochemistry*; Vol 11, Elsevier.
- Carranza, E.J.M. (2009b). Objective selection of suitable unit cell size in data-driven modeling of mineral prospectivity. *Comput. Geosci.* 35:2032-2046.
- Carranza, E.J.M. (2009c). Controls on mineral deposit occurrence inferred from analysis of their spatial pattern and spatial association with geological features. *Ore Geology Reviews* 35: 383–400.
- Carranza, E.J.M. (2011). Geocomputation of mineral exploration targets. *Comput. Geosci.* 37:1907-1916.
- Carranza, E.J.M. and Hale, M. (2000). Geologically constrained probabilistic mapping of gold potential, Baguio District, Philippines. *Nat. Resour. Res.* 9 :237-253.
- Carranza, E.J.M. and Hale, M. (2002a). Evidential belief functions for data-driven geologically constrained mapping of gold potential, Baguio district, Philippines. *Ore Geol. Rev.* 22:117-132.
- Carranza, E.J.M. and Hale, M. (2002b). Mineral mapping with Landsat Thematic Mapper data for hydrothermal alteration mapping in heavily vegetated terrain. *Int. J. Remote Sens.* 23 (22): 4827–4852.
- Carranza, E.J.M. and Sadeghi, M. (2010). Predictive mapping of prospectivity and quantitative estimation of undiscovered VMS deposits in Skellefte district (Sweden). *Ore Geology Reviews* 38: 219–241.
- Carranza, E.J.M., Hale, M. and Faassen, C. (2008). Selection of coherent deposit-type locations and their application in data-driven mineral prospectivity mapping. *Ore Geol. Rev.* 33:536-558.
- Carranza, E.J.M., Mangaoang, J.C. and Hale, M. (1999). Application of mineral exploration models and GIS to generate mineral potential maps as input for optimum land-use planning in the Philippines. *Natural Resources Research* 8(2): 165-173.
- Chang, K. (2010). *Introduction to geographic information systems*. McGraw-Hill International Edition.
- Cheng, Q. and Agterberg, F.P. (1999). Fuzzy weights of evidence and its application in mineral potential mapping. *Natural resources research*, 8 (1): 27-35.
- Chu, L.H. (1980). Preliminary report on geochemistry, Anomaly 3503, Kelantan. Geological Survey Malaysia Report (unpublished).

- Chu, L.H. (1983). Exploration in the Sok Base Metal Prospect, Kelantan. Geological Survey Malaysia Report (unpublished).
- Chu, L.H., Muntanion, H., Sidek, A., Chand, F. and Troup, A. (1983). Regional geochemistry of South Kelantan. Geological Survey Malaysia.
- Chu, L.H. (2000). Malaysia's mineral resource availability and potential. Prosiding Asia Pacific Mining and Quarrying. Kuala Lumpur.
- Chung, C.F. (1978). Computer program for the logistic model to estimate the probability of occurrence of discrete events. Geological Survey of Canada, Paper 78 (12).
- Chung, C.F. and Agterberg, F.P. (1980). Regression models for estimating mineral resources from geological map data. *Math. Geol.* 12:473-488.
- Chung, C.F. and Fabbri, A.G. (2003). Validation of spatial prediction models for landslide hazard mapping. *Nat. Hazards*, 30: 451-472.
- Clarke, K.C. (1986). Advances in geographic information systems, computers, environment and urban systems, 10: 175-184.
- Cohen, P.R. (1985). *Heuristic Reasoning About Uncertainty: An Artificial Intelligence Approach*, Pitman, London.
- Dempster, A.P. (1967). Upper and lower probabilities induced by a multivalued mapping. *Annals of Mathematical Statistics* 38(2): 325-339.
- Dempster, A.P. (1968). A generalization of Bayesian inference. *Journal of the Royal Statistical Society* 30(2): 205-247.
- Dickson, B.L., Fraser, S.L. and Kinsey-Henderson, A. (1996). Interpreting aerial gamma-ray surveys utilizing geomorphological and weathering models. *J. Geochem. Explor.*, 57: 75-88.
- Dodge, N. N. (1977). Mineral production on the east coast of Malaya in the Nineteenth Century. *Journal Malayan Branch Royal Asiatic Soc.* 50(2):89-108.
- Eddy, B.G., Bonham-Carter, G.F. and Jefferson, C.W. (1995). Mineral resource assessment of the Parry Islands, high Arctic, Canada: A GIS-base fuzzy logic model, Proceedings of Canadian Conference on GIS, CD ROM session C3, Paper 4, Ottawa.
- Emmons, J. (1933). On the mechanism of the deposition of certain metalliferous lode systems associated with granitic batoliths, Ore deposits of the Western United States. *Amer. Inst. Min. Metall. Petrol. Engineers, Lindgren Volume*: 327-439.
- Ernowo. and Oktaviani, P. (2010). Regional probabilistic of gold-silver potential mapping using likelihood ratio models in Flores Island. Proceeding of the 39th IAGI Annual Convention and Exhibition, Lombok.

- Evan, M.E. (1987). An introduction to ore geology. Blackwell scientific publications, Oxford.
- Evans, M.E. (1993). Ore Geology and Industrial Minerals; an Introduction, Blackwell Scientific Publications, Oxford.
- Franca-Rocha, W., Bonham-Carter, G. and Misi, A. (2003). GIS modeling for mineral potential mapping of carbonate-hosted pb-zn deposits. *Revista Brasileira de Geociências*, 33: 191-196.
- Galvão, L.S., Almeida-Filho, R. and Vitorello, I. (2005). Spectral discrimination of hydrothermally altered materials using ASTER short-wave infrared bands: evaluation in a tropical savannah environment. *Int. J. Appl. Earth Obs. Geoinformation* 7: 107–114.
- Gettings, M.E. and Bultman, M.W. (1993). Quantifying favorableness for occurrence of a mineral deposit type using fuzzy logic-an examples from Arizona. USGS Open-File Report 93-392.
- Geological Survey Malaysia. (1983). Airbone spectrometric and magnetic survey, Central Belt Project.
- Goh, S.H., Teh, G.H., Hassan, W.F.W. (2006). Gold mineralization and zonation in the State of Kelantan. *Geological Society of Malaysia Bulletin*, 52: 129-135.
- Groves, D.I., Goldfarb, R.J., Gebre_Mariam, M., Hagemann, S.G. and Robert, F. (1998). Orogenic gold deposits: a proposed classification in the context of their crustal distribution and relationships to other gold deposits types. *Ore Geology Reviews*, 13: 7-27.
- Hamzah, A. H. and Hamzah, M. (1989). Reconnaissance geochemical survey of the Tanah Merah area, Sheet 22, Kelantan. Laporan EMR 11/1989 Jabatan Penyiasatan Kajibumi.
- Harris, D.P. (1965). An application of multivariate statistical analysis to mineral exploration. Unpublished PhD. Dissertation, Pennsylvania State University.
- Harris, D.P. (1969). Alaska's base and precious metals resources: a probabilistic regional appraisal. *Quarterly of the Colorado School of Mines* 64: 295–327.
- Harris, J.R. (1989). Data integration for gold exploration in eastern Nova Scotia using a GIS. In: *Remote Sensing for Exploration Geology*, Proc. Calgary, Alberta: 233–249.
- Harris, J.R., Wilkinson, L. and Grunsky, E.C. (2000). Effective use and interpretation of lithogeochemical data in regional exploration programs. *Ore Geol. Rev.* 16:107-143.
- Harris, D.P. and Pan, G.C. (1999). Mineral favorability mapping: a comparison of artificial neural networks, logistic regression and discriminate analysis. *Natural Resources Research* 8: 93–109.

- Hassan, W. F. W., Almashoor, S. S., Tan, T. H., Mohamad, H., Wood, K. H., Hamzah, M. S. (1997). Kajian status potensi emas dalam batuan sedimen di Semenanjung Malaysia. Laporan akhir penyelidikan IRPA 4-07-03-025, tidak diterbitkan, Jabatan Geologi Universiti Kebangsaan Malaysia.
- Hassan, W.F.W. (2008). The mineralogy of gold mineralization of the Ajmal Mine, Kechau Tui, Pahang Darul Makmur. Bulletin of the Geological Society of Malaysia 54: 59-62.
- Hassan W.F.W.(2007). Khersonese emas geologi emas Semenanjung Malaysia. Universiti Kebangsaan Malaysia: 30-36.
- Henley, R.W. and Adams, D.P.M. (1992). Strike-slip fault reactivation as a control on epithermal vein-style gold mineralization. *Geology* 20: 443–446.
- Hutchison, C.S. (1975). Ophiolite in Southeast Asia. *Geological Society of America Bulletin*, 1975, 86: 797-806.
- Ho, S.E., Groves, D.I. and Bennett, J.M. (Eds.) (1990). Gold deposits of the Archaean Yilgarn Block, Western Australia: nature, genesis and exploration guides: Geology Department and University Extension, University of Western Australia Publication, 20.
- JMG (2012a). Malaysian Mineral Yearbook 2012.
- JMG (2012b). Malaysian Mining Industry 2012.
- Karimi, M., Menhaj, M.B. and Mesgari, M.S. (2008). Preparing mineral potential map using fuzzy logic in GIS environment. The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences. Vol. XXXVII. Part B8, Beijing.
- Kearey, P. and Brooks, M. (1984). An introduction to geophysical prospecting, 2nd Edn., Blackwell Scientific Publication, Oxford.
- Kendall, M.G. and Buckland, W.R. (1982). A dictionary of statistical terms, 4th Ed. Longmans, London.
- Knox-Robinson, C.M. (2000). Vectorial fuzzy logic: a novel technique for enhanced mineral prospectivity mapping, with reference to the orogenic gold mineralisation potential of the Kalgoorlie Terrane, Western Australia. *Aust. J. of Earth Sci.* 47:929-941.
- Kia, M.B., Pirasteh, S., Pradhan, B., Mahmud, A.R., Sulaiman, W.N.A. and Moradi, A. (2012). An artificial neural network model for flood simulation using GIS: Johor River Basin, Malaysia. *Environmental Earth Sciences* 67 (1): 251-264.
- Lee, S. (2009). Probabilistic integration of integrated gold mineral potential maps using GIS. Proceeding of 7th Symposium On Stepping Stones To The Future: Strategies, Architectures, Concepts and Technologies, Daejeon, Korea.

- Lee, S. (2011). Integration of mineral potential maps from various geospatial models. Proceedings of the 2nd International Conference on Computing for Geospatial Research & Applications, 52.
- Lee, S. and Pradhan, B. (2006). Probabilistic landslide hazards and risk mapping on Penang Island, Malaysia. *J. Earth Syst. Sci.*, 2006, 6: 1-12.
- Lee, S. and Pradhan, B. (2007). Landslide hazard mapping at Selangor, Malaysia using frequency ratio and logistic regression model. *Landslides*, 4: 33-41.
- Liu, Y., Xia, Q., Cheng, Q. and Wang, X. (2013). Application of singularity theory and logistic regression model for tungsten polymetallic potential mapping. *Nonlin. Processes Geophys.*, 20: 445-453.
- Low, V. F. (1921). Kelantan and its natural resources. *Mining magazine*. 24:11-20.
- Lowell, J.D. (1991). The discovery of the La Escondida Orebody. In: R.W. Hutchinson and R.I. Grauch (Eds.), *Historical perspective of genetic concepts and case history of famous discoveries*, Economic Geology Monograph 8, Economic Geology Publishing Company.
- Macdonald, S. (1967). Geology and mineral resources of North Kelantan and North Terengganu. *Geology Survey Malaysia District Memoir* 10.
- Manap, M.A., Nampak, H., Pradhan, B., Lee, S., Sulaiman, W.N.A. and Ramli, M.F. (2012). Application of probabilistic-based frequency ratio model in groundwater potential mapping using remote sensing data and GIS. *Arab J Geosci*. Doi: 10.1007/s12517-012-0795-z.
- McCuaig, T.C., Kreuzer, O.P. and Brown, W.M. (2007). Fooling ourselves-dealing with model uncertainty in a mineral systems approach to exploration. *Mineral Exploration and Research-Digging Deeper*. 9th Biennial SGA Meeting, Dublin, Proceedings: 1435-1438.
- McCuaig, T.C., Porwal, A. and Gessner, K. (2009). Fooling ourselves: recognizing uncertainty and bias in exploration targeting. *Centre for Exploration Targeting Quarterly News*, The University of Western Australia, 2(7), 1 and 5-8.
- Middlebrook, S. M. (1933). Pulai: An early Chinese settlement in Kelantan. *Journal Malayan Branch Royal Asiatic Soc.* 11(2):151-156.
- Moon, W.M. (1989). Integration of remote sensing and geological/geophysical data using Dempster-Shafer approach. *Proceedings of International Geoscience and Remote Sensing Symposium (IGARSS)*: 838-841.
- Moon, W.M. (1990). Integration of geophysical and geological data using evidential belief function. *IEEE Transactions on Geoscience and Remote Sensing* 28(4): 711-720.
- Moon, W.M., Chung, C.F. and An, P. (1991). Representation and Integration of geological, geophysical and remote sensing data. *Geoinformatics* 2:177-182.

- Nicolais, S.M. (1974). Mineral exploration with ERTS imagery: Third ERTS-1 Symposium, NASA SP-351 (1): 785–796.
- Oh, H.J. and Lee, S. (2011). Application of artificial neural network for gold–silver deposits potential mapping: A Case Study of Korea. *Nat. Resour. Res.* 19:103–124.
- Oh, H.J. and Lee, S. (2008). Regional probabilistic and statistical mineral potential, mapping of gold-silver deposits using GIS in the Gangreung area, Korea. *Resour. Geol.* 58:171–187.
- Pan, G.C. and Harris, D.P. (2000). Information synthesis for mineral exploration. Oxford University Press, New York.
- Park, C.F. and MacDiarmid, R. A. (1975). Ore deposits, 3rd Ed. Freeman, San Francisco.
- Partington, G.A. (2010). Exploration targeting using GIS: more than a digital light table. Geo Computing Conference, Brisbane, Australia.
- Pirajno, F. (1992). Hydrothermal mineral deposits, principles and fundamental concepts for the exploration geologists, Springer-Verlag, Berlin.
- Pour, A.B., Hashim, M. and Genderen, J.V. (2013). Detection of hydrothermal alteration zones in a tropical region using satellite remote sensing data: Bau goldfield, Sarawak, Malaysia *Ore Geology Reviews* 54:181–196.
- Pourghasemi, H.R., Pradhan, B., Gokceoglu, C., Mohammadi, M. and Moradi, H.R. (2013). Application of weights-of-evidence and certainty factor models and their comparison in landslide susceptibility mapping at Haraz watershed, Iran. *Arabian Journal of Geosciences* 6 (7): 2351–2365.
- Porwal, A., Carranza, E.J.M. and Hale, M. (2006). A hybrid fuzzy weights-of-evidence model for mineral potential mapping. *Natural Resources Research*, 15 (1): 1–14.
- Porwal, A. and Kreuzer, O.P. (2010). Introduction to the Special Issue: Mineral prospectivity analysis and quantitative. *Ore Geology Reviews* 38:121–127.
- Pradhan, B. (2013). A comparative study on the predictive ability of the decision tree, support vector machine and neuro-fuzzy models in landslide susceptibility mapping using GIS. *Comput. Geosci.*, 51: 350–365.
- Pradhan, B. (2011). Manifestation of an advanced fuzzy logic model coupled with Geo-information techniques to landslide susceptibility mapping and their comparison with logistic regression modelling. *Environmental and Ecological Statistics* 18 (3): 471–493.
- Pradhan, B. (2010). Landslide susceptibility mapping of a catchment area using frequency ratio, fuzzy logic and multivariate logistic regression approaches. *J. Indian Soc. Remote Sens.*, 38: 301–320.

- Pradhan, B. (2009). Groundwater potential zonation for basaltic watersheds using satellite remote sensing data and GIS techniques. *Central European Journal of Geoscience* 1(1):120–129.
- Pradhan, B. and Lee, S. (2010a). Landslide susceptibility assessment and factor effect analysis: backpropagation artificial neural networks and their comparison with frequency ratio and bivariate logistic regression modelling. *Environ. Modell Softw.*, 25: 747-759.
- Pradhan, B. and Lee, S. (2010b). Delineation of landslide hazard areas on Penang Island, Malaysia, by using frequency ratio, logistic regression, and artificial neural network models. *Environ Earth Sci* 60:1037–1054.
- Pradhan, B., Latif, Z.A. and Aman S.N.A. (2012). Application of airborne LiDAR-derived parameters and probabilistic-based frequency ratio model in landslide susceptibility mapping. *Applied Mechanics and Materials* 225: 442-447.
- Pradhan, B., Suliman, M.D.H. and Awang, M.A. (2007). Forest fire susceptibility and risk mapping using remote sensing and geographical information systems (GIS). *Disaster Prevention and Management* 16:344–352.
- Purwanto, H.S. (2002). Gold mineralisation and structural control in the Penjom area, Pahang and Lubuk Mandi area, Terengganu, Peninsular Malaysia. Unpublished PhD. Thesis, Universiti Kebangsaan Malaysia.
- Raines, G.L. (1999). Evaluation of weights of evidence to predict epithermal gold deposits in the Great Basin of the western United States: *Nat. Resour. Res.* 8:257-276.
- Reimann, C. and Filzmoser, P. (2000). Normal and lognormal data distribution in geochemistry: death of a myth. *Consequences for the statistical treatment of geochemical and Environmental Geology*, 39: 1001-1014.
- Rencz, A.N., Harris, J.R., Watson, G.P. and Murphy, B. (1994). Data integration for mineral exploration in the Antigonish Highlands, Nova Scotia. *Can. Jour. Remote Sensing* 20:258-267.
- Rose, A.W., Hawkes, H.E. and Webb, J.S. (1979). *Geochemistry in mineral exploration*, 2nd Ed., Academic Press, London.
- Rowan, L.C. and Wetlaufer, P.H. (1975). Iron-absorption band analysis for the discrimination of iron-rich zones. U.S. Geol. Surv. Type III Final Report, Contract S-70243-AG.
- Rowan, L.C. and Bowers, T.L. (1995). Analysis of linear features mapped in Landsat thematic mapper and side-looking airborne radar images of the Reno, Nevada 18 by 28 quadrangle, Nevada and California-implications for mineral resource studies. *Photogramm. Eng. Remote Sensing* 61: 749–759.
- Sabins, F.F. (1997). *Remote sensing-principles and interpretation*, 3rd edn., W.H. Freeman, New York, NY.

- Sabins, F.F. (1999). Remote sensing for mineral exploration. *Ore Geology Reviews* 14:157–183.
- Schowengerdt and Robert, A. (2007). Remote sensing: models and methods for image processing (3rd ed.). Academic Press. p.2.ISBN978-0-12-369407-2.
- Sentz, K., Ferson, S. and Laboratories, S.N. (2002). Combination of Evidence in Dempster–Shafer Theory, Citeseer.
- Shafer, G. (1976). A mathematical theory of evidence. Princeton University Press, Princeton, N.J.
- Sinclair, A.J. and Woodsworth, G.L. (1970). Multiple regression as a method of estimating exploration potential in an area near Terrace, B.C. *Economic Geology* 65: 998–1003.
- Singer, D.A. (1972). Multivariate statistical analysis of the unit regional value of mineral resources. Unpublished PhD. Dissertation, Pennsylvania State University.
- Silva, E.C., Silva, A.M., Toledo, C.L.B., Mol, A.G., Otterman, D.W. and Souza, S.R.C. (2013). Mineral potential mapping for orogenic gold deposits in the Rio Maria Granite Greenstone Terrane, Southeastern Pará State, Brazil.
- Singer, D.A. and Kouda, R. (1996). Application of a feed forward neural network in the search for Kuruko deposits in the Hokuroku district, Japan. *Mathematical Geology* 28(8): 1017-1023.
- Surip, N., Hamzah, A.H., Zakaria, M.R., Napiah, A. and Talib, J.A. (2007). Mapping of gold in densely vegetated area using remote sensing and GIS techniques in Pahang, Malaysia. *Proceeding of Asian Conference on Remote Sensing (ACRS)*, Kuala Lumpur.
- Stanley, C.R. (2006). Numerical transformation of geochemical data: 1. maximizing geochemical contrast to facilitate information extraction and improve data presentation. *Geochemistry-Exploration Environment Analysis* 6: 69-78.
- Taylor, R.B. and Steven, T.A. (1983). Definition of mineral resource potential: *Economic Geology*, 78(6): 1268-1270.
- Tehrany, M.S., Pradhan, B. and Jebur M.N. (2014). Flood susceptibility mapping using a novel ensemble weights-of-evidence and support vector machine models in GIS. *Journal of Hydrology*, 512: 332-343.
- Teoh, L.H., Lee, A.K and Foo, K.Y. (1987). Gold mineralization and prospects in Kelantan, Peninsular Malaysia. Geological Survey Malaysia Regional Mineral Exploration Project report.
- Teoh, L.H. and Hamzah, A.H. (1986). Follow-up investigation of the Bukit Kedah gold anomaly, Tanah Merah Kelantan. Laporan Eksplorasi Mineral EMR No. 04/1986. Laporan Jabatan Penyiasatan Kajibumi Malaysia.

- Turekian, K. K., Wedepohl, K. H. (1961). Distribution of the elements in some major units of the earth's crust. *Bull. Geol. Soc. Am.* 72:175-192.
- USGS. (2007). What is a GIS?
http://webgis.wr.usgs.gov/globalgis/tutorials/what_is_gis.htm. Retrieved 2 July 2010.
- Umar, Z., Pradhan, B., Ahmad, A., Jebur, M.N and Tehrany M.S. (2014). Earthquake induced landslide susceptibility mapping using an integrated ensemble frequency ratio and logistic regression models in West Sumatera Province, Indonesia. *Catena* 118:124-135.
- Vicente, L.E. and Filho, C.R.S. (2011). Identification of mineral components in tropical soils using reflectance spectroscopy and advanced spaceborne thermal emission and reflection radiometer (ASTER) data. *Remote Sens. Environ.* 115: 1824–183.
- Wally, P. (1987). Belief function representation of statistical evidence. *Annals Statistics* 15: 1439–1485.
- Wheatley, P. (1961). *The Golden Khersonese: studies in the historical geography of the Malay Peninsula before A. D. 1500*. University of Malaya Press.
- World Gold Council. (2013a). Gold mining.
<http://www.gold.org/gold-mining>. Retrieved on 12 September 2013.
- World Gold Council (2013b). Supply and demand.
<http://www.gold.org/supply-and-demand/gold-demand-trends>. Retrieved on 12 September 2013.
- Wright, D.F. and Bonham-Carter, G.F. (1996). VHMS favourability mapping with GIS based integration models, Chisel Lake-Anderson Lake area. In: G.F. Bonham-Carter, A.G. Galley and G.E.M. Hall (Eds.), *EXTECH I: A multidisciplinary approach to massive sulphide research in the Rusty Lake-Snow Lake Greenstone Belts, Manitoba*. Geological Survey of Canada, Bulletin 426: 339-376, 387-401.
- Yin, E.H. (1965). Provisional draft report on the geology and mineral resources of the Gua Musang area, Sheet 45, South Kelantan. Geological Survey of Malaysia.
- Yin, E.H. (1988). Geological map of Peninsular Malaysia, 8th edition, Geological Survey of Malaysia, Kuala Lumpur.
- Yousefi, M., Carranza, E.J.M. and Kamkar-Rouhani, A. (2013). Weighted drainage catchment basin mapping of geochemical anomalies using stream sediment data for mineral potential modeling.
- Zadeh, L. (1965). "Fuzzy sets", *Information and Control*, 8: 338–353.
- Zaw, K., Peters, S.G., Cromie, P., Burrett, C. and Hou, Z. (2007). Nature, diversity of deposit types and metallogenic relations of South China. *Ore Geology Reviews* 31: 3–47.

Ziaei, M., Carranza, E.J.M. and Ziaei, M. (2011). Application of geochemical zonality coefficients in mineral prospectivity mapping.

