



***EFFECTS OF A HOME-BASED PROGRAMME ON PHYSIOLOGICAL
VARIABLES, PHYSICAL ACTIVITY, PSYCHOLOGICAL DISTRESS
AND ENERGY INTAKE AMONG UNDERGRADUATE FEMALE
STUDENTS IN IRAQ***

JIAN ABDULLAH NOORI

FPP 2016 28



**EFFECTS OF A HOME-BASED PROGRAMME ON PHYSIOLOGICAL
VARIABLES, PHYSICAL ACTIVITY, PSYCHOLOGICAL DISTRESS
AND ENERGY INTAKE AMONG UNDERGRADUATE
FEMALE STUDENTS IN IRAQ**

By

JIAN ABDULLAH NOORI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

May 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



DEDICATED

To

The person who encouraged and supported me by all means

“Omar”



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Doctor of Philosophy

EFFECTS OF A HOME-BASED PROGRAMME ON PHYSIOLOGICAL VARIABLES, PHYSICAL ACTIVITY, PSYCHOLOGICAL DISTRESS AND ENERGY INTAKE AMONG UNDERGRADUATE FEMALE STUDENTS IN IRAQ

By

JIAN ABDULLAH NOORI

May 2016

Chairman : Associate Professor Soh Kim Geok, PhD
Faculty : Educational Studies

The study aimed to evaluate the effect of home-based programme to improve physiological variables, physical activity (PA), psychological distress (PD), and energy intake (EI) among undergraduate female students in Iraq. Besides, aimed to evaluate which variable was the most affected by the home based intervention programme. The participants were Iraqi sedentary undergraduate female students (N=44) who were assigned to two groups consisted of experimental group (N=22) and control group (N=22) respectively. The experimental group received a 12-week home based intervention programme which focuses on (PA and dietary awareness), while the control group were maintaining their usual life. Measurements for all the variables were taken prior to the intervention (pre-test), at week 6 (post-test1), and after 12 weeks (post-test2). Mixed between-within subjects analysis of variance was used to analyse physiological variables and PA, while, two way repeated measure of ANOVA was used to analyse PD and EI. In addition, Cohen's d was used to converting the Eta squared to evaluate which of the variables was more affected by the intervention programme.

The results of Mixed between-within subjects analysis of variance shows a statistically significant between the mean test scores in the pre-test, post-test1, and post-test2 measurements of certain physiological variables and PA in the experimental group. The variables were diastolic blood pressure: $F(2, 84) = 3.507, p < 0.05$, blood glucose: $F(2, 84) = .917, p < 0.05$, body fat: $F(2, 84) = 6.246, p < 0.05$, $VO_2\text{max}$: $F(2, 84) = 57.277, p < 0.05$, and vital capacity: $F(1.654, 69.485) = 16.839, p < 0.05$, and PA: $F(2, 84) = 731.01, p < 0.05$ respectively. Two way repeated measures of ANOVA shows a statistically significant between the mean test scores in the pre-test and post-test measurements of participants' PD and EI in the experimental group (i.e. total PD: $F(1,42) = 80.838, p < 0.05$, depression: $F(1,42) = 61.105, p < 0.05$, anxiety: $F(1,42) = 39.613, p < 0.05$, stress: $F(1,42) = 91.271, p < 0.05$, EI: $F(1,42) = 88.332, p < 0.05$) respectively.

Analysis of the post-test² scores of the experimental and control groups revealed that the difference between the mean scores of the two groups is statistically significant in certain physiological variables: diastolic blood pressure: $F(1, 42)=4.948$, $p<0.05$, blood glucose: $F(1,42)=8.783$, $p<0.05$, $VO_2\text{max}$: $F(1, 42)=52.034$, $p<0.005$, vital capacity: $F(1, 42)=14.047$, $p<0.05$, and PA $F(2, 42)=2260.951$, $p<0.05$ respectively. Meanwhile, the analysis of the post-test scores of the experimental and control groups revealed that the difference between the mean scores of the two groups is statistically significant in PD: total PD: $F(1,42)=47.452$, $p<0.05$, depression: $F(1, 42)=27.722$, $p<0.05$, anxiety: $F(1, 42)= 33.156$, $p<0.05$, stress: $F(1, 42)=37,310$, <0.05), and EI ($F(1, 42)= 40.289$, $p<0.05$) respectively. Finally, the Eta squared showed that certain physiological variables were the more effected by the intervention programme blood cholesterol=18.147 and $VO_2\text{max}=4.467$ follow by PA= 18.147, then EI =3.701. While PD =2.774 was the lowest effected by the home based intervention programme.

This study provided evidence that home-based intervention programme which focused on PA and dietary awareness had a significant effect on improving physiological variables, PA, PD, and EI. However, to achieve these benefits, exercise and dietary programme need to be carried out regularly as a life routine.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**KESAN PROGRAM BERASA DI RUMAH TERHADAP PEMBOLEH UBAH
PSIKOLOGIKAL, AKTIVITI FIZIKAL, DISTRES PIKOLOGIKAL DAN
DAPATAN TENAGA DALAM KALANGAN PELAJAR PEREMPUAN DI IRAQ**

Oleh

JIAN ABDULLAH NOORI

Mei 2016

Pengerusi : Profesor Madya Soh Kim Geok, PhD
Fakulti : Pengajian Pendidikan

Kajian ini bertujuan untuk menilai keberkesanan program intervensi berasas di rumah untuk meningkatkan pemboleh ubah fisiologi, aktiviti fizikal (PA), distress psikologikal (PD) dan daoatan tenaga (EI) dalam kalangan pelajar perempuan di Iraq. Selain itu, bertujuan untuk menilai pemboleh ubah yang paling dipengaruhi oleh program intervensi berasas di rumah. Peserta adalah pelajar perempuan yang mengamalkan gaya hidup sendentari di Iraq (N = 44) yang dibahagikan kepada kumpulan eksperimen (N=22) dan kumpulan kawalan (N=22). Kumpulan eksperimen mengikuti program intervensi berasas di rumah selama 12 minggu berfokus kepada (aktiviti fizikal dan kesedaran pemakanan), manakala kumpulan kawalan mengamalkan gaya hidup yang normal. Pengukuran terhadap pemboleh ubah dilakukan sebelum intervensi (Ujian pra), minggu ke 6 (Ujian pasca 1) dan selepas minggu ke 12 (Ujian pasca 2). Mixed Between-within Subjects Analysis of Variance digunakan untuk menganalisis pemboleh ubah fisiologi dan PA, manakala Two Way Repeated Measure of ANOVA digunakan untuk menganalisis PD dan EI. Selain itu, Cohen d digunakan untuk menukarkan kuasa dua Eta bagi menilai pemboleh ubah yang lebih dipengaruhi oleh program intervensi.

Keputusan analisis Mixed Between-within Subjects Analysis of Variance menunjukkan terdapat keputusan yang signifikan secara statistik di antara skor ujian min dalam Ujian pra, Ujian pasca 1 dan Ujian pasca 2 bagi pemboleh ubah fisiologi tertentu dan aktiviti fizikal dalam kumpulan eksperimen. Pemboleh ubah-pemboleh ubah berkenaan adalah tekanan darah diastolik : $F(2, 84) = 3.507, p < 0.05$, glukosa darah : $F(2, 84) = .917, p < 0.05$, lemak badan : $F(2, 84) = 6.246, p < 0.05$, $VO_2\text{max}$: $F(2, 84) = 57.277, p < 0.05$, dan vital capacity : $F(1.654, 69.485) = 16.839, p < 0.05$, dan PA : $F(2, 84) = 731.01, p < 0.05$. Two way repeated measures of ANOVA menunjukkan terdapat signifikan secara statistik min skor Ujian pra dan Ujian pasca PD dan EI peserta kumpulan eksperimen (iaitu jumlah PD : $F(1,42) = 80.838, p < 0.05$, kemurungan : $F(1,42) = 61.105, p < 0.05$, kebimbangan : $F(1,42) = 39.613, p < 0.05$, tekanan : $F(1,42) = 91.271, p < 0.05$, EI : $(1,42) = 88.332, p < 0.05$).

Analisis skor Ujian pasca 2 bagi kumpulan eksperimen dan kumpulan kawalan menunjukkan terdapat perbezaan statistik yang signifikan skor min kedua-dua kumpulan terhadap pemboleh ubah fisiologi yang tertentu : tekanan darah diastolik: $F(1, 42) = 4.948$, $p < 0.05$, glukosa darah: $F(1, 42) = 8.783$, $p < 0.05$, $VO_2\text{max}$: $F(1, 42) = 52.034$, $p < 0.05$, kapasiti vital: $F(1, 42) = 14.047$, $p < 0.05$ dan PA: $F(2, 42) = 2260.951$, $p < 0.05$. Manakala, analisis skor Ujian pasca kumpulan eksperimen dan kumpulan kawalan menunjukkan bahawa terdapat perbezaan yang signifikan di antara skor min PD bagi kedua-dua kumpulan (PD keseluruhan: $F(1, 42) = 47.452$, $p < 0.05$, kemurungan: $F(1, 42) = 27.722$, $p < 0.05$, kebimbangan: $F(1, 42) = 33.156$, $p < 0.05$, tekanan: $F(1, 42) = 37.310$, $p < 0.05$ dan EI ($F(1, 42) = 40.289$, $p < 0.05$). Akhir sekali, kuasa dua Eta menunjukkan bahawa pemboleh ubah fisiologi yang tertentu adalah lebih dipengaruhi oleh program intervensi bagi kolesterol dalam darah = 18.147 dan $VO_2\text{max}$ = 4.467 diikuti oleh PA = 18.147 dan seterusnya EI = 3.701. Manakala, PD (2.774) adalah kurang dipengaruhi oleh program intervensi berasas di rumah.

Kajian ini menunjukkan program intervensi berasas di rumah mempunyai kesan signifikan untuk meningkatkan pemboleh ubah fisiologi, PA, PD dan EI. Walau bagaimanapun, manfaat ini boleh dicapai dengan senaman dan program pemakanan yang perlu dilakukan secara kerap sebagai rutin kehidupan.

ACKNOWLEDGEMENTS

I would like to take this opportunity to express my gratefulness to the most high ALLAH (sw), who gives me strength, courage, inspiration, and love to be able to go through all the days of my life and also afford me great understanding and wisdom to complete my thesis, all the glory of his name.

First of all, I would like to acknowledgment the sincerest gratitude to my supervisor Associate Professor Dr. Soh Kim Geok for her care, wisdom, careful supervision and pleasant support during this investigation. I appreciate her invaluable guidance and the precious time she spend on reading the draft of the whole work. She always support me when I need it the most in academic as well as research. For this, I am truly grateful.

I wish to extend deepest and the most profound gratitude to my committee members, Associated Professor Dr. Norhaizan Mohd. Esa, Associated Professor Dr. Rohani Ahmad Tarmizi, Professor Dr. Nabeel Abdulwahab Ahmed for their professional guidance and valuable advice, which have contributed to my success of this study.

Of course, I would not be where I am today if it were not for my family, I would like to express my appreciation to them of their continual support. I am especially grateful to my husband "Omar" and my children "Aska" and "Yahya" for their continuing love, support, and understanding throughout my academic career.

My sincere appreciation to all the participants involved in this experimental study. Without their cooperation, this research could not have been possible. My utmost appreciation goes to the Ministry of Higher Education and Scientific Research-Iraq, the dean, the lecturers, and all freshmen female students in the Faculty of Education- Soran University in Erbil, and the jury experts in the field. My deepest appreciation also goes to my friends Dr. Arporn Popa, Dr. Chun Cheng Chuan, Dr. Zeinab Ghiami, for their concern, encouragement, and continuous support during the wonderful four years in UPM.

Last but not least, thanks to those individuals whose name not mentioned, but certainly they have a place in my heart; surly ALLAH (sw) will reward them.

I certify that a Thesis Examination Committee has met on 24 May 2016 to conduct the final examination of Jian Abdullah Noori on her thesis entitled "Effects of a Home-Based Programme on Physiological Variables, Physical Activity, Psychological Distress and Energy Intake among Undergraduate Female Students in Iraq" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Aminuddin bin Yusof, PhD

Professor
Faculty of Educational Studies
Universiti Putra Malaysia
(Chairman)

Mohd Nasir bin Mohd Taib, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Tengku Fadilah binti Tengku Kamalden, PhD

Senior Lecturer
Faculty of Educational Studies
Universiti Putra Malaysia
(Internal Examiner)

Lateef Oluwole Amusa, PhD

Professor
University of Venda
South Africa
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 23 August 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Soh Kim Geok, PhD

Associate Professor
Faculty of Educational Studies
Universiti Putra Malaysia
(Chairman)

Norhaizan Mohd. Esa, PhD

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Rohani Ahmad Tarmizi, PhD

Associate Professor
Institute for Mathematical Research
Faculty of Educational Studies
Universiti Putra Malaysia
(Member)

Nabeel Abdulwahab Ahmed, PhD

Professor
Faculty of Basic Education
Al-Mustansiriya University- Iraq
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software

Signature: _____ Date: _____

Name and Matric No.: Jian Abdullah Noori (GS33395)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____

Name of Chairman
of Supervisory
Committee:

Associate Professor Dr. Soh Kim Geok

Signature: _____

Name of Member
of Supervisory
Committee:

Associate Professor Dr. Norhaizan Mohd. Esa

Signature: _____

Name of Member
of Supervisory
Committee:

Associate Professor Dr. Rohani Ahmad Tarmizi

Signature: _____

Name of Member
of Supervisory
Committee:

Professor Dr. Nabeel Abdulwahab Ahmed

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xviii
CHAPTER	
1 INTRODUCTION	1
1.1 Overview	1
1.2 Background of the Study	1
1.3 Problem Statement	4
1.4 Objectives	5
1.5 Hypotheses	6
1.5.1 Hypothesis 1-8 for Objective 1	6
1.5.2 Hypothesis 9 for Objective 2	6
1.5.3 Hypothesis 10-13 for Objective 3	7
1.5.4 Hypothesis 14 for Objective 4	7
1.5.5 Research Question for Objective 5	7
1.6 Significance of the Study	7
1.7 Delimitation of the Study	8
1.8 Limitation of the Study	8
1.9 Definition of the Terms	9
1.9.1 Physiological Variables	9
1.9.2 Physical Activity (PA)	9
1.9.3 Psychological distress	9
1.9.4 Energy Intake	10
1.9.5 Iraqi Undergraduate Level	10
1.9.6 Home Based Intervention Programme	10
2 LITERATURE REVIEW	11
2.1 Overview	11
2.2 Introduction	11
2.3 Physical Activity	11
2.4 Energy Intake	13
2.5 Benefits of physical activity and dietary intervention programme	14
2.6 The benefits of intervention programme on physiological variables	16
2.7 The benefits of intervention programme on physical activity	18
2.8 The benefits of intervention programme on psychological distress	19
2.9 The benefits of intervention programme on Energy intake	21
2.10 Conclusion of literature review	22

3	METHODOLOGY	24
3.1	Overview	24
3.2	Research Design	24
3.3	Research Framework	25
3.4	Population and Sampling Procedures	27
3.5	Instrumentation	28
3.5.1	Physiological Variables Practical Tests	28
3.5.2	Practical Tests for Measuring Physical Activity Variable	32
3.5.3	Psychological Distress Rating Depression Anxiety Stress Scales (DASS) Questionnaire	32
3.5.4	24-Hour Dietary Record	33
3.6	Validity	33
3.7	Pilot Study	33
3.8	Reliability	34
3.9	The Intervention Programme	34
3.10	Research Procedure	41
3.11	Data Analysis	42
4	DATA ANALYSIS	44
4.1	Overview	44
4.2	Descriptive Statistics	47
4.3	Inferential Statistics	51
4.3.1	Null hypothesis 1	52
4.3.2	Null hypothesis 2	56
4.3.3	Null hypothesis 3	58
4.3.4	Null hypothesis 4	61
4.3.5	Null hypothesis 5	63
4.3.6	Null hypothesis 6	66
4.3.7	Null hypothesis 7	68
4.3.8	Null hypothesis 8	70
4.3.9	Null hypothesis 9	73
4.3.10	Null hypothesis 10	75
4.3.11	Null hypothesis 11	77
4.3.12	Null hypothesis 12	79
4.3.13	Null hypothesis 13	81
4.3.14	Null hypothesis 14	83
4.3.15	Research Question	85
5	DISCUSSIONS, CONCLUSIONS, IMPLICATION, AND RECOMMENDATIONS FOR FUTER RESULTS	87
5.1	Overview	87
5.2	Discussion	87
5.2.1	Discussions on Physiological variables	87
5.2.2	Discussions on Physical Activity	93
5.2.3	Discussions on Psychological Distress	95
5.2.4	Discussions on energy intake	96
5.2.5	Discussions on the effectiveness of the intervention programme	97
5.3	Conclusion	99

5.4	Implications	100
5.5	Recommendations for Future Research	101
REFERENCES		102
APPENDICES		123
BIODATA OF STUDENT		179
LIST OF CONFERANCES		180



LIST OF TABLES

Table		Page
3.1	Research design	25
3.2	G-power method for finding total sample size	28
3.3	Reliability statistics for research variables	34
3.4a	The intervention schedule for the physical activity programme	36
3.4b	The intervention schedule for the dietary awareness programme	40
3.5	The analysis of the hypotheses and research question of the study	42
4.1	Independent sample test of pre-test	45
4.2	Normality test for the dependent variables of the study	46
4.3	Levene's test of equality of error variances between two groups based on mean	47
4.4	Frequency and percentage distribution by groups.	48
4.5	Descriptive statistic for the students' demographic variables in both experimental and control groups	49
4.6	Descriptive statistics score for physiological variables and physical activity for experimental and control group	50
4.7	Descriptive statistics score for psychological distress and energy intake for experimental and control group	51
4.8	Levene's test of equality of error variances	52
4.9	Dia. blood pressure mean difference between experimental and control groups in pre, post1 and post-test2	53
4.10	Sys. blood pressure mean difference between experimental and control groups in pre, post1 and post-test2	53
4.11	The difference of Dia. blood pressure mean scores between tests in experimental and control groups	54
4.12	The difference of Sys. Blood pressure mean scores between tests in experimental and control groups	54
4.13	Levene's test of equality of error variances	57

4.14	Blood cholesterol mean difference between experimental and control groups in pre-test, post-test1 and post-test2	57
4.15	The difference of B.Ch. mean scores between tests in Experimental and Control Groups	58
4.16	Levene's test of equality of error variances	59
4.17	Blood glucose mean difference between experimental and control groups in pre-test, post-test1 and post-test2	60
4.18	The difference of blood glucose between tests in experimental and control groups	60
4.19	Levene's test of equality of error variances	61
4.20	BMI mean difference between experimental and control groups in pre, post-test1 and post-test2	62
4.21	The difference of BMI mean scores between tests in experimental and control groups	63
4.22	Levene's test of equality of error variances	64
4.23	WHR mean difference between experimental and control groups in pre, post-test1 and post-test2	64
4.24	The difference of WHR scores between tests in experimental and control groups	65
4.25	Levene's test of equality of error variances	66
4.26	BF mean difference between experimental and control groups in pre, post-test1 and post-test2	67
4.27	The difference of BF scores between tests in experimental and control groups	67
4.28	Levene's Test of Equality of Error Variances	68
4.29	VO ₂ max mean difference between experimental and control groups in pre, post-test1 and post-test2	69
4.30	The difference of VO ₂ max scores between tests in experimental and control groups	70
4.31	Levene's test of equality of error variances	71
4.32	V.C. mean difference between experimental and control groups in pre, post-test1 and post-test2	71

4.33	The difference of V.C. scores between tests in experimental and control groups	72
4.34	Levene's test of equality of error variances	73
4.35	Physical activity difference between experimental and control groups in pre-test, post-test1 and post-test2	74
4.36	The difference of physical activity scores between tests in experimental and control groups	74
4.37	Levene's test of equality of error variances	76
4.38	Psychological distress mean difference between experimental and control groups in pre-test and post-test	76
4.39	The difference of psychological distress scores between tests in experimental and control groups	77
4.40	Levene's test of equality of error variances	78
4.41	Depression mean difference between experimental and control groups in pre-test and post-test	78
4.42	The difference of depression scores between tests in experimental and control groups	79
4.43	Levene's test of equality of error variances	80
4.44	Anxiety mean difference between experimental and control groups in pre-test and post-test	80
4.45	The difference of anxiety scores between tests in experimental and control groups	81
4.46	Levene's test of equality of error variances	82
4.47	Stress mean difference between experimental and control groups in pre-test and post-test	82
4.48	The difference of stress scores between tests in experimental and control groups	83
4.49	Levene's test of equality of error variances	84
4.50	Energy intake mean difference between experimental and control groups in pre-test and post-test	84
4.51	The difference of energy intake scores between tests in Experimental and Control Groups	85



LIST OF FIGURES

Figure		Page
3.1	Research Framework	26
3.2	BMI equation	30
4.1	Mean Plot of Dia. Blood Pressure in Both Experimental (1) and Control (2) Groups	55
4.2	Mean Plot of Sys. Blood Pressure in Both Experimental (1) and Control (2) Groups	56
4.3	Mean plot of B.Ch. in Both Experimental (1) and Control (2) Groups	58
4.4	Mean Plot of B.G. in Both Experimental (1) and Control (2) Groups	61
4.5	Mean Plot of BMI in Both Experimental (1) and Control (2) Groups	63
4.6	Mean Plot of WHR in Both Experimental (1) and Control (2) Groups	65
4.7	Mean Plot of BF in Both Experimental (1) and Control (2) Groups	68
4.8	Mean plot of VO ₂ max in both experimental (1) and control (2) groups	70
4.9	Mean plot of V.C. in both experimental (1) and control (2) groups	72
4.10	Mean Plot of physical activity score in both experimental (1) and control (2) groups	75
4.11	Mean plot of psychological distress in both experimental (1) and control (2) groups	77
4.12	Mean plot of depression in both experimental (1) and control (2) groups	79
4.13	Mean plot of anxiety in both experimental (1) and control (2) groups	81
4.14	Mean plot of stress in both experimental (1) and control (2) groups	83
4.15	Mean plot of energy intake in both experimental (1) and control (2) groups	85

LIST OF ABBREVIATIONS

WHO	World Health Organization
UN	United nation
MOHERS	Ministry of Higher Education and Scientific Research
BP	Blood Pressure
DIA.BP	Diastolic Blood Pressure
SYS.BP	Systolic Blood Pressure
BCH	Blood Cholesterol
BG	Blood Glucose
BMI	Body Mass Index
BF	Body Fat
WHR	Waist Hip Ratio
VO ₂ max	Maximal Oxygen Consumption
VC	Vital Capacity
PA	Physical Activity
PD	Psychological Distress
DEP.	Depression
ANX	Anxiety
STR	Stress
DASSQ	Depression, Anxiety, and Stress Questionnaire
EI	Energy Intake

CHAPTER 1

INTRODUCTION

1.1 Overview

The first chapter introduces the background and the nature of the present study. In particular, the role of physical activity (PA) and dietary knowledge related to the condition of Iraqis' daily life and its effect on females, especially on those who are attending Iraqi universities are presented. The chapter starts with a detailed description of obstacles and restrictions in Iraqi society in the light of the current situation and their impact on the decline in PA and dietary awareness in the stage of university, which leads to the establishment of an active and healthy society. Therefore, the problem statement, significance, limitation and delimitation of the study are also presented. The last part of this chapter presents the operational definitions of key terms used in this study.

1.2 Background of the Study

Iraq is going through a critical period in the present time in which terrorist attacks continue to occur on an almost daily basis, which have directly affected most aspects of life around the country. This situation has led to further deterioration of the security situation, especially in the capital Baghdad, through repeated security breaches of terrorist bombings, improvised explosive devices and car bombs and assassinations that have continued until the present time and the majority of these attacks were directed at civilians (Habib, 2007). Women and other vulnerable groups in Iraq still suffer from discrimination, economic and social barriers, and the attacks against them. According to the latest report by the United Nation (UN) in 2013, the human rights situation in Iraq, despite some progress, is threatened by the rise of armed violence, which has killed at least 3,238 civilians and injured 10,379 others in 2012 (UN, 2013). While the number of civilians wounded or killed reached to 55,856 during the years 2014 to 2015 (Kubiš & Al Hussein, 2016).

All of these conditions have directly influenced the lifestyle and daily behaviour of the people. The unsuitable conditions which Iraq is currently facing have made it difficult to provide the most basic means to improve the quality of life in general. Especially, women are not able to enjoy their freedom, economical independence and social activity. They are not only under oppression from the government, but they are also being oppressed and discriminated against by the community and men in particular (Madi, 2007). The new civil constitution of Iraq still lacks the women's basic rights to enhance the poor state being currently faced by Iraqi women. They are not only confined to the house, and kitchen and cloak, but they are also trapped mostly by male terrorists when they are out of the house. Most Iraqi women must suffer from great ignorance and misconceptions due to religious beliefs and the law by males against their freedom. For example, there is a fear of taking part in sports activities among young women since they believe that physical exercise would turn their bodies into more masculine bodies, and the society would view them as more masculine individuals (Habib, 2007). All this shows an actual decline in the role of

women in society, cultural events, sports, and intellectual life as well as the positive impact of this involvement. This concern is confirmed by the World Health Organization (WHO)'s reports indicating that the percentage of physically inactive and overweight females in Iraq is 51.3% and 65.1%, respectively (Al-Tamimi, Armstrong, Cowan, & Riley, 2011).

School system at the forefront of educational institutions has also deteriorated. Approximately 80% of schools in the capital city of Baghdad are neglected by the authorities (ARFAD, 2008). Higher Education offered in universities and colleges has also faced the same fate. Iraqi universities in the current stage suffer from the lack of development, still following the traditional rules which does not fit the principles of modern education and theories. Especially after the change of the former regime in April, 2003, due to the destruction of the most educational institutions, the poor infrastructure of universities negatively impact the involvement of women in PA and sports (Al-Haidari, 2011).

Lifestyle and behavioural factors such as daily PA play a key role in the prevention of chronic diseases including cardiovascular disease, diabetes, and obesity (Cooper & Hancock, 2011; Hardin, Hebert, Bayden, Dehart, & Mazur, 1997; Rowlands, Eston, & Ingledew, 1999; Strong et al., 2005). PA promotes many values such as self-confidence and community spirit, communication and integration, discipline and respect, in addition to the psychological gains and disposal of depression and improving concentration (Al-Tamimi, 2007). Hence, being physically active plays an essential role in increasing health and well-being. A large number of researchers investigated the benefits of PA on many parts of the body such as heart, skeletal muscles, bones, blood (i.e. cholesterol levels), the immune system and the nervous system (Cooper & Hancock, 2011; Soroush et al., 2013; Stefanick et al., 1998; Stone, McKenzie, Welk, & Booth, 1998; Strong et al., 2005). However, to achieve all of these benefits, specific guidelines were recommended to improve physiological variables, PA, PD, and EI for adults starting from the minimum amount of PA and increasing dietary knowledge (Agriculture & Services, 2010; P. A. G. A. Committee, 2008b).

Specifically, the 2008 PA Guidelines recommend 150 minutes of moderate and/or 75 minutes of vigorous PA per week to reduce the risk of obesity, cardiovascular diseases (CVD) and type 2 diabetes (Hamilton, Healy, Dunstan, Zderic, & Owen, 2008). Moreover, moderate intensity PA includes walking which is a popular, accessible and acceptable form of activity, particularly among populations who are the most physically inactive. Walking programmes are considered a safe and effective way to increase the physical fitness. Also, walking does not require special equipment and has a low risk of injuries. Furthermore, relevant research found that increased walking leads to increased fitness, decreased body weight, body mass index, and percentage of body fat and resting diastolic blood pressure in previously sedentary adults (P. A. G. A. Committee, 2008b; Cooper & Hancock, 2011; Kassavou, Turner, & French, 2013). Fitness is important in daily life, and low levels of PA and less physical fitness are independent risk factors for chronic diseases and premature mortality among adults (Katzmarzyk, 2010). Therefore, introducing walking as an important and easily accessible fitness activity to the

individuals' daily activities seems to be necessary and helpful in improving the overall fitness and health of an individual.

There are considerable published data to strongly support the benefits of the intervention programmes which consist of PA and dietary changes as a means to decrease the morbidity and mortality from cardiovascular disease and stroke in adults (Artinian et al., 2010; Blumenthal et al., 2010; T. Brown & Summerbell, 2009; Burke et al., 2013; Butler, Black, Blue, & Gretebeck, 2004). Moreover, there is a strong evidence that obesity is related to energy content of diet and an increasingly sedentary lifestyle because obesity in childhood is known to be an independent risk factor for adult obesity (Flegal, Carroll, Ogden, & Curtin, 2010; Ogden & Carroll, 2010). Therefore, there is a need to develop interventions to reduce the prevalence of obesity among the youth. These interventions should focus on changing these behaviours. The effectiveness of interventions that focus on improving diet and PA on the individuals' health has been shown in different studies (T. Brown & Summerbell, 2009; Coghill & Cooper, 2008; Collins et al., 2011; Danielsen, Svendsen, Mæhlum, & Sundgot-Borgen, 2013). However, there is still a need for further investigations to determine the effects of such interventions on the physical and PD of the individuals, especially among young people in Iraq.

It has become well acceptable that the type of environment and lifestyle of individuals have a strong influence on health. New evidence indicates that good dietary elements have a strong impact on the health of the body and mind. In particular, insufficient levels of dietary intake which is typical of our modern society can be considered as risk factors for several modern diseases and various mental diseases such as Alzheimer's disease, as well as psychiatric disorders such as depression (Gomez-Pinilla, 2011). Many researchers have shown that inactive lifestyle with unhealthy eating habits leads to overweight, obesity, and health problems (Bauer, Neumark-Sztainer, Fulkerson, Hannan, & Story, 2011; Blair, Kohl, & Barlow, 1993; Boyle, Jones, & Walters, 2010; Cecchini et al., 2010). According to Al-Tamimi (2007) the World Health Organization (WHO) indicated that the proportion of death from non-communicable diseases is 60% of the total death. The most important reasons are unhealthy and proper diets and lack of exercise.

Non-communicable diseases related with nutrition cause morbidity and mortality in most countries in the Eastern Mediterranean region, specifically cardiovascular disease, diabetes, and cancer. The key risk factors for non-communicable diseases include high blood pressure, high concentration of serum cholesterol, tobacco smoking, unhealthy eating habits, overweight or obesity, and physical inactivity (Musaiger & Al-Hazzaa, 2012). These facts clearly demonstrate the lack of dietary awareness of the majority of people which is reflected negatively on their EI in the daily life and shows that they are unlikely to follow healthy eating habits. Hence, intervention programme including dietary awareness along with PA was found to be more beneficial to prevent diseases and maintain life (Daubenmier et al., 2007; G. A. Kelley, Kelley, Roberts, & Haskell, 2012a, 2012b; Söderlund, Fischer, & Johansson, 2009).

1.3 Problem Statement

Physical inactivity and lack of dietary awareness have been identified as important public health concerns for the youth. According to WHO's reports, the lack of exercise and inadequate good diet cause an increase in the rate of global deaths from non-infectious diseases by an estimated 60% (Al-Tamimi, 2007). Poor physical inactivity and diet cause 310,000 to 580,000 deaths per year and are the major contributors to disabilities that result from diabetes, osteoporosis, obesity, and stroke (Pribis, Burtnack, McKenzie, & Thayer, 2010). This phenomenon has highlighted the importance of PA and dietary awareness in our daily life behaviour and how the lack of PA and dietary awareness can be dangerous, especially to youth life. Accordingly, Mirkin (2010) indicated that Iraq has approximately 50% of the population under the age of 19. However, it is difficult for people in this country to keep healthy and live actively due to poor security conditions and the instability of safety

In addition, Humairi (2015) indicated the low level of nutrition and dietary awareness for Iraqi rural women with an average of 19.3 equivalent to 55%. Blood pressure, blood glucose and obesity rates among females aged 15 and above in Iraq are higher compared to their counterparts in the Eastern Mediterranean Region that (12.5 vs. 11.6 in BP, 28.7 vs. 29.1 in BG, and 36.2 vs. 26.5 in obesity) due to inactive lifestyle and bad EI (GHO, 2013). All those problems showed the need to increase dietary awareness and female knowledge to help prevent diseases as reported by WHO on Iraqi health profile.

Different studies conducted on PA among adolescents and adults show a significant decline in the rate of PA in adolescence (Caspersen, Pereira, & Curran, 2000; Malina, 2001). This rate continues to decline throughout adulthood (Dwyer, Wilson, Limarzi, Callaghan, & Croskery, 2013; Malina, 2001). Furthermore, research has shown that the level of PA has declined among college students in recent years. It was reported that up to 50% of college students are not physically active at the recommended levels (Jackson & Howton, 2008; Leslie, Fotheringham, Owen, & Bauman, 2001). Al Subaie (2005) revealed a low percentage of students (5.8%) involved in physical activities in Saudi Arabia during the period of university study. In tracking PA participation in the early college years, Racette, Deusinger, Strube, Highstein, and Deusinger (2005) found that 30% of students did not do physical exercise during their freshman year.

Ismail and Shihab (1994) showed the negative trends and misconceptions among educators, teaching staff, and management in schools in Iraq, who consider PA as an outsider subject in education. In addition, the evidence indicates that physical education is not properly acknowledged in education system due to the lack of interest in physical education lessons, lack of facilities and equipment, insufficient time allocated to physical education lessons, and lack of interest in school administration to support PA.

Despite the significant number of evidence regarding the benefits of being physically active and improving students physiologically, physically and psychologically, there is no physical education course in the curriculum of non-sports colleges in Iraqi universities. This problem is faced by college students to engage in PA during their undergraduate

years. Moreover, the lack of adequate encouragement that students receive from faculty members, the congestion of university lectures, and the lack of credits for PA in the educational calendar of universities, are the factors leading students' poor participation in sports activities in the university. Furthermore, the unsecure society (e.g. students to do exercises at home with lack of time and poor application of the lessons on physical education) is the key factor that the very few students receive proper physical and dietary information.

Meanwhile, parents do not usually encourage their children, especially their daughters to engage in sports teams and social clubs. Some parents do not allow their daughters to participate in sports activities due to the security situation which could endanger the lives of their children. This unwillingness is heightened by the wrong society's perception of sports, especially for girls as pointed out by a representative of the Olympic Committee in the province of Nineveh who said: *"Social conditions and the reluctance of families on the participation of their daughters in sports and increasing number of girls who wear hijab because of customs and traditions, formed all the straw that broke the women's sport"* explaining to the UNAMI (United Nations Assistance Mission for Iraq) in 2013. In addition, in several reports, UNAMI indicated that in terms of society's perception, women's sports involvement is an indecent phenomenon in the society (Aoda, 2013).

The above mentioned issues are the reasons behind the manifestation of potentially inactive and uneducated healthy generation of young people. Phenomenon of inactivity and nutrition illiteracy is prevalent, especially among female members of the society, who are prone to various physiological, physical, and mental problems as the consequence of not engaging in any type of PA or sports and not receiving sufficient dietary awareness. Therefore, there seems to be an urgent need to design a home-based intervention programme to help the sedentary young Iraqi female students improve their health physiologically, physically, and mentally as well as their EI. This intervention should include physical activities, simple exercises, and dietary awareness tailored for these inactive young women, which can be carried out easily at home where most of these students spend a great deal of their free time. In addition, it is important to know which health indicators (i.e. physiologically, physically, mentally or EI) is mostly affected by this intervention programme. Hopefully, this study will help to reduce obesity, cardiovascular diseases, and diabetes as well as increase physically active individuals and improve health in the society.

1.4 Objectives

The main objective for this research is to investigate the effect of a home-based intervention programme on the physiological variables, PA, PD and EI among undergraduate female students in Iraq. The specific objectives are presented as follows:

1. To investigate the effect of the 12 week home based intervention programme on the physiological variables (blood pressure, blood cholesterol, blood glucose, body mass index, waist hip ratio, body fat, VO₂max and vital capacity) among undergraduate female students in Iraq.

2. To investigate the effect of the 12 week home based intervention programme on the physical activity among undergraduate female students in Iraq.
3. To investigate the effect of the 12 week home based intervention programme on the psychological distress among undergraduate female students in Iraq.
4. To investigate the effect of the 12 week home based intervention programme on the energy intake undergraduate female students in Iraq.
5. To evaluate which variables (physiological variables, physical activity, psychological distress and energy intake) are the most affected by the intervention programme.

1.5 Hypotheses

In order to achieve the objectives of the study, the hypotheses are addressed based on the variables measured. However, for objective five, research question is posed to evaluate which variables are the most affected by the intervention programme. The specific hypotheses and research question based on research objectives are listed below:

1.5.1 Hypothesis 1-8 for Objective 1

H₀₁ There are no significant differences in blood pressure across pre-test, post-test1, and posttest2 between experimental and control groups.

H₀₂ There are no significant differences in blood cholesterol across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₃ There are no significant differences in blood glucose across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₄ There are no significant differences in body mass index across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₅ There are no significant differences in waist hip ratio across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₆ There are no significant differences in body fat across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₇ There are no significant differences in VO₂max across pre-test, post-test1, and post-test2 between experimental and control groups.

H₀₈ There are no significant differences in vital capacity across pre-test, post-test1, and post-test2 between experimental and control groups.

1.5.2 Hypothesis 9 for Objective 2

H₀₉ There are no significant differences in physical activity across pre-test, post-test1, and post-test2 between experimental and control groups.

1.5.3 Hypothesis 10-13 for Objective 3

Ho10 There are no significant differences in psychological distress across pre-test and post-test between experimental and control groups.

Ho11 There are no significant differences in depression across pre-test and post-test between experimental and control groups.

Ho12 There are no significant differences in anxiety across pre-test and post-test between experimental and control groups.

Ho13 There are no significant differences in stress across pre-test and post-test between experimental and control groups.

1.5.4 Hypothesis 14 for Objective 4

Ho14 There are no significant differences in energy intake cross pre-test and post-test between experimental and control groups.

1.5.5 Research Question for Objective 5

Which variables (physiological variables, physical activity, psychological distress_and energy intake) are the mostly affected by the intervention programme?

1.6 Significance of the Study

This study aimed to investigate the effect of a 12 week home-based intervention programme which focused on simple exercises and information on dietary awareness. The intervention is characterised by its easiness and availability to perform at home without the need for female students to participate in sports or physical activities outside of their homes. Considering society's negative perceptions and sensitivity towards female's engagement in sports events and physical activities in public, doing physical activities at home would provide sedentary young females with opportunities to become more physically active and improve their health. Moreover, it will be a safer way to stay healthy and active despite the current unstable security situation in Iraq.

Additionally, the current study will contribute to a better understanding of the practical application of PA and dietary awareness programme through utilization of an experimental design. The findings of this study could be considered by the Ministries of Education and Higher Education in the curriculum development process so that physical education courses are introduced to schools and university students in Iraq. Universities and colleges may use the intervention discussed in the current study to help overcome the phenomenon of physical inactivity among female university students and encourage them to engage in various physical activities at home. Indeed, this will lead to better health among young female students in Iraq.

The findings of this study will be a useful reference for educators, researchers, and lecturers in universities in terms of designing new methods to help sedentary female students become more physically active and more aware of their daily food intake culminating in a healthier generation. College administrators or health promoters may prefer to incorporate eating regulation skills into female programmes, teach information about food choices, and enhance social supports for healthy attitudes and behaviour during this critical developmental period due to the importance of increasing knowledge and awareness of good nutrients and EI that contribute to health maintenance. Moreover, the finding of this study will be a useful for the social counsellors in universities in terms of decreasing the PD phenomenon. Indeed, practising PA and improving daily diet habits will reduce depression, anxiety, and stress which is popular among female students at this age due to the conditions mentioned that Iraqi society going through, which led to increasing the negative sentiment for life.

1.7 Delimitation of the Study

The present study has a number of delimitations as mentioned below:

First, the study was conducted in the College of Education, Soran University, Erbil, in the Northern Region of Iraq due to the lack of security and stability in the capital city, Baghdad and southern areas of Iraq.

Although there are many physiological variables that can be adopted and tested, this study only selected nine physiological variables (blood pressure, blood cholesterol, blood glucose, BMI, body fat, WHR, VO_2 max and vital capacity). Indeed, these variables are more related to general health (Danielsen et al., 2013; Daubenmier et al., 2007; Dunn et al., 1999; Esposito et al., 2003). Therefore, according to Goodpaster et al. (2010), Islam et al. (2013), and J ärvel äet al. (2012), the selected variables are the most related variables that can show the relevance of this variables to the improvement in health.

1.8 Limitation of the Study

There are some limitations of the current study. The official age of the students for entering university is 18 years old in Iraq. However, due to the possibility that some students dropped out or left school year because of security problems in the country and restarted school after several years, Ministry of Higher Education in Iraq determined the appropriate age range of 18-22 for accepting the students in universities. As a result, the selection of a homogeneous sample regarding the students' age would not be possible in the current study.

Studies have showed that the laboratory test is the most reliable way to get the results. However, due to the insecurity situation, difficulty in mobility and to ensure the safety of the students in this research, the field tests were used. However, the selected field tests were reported to have high validity and reliability as compared to the lab test. Examples include using step test instead of the lab treadmill test to measure the VO_2 max (Chatterjee,

Chatterjee, & Bandyopadhyay, 2005; Mcardle, Katch, Pechar, Jacobson, & Ruck, 1971). VO₂max step test with reliability score of .082 (Johnson & Nelson, 1969), using the caliper to measure body fat percentage instead of hydrostatic weighing test (Agarwal, Bills, & Light, 2010; Durnin & Womersley, 1974). Reliability for measuring body fat by Calipers were reported to be 0.79 to 0.91 (Kispert & Merrifield, 1987).

1.9 Definition of the Terms

The following definitions have been given to contribute to a better understanding of the basic terms used in the current study.

1.9.1 Physiological Variables

The physiological variables are the amount that can change frequently in the bodies of the human beings, animals or plants (Ting & Education, 2009). In this study, physiological variables include subjects' blood pressure, blood cholesterol, blood glucose, body fat, body mass index and waist hip ratio, VO₂max, vital capacity are the dependent variables and will be tested 3 times during the experiment. These variables are suggested because they are good indicators for health (Tjøanna et al., 2013; Tudor-Locke, 2010; Wing, Venditti, Jakicic, Polley, & Lang, 1998).

1.9.2 Physical Activity (PA)

WHO (2004b) defined the physical activity as any bodily movement produced by skeletal muscles that requires EI. Regular moderate intensity PA such as walking, cycling, or participating in sports, which has positive effect on health. For instance, it can reduce the risk of cardiovascular diseases, diabetes, colon and breast cancer, and depression. Moreover, adequate levels of PA decreases the risk of a hip or vertebral fracture and help control weight (Kushi et al., 2012; Lakoski et al., 2011; Lee et al., 2012). In this study, PA includes simple exercises used in the intervention programme tailored to be done at home, which is measured by counting the steps average in three days by pedometer.

1.9.3 Psychological distress

PD is defined as a general term used to describe unpleasant feelings or emotions that impact your level of functioning. In other words, it is psychological discomfort that interferes with your activities of daily living. PD can result in negative views of the environment, others, and the self. Sadness, anxiety, distraction, and symptoms of mental illness are manifestations of PD (Karkhanis & Mathur, 2016; Tiwari, 2016). The focus should be on promoting PD throughout the lifespan to ensure a healthy start in life and to prevent mental disorder (WHO, 2007). In this study, PD of depression, anxiety, and stress signs were measured in two different times: before and at the end of the study using DASS Questionnaire. The DASS depression, anxiety, and stress scales were developed by researchers at the University of New South Wales-Australia. A 42-item questionnaire includes three self-report scales designed to measure the negative emotional states of

depression, anxiety and stress. Each of the three scales contains 14 items, divided into subscales of 2-5 items with similar content (Lovibond & Lovibond, 1995).

1.9.4 Energy Intake

According to Bender (2009), EI is the total energy cost of maintaining constant conditions in the body, i.e. homeostasis (basal metabolism, BMR) plus the energy cost of PA. The average total EI in Western countries is about 1.4 times BMR; a desirable level of PA is about 1.7 times BMR. Energy intake plays a significant role in the quality of life, health and longevity (Marmot et al., 2010). In this study, the EI was measured depending on the amount of kilocalories (Kcal) intake per day in two different times: before and at the end of the study by using 24-hours food record (Yang et al., 2010).

1.9.5 Iraqi Undergraduate Level

The undergraduate level is defined as the level which students are admitted after completing the secondary or high school, which generally lasts for four years. In order to pursue their studies at undergraduate level in Iraq. The applicants, who must be 18 years old or above, register in colleges and universities to pursue their studies in various field such as arts, sciences, medicine, engineering so on and so forth. Bachelor's degrees in the fields of Education and Science are awarded upon the completion of four-year programmes (WES, 2004). In the current study, the first-year female students studying at the faculty of Education under a 4-year programme in the 18-22 age range were selected as the participants in the experiment.

1.9.6 Home Based Intervention Programme

An intervention programme should be a system that helps develop the individuals' skills, potentials, and capabilities, whether physical or psychological, and to reform their daily habits and lifestyle, and it may consist of different groups of programmes, tables, or schedules related to the type of target to be achieved (Guralnick & Bricker, 1987). In this study, the home-based intervention programme of contained PA combined with dietary awareness sessions in the form of a booklet including instructions and schedules for moderate and /or high intensity physical activities (5 days per week, 30 minutes per day) with paragraphs of nutrition knowledge and dietary awareness (2 days per week, 15 minutes per day), which it tailored to could be done easily at home. The study lasted for 12 weeks adapted from the 2008 Physical Activity Guidelines for Americans (P. A. G. A. Committee, 2008b) and from the Dietary Guidelines for Americans, 2010 (Agriculture & Services, 2010). The effects of the intervention on physiological, physical, PD and EI among undergraduate female students were measured through pre-test, post-test1 and post-test2.

REFERENCES

- Adams, M. A. (2009). *A pedometer-based intervention to increase physical activity: Applying frequent, adaptive goals and a percentile schedule of reinforcement.*
- Agarwal, R., Bills, J. E., & Light, R. P. (2010). Diagnosing obesity by body mass index in chronic kidney disease an explanation for the "Obesity Paradox?". *Hypertension Journal*, 56(5), 893-900.
- Agriculture, U. S. D. o., & Services, U. S. D. o. H. a. H. (2010). *Dietary Guidelines for Americans, 2010.* Washington, DC: U.S. Government Printing Office Retrieved from www.dietaryguidelines.gov
- Al-Haidari, A. A. J. (2011). The Reality of Higher Education in Iraq Between The Challenges and Necessity. *Modern Discussion* Retrieved 3573, from <http://www.ahewar.org/search/Dsearch.asp?nr=3573>
- Al-Haifi, A. R., Al-Fayez, M. A., Al-Athari, B. I., Al-Ajmi, F. A., Allafi, A. R., Al-Hazaa, H. M., & Musaiger, A. O. (2013). Relative contribution of physical activity, sedentary behaviors, and dietary habits to the prevalence of obesity among kuwaiti adolescents. *Food & Nutrition Bulletin Journal*, 34(1), 6-13.
- Al-Hazaa, H. M. (2005). Measuring physical activity and energy spent in humans: A brief review. *Arab Journal of Food and Nutrition*, 6(13), 26-50.
- Al-Tamimi, Y. A. (2007). Sports and social development goals. *Journal of Physical Education Sciences*, 2(5), 143-145.
- Al-Tamimi, Y. A., Armstrong, T., Cowan, M., & Riley, L. (2011). Noncommunicable Diseases Country Profiles 2011. France: World Health Organization.
- Al Subaie, K. B. S. M. (2005). The factors leading to poor student participation in student activities and means of overcoming them from the perspective of students at King Saud University. *Message Arabian Gulf Journal*, 94(1), 83.
- Alderson, J. C., Clapham, C., & Wall, D. (1995). *Language Test Construction and Evaluation* (1st ed.). Cambridge University Press Ernst Klett Sprachen.
- Allender, S., Cowburn, G., & Foster, C. (2006). Understanding participation in sport and physical activity among children and adults: A review of qualitative studies. *Health Education Research Journal*, 21(6), 826-835.
- Amine, E., Baba, N., Belhadj, M., Deurenbery-Yap, M., Djazayery, A., Forrester, T., . . . MBuyamba, J. (2002). Diet, Nutrition and the Prevention of Chronic Diseases: Report of a Joint WHO/FAO Expert Consultation. Switzerland, Geneva: World Health Organization.
- Ängeby, L. (2014). Dietary Habits. *SE-171 77*, from <http://ki.se/en/education/dietary-habits>

- Aoda, S. (2013). Women's Sport in Mosul, "Crime" Behind Closed Doors. <http://unami.unmissions.org/>.
- ARFAD. (2008). Education in Iraq Between Past and Present, from <http://www.irfad.org/ar/?s=%D8%A7%D9%84%D8%AA%D8%B9%D9%84%D9%8A%D9%85+%D9%81%D9%8A+%D8%A7%D9%84%D8%B9%D8%B1%D8%A7%D9%82+%D8%A8%D9%8A%D9%86+%D8%A7%D9%84%D9%85%D8%A7%D8%B6%D9%8A+%D9%88%D8%A7%D9%84%D8%AD%D8%A7%D8%B6%D8%B1>
- Arroll, B., & Beaglehole, R. (1992). Does physical activity lower blood pressure: A critical review of the clinical trials. *Journal of Clinical Epidemiology*, 45(5), 439-447.
- Artinian, N. T., Fletcher, G. F., Mozaffarian, D., Kris-Etherton, P., Van Horn, L., Lichtenstein, A. H., . . . Redeker, N. S. (2010). Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults a scientific statement from the american heart association. *Circulation Journal*, 122(4), 406-441.
- Ary, D., Jacobs, L. C., & Sorensen, C. (1990). *Introduction to Research in Education* (4th ed.). Chicago: Holt, Rinehart, and Winston.
- Ary, D., Jacobs, L. C., & Sorensen, C. (2010). *Introduction to Research in Education* (8th ed.): Belmont, CA: Wadsworth.
- Azevedo, K., Luiz, R., Rocco, P., & Conde, M. (2012). Vital capacity and inspiratory capacity as additional parameters to evaluate bronchodilator response in asthmatic patients: A cross sectional study. *BMC Pulmonary Medicine Journal*, 12(1), 49.
- Bandini, L. G., Anderson, S. E., Curtin, C., Cermak, S., Evans, E. W., Scampini, R., . . . Must, A. (2010). Food selectivity in children with autism spectrum disorders and typically developing children. *The Journal of Pediatrics*, 157(2), 259-264.
- Barbosa, L., Chaves, O. C., & Ribeiro, R. d. C. L. (2012). Anthropometric and body composition parameters to predict body fat percentage and lipid profile in schoolchildren. *Revista Paulista de Pediatria Journal*, 30(4), 520-528.
- Bartsch, R., Kantelhardt, J. W., Penzel, T., & Havlin, S. (2007). Experimental evidence for phase synchronization transitions in the human cardiorespiratory system. *Physical Review Letters*, 98(5), 4.
- Bauer, K. W., Neumark-Sztainer, D., Fulkerson, J. A., Hannan, P. J., & Story, M. (2011). Familial correlates of adolescent girls' physical activity, television use, dietary intake, weight, and body composition. *International Journal of Behavioral Nutrition and Physical Activity* 8(1), 1-10.
- Bender, D. A. (2009). *A Dictionary of Food and Nutrition* (3rd ed.). United States: OUP Oxford.

- Blair, S. N., Kohl, H. W., & Barlow, C. E. (1993). Physical activity, physical fitness, and all-cause mortality in women: Do women need to be active? *Journal of the American College of Nutrition*, 12(4), 368-371.
- Blissmer, B., Riebe, D., Dye, G., Ruggiero, L., Greene, G., & Caldwell, M. (2006). Health-related quality of life following a clinical weight loss intervention among overweight and obese adults: Intervention and 24 month follow-up effects. *Health & Quality of Life Outcomes Journal* 4(1), 43.
- Blumenthal, J. A., Babyak, M. A., Hinderliter, A., Watkins, L. L., Craighead, L., Lin, P.-H., . . . Sherwood, A. (2010). Effects of the DASH diet alone and in combination with exercise and weight loss on blood pressure and cardiovascular biomarkers in men and women with high blood pressure: The ENCORE study. *Archives of Internal Medicine Journal*, 170(2), 126-135.
- Bogaerts, A., Devlieger, R., Nuyts, E., Witters, I., Gyselaers, W., & Van den Bergh, B. (2013). Effects of lifestyle intervention in obese pregnant women on gestational weight gain and mental health: A randomized controlled trial. *International Journal of Obesity*, 37(6), 814-821.
- Booth, J. (1977). A short history of blood pressure measurement. *Proceedings of The Royal Society of Medicine Journal* 70(11), 793.
- Bouchard, C., Shephard, R. J., Stephens, T., Sutton, J., & McPherson, B. (1990). *Exercise, Fitness, and Health: A Consensus of Current Knowledge: Proceedings of The International Conference on Exercise, fitness, and health, May 29-June 3, 1988, Toronto, Canada.*
- Boyle, S. E., Jones, G. L., & Walters, S. J. (2010). Physical activity, quality of life, weight status and diet in adolescents. *Quality of Life Research Journal*, 19(7), 943-954.
- Brown, Potter, J. D., Jacobs Jr, D. R., Kopher, R. A., Rourke, M. J., Barosso, G. M., . . . Schmid, L. A. (1996). Maternal waist-to-hip ratio as a predictor of newborn size: results of the Diana Project. *Epidemiology*, 7(1), 62-66.
- Brown, T., & Summerbell. (2009). Systematic review of schoolbased interventions that focus on changing dietary intake and physical activity levels to prevent childhood obesity: An update to the obesity guidance produced by the National Institute for Health and Clinical Excellence. *Obesity Reviews Journal*, 10(1), 110-141.
- Burke, L., Lee, A. H., Jancey, J., Xiang, L., Kerr, D. A., Howat, P. A., . . . Anderson, A. S. (2013). Physical activity and nutrition behavioural outcomes of a home-based intervention program for seniors: A randomized controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), 14.
- Butler, S. M., Black, D. R., Blue, C. L., & Gretebeck, R. J. (2004). Change in diet, physical activity, and body weight in female college freshman. *American Journal of Health Behavior*, 28(1), 24-32.

- Byrne, B. M. (2013). *Structural Equation Modeling With AMOS: Basic Concepts, Applications, and Programming* (2nd ed.). New York: Routledge.
- Calfas, K. J., Sallis, J. F., Nichols, J. F., Sarkin, J. A., Johnson, M. F., Caparosa, S., . . . Alcaraz, J. E. (2000). Project GRAD: Two-year outcomes of a randomized controlled physical activity intervention among young adults. *American Journal of Preventive Medicine*, 18(1), 28-37.
- Caro, C. G., Pedley, T., Schroter, R., & Seed, W. (2012). *The Mechanics of The Circulation* (2nd ed.). England, Cambridge: Cambridge University Press.
- Caspersen, C. J., Pereira, M. A., & Curran, K. M. (2000). Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Medicine and Science in Sports and Exercise Journal*, 32(9), 1601-1609.
- Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and physical fitness: Definitions and distinctions for health-related research. *Public Health Reports Journal*, 100(2), 126.
- Cecchini, M., Sassi, F., Lauer, J. A., Lee, Y. Y., Guajardo-Barron, V., & Chisholm, D. (2010). Tackling of unhealthy diets, physical inactivity, and obesity: Health effects and cost-effectiveness. *The Lancet Journal*, 376(9754), 1775-1784.
- Chatterjee, S., Chatterjee, P., & Bandyopadhyay, A. (2005). Validity of queen's college step test for estimation of maximum oxygen uptake in female students. *Indian Journal of Medical Research*, 121(1), 32-35.
- Chobanian, A. V., Bakris, G. L., Black, H. R., Cushman, W. C., Green, L. A., Izzo Jr, J. L., . . . Wright Jr, J. T. (2003). The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: The JNC 7 report. *Jama Journal*, 289(19), 2560-2571.
- Christy, S. M., Mosher, C. E., Sloane, R., Snyder, D. C., Lobach, D. F., & Demark-Wahnefried, W. (2011). Long-term dietary outcomes of the FRESH START intervention for breast and prostate cancer survivors. *Journal of The American Dietetic Association*, 111(12), 1844-1851.
- Church, T. S., Earnest, C. P., Skinner, J. S., & Blair, S. N. (2007). Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure: A randomized controlled trial. *Jama Journal*, 297(19), 2081-2091.
- Coghill, N., & Cooper, A. R. (2008). The effect of a home-based walking program on risk factors for coronary heart disease in hypercholesterolaemic men.: A randomized controlled trial. *Preventive Medicine Journal*, 46(6), 545-551.
- Cohen, A. D. (1980). *Testing Language Ability in The Classroom* (1st ed.). United States, New York: Newbury House Pub.

- Cohen, J. (1988). *Statistical Power Analysis for The Behavioral Sciences* (2nd ed.). United States: L. Erlbaum Associates.
- Collins, T. C., Lunos, S., Carlson, T., Henderson, K., Lightbourne, M., Nelson, B., & Hodges, J. S. (2011). Effects of a home-based walking intervention on mobility and quality of life in people with diabetes and peripheral arterial disease a randomized controlled trial. *Diabetes Care Journal*, 34(10), 2174-2179.
- Committee, D. G. A. (2010). Report of the dietary guidelines advisory committee on the dietary guidelines for Americans, 2010, to the Secretary of Agriculture and the Secretary of Health and Human Services. *Agricultural Research Service Journal*.
- Committee, P. A. G. A. (2008a). Physical activity guidelines advisory committee report, 2008. *Washington, DC: US Department of Health and Human Services, 2008*.
- Committee, P. A. G. A. (2008b). Physical activity guidelines for Americans. *Washington, DC: US Department of Health and Human Services* 15-34.
- Cooper, K., & Hancock, C. (2011). Review: The Benefits of Physical Activity for Health and Well-being. UK: C3 Collaborating for Health.
- Council, N. H. a. M. R. (2013). *Australian Dietary Guidelines*. Canberra: National Health and Medical Research Council.
- Crawford, J. R., & Henry, J. D. (2003). The Depression Anxiety Stress Scales (DASS): Normative data and latent structure in a large non-clinical sample. *British Journal of Clinical Psychology*, 42(2), 111-131.
- Crawford, P. B., Obarzanek, E., Morrison, J., & Sabry, Z. (1994). Comparative advantage of 3-day food records over 24-hour recall and 5-day food frequency validated by observation of 9-and 10-year-old girls. *Journal of The American Dietetic Association*, 94(6), 626-630.
- Danielsen, K. K., Svendsen, M., Mæhlum, S., & Sundgot-Borgen, J. (2013). Changes in body composition, cardiovascular disease risk factors, and eating behavior after an intensive lifestyle intervention with high volume of physical activity in severely obese subjects: A prospective clinical controlled trial. *Journal of Obesity*, 2013, 12. doi: 10.1155/2013/325464
- Daubenmier, J. J., Weidner, G., Sumner, M. D., Mendell, N., Merritt-Worden, T., Studley, J., & Ornish, D. (2007). The contribution of changes in diet, exercise, and stress management to changes in coronary risk in women and men in the multisite cardiac lifestyle intervention program. *Annals of Behavioral Medicine Journal*, 33(1), 57-68.
- Davies, A. (1999). *Dictionary of Language Testing* (1st ed. Vol. 7). UK: Cambridge University Press.

- Denke, M. A., Sempos, C. T., & Grundy, S. M. (1993). Excess body weight: An underrecognized contributor to high blood cholesterol levels in white American men. *Archives of Internal Medicine Journal*, 153(9), 1093.
- Dlugosz, E. M., Chappell, M. A., Meek, T. H., Szafrńska, P. A., Zub, K., Konarzewski, M., . . . Careau, V. (2013). Phylogenetic analysis of mammalian maximal oxygen consumption during exercise. *The Journal of Experimental Biology*, 216(24), 4712-4721.
- Dorothy J. Klimis-Zacas. (2001). *Nutrition 01/02* (13th ed.). USA: Guilford, CT: McGraw-Hill/Dushkin.
- Dunlap, A. N. (2012). *The Impact of a Intervention Program on The Knowledge and Behaviors of School-age Children in Alabama Regarding Nutrition and Physical activity*. Ph.D Graduate Theses and Dissertations, Iowa State University, Ames, Iowa.
- Dunn, A. L., Marcus, B. H., Kampert, J. B., Garcia, M. E., Kohl III, H. W., & Blair, S. N. (1999). Comparison of lifestyle and structured interventions to increase physical activity and cardiorespiratory fitness: A randomized trial. *Jama Journal*, 281(4), 327-334.
- Durnin, J., & Womersley, J. (1974). Body fat assessed from total body density and its estimation from skinfold thickness: Measurements on 481 men and women aged from 16 to 72 years. *British Journal of Nutrition*, 32(01), 77-97.
- Dwyer, J. J. M., Wilson, K., Limarzi, L., Callaghan, B., & Croskery, L. (2013). Physical activity among female adolescents of indian and polish origin in mississauga, ontario: An examination of shared and ethno-cultural barriers. *Revue ph éEPS/PHEnex Journal*, 4(3), 14.
- Eaton, C. B., Lapane, K. L., Garber, C. A., Assaf, A. R., Lasater, T. M., & Carleton, R. A. (1995). Sedentary lifestyle and risk of coronary heart disease in women. *Medicine and Science in Sports and Exercise Journal*, 27(11), 1535-1539.
- Elliot, C. A., Kennedy, C., Morgan, G., Anderson, S. K., & Morris, D. (2012). Undergraduate physical activity and depressive symptoms: A national study. *American Journal of Health Behavior*, 36(2), 230-241.
- Ellis, P. D. (2010). *The Essential Guide to Effect Sizes: Statistical Power, Meta-Analysis, and The Interpretation of Research Results* (1st ed.). United Kingdom: University Press Cambridge.
- Epstein, L. H., Wing, R. R., Thompson, J. K., & Griffin, W. (1980). Attendance and fitness in aerobics exercise the effects of contract and lottery procedures. *Behavior Modification Journal*, 4(4), 465-479.

- Esposito, K., Pontillo, A., Di Palo, C., Giugliano, G., Masella, M., Marfella, R., & Giugliano, D. (2003). Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: A randomized trial. *Jama Journal*, 289(14), 1799-1804.
- Ethington, M. D. (2009). *Short-Term Effects of a Nutrition Education Program on Food Choices in Adolescents at Risk for Type 2 Diabetes*. Ph.D, University of Texas Medical Branch, Galveston, Texas, United States.
- Fanelli, M., & Stevenhagen, K. (1986). Consistency of energy and nutrient intakes of older adults: 24-hour recall vs. 1-day food record. *Journal of the American Dietetic Association (USA)*, 86(5.), 665 - 667.
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods Journal*, 39(2), 175-191.
- Ferreira, M., Matsudo, S., Matsudo, V., & Braggion, G. (2005). Effects of an intervention program of physical activity and nutrition orientation on the physical activity level of physically active women aged 50 to 72 years old. *Revista Brasileira de Medicina do Esporte Journal*, 11(3), 172-176.
- Fink, A. (2012). *Evidence-based public health practice*: Sage Publications.
- Fitzgerald, A., Heary, C., Nixon, E., & Kelly, C. (2010). Factors influencing the food choices of Irish children and adolescents: A qualitative investigation. *Health Promotion International Journal*, 25(3), 289-298.
- Flegal, K. M., Carroll, M. D., Ogden, C. L., & Curtin, L. R. (2010). Prevalence and trends in obesity among US adults, 1999-2008. *Jama Journal*, 303(3), 235-241.
- Fletcher, G. F., Balady, G., Blair, S. N., Blumenthal, J., Caspersen, C., Chaitman, B., . . . Pina, I. L. (1996). Statement on exercise: Benefits and recommendations for physical activity programs for all Americans a statement for health professionals by the committee on exercise and cardiac rehabilitation of the council on clinical cardiology, American heart association. *Circulation Journal*, 94(4), 857-862.
- Ford, E. S., Bergmann, M. M., Boeing, H., Li, C., & Capewell, S. (2012). Healthy lifestyle behaviors and all-cause mortality among adults in the United States. *Preventive Medicine Journal*, 55(1), 23-27.
- Fox, K. R. (1999). The influence of physical activity on mental well-being. *Public Health Nutrition Journal*, 2(3a), 411-418.
- Freak-Poli, R., Wolfe, R., Brand, M., Courten, M., & Peeters, A. (2013). Eight-month postprogram completion: Change in risk factors for chronic disease amongst participants in a 4-month pedometer-based workplace health program. *Obesity Research Journal*, 21(9), E360-E368.

- Freedman, M. R., King, J., & Kennedy, E. (2001). Executive summary. *Obesity Research Journal*, 9(S3), 1S-5S.
- Ganasegeran, K., Al-Dubai, S., Qureshi, A. M., Al-abed, A.-a. A., & Aljunid, S. M. (2012). Social and psychological factors affecting eating habits among university students in a Malaysian medical school: A cross-sectional study. *Nutrition Journal*, 11(48), 107.
- Garson, G. (2010). Testing of Assumptions: Statnotes, from North Carolina State University. *Public Administration Program. Online*, 21.
- George, D. (2003). *SPSS for Windows Step by Step: A Simple Study Guide and Reference, 17.0 update, 10/e* (4th ed.). India: Pearson Education
- GHO, G. H. O. (2013). Iraq: Country Profiles. world health organization.
- Gibney, M., Margetts, B., Kearney, J., & Arab, L. (2004). *Public Health Nutrition: The Nutrition Society Textbook Series*. United States: Oxford: Blackwell Science.
- Gomez-Pinilla, F. (2011). The combined effects of exercise and foods in preventing neurological and cognitive disorders. *Preventive Medicine Journal*, 52(1), S75-S80.
- Goodpaster, B. H., DeLany, J. P., Otto, A. D., Kuller, L., Vockley, J., South-Paul, J. E., . . . Hames, K. C. (2010). Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults a randomized trial. *Jama Journal*, 304(16), 1795-1802.
- Government, H. (2011). *No health without mental health: A cross-government mental health outcomes strategy for people of all ages*. London: Department of Health Retrieved from www.dh.gov.uk/mentalhealthstrategy.
- Greaves, C. J., Sheppard, K. E., Abraham, C., Hardeman, W., Roden, M., Evans, P. H., & Schwarz, P. (2011). Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health Journal*, 11(1), 119.
- Groeneveld, I. F., Proper, K. I., Van der Beek, A. J., Hildebrandt, V. H., & Van Mechelen, W. (2011). Short and long term effects of a lifestyle intervention for construction workers at risk for cardiovascular disease: A randomized controlled trial. *BMC Public Health Journal*, 11(1), 836.
- Group, D. P. P. R. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *The New England Journal of Medicine*, 346(6), 393.
- Guelinckx, I., Devlieger, R., Mullie, P., & Vansant, G. (2010). Effect of lifestyle intervention on dietary habits, physical activity, and gestational weight gain in obese pregnant women: A randomized controlled trial. *The American journal of clinical nutrition*, 91(2), 373-380.

- Gulliford, M. C., Mahabir, D., & Roche, B. (2003). Food insecurity, food choices, and body mass index in adults: nutrition transition in Trinidad and Tobago. *International Journal of Epidemiology*, 32(4), 508-516.
- Guralnick, M. J., & Bricker, D. (1987). The Effectiveness of Early Intervention for Children with Cognitive and General Developmental Delays (pp. 115-173). San Diego, CA, US: Academic Press.
- Habib, K. (2007). The Status of Women in Iraqi Society and The Necessities of Change. *The reality of women in Iraq after the change, freedom and women's equality is an essential part of the democratic values of civil society 4*. Retrieved from <http://www.ahewar.org/debat/show.art.asp?aid=91148>
- Hamilton, M. T., Healy, G. N., Dunstan, D. W., Zderic, T. W., & Owen, N. (2008). Too little exercise and too much sitting: Inactivity physiology and the need for new recommendations on sedentary behavior. *Current Cardiovascular Risk Reports Journal*, 2(4), 292-298.
- Hardin, D. S., Hebert, J. D., Bayden, T., Dehart, M., & Mazur, L. (1997). Treatment of childhood syndrome X. *Journal of Pediatrics*, 100(2), 6.
- Harrison, R. L. (2007). *Weigh to Life: A Combined Lifestyle Intervention of Diet, Physical Activity, and Behavioral Strategies*. University of Missouri--St. Louis.
- Haviland, W., Prins, H., McBride, B., & Walrath, D. (2010). *Cultural Anthropology: The Human Challenge* (13th ed.). United states: Cengage Learning.
- Health, U. D. o., & Services, H. (1996). Physical activity and health: a report of the surgeon general. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention. *National Center for Chronic Disease Prevention and Health Promotion*, 147.
- Health, U. D. o., Services, H., General, U. S. P. H. S. O. o. t. S., Prevention, N. C. f. C. D., Promotion, H., Fitness, P. s. C. o. P., & Sports. (1996). *Physical Activity and Health: A Report of The Surgeon General* (1st ed.). UK: International Medical Pub.
- Heywood, V. (2002). *Advanced Fitness Assessment and Exercise Prescription* (5th ed.). United States: Champaign, IL: Human Kineticspub Europe Ltd.
- Honkola, A., Forsen, T., & Eriksson, J. (1997). Resistance training improves the metabolic profile in individuals with type 2 diabetes. *Acta Diabetologica Journal*, 34(4), 245-248.
- Humairi, A. (2015). Nutrition awareness of rural women and it's relationship with some personal and social – economical factors. *Journal of The University of Babylon*, 23(1), 15.

- Imayama, I., Alfano, C. M., Kong, A., Foster-Schubert, K. E., Bain, C. E., Xiao, L., . . . Blackburn, G. L. (2011). Dietary weight loss and exercise interventions effects on quality of life in overweight/obese postmenopausal women: A randomized controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, 8(118), 1-12.
- Islam, N. S., Zanolwiak, J. M., Wyatt, L. C., Chun, K., Lee, L., Kwon, S. C., & Trinh-Shevrin, C. (2013). A randomized-controlled, pilot intervention on diabetes prevention and healthy lifestyles in the New York City Korean community. *Journal of community health*, 38(6), 1030-1041.
- Ismail, Y. A., & Shihab, I. H. (1994). Trends of female teachers and male teachers of various subjects about physical education depending on the size of the practice of sports. *Journal of The Science of Physical Education*, 1(7), 8.
- Iwane, M., Arita, M., Tomimoto, S., Satani, O., Matsumoto, M., Miyashita, K., & Nishio, I. (2000). Walking 10,000 steps/day or more reduces blood pressure and sympathetic nerve activity in mild essential hypertension. *Hypertension research: Official Journal of The Japanese Society of Hypertension*, 23(6), 573-580.
- Jacka, F. N., Mykletun, A., Berk, M., Bjelland, I., & Tell, G. S. (2011). The association between habitual diet quality and the common mental disorders in community-dwelling adults: The Hordaland health study. *Psychosomatic Medicine Journal*, 73(6), 483-490.
- Jackson, & Howton, A. (2008). Increasing walking in college students using a pedometer intervention: Differences according to body mass index. *Journal of American College Health* 57(2), 159-164.
- Jackson, A., Stanforth, P., Gagnon, J., Rankinen, T., Leon, A., Rao, D., . . . et, a. (2002). The effect of sex, age and race on estimating percentage body fat from body mass index: The Heritage Family Study. "True" *International Journal of Obesity*, 26(6), 789-796.
- Jackson, A. S., Pollock, M. L., & Ward, A. (1979). Generalized equations for predicting body density of women. *Medicine and science in sports and exercise*, 12(3), 175-181.
- Jaggers, J. R., Prasad, V., Dudgeon, W. D., Hand, G. A., Burgess, S., Kalinski, M. I., & Blair, S. N. (2014). *Results of a home-based exercise intervention to increase physical activity among people living with HIV*. Paper presented at the Medicine and Science in Sport and Exercise.
- Jakicic, J. M., Winters, C., Lang, W., & Wing, R. R. (1999). Effects of intermittent exercise and use of home exercise equipment on adherence, weight loss, and fitness in overweight women: A randomized trial. *Jama Journal*, 282(16), 1554-1560.

- Järvelä L. S., Kempainen, J., Niinikoski, H., Hannukainen, J. C., Lähteenmäki, P. M., Kapanen, J., . . . Heinonen, O. J. (2012). Effects of a home-based exercise program on metabolic risk factors and fitness in long-term survivors of childhood acute lymphoblastic leukemia. *Pediatric Blood & Cancer Journal*, 59(1), 155-160.
- Jednacz, E., & Rutkowska-Sak, L. (2015). Assessment of the body composition and parameters of the cardiovascular risk in juvenile idiopathic arthritis. *BioMed Research International Journal*, 2015, 8.
- Johnson, B. L., & Nelson, J. K. (1969). *Practical Measurements for Evaluation in Physical Education* (4th ed.). University of Michigan: Burgess Pub.
- Karkhanis, R., & Mathur, K. (2016). Impact of physical distress and psychological distress in women passing through different stages of menopause. *Indian Journal of Health and Wellbeing*, 7(1), 37.
- Karvetti, R., & Knuts, L. R. (1985). Validity of the 24-hour dietary recall. *Journal of the American Dietetic Association*, 85(11), 1437-1442.
- Kassavou, A., Turner, A., & French, D. P. (2013). Do interventions to promote walking in groups increase physical activity? A meta-analysis. *International Journal of Behavioral Nutrition and Physical Activity*, 10(1), 18.
- Katzmarzyk, P. T. (2010). Physical activity, sedentary behavior, and health: Paradigm paralysis or paradigm shift? *Diabetes Journal*, 59(11), 2717-2725.
- Kelley, & Kelley, K. S. (2009). Impact of progressive resistance training on lipids and lipoproteins in adults: A meta-analysis of randomized controlled trials. *Preventive Medicine Journal*, 48(1), 9-19.
- Kelley, Kelley, K. S., & Tran, Z. V. (2004). Aerobic exercise and lipids and lipoproteins in women: A meta-analysis of randomized controlled trials. *Journal of Women's Health* 13(10), 1148-1164.
- Kelley, & Preacher, K. J. (2012). On effect size. *Psychological Methods Journal*, 17(2), 137-152.
- Kelley, G., & McClellan, P. (1994). Antihypertensive effects of aerobic exercise a brief meta-analytic review of randomized controlled trials. *American Journal of Hypertension*, 7(2), 115-119.
- Kelley, G. A., Kelley, K. S., Roberts, S., & Haskell, W. (2012a). Combined effects of aerobic exercise and diet on lipids and lipoproteins in overweight and obese adults: A meta-analysis. *Journal of Obesity*, 2012, 16.
- Kelley, G. A., Kelley, K. S., Roberts, S., & Haskell, W. (2012b). Comparison of aerobic exercise, diet or both on lipids and lipoproteins in adults: A meta-analysis of randomized controlled trials. *Clinical Nutrition Journal*, 31(2), 156-167.

- Kispert, C. P., & Merrifield, H. H. (1987). Interrater reliability of skinfold fat measurements. *Physical Therapy Journal*, 67(6), 917-920.
- Kittler, P. G., & Sucher, K. P. (1998). *Food and Culture in America: A Nutrition Handbook* (2nd ed.). USA, Belmont: Wadsworth Publishing Company, Inc.
- Kubiš, J., & Al Hussein, Z. R. a. (2016). Report on the Protection of Civilians in the Armed Conflict in Iraq: 1 May – 31 October 2015 (pp. 46). Baghdad-Iraq United Nations Assistance Mission for Iraq (UNAMI) & the Office of the United Nations High Commissioner for Human Rights (OHCHR).
- Kushi, L. H., Doyle, C., McCullough, M., Rock, C. L., Demark-Wahnefried, W., Bandera, E. V., . . . Gansler, T. (2012). American cancer society guidelines on nutrition and physical activity for cancer prevention. *CA: A Cancer Journal for Clinicians*, 62(1), 30-67.
- Labrosse, E. (2008). Master your pedometer: A study examining the effects of wearing a hip pedometer on the activity behaviours of grade five and six children. *Masters Abstracts International Journal*, 47(03), 6.
- Lakoski, S. G., Barlow, C. E., Farrell, S. W., Berry, J. D., Morrow Jr, J. R., & Haskell, W. L. (2011). Impact of body mass index, physical activity, and other clinical factors on cardiorespiratory fitness (from the Cooper Center longitudinal study). *The American Journal of Cardiology*, 108(1), 34-39.
- Leavey, V., Sandrey, M., & Dahmer, G. (2010). Comparative effects of 6-week balance, gluteus medius strength, and combined programs on dynamic postural control. *Journal of Sport Rehabilitation*, 19(3).
- Lee, I.-M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., Katzmarzyk, P. T., & Group, L. P. A. S. W. (2012). Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *The Lancet Journal*, 380(9838), 219-229.
- Leslie, E., Fotheringham, M. J., Owen, N., & Bauman, A. (2001). Age-related differences in physical activity levels of young adults. *Medicine and Science in Sports and Exercise Journal*, 33(2), 255-258.
- Lindström, J., Louheranta, A., Mannelin, M., Rastas, M., Salminen, V., Eriksson, J., . . . Tuomilehto, J. (2003). The finnish diabetes prevention study (DPS) lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care Journal*, 26(12), 3230-3236.
- Lock, R. S. (1990). College women's decision-making skills relating to voluntary participation in physical activity during leisure time. *Perceptual and Motor Skills Journal*, 71(1), 141-146.
- Loprinzi, P. D., Smit, E., & Mahoney, S. (2014). Physical activity and dietary behavior in US adults and their combined influence on health. *Perceptual and Motor Skills Journal*, 89(22), 190-198.

- Lotrič, M. B., & Stefanovska, A. (2000). Synchronization and modulation in the human cardiorespiratory system. *Physica A: Statistical Mechanics and its Applications*, 283(3), 451-461.
- Lovibond, P. F. (1998). Long-term stability of depression, anxiety, and stress syndromes. *Journal of Abnormal Psychology*, 107(3), 520-526.
- Lovibond, P. F. (2014, Nov 10, 2014). Depression Anxiety Stress Scales (DASS), from <http://www2.psy.unsw.edu.au/dass/>
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the depression anxiety stress scales (DASS) with the beck depression and anxiety inventories. *Behaviour Research and Therapy Journal*, 33(3), 335-343.
- Lowenberg, M. E., Todhunter, E. N., Wilson, E. D., Savage, J. R., & Lubawski, J. (1979). *Food and People* (3rd ed.). USA: John Wiley and Sons.
- Macera, C. A. (2003). Promoting healthy eating and physical activity for a healthier nation. *Center for Disease Control and Prevention*. Viewed at <http://www.cdc.gov/healthyouth/publications/pdf/pp-ch7.pdf>.
- Madi, A. (2007). *Is there a woman's role in Iraq after the change?* (1st ed. Vol. 4). Iraq, Baghdad: Civilized dialogue Foundation.
- Malina, R. M. (2001). Physical activity and fitness: Pathways from childhood to adulthood. *American Journal of Human Biology*, 13(2), 162-172.
- Malina, R. M., Bouchard, C., & Bar-Or, O. (2004). *Growth, Maturation, and Physical Activity* (2nd ed.). United States: Human Kinetics.
- Mann, J., & Truswell, S. (2012). *Essentials of Human Nutrition* (4th ed.). United States: Oxford University Press.
- Marcus, B. H., & Forsyth, L. (2003). *Motivating People to Be Physically Active* (2nd ed.). United States of America Human Kinetics.
- Marmot, M., Atinmo, T., Byers, T., Cavill, N., Chen, J., Hirohata, T., . . . Leitzmann, C. (2010). Executive Summary Policy and Action for Cancer Prevention Food, Nutrition, and Physical Activity. . Washington, DC: World Cancer Research Fund and American Institute for Cancer Research.
- Mcardle, W. D., Katch, F. I., Pechar, G. S., Jacobson, L., & Ruck, S. (1971). Reliability and interrelationships between maximal oxygen intake, physical work capacity and step-test scores in college women. *Medicine and Science in Sports Journal*, 4(4), 182-186.

- McDowell, M. A., Fryar, C. D., Ogden, C. L., & Flegal, K. M. (2008). *Anthropometric Reference Data for Children and Adults: United States, 2003-2006*. United States: Hyattsville, MD: National Center for Health Statistics Retrieved from <http://www.cdc.gov/nchs/data/nhsr/nhsr010.pdf>.
- McGinnis, J. M., & Foege, W. H. (1993). Actual causes of death in the United States. *JAMA: The Journal of the American Medical Association*, 270(18), 2207-2212.
- McKinney, C., Bishop, V., Cabrera, K., Medina, R., Takawira, D., Donate, N., . . . Guevara, B. (2014). NuFit: Nutrition and fitness CBPR program evaluation. *Journal of Prevention & Intervention in The Community*, 42(2), 112-124.
- McMartin, S. E., Kuhle, S., Colman, I., Kirk, S. F., & Veugelers, P. J. (2012). Diet quality and mental health in subsequent years among Canadian youth. *Public Health Nutrition Journal*, 15(12), 2253-2258.
- Mcmurray, R. G., Harrell, J. S., Bangdiwala, S. I., Bradley, C. B., Deng, S., & Levine, A. (2002). A school-based intervention can reduce body fat and blood pressure in young adolescents. *Journal of Adolescent Health*, 31(2), 125-132.
- Meyers, L. S., & Grossen, N. E. (1974). *Behavioral Research: Theory, Procedure, and Design* (2nd ed.). New York: WH Freeman.
- Miller, P. B., Anthony, D., & Yarrish, K. K. (2013). Exploration of healthy eating education and consumption of various foods of college students in a small liberal arts college. *Contemporary Issues in Education Research (CIER) Journal*, 6(1), 67-72.
- Minhas, M. (2013a). Interaction of Physical Activity, Diet, Health Locus of Control and Quality of Life among Finnish University Students.
- Minhas, M. (2013b). *Interaction of Physical Activity, Diet, Health Locus of Control and Quality of Life Among Finnish University Students*. MSc, University of Jyväskylä Finland.
- Mirkin, B. (2010). *Population Levels, Trends and Policies in The Arab Region: Challenges and Opportunities*. UN United Nations Development Programme, Regional Bureau for Arab States.
- Mitchell, M., & Jolley, J. (2004). *Research Design Explained* (5th ed.). Belmont, CA Wadsworth.
- Montoye, H. J. (1975). *Physical Activity and Health: An Epidemiologic Study of an Entire Community*. United States: Prentice-Hall Englewood Cliffs, NJ.
- Morey, M. C., Snyder, D. C., Sloane, R., Cohen, H. J., Peterson, B., Hartman, T. J., . . . Demark-Wahnefried, W. (2009). Effects of home-based diet and exercise on functional outcomes among older, overweight long-term cancer survivors: RENEW: a randomized controlled trial. *Jama*, 301(18), 1883-1891.

- Morgan, A. Z., Keiley, M. K., Ryan, A. E., Radomski, J. G., Gropper, S. S., Connell, L. J., . . . Ulrich, P. V. (2012). Eating regulation styles, appearance schemas, and body satisfaction predict changes in body fat for emerging adults. *Journal of Youth and Adolescence*, 41(9), 1127-1141.
- Mostert, S., & Kesselring, J. (2002). Effects of a short-term exercise training program on aerobic fitness, fatigue, health perception and activity level of subjects with multiple sclerosis. *Multiple Sclerosis Journal*, 8(2), 161-168.
- Musaiger, A. (2013). Nutrition and Its Relationship to Physical Education and Sports. 2. Retrieved from <http://www.acnut.com/v/>
- Musaiger, A., & Al-Hazzaa, H. M. (2012). Prevalence and risk factors associated with nutrition-related noncommunicable diseases in the Eastern Mediterranean region. *International Journal of General Medicine*, 5, 199-217.
- Mytton, O. T., Clarke, D., & Rayner, M. (2012). Taxing unhealthy food and drinks to improve health. *BMJ Journal*, 2012(344), 7.
- Nemet, D., Barkan, S., Epstein, Y., Friedland, O., Kowen, G., & Eliakim, A. (2005). Short-and long-term beneficial effects of a combined dietary-behavioral-physical activity intervention for the treatment of childhood obesity. *The Journal of Pediatrics*, 115(4), e443-e449.
- Newby, P. K., Muller, D., Hallfrisch, J., Qiao, N., Andres, R., & Tucker, K. L. (2003). Dietary patterns and changes in body mass index and waist circumference in adults. *The American Journal of Clinical Nutrition*, 77(6), 1417-1425.
- Ogden, C. L., & Carroll, M. D. (2010). Prevalence of overweight, obesity, and extreme obesity among adults: United States, trends 1960-1962 through 2007-2008. *National Center for Health Statistics Journal*, 6, 1-6.
- Ohlhaut, A. (2012). *Effects of Instruction Method on Vital Capacity and Maximum Sustained Phonation in Adult Female Controls*. M.S., East Carolina University. Retrieved from <http://hdl.handle.net/10342/3905>
- Paffenbarger, R. S., Wing, A. L., & Hyde, R. T. (1978). Physical activity as an index of heart attack risk in college alumni. *American Journal of Epidemiology*, 108(3), 161-175.
- Panel, E. (2010). American college of sports medicine roundtable on exercise guidelines for cancer survivors. *ACSM Journal*, 1409-1426.
- Patel, A., Keogh, J. W., Kolt, G. S., & Schofield, G. M. (2013). The long-term effects of a primary care physical activity intervention on mental health in low-active, community-dwelling older adults. *Aging and Mental Health Journal* 17(6), 766-772.

- Patrick, K., Calfas, K. J., Norman, G. J., Zabinski, M. F., Sallis, J. F., Rupp, J., . . . Cella, J. (2006). Randomized controlled trial of a primary care and home-based intervention for physical activity and nutrition behaviors: PACE+ for adolescents. *Archives of Pediatrics and Adolescent Medicine Journal*, *160*(2), 128-136.
- Pekmezi, D. W., & Demark-Wahnefried, W. (2011). Updated evidence in support of diet and exercise interventions in cancer survivors. *Acta Oncologica Journal*, *50*(2), 167-178.
- Peluso, M. A. M., & Andrade, L. H. S. G. d. (2005). Physical activity and mental health: The association between exercise and mood. *Clinics Journal*, *60*(1), 61-70.
- Penedo, F. J., & Dahn, J. R. (2005). Exercise and well-being: A review of mental and physical health benefits associated with physical activity. *Current Opinion in Psychiatry Journal*, *18*(2), 189-193.
- Petajan, J. H., Gappmaier, E., White, A. T., Spencer, M. K., Mino, L., & Hicks, R. W. (1996). Impact of aerobic training on fitness and quality of life in multiple sclerosis. *Annals of Neurology Journal*, *39*(4), 432-441.
- Peterson, J. A., Yates, B. C., Atwood, J. R., & Hertzog, M. (2005). Effects of a physical activity intervention for women. *Western Journal of Nursing Research*, *27*(1), 93-110.
- Pickering, T. G., Hall, J. E., Appel, L. J., Falkner, B. E., Graves, J., Hill, M. N., . . . Roccella, E. J. (2005). Recommendations for blood pressure measurement in humans and experimental animals part 1: Blood pressure measurement in humans: A statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension Journal*, *45*(1), 142-161.
- Pinto, B. M., Frierson, G. M., Rabin, C., Trunzo, J. J., & Marcus, B. H. (2005). Home-based physical activity intervention for breast cancer patients. *Journal of Clinical Oncology*, *23*(15), 3577-3587.
- Ponichtera-Mulcare, J. A., Mathews, T., Barrett, P. J., & Gupta, S. C. (1997). Change in aerobic fitness of patients with multiple sclerosis during a 6-month training program. *Research in Sports Medicine: An International Journal*, *7*(3-4), 265-272.
- Popkin, B. M., Adair, L. S., & Ng, S. W. (2012). Global nutrition transition and the pandemic of obesity in developing countries. *Nutrition Reviews Journal*, *70*(1), 3-21.
- Powers, A. R., Struempfer, B. J., Guarino, A., & Parmer, S. M. (2005). Effects of a nutrition education program on the dietary behavior and nutrition knowledge of second-grade and third-grade students. *Journal of School Health*, *75*(4), 129-133.

- Pribis, P., Burtnack, C. A., McKenzie, S. O., & Thayer, J. (2010). Trends in body fat, body mass index and physical fitness among male and female college students. *Nutrients Journal*, 2(10), 1075-1085.
- Racette, S. B., Deusinger, S. S., Strube, M. J., Highstein, G. R., & Deusinger, R. H. (2005). Weight changes, exercise, and dietary patterns during freshman and sophomore years of college. *Journal of American College Health*, 53(6), 245-251.
- Rangul, V., Bauman, A., Holmen, T. L., & Midthjell, K. (2012). Is physical activity maintenance from adolescence to young adulthood associated with reduced CVD risk factors, improved mental health and satisfaction with life: The HUNT Study, Norway. *International Journal of Behavioral Nutrition and Physical Activity*, 9(144), 27.
- RehmanArshad, A. (2013). Accuracy of accutrend GCT meter for the measurement of blood cholesterol levels. *Rawal Medical Journal*, 38(4), 354-357.
- Richardson, C. R., Faulkner, G., McDevitt, J., Skrinar, G. S., Hutchinson, D. S., & Piette, J. D. (2014). Integrating physical activity into mental health services for persons with serious mental illness. *Psychiatric Services Journal*, 56(3), 8.
- Rippe, J. M., Price, J. M., Hess, S. A., Kline, G., DeMers, K. A., Damitz, S., . . . Freedson, P. (1998). Improved psychological well-being, quality of life, and health practices in moderately overweight women participating in a 12-week structured weight loss program. *Obesity Research Journal*, 6(3), 208-218.
- Rowlands, A. V., Eston, R. G., & Ingledew, D. K. (1999). Relationship between activity levels, aerobic fitness, and body fat in 8-to 10-yr-old children. *Journal of Applied Physiology*, 86(4), 1428-1435.
- Sakamaki, R., Toyama, K., Amamoto, R., Liu, C.-J., & Shinfuku, N. (2005). Nutritional knowledge, food habits and health attitude of Chinese university students: A cross sectional study. *Nutrition Journal*, 4(1), 4.
- Sallis, J. F., Calfas, K. J., Nichols, J. F., Sarkin, J. A., Johnson, M. F., Caparosa, S., . . . Alcaraz, J. E. (1999). Evaluation of a university course to promote physical activity: Project GRAD. *Research Quarterly for Exercise and Sport Journal*, 70(1), 1-10.
- Samuelson, G. (1997). Physical status: The use and interpretation of anthropometry. WHO technical report series. *Acta Paediatrica Journal* 86(3), 280-280.
- Santo, A. S., & Golding, L. A. (2003). Predicting maximum oxygen uptake from a modified 3-minute step test. *Research Quarterly for Exercise and Sport Journal*, 74(1), 110-115.
- Sapienza, C. M., & Ruddy, B. H. (2009). *Voice Disorders* (1st ed.). United States, San Diego: Plural.

- Schlosser, E. (2012). *Fast Food Nation: The Dark Side of The All-American Meal* (1st ed.). United States of America, New York: Houghton Mifflin Harcourt.
- Sengupta, P., Chaudhuri, P., & Bhattacharya, K. (2014). Screening obesity by direct and derived anthropometric indices with evaluation of physical efficiency among female college students of Kolkata. *Annals of Medical and Health Sciences Research Journal*, 3(4), 517-522.
- Services, U. D. o. H. H., & Agriculture, U. D. o. (2010). *Dietary guidelines for Americans*, 2010: US Government Printing Office Washington, DC.
- Services, U. S. D. o. H. a. H. (1996). *Physical Activity and Health: A report of the Surgeon General*. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion.
- Severson, K. (2010). Told to eat its vegetables, America orders fries. *New York Times*, 24.
- Shepherd, R. (1999). Social determinants of food choice. *Proceedings of the Nutrition Society Journal*, 58(04), 807-812.
- Sherwood, N. E., & Jeffery, R. W. (2000). The behavioral determinants of exercise: Implications for physical activity interventions. *Annual Review of Nutrition Journal*, 20(1), 21-44.
- Shruthi, B., Hassan, A., & Reddy, B. (2013). The effect of Ramadan fasting on the body composition, blood pressure, heart rate of healthy young adults. *International Journal of Recent Trends in Science And Technology*, 8(1), 31-35.
- Siri, W. E. (1961). Body composition from fluid spaces and density: Analysis of methods. *Techniques for measuring body composition*, 61, 223-244.
- Söderlund, A., Fischer, A., & Johansson, T. (2009). Physical activity, diet and behaviour modification in the treatment of overweight and obese adults: A systematic review. *Perspectives in Public Health Journal*, 129(3), 132-142.
- Soroush, A., Der Ananian, C., Ainsworth, B. E., Belyea, M., Poortvliet, E., Swan, P. D., . . . Yngve, A. (2013). Effects of a 6-month walking study on blood pressure and cardiorespiratory fitness in US and Swedish adults: ASUKI step study. *Asian Journal of Sports Medicine*, 4(2), 114.
- Spark, L. C., Reeves, M. M., Fjeldsoe, B. S., & Eakin, E. G. (2013). Physical activity and/or dietary interventions in breast cancer survivors: A systematic review of the maintenance of outcomes. *Journal of Cancer Survivorship*, 7(1), 74-82.
- Stathopoulou, G., Powers, M. B., Berry, A. C., Smits, J. A., & Otto, M. W. (2006). Exercise interventions for mental health: A quantitative and qualitative review. *Clinical Psychology: Science and Practice Journal*, 13(2), 179-193.

- Stefanick, M. L., Mackey, S., Sheehan, M., Ellsworth, N., Haskell, W. L., & Wood, P. D. (1998). Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *New England Journal of Medicine*, 339(1), 12-20.
- Stone, E. J., McKenzie, T. L., Welk, G. J., & Booth, M. L. (1998). Effects of physical activity interventions in youth: Review and synthesis. *American Journal of Preventive Medicine*, 15(4), 298-315.
- Strong, Malina, Blimkie, Daniels, Dishman, Gutin, . . . Pivarnik. (2005). Evidence based physical activity for school-age youth. *The Journal of Pediatrics*, 146(6), 732-737.
- Strycker, L. A., Duncan, S. C., Chaumeton, N. R., Duncan, T. E., & Toobert, D. J. (2007). Reliability of pedometer data in samples of youth and older women. *International Journal of Behavioral Nutrition and Physical Activity*, 4(1), 4.
- Sugiyama, T., Healy, G. N., Dunstan, D. W., Salmon, J., & Owen, N. (2008). Joint associations of multiple leisure-time sedentary behaviours and physical activity with obesity in Australian adults. *International Journal of Behavioral Nutrition and Physical Activity*, 5(1), 35.
- Swendseid, M. E. (1981). *Essential Amino Acid Requirements: A Review*. Rome: food and agriculture organization of the united nation (FAO) Retrieved from <http://www.fao.org/DOCREP/MEETING/004/M2772E/M2772E00.HTM>.
- Talbot, L. A., Metter, E. J., Morrell, C. H., Frick, K. D., Weinstein, A. A., & Fleg, J. L. (2011). A pedometer-based intervention to improve physical activity, fitness, and coronary heart disease risk in national guard personnel. *Military Medicine Journal*, 176(5), 592-600.
- Taylor, H. L., Jacobs Jr, D. R., Schucker, B., Knudsen, J., Leon, A. S., & Debacker, G. (1978). A questionnaire for the assessment of leisure time physical activities. *Journal of Chronic Diseases*, 31(12), 741-755.
- Ting, L. Y., & Education, P. (2009). *A Comparison on Physiological Variables Between Traditional and Latest Models of Elliptical Cross Trainers in University Female Students*. BA, Hong Kong Baptist University China.
- Tiwari, M. B. (2016). The rubrics. *Journal of Interdisciplinary Studies*, 2(2), 10.
- Tjønnå, A. E., Leinan, I. M., Bartnes, A. T., Jenssen, B. M., Gibala, M. J., Winett, R. A., & Wisløff, U. (2013). Low-and high-volume of intensive endurance training significantly improves maximal oxygen uptake after 10-weeks of training in healthy men. *Plos One Journal*, 8(5), 5.
- Tudor-Locke, C. (2010). Steps to better cardiovascular health: How many steps does it take to achieve good health and how confident are we in this number? *Current Cardiovascular Risk Reports Journal*, 4(4), 271-276.

- Tully, M. A., & Cupples, M. E. (2011). UNISTEP (university students exercise and physical activity) study: A pilot study of the effects of accumulating 10,000 steps on health and fitness among university students. *Journal of Physical Activity and Health*, 8(5), 663-667.
- Tyson, P., Wilson, K., Crone, D., Brailsford, R., & Laws, K. (2010). Physical activity and mental health in a student population. *Journal of Mental Health* 19(6), 492-499.
- UN. (2013). Escalating Violence Threatens Iraq's Progress in the Field of Human Rights. United States, New York: UNAMI.
- Uthaymeen, A.-A. (2005). Dietary Habits and Their Role in The Pathogenesis and Prevention of Diseases. (1st ed). from <http://albalsem.info/enutrition/main.htm>
- Uusitupa, M., Louheranta, A., Lindström, J., Valle, T., Sundvall, J., Eriksson, J., & Tuomilehto, J. (2000). The finnish diabetes prevention study. *British Journal of Nutrition*, 83(S1), S137-S142.
- Von Gruenigen, V. E., Courneya, K. S., Gibbons, H. E., Kavanagh, M. B., Waggoner, S. E., & Lerner, E. (2008). Feasibility and effectiveness of a lifestyle intervention program in obese endometrial cancer patients: A randomized trial. *Gynecologic Oncology Journal*, 109(1), 19-26.
- WES, W. E. S. (2004). Iraq - Higher Education. *Higher Education*, from <http://www.wes.org/ca/wedb/iraq/izhigher.htm>
- Whittemore, R., Melkus, G., Sullivan, A., & Grey, M. (2003). A nurse-coaching intervention for women with type 2 diabetes. *The Diabetes Educator Journal*, 30(5), 795-804.
- WHO. (2004a). Diet. *Global Strategy on Diet, Physical Activity and Health* (1st ed). from <http://www.who.int/dietphysicalactivity/diet/en/>
- WHO. (2004b). Physical Activity. *Global Strategy on Diet, Physical Activity and Health* (2nd ed). from <http://who.int/dietphysicalactivity/pa/en/>
- WHO. (2004c). Promoting Fruit and Vegetable Consumption Around The World. *Global Strategy on Diet, Physical Activity and Health* (1st ed). from <http://www.who.int/dietphysicalactivity/fruit/en/>
- WHO. (2007). Mental Health. *Mental Health* (1st ed). from <http://www.who.int/en/>
- WHO. (2011). Waist Circumference and Waist–Hip Ratio: Report of a WHO Expert Consultation (pp. 47). Switzerland, Geneva: World Health Organization (WHO).
- WHO. (2013). Blood Pressure *Questions and answers about hypertension*, from <http://www.who.int/features/qa/82/ar/>
- WHO, & Consultation, F. E. (2003). Diet, Nutrition and the Prevention of Chronic Diseases (Vol. 916, pp. 160): World Health Organ Tech Rep Ser.

- Wiklund, P., Alen, M., Munukka, E., Cheng, S. M., Yu, B., Pekkala, S., & Cheng, S. (2014). Metabolic response to 6-week aerobic exercise training and dieting in previously sedentary overweight and obese pre-menopausal women: A randomized trial. *Journal of Sport and Health Science*, 3(3), 217-224.
- Wilkinson, L. (1999). Statistical methods in psychology journals: Guidelines and explanations. *American psychologist Journal*, 54(8), 594.
- Willett, W. (2011). *Eat, Drink, and be Healthy: The Harvard Medical School Guide to Healthy Eating*. USA, New York City: Simon and Schuster.
- Wing, R. R., Matthews, K. A., Kuller, L. H., Meilahn, E. N., & Plantinga, P. (1991). Waist to hip ratio in middle-aged women. Associations with behavioral and psychosocial factors and with changes in cardiovascular risk factors. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 11(5), 1250-1257.
- Wing, R. R., Venditti, E., Jakicic, J. M., Polley, B. A., & Lang, W. (1998). Lifestyle intervention in overweight individuals with a family history of diabetes. *Diabetes Care Journal*, 21(3), 350-359.
- Winner, B. C. (2013). *Effects of an Empirically-Based Physical Activity Intervention Aimed to Increase Moderate-to-Vigorous Physical Activity and Improve Body Composition and Blood Pressure in Appalachian Children*. M.Sc., Ohio University, USA. Retrieved from <https://etd.ohiolink.edu/>
- Wood, P. D., Stefanick, M. L., Williams, P. T., & Haskell, W. L. (1991). The effects on plasma lipoproteins of a prudent weight-reducing diet, with or without exercise, in overweight men and women. *New England Journal of Medicine*, 325(7), 461-466.
- Wu, X. Y., Ohinmaa, A., & Veugelers, P. J. (2012). Diet quality, physical activity, body weight and health-related quality of life among grade 5 students in Canada. *Public Health Nutrition Journal*, 15(01), 75-81.
- Yang, Y. J., Kim, M. K., Hwang, S. H., Ahn, Y., Shim, J. E., & Kim, D. H. (2010). Relative validities of 3-day food records and the food frequency questionnaire. *Nutrition Research and Practice Journal*, 4(2), 142-148.