



***EVALUATION OF POLYPHENOL RICH PAPAYA (CARICA PAPAYA L.)
LEAF AND ITS EXTRACTS ON RUMEN MICROBES,
BIOHYDROGENATION AND METHANE PRODUCTION***

SAEID JAFARI

FPV 2016 16



**EVALUATION OF POLYPHENOL RICH PAPAYA (*CARICA PAPAYA* L.)
LEAF AND ITS EXTRACTS ON RUMEN MICROBES,
BIOHYDROGENATION AND METHANE PRODUCTION**

By
SAEID JAFARI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

March 2016

© COPYRIGHT UPM



COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright©Universiti Putra Malaysia



DEDICATION

I dedicate this work to my family. This dissertation exists because of their love, support and prayers.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**EVALUATION OF POLYPHENOL RICH PAPAYA (*CARICA PAPAYA* L.)
LEAF AND ITS EXTRACTS ON RUMEN MICROBES,
BIOHYDROGENATION AND METHANE PRODUCTION**

By

SAEID JAFARI

March 2016

Chairman : Associate Professor Goh Yong Meng, PhD
Faculty : Veterinary Medicine

Methane (CH₄) emission from enteric fermentation is one of the causes of climate changes and global warming. Rumen biohydrogenation (BH) is also detrimental to human health because of increasing the risk of cardiovascular diseases. The effect of increasing inclusion levels of papaya leaf (PL), PL extract (PLE) and PL with different solvent fractions (PLFs) on biohydrogenation (BH) of polyunsaturated fatty acids (PUFA), microbial quantification and fermentation characteristics were studied.

For three *in vitro* studies, PL at different concentrations: 0 (control; CON, 50% alfalfa hay (AH) + 50% concentrate), 15% of AH in substrate replaced by PL (18.75 mg /250 mg dry matter, DM), 25% of AH in substrate replaced by PL (31.25 mg/250 mg DM) and 50% of AH in substrate replaced by PL (62.50 mg/250 mg DM) were mixed with 30 ml of buffered rumen fluid and were incubated for 24 h. CH₄ production (mL/250mg DM) declined at a decreasing rate (linear, $p = 0.03$) with increasing levels of PL. Supplementation of the diet with PL significantly ($P < 0.05$) decreased the rate of BH of LA and C18-PUFA after 24 h of incubation, especially at the higher inclusion rate, resulting in an increased production of rumen BH intermediates such as conjugated linoleic acid (CLA), whereas saturated fatty acid (SFA) decreased in the rumen fluid after 24 h of incubation. Total methanogens ($P < 0.05$) were lower in MPL (-12%) and HPL (-16%) compared to CON. Moreover, there were increases in total bacterial population (HPL: +5%) and *Butyrivibrio fibrisolvens* (MPL: +20; HPL: +23%) among PL treatment groups compared to the CON group.

PLE at different concentrations with no addition of PLE (CON, 0), 5 mg /250mg DM (low leaf extract: LLE), 10 mg /250mg DM (medium leaf extract: MLE) and 15 mg /250mg DM (high leaf extract: HLE) was mixed with 30 ml of buffered rumen fluid and were incubated for 24 h. CH₄ production decreased ($P < 0.05$) with increasing levels of PLE. Supplementation of the diet with PLE significantly ($P < 0.05$) decreased the rate of BH of oleic acid (OA), linoleic acid (LA), α -linolenic acid (LNA) and C18-PUFA compared to CON after 24 h incubation resulting in increased concentrations of BH

intermediates such as vaccenic acid (VA), rumenic acid (RA) and t10c12 CLA. Total protozoa and methanogen population in HLE decreased ($P < 0.05$) compared to CON, but the total bacteria and *Butyrivibrio fibrisolvens* population were higher ($P < 0.05$) in CON compared to the PLE treatment groups.

Different solvent fractions of PL (PLFs; Hexane fraction, Chloroform fraction, Ethyl acetate fraction, Butanol fraction and water fraction) at a concentration of CON with no PLFs (0) and 15mg /250 mg DM which were equivalent to 0 and 60 g/kg DM, respectively were mixed with 30 mL of buffered rumen fluid and were incubated for 24 h. CH_4 production was highest ($P < 0.05$) for CON (7.65) and lowest for chloroform fraction (5.41) compared to other PLFs at 24 h of incubation. Supplementation of the diet with PLFs significantly ($P < 0.05$) decreased the rate of BH of OA, LA and LNA compared to CON after 24 h of incubation. The chloroform fraction had the lowest BH of OA (45.01%) and LA (54.65%) among treatments. Total protozoa and total methanogen populations in PLFs decreased ($P < 0.05$) compared to CON, but the population of *Butyrivibrio fibrisolvens* was higher ($P < 0.05$) in CON compared to PLFs treatment groups. These *in vitro* results indicate that PL, PLE and PLFs supplementation as the substrate decreased BH of PUFA, inhibited rumen methanogenesis without adversely affecting other rumen fermentation characteristics in rumen fermentation liquid.

For *in vivo* study, four fistulated adult Kajang cross-bred goats (39.0 ± 0.75 kg mean \pm standard error) were used in a cross-over design study to test the effect of PL on rumen fermentation characteristics, rumen microbial population and FA composition of rumen fluid obtained at different times of sampling (0, 2, 4, 6 and 8 h). Goats were fed twice daily to the following diets: (CON, 50 % concentrate + 50% AH), 25% AH in basal diet replaced by PL (MPL), and 50% AH in basal diet replaced by PL (HPL). Rumen samples were collected at the end of the experiment (3rd weeks). Supplementation of the diet with PL significantly ($P < 0.05$) decreased the rate of rumen BH of LA after different hours of feeding at the higher inclusion of PL (HPL), resulting in an increased concentration of rumen intermediate products such as CLA e.g. t10 c12, whereas SFA and stearic acid (SA) decreased ($P < 0.05$) in the rumen fluid and blood plasma, respectively. The concentration of LNA was higher ($P < 0.05$) in the blood of goats fed PL especially HPL, compared to the goats receiving only the basal diet with no PL inclusion (CON). The *Butyrivibrio fibrisolvens* population as determined by real-time PCR at the end of the feeding trial was higher ($P < 0.05$) for MPL (+8%) and HPL (+16 %) in the rumen of goats compared to the CON. Methanogenic bacteria (\log_{10} cell/L) decreased ($P < 0.05$) in PL treatment groups.

In summary, this research revealed that supplementation of diets with different levels of PL even as a whole or as the extract (PLE and PLFs) plays an important role in affecting rumen microbial fermentation characteristics resulting in a reduction of CH_4 production (*in vitro*) and BH of FA (*in vitro* and *in vivo*).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

PENILAIAN POLIFENOL YANG TERDAPAT DALAM DAUN POKOK BETIK (*CARICA PAPAYA L.*) DAN EKSTRAK DALAM RUMEN MIKROB, RUMEN BIOHYDROGENATION DAN RUMEN PENGHASILAN METANA

Oleh

SAEID JAFARI

Mac 2016

Pengerusi : Profesor Madya. Goh Yong Meng, PhD
Fakulti : Perubatan Veterinar

Daun betik (PL) telah menjadi makanan haiwan yang berkhasiat dan mengandungi kepelbagaian spektrum metabolit sekunder. Kesan daripada peningkatan tahap kemasukan PL, ekstrak metanol PL (PLE) dan PL dengan pecahan pelarut yang berbeza (PLFs) di biohydrogenation (BH) asid lemak poli tak tepu (PUFA), kuantifikasi dan penapaian ciri-ciri mikrob telah dikaji.

Dalam kajian vitro, PL pada kepekatan yang berbeza PL: 0 (control; CON, 50% alfalfa hay (AH) + 50% pekat), 15% daripada AH dalam substrat digantikan dengan PL (18.75 mg / 250 mg bahan kering, DM), 25% daripada AH dalam substrat digantikan dengan PL (31.25 mg / 250 mg PTT) dan 50% daripada AH dalam substrat digantikan dengan PL (62.50 mg / 250 mg PTT) telah dicampur dengan 30 ml larutan cecair rumen dan dieram selama 24 jam. Pengeluaran metana (CH₄) (mL / 250mg DM) merosot pada kadar yang berkurangan (linear $P = 0.03$) dengan tahap peningkatan PL.

Suplemen diet dengan PL ketara ($P < 0.05$) mengurangkan kadar BH LA dan C18-PUFA selepas 24 jam pengeraman, terutama pada kadar kemasukan yang lebih tinggi, menyebabkan peningkatan penghasilan perantaraan rumen BH seperti asid linoleik (PTK), manakala asid lemak tepu (SFA) menurun dalam cecair rumen selepas 24 jam pengeraman. Jumlah methanogens ($P < 0.05$) adalah lebih rendah pada MPL (-12%) dan HPL (-16%) berbanding dengan CON. Selain itu, terdapat peningkatan dalam jumlah bilangan bakteria (HPL: + 5%) dan fibrisolvens Butyrivibro (MPL: 20; HPL: + 23%) di kalangan kumpulan rawatan PL berbanding dengan kumpulan CON itu. PLE pada kepekatan yang berbeza tanpa penambahan PLE (CON, 0), 5 mg / DM 250mg (ekstrak daun rendah LLE), 10 mg / 250mg DM (ekstrak daun sederhana: MLE) dan 15 mg / 250mg DM (ekstrak daun tinggi: HLE) telah dicampur dengan 30 mL larutan cecair rumen dan dieram selama 24 jam. Pengeluaran CH₄ menurun ($P < 0.05$) dengan peningkatan tahap PLE. Suplemen diet dengan PLE ketara ($P < 0.05$) mengurangkan kadar BH asid oleik (OA), asid linoleik (LA), α -linolenik (LNA) dan C18-PUFA berbanding CON selepas 24 jam pengeraman mengakibatkan peningkatan kepekatan

perantaraan BH seperti asid vaccenic (VA), asid rumenic (RA) dan t 10 c 12 CLA. Jumlah protozoa dan populasi methanogen dalam HLE menurun ($P < 0.05$) berbanding CON, tetapi jumlah bakteria dan populasi *Butyrivibrio fibrisolvens* adalah lebih tinggi ($P < 0.05$) dalam CON berbanding dengan kumpulan rawatan PLE.

Pecahan yang berbeza pelarut PL (PLFs; pecahan heksana, pecahan Chloroform, pecahan Ethyl asetat, pecahan Butanol dan pecahan air) pada kepekatan CON tanpa PLFs (0) dan 15mg / 250 mg DM yang bersamaan dengan 0 dan 60 g / kg DM, masing-masing telah dicampur dengan 30 mL larutan cecair rumen dan dieram selama 24 jam. Antara rawatan PLFs, pecahan kloroform tidak jauh berbeza ($P > 0.05$) berbanding dengan rawatan tanpa penambahan PLFs (CON: 50% AH + 50% pekat) dari segi pengeluaran jumlah gas (mL) dan kadar pengeluaran gas (mL / h).

Pengeluaran CH₄ adalah paling tinggi ($P < 0.05$) bagi CON (7.65) dan terendah untuk pecahan kloroform (5.41) berbanding PLFs lain pada 24 jam penderaman. Suplemen diet dengan PLFs ketara ($P < 0.05$) mengurangkan kadar BH OA, LA dan LNA berbanding CON selepas 24 jam penderaman. Pecahan kloroform mempunyai BH terendah OA (45.01%) dan LA (54.65%) di kalangan rawatan. Jumlah protozoa dan jumlah populasi methanogen dalam PLFs menurun ($P < 0.05$) berbanding CON, tetapi populasi *fibrisolvens Butyrivibrio* adalah lebih tinggi ($P < 0.05$) dalam CON berbanding kumpulan rawatan PLFs. Keputusan dalam vitro ini menunjukkan bahawa PL, PLE dan PLFs suplemen sebagai substrat menurun BH daripada PUFA, menghalang rumen methanogenesis tanpa menjejaskan ciri-ciri rumen penapaian lain dalam cecair rumen penapaian.

Dalam kajian vivo, empat ekor kambing dewasa yang difistula telah digunakan dalam kajian reka bentuk cross-over untuk menguji kesan PL kepada ciri-ciri rumen penapaian, populasi rumen mikrob dan komposisi FA cecair rumen diperolehi pada masa yang berlainan persampelan (0, 2, 4, 6 dan 8 jam). Kambing dengan berat badan (BW) dari (39.0 ± 0.75 kg min \pm SE) telah diberi makan dua kali sehari pada pukul 8:00 dan 5:00 untuk diet yang berikut: (CON, 50% pekat + 50% Hijrah), 25% AH dalam diet basal digantikan dengan PL (MPL), dan 50% AH dalam diet basal digantikan dengan PL (HPL). Darah dan rumen sampel telah dikumpulkan pada akhir eksperimen (minggu ke-3). Keputusan kami menunjukkan bahawa pH cecair rumen adalah lebih rendah ($P < 0.05$) dalam HPL berbanding CON terutama pada 0 dan 6 jam rumen persampelan. Suplemen diet dengan PL ketara ($P < 0.05$) mengurangkan kadar rumen BH LA selepas waktu yang berbeza memberi makan pada kemasukan yang lebih tinggi PL (HPL), mengakibatkan peningkatan kepekatan produk perantaraan rumen seperti CLA contohnya t10 C12, manakala SFA dan asid stearik (SA) menurun ($P < 0.05$) di dalam cecair rumen dan plasma darah, masing-masing. Kepekatan LNA adalah lebih tinggi ($P < 0.05$) dalam membawa darah pada kambing jantan diberi makan PL terutama HPL, berbanding kambing hanya menerima diet basal tanpa kemasukan PL (CON). Populasi *Butyrivibrio fibrisolvens* seperti yang ditentukan oleh masa sebenar PCR pada akhir percubaan makan adalah lebih tinggi ($P < 0.05$) bagi MPL (+ 8%) dan HPL (16%) dalam rumen kambing berbanding CON itu. Bakteria metanogen (\log_{10} cell / L) menurun ($P < 0.05$) dalam kumpulan rawatan PL.

Secara ringkasnya, kajian mendapati suplemen diet dengan pelbagai peringkat PL walaupun secara keseluruhannya atau secara ekstrak (PLE dan PLFs) memainkan peranan yang penting dalam mempengaruhi fermentasi rumen mikrob yang menyebabkan penurunan pengeluaran CH₄ dan BH bagi FA.



ACKNOWLEDGEMENTS

I would like to thank God who until now makes things possible for me. He always protects and helps me and I am able to make this work with him.

I would like to express my sincere appreciation to my supervisory committee, Associate Professor Dr. Goh Yong Meng as chairman and Professor Dr. Mohamed Ali Rajion, Dr Mahdi Ebrahimi and Dr Yusof Hamali as members for their motivation and technical support throughout this research.

I wish to express my sincere gratitude to the Deans of the Faculty of Veterinary Medicine and Institute of Tropical Agriculture, Universiti Putra Malaysia for the use of their facilities and the unlimited assistance from their staff during the course of this study. I would also like to thank staff members of the laboratory of physiology in the Faculty of Veterinary Medicine.

I want to express my appreciation to my friends Dr. Abdoreza Soleimani Farjam, Dr. Mohammad Feseleh Jahromi and Dr. Saied Nikbin.

I would like to thank my dear parents who provided me the opportunity for continuation of my study, and special thanks to my brother and sister. Their love, support and patience have made this work possible.

I certify that a Thesis Examination Committee has met on 29 March 2016 to conduct the final examination of Saeid Jafari on his thesis entitled "Evaluation of Polyphenol Rich Papaya (*Carica papaya* L.) Leaf and its Extracts on Rumen Microbes, Biohydrogenation and Methane Production" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Abd. Wahid bin Haron, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Liang Juan Boo, PhD

Associate Professor
Institute of Tropical Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Md Zuki bin Abu Bakar @ Zakaria, PhD

Professor
Institute of Bioscience
Universiti Putra Malaysia
(Internal Examiner)

Dewi Apri Astuti, PhD

Professor Ir.
Bogor Agricultural University
Indonesia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 25 May 2016

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Goh Yong Meng, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Mohamed Ali Rajion, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Mahdi Ebrahimi, PhD

Postdoctoral Fellow
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Yussof Hamali, PhD

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- This thesis is my original work;
- Quotation, illustrations and citations have been duly referenced;
- This thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- Intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journal, modules, proceeding, popular writing, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No: Saeid Jafari, GS32335

Declaration by Members of Supervisory committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman of
Supervisory

Committee: Associate Professor Goh Yong Meng

Signature: _____

Name of
Member of
Supervisory

Committee: Professor Mohamed Ali Rajion

Signature: _____

Name of
Member of
Supervisory

Committee: Dr. Mahdi Ebrahimi

Signature: _____

Name of
Member of
Supervisory

Committee: Dr. Yussof Hamali

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xix
CHAPTER	
1. GENERAL INTRODUCTION	1
1.1 Background	1
1.2 Hypothesis Statements	2
1.3 Objectives	3
2. LITERATURE REVIEW	4
2.1 The Use of Agricultural Residues in Ruminant Nutrition	4
2.2 Papaya Leaf as an Agricultural By-Product in Ruminant Nutrition	4
2.3 Plant secondary metabolites contents of Papaya leaf	5
2.4 Antibacterial and antioxidant effect of papaya leaf	5
2.5 Plant Containing Secondary Metabolites (PSM) In Agricultural Residues	6
2.6 Phytochemical Screening and Extraction of PSM	6
2.7 Rumen Biohydrogenation in Ruminants	6
2.8 Nutritional Strategies For Reducing Rumen Biohydrogenation	7
2.9 Effect Of Plant Containing Secondary Metabolites on Rumen Biohydrogenation	7
2.10 Rumen Methanogenesis in Ruminants	8
2.11 Nutritional Strategies For Reducing Rumen methanogenesis	8
2.12 Effect of Plant Containing Secondary Metabolites On Rumen Methanogenesis	9
2.13 Effect of Plant Containing Secondary Metabolites on the Fermentation Characteristics of Ruminants	12
2.14 Effects of plants containing secondary metabolites on the rumen microbial population	15
2.15 The role of rumen microorganisms in rumen biohydrogenation and rumen methanogenesis	17
2.16 Effects of plants containing secondary metabolites on antioxidant activity and biochemistry of blood plasma	17
3. EFFECTS OF PAPAYA LEAF, PAPAYA LEAF EXTRACT AND PAPAYA LEAF FRACTIONS ON <i>IN VITRO</i> RUMEN METHANOGENESIS AND RUMENBIOHYDROGENATION	19
3.1 Introduction	19
3.2 Materials and methods	20
3.2.1 Experimental samples and their chemical analyses	

3.2.2	Preparation of fistulated animals	21
3.2.3	Experimental animals	21
3.2.4	Extraction-Fractionation Procedure	23
3.2.5	Phenolic composition of papaya leaf	24
3.2.6	<i>In vitro</i> Incubation	24
3.2.7	Post Incubation Analysis	25
3.2.8	Fatty Acid Analysis and Biohydrogenation Calculation	25
3.2.9	Ruminal Microbial Quantification Using Real Time Polymerase Chain Reaction	26
3.3	Statistical Analysis	27
3.4	Results	27
3.4.1	Effect of PL on Gas and CH ₄ Production	27
3.4.2	Effect of PLE on Gas and CH ₄ Production	28
3.4.3	Effect of PLFs on Gas and CH ₄ Production	29
3.4.4	Rumen Fermentation Parameters	30
3.4.5	Effect of PL on Rumen Fermentation Parameters	31
3.4.6	Effect of PLE on Rumen Fermentation Parameters	31
3.4.7	Effect of PLFs on Rumen Fermentation Parameters	33
3.4.8	Effect of PL on Rumen Concentration of Fatty acids and Biohydrogenation of PUFA	36
3.4.9	Effect of PLE on Rumen Concentration of Fatty acids and Biohydrogenation of	39
3.4.10	Effect of PLFs on Rumen Concentration of Fatty acids and Biohydrogenation of	43
3.4.11	Effect of PL on Rumen Bacteria and Protozoa Population	44
3.4.12	Effect of PLFs on Rumen Bacteria and Protozoa Population	44
3.5	Discussion	47
3.5.1	Effect of PL on Gas and CH ₄ Production	47
3.5.2	Effect of PLE on Gas and CH ₄ Production	48
3.5.3	Effect of PLFs on Gas and CH ₄ Production	49
3.5.4	Effect of PL on Rumen Fermentation Parameters	50
3.5.5	Effect of PLE on Rumen Fermentation Parameters	50
3.5.6	Effect of PLFs on Rumen Fermentation Parameters	51
3.5.7	Effect of PL on Rumen Concentration of Fatty acids and Biohydrogenation of PUFA	51
3.5.8	Effect of PLE on Rumen Concentration of Fatty acids and Biohydrogenation of PUFA	52
3.5.9	Effect of PLFs on Rumen Concentration of Fatty acids and Biohydrogenation	53
3.5.10	Effect of PL on Rumen Bacteria and Protozoa Population	54
3.5.11	Effect of PLE on Rumen Bacteria and Protozoa Population	54
3.5.12	Effect of PLFs on Rumen Bacteria and Protozoa Population	55
3.6	Conclusions	56

4.	IN VIVO STUDY OF PAPAYA (CARICA PAPAYA) LEAF AS A POSSIBLE NATURAL ALTERNATIVE FOR MODULATING RUMEN FERMENTATION CHARACTERISTICS, RUMEN MICROBIAL POPULATION AND FOR REDUCING RUMINAL BIOHYDROGENATION OF C18-POLYUNSATURATED FATTY ACIDS IN GOATS	57
4.1	Introduction	57
4.2	Materials and Methods	58
4.2.1	Experimental Animals and Diets	58
4.2.2	Rumen Sample Collection and Laboratory Analysis	59
4.2.3	Chemical Composition of Diets	59
4.2.4	Fatty Acid Profile in Diet, Rumen fluid and Blood Plasma	61
4.2.5	Calculation of Disappearance of C18-PUFA	61
4.2.6	Rumen Microbial Quantification by Real Time PCR	61
4.2.7	Blood Biochemical Parameters and Antioxidant Activity	61
4.3	Statistical Analysis	62
4.4	Results	62
4.4.1	Rumen Fermentation Parameters	62
4.4.2	Fatty Acid Profile of Rumen fluid and Biohydrogenation of C18-PUFA	64
4.4.3	Fatty acid composition of blood plasma	75
4.4.4	Rumen Microbial Quantification	76
4.4.5	Blood plasma Biochemical Profile and Antioxidant Activity	78
4.5	Discussion	79
4.5.1	Rumen Fermentation Parameters	79
4.5.2	Fatty Acid Profile of the Rumen fluid and Biohydrogenation of C18-PUFA	80
4.5.3	Fatty acid Composition of blood plasma	81
4.5.4	Microbial and Protozoal Population	81
4.5.5	Blood Biochemical Profile and Antioxidant Activity	82
4.6	Conclusions	83
5.	GENERAL DISCUSSION	84
6.	SUMMARY, CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	86
	REFERENCES	88
	APPENDIX	102
	BIODATA OF STUDENT	103
	LIST OF PUBLICATIONS	104

LIST OF TABLES

Table	Page
1. Effects of plant secondary metabolites on methane production in the Rumen	11
2. Effects of plant secondary metabolites on fermentation parameters in the Rumen	14
3. Effects of plants secondary metabolites on microbial populations in Rumen	16
4. Ingredients and chemical composition and fatty acid content of substrates used for the <i>in vitro</i> incubation	22
5. Phenolic content of PL	24
6. Polymerase chain reaction primers used for amplifying target bacteria in the Rumen	27
7. Effect of the addition of different levels of PL on rumen CH ₄ production after <i>in vitro</i> incubation with rumen fluid from goats	28
8. Effect of the addition of different levels of PLE on rumen CH ₄ production after <i>in vitro</i> incubation with rumen fluid from goats	29
9. Effect of PLFs on gas (mL/250mg DM) and CH ₄ (mL/250mg DM) production at 24 h of incubation <i>in vitro</i>	30
10. Effect of the addition of different levels of PL on rumen fermentation parameters after <i>in vitro</i> incubation with rumen fluid from goats	30
11. Effect of the addition of different levels of PLE on rumen fermentation parameters after <i>in vitro</i> incubation with rumen fluid from goats	31
12. Effect of the addition of different levels of PLFs on rumen fermentation parameters after <i>in vitro</i> incubation with rumen fluid from goats	32
13. Effect of the addition of different levels of PL on rumen concentration of fatty acids at 0 h and 24 h of <i>in vitro</i> incubation with rumen fluid from goat.	34
14. Effect of the addition of different levels of PLE on rumen concentration of fatty acids at 0 h and 24 h of <i>in vitro</i> incubation with rumen fluid from goat.	37

15.	Effect of the addition of different levels of PLFs on rumen concentration of fatty acids at 0 h and 24 h of <i>in vitro</i> incubation with rumen fluid from goat.	41
16.	Effect of PL on the population of bacteria and protozoa at 24 h of <i>in vitro</i> incubation	44
17.	Effect of PLE on the population of bacteria and protozoa at 24 h of <i>in vitro</i> incubation	44
18.	Cross-over feeding design	59
19.	Ingredients and chemical composition of the experimental diets	60
20.	<i>In vivo</i> Effect of PL on total VFA at different times of measurements	64
21.	Effect of papaya leaf on blood plasma fatty acid composition of goats (% of total identified fatty acids)	75
22.	<i>In vivo</i> effect of different inclusion level of PL on blood biochemical parameters of goat	75

LIST OF FIGURES

Figure	Page
1. A schematic diagram depicting the extraction - fractionation of PL	23
2. <i>In vitro</i> gas production of control (CON) and different inclusion levels of PL at different times of incubation.	28
3. <i>In vitro</i> gas production of control (CON) and different inclusion levels of PLE at different times of incubation.	29
4. Effect of incremental levels of PL on percentage of BH after 24 h <i>in vitro</i> incubation with rumen inocula from the goats.	35
5. Effect of incremental levels of PL on production of BH intermediate products after 24 h <i>in vitro</i> incubation with rumen inocula from goats.	35
6. Effect of incremental levels of PLE on the percentage of BH after 24 h <i>in vitro</i> incubation with rumen inocula from the goats.	38
7. Effect of incremental levels of PLE on production of biohydrogenation intermediate products after 24 h <i>in vitro</i> incubation with rumen inocula from goats	38
8. Effect of PLFs on rumen biohydrogenation of OA at 24 h of incubation <i>in vitro</i>	39
9. Effect of PLFs on rumen biohydrogenation of LA at 24 h of incubation <i>in vitro</i>	40
10. Effect of PLFs on rumen biohydrogenation of linolenic acid (LNA) at 24 h of incubation <i>in vitro</i>	40
11. Effect of PLFs on the production of rumen biohydrogenation intermediates after 24 h of incubation <i>in vitro</i>	43
12. <i>In vitro</i> effect of PLFs on abundance of total protozoa at 24 h of incubation.	45
13. <i>In vitro</i> effect of PLFs on abundance of total bacteria at 24 h of Incubation	46
14. <i>In vitro</i> effect of PLFs on abundance of <i>Butyrivibrio fibrisolvens</i> at 24 h of incubation.	46
15. <i>In vitro</i> effect of PLFs on abundance of total methanogens at 24 h of Incubation	47

16.	<i>In vivo</i> Effect of PL on rumen pH at different times of measurements.	63
17.	<i>In vivo</i> effect of PL on NH ₃ N concentration at different times of rumen Fermentation	63
18.	<i>In vivo</i> effect of PL on the concentration of C14:0 at different times of sampling.	65
19.	<i>In vivo</i> effect of PL on the concentration of C15:0 at different times of sampling.	65
20.	<i>In vivo</i> effect of PL on the concentration of C15:1 at different times of sampling.	66
21.	<i>In vivo</i> effect of PL on the concentration of C16:0 at different times of sampling.	66
22.	<i>In vivo</i> effect of PL on the concentration of C16:1n-7 at different times of sampling.	67
23.	<i>In vivo</i> effect of PL on the concentration of C18:0 at different times of sampling.	67
24.	<i>In vivo</i> effect of PL on the concentration of C18:1 t-11 at different times of sampling.	68
25.	<i>In vivo</i> effect of PL on the concentration of C18:1 n-9 at different times of sampling.	68
26.	<i>In vivo</i> effect of PL on the concentration of C18:2 n-6 at different times of sampling.	69
27.	<i>In vivo</i> effect of PL on the concentration of C18:3 n-3 at different times of sampling.	69
28.	<i>In vivo</i> effect of PL on the concentration of CLA c9-t11 at different times of sampling.	70
29.	<i>In vivo</i> effect of PL on the concentration of CLA t10c12 at different times of sampling.	70
30.	<i>In vivo</i> effect of PL on the concentration of total SFA at different times of sampling.	71
31.	<i>In vivo</i> effect of PL on the concentration of total MUFA at different times of sampling.	71
32.	<i>In vivo</i> effect of PL on the concentration of n-6 PUFA at different times of sampling.	72

33.	<i>In vivo</i> effect of PL on the concentration of n-3 PUFA at different times of sampling.	72
34.	<i>In vivo</i> effect of PL on the concentration of C18 PUFA at different times of sampling.	73
35.	<i>In vivo</i> effect of PL on the biohydrogenation of OA (%) at different times of Sampling	73
36.	<i>In vivo</i> effect of PL on the biohydrogenation of LA (%) at different times of Sampling	74
37.	<i>In vivo</i> effect of PL on the biohydrogenation of LNA (%) at different times of Sampling	74
38.	<i>In vivo</i> effect of PL on the population of total methanogens	76
39.	<i>In vivo</i> effect of PL on the population of total protozoa	77
40.	<i>In vivo</i> effect of PL on total bacterial population	77
41.	<i>In vivo</i> effect of PL on <i>Butyriovibro fibrisolvens</i> population	78
42.	<i>In vivo</i> effect of PL on the concentration of malonaldehyde (MDA) in blood Plasma	78

LIST OF ABBREVIATIONS

ADF	acid detergent fiber
ANOVA	analysis of variance
BH	biohydrogenation
°C/min	degree centigrade per minute
CH ₄	Methane
CLA	conjugated linoleic acid
CT	condensed tannin
DM	dry matter
FAME	fatty acid methyl esters
G	Gram
GLM	general linear model
GHG	green house gas
H	Hour
HT	hydrolyzed tannin
Kcal	kilo calories
Kg	Kilogram
LA	linoleic acid
LNA	α -linolenic acid
Mm	Millimeter
MDA	Malondialdehyde
ME	metabolizable energy
Min	Minute
mmol/L	millimoles per liter
μ mol/L	micromoles per liter
mg/L	milligrams per liter

MI	Milliliter
MUFA	monounsaturated fatty acids / monoenoic fatty acids
NDF	neutral detergent fiber
OA	oleic acid
PL	papaya leaf
PLE	papaya leaf extract
PLFs	papaya leaf fractions
PSM	plant secondary metabolite
PUFA	polyunsaturated fatty acids
RA	rumenic acid
SA	stearic acid
SEM	standard error of mean
SFA	saturated fatty acids
UFA	unsaturated fatty acids
VA	vaccenic acid
VFA	volatile fatty acid

CHAPTER 1

GENERAL INTRODUCTION

1.1 Background

Animals, particularly ruminants, produce methane (CH₄) from anaerobic fermentation in their gastrointestinal tracts as a pathway for the disposal of metabolic hydrogen produced during microbial metabolism (Jayanegara *et al.*, 2011). Methane accounts for 2-12% loss of dietary gross energy in ruminants and is a potent greenhouse gas (GHG) with a global warming potential 23 times higher than that of carbon dioxide (CO₂) in trapping the heat (Bhatta *et al.*, 2013; IPCC, 2001). Therefore, reducing ruminal CH₄ not only improves the efficiency of nutrient utilization but also helps to protect the environment from global warming. Manipulating the rumen microbial ecosystem to reduce CH₄ emission and nitrogen excretion by ruminants to improve their performance is an important issue for animal nutritionists. There is a need for identifying feed additives with the potential to modify rumen fermentation for enhancing the efficiency of utilization of feed energy while decreasing rumen methanogenesis (Bhatta *et al.*, 2012).

The production of healthy food is an important issue in current ruminant nutrition research. The conversion of dietary polyunsaturated fatty acid (PUFA) to saturated fatty acid (SFA) by ruminants, which is called rumen biohydrogenation (BH) has important health implications for human health. In fact, cardiovascular diseases due to using animal products with high SFA are known to be one of the important causes of death in human populations worldwide (WHO, 2011). Moreover, the transformation of PUFA to SFA results in an increased accumulation of several hydrogenation intermediates in ruminant meat and milk, such as rumenic acid (RA) and vaccenic acid (VA), which are well known for their anti-carcinogenic, anti-atherogenic and anti-oxidative health promoting properties (Durmic *et al.*, 2008). Increased conjugated linoleic acid (CLA) levels in ruminant-derived food products can be achieved by nutritional and managerial practices that facilitate the higher forestomach output of CLA and VA (Chilliard *et al.*, 2007).

Feed additives ranging from antibiotics, CH₄ inhibitors and defaunating agents have been used for years in ruminant nutrition to improve performance and decrease CH₄ production (Benchaar *et al.*, 2008). However, negative consequences of these feed additives have encountered nutritionists with big problems and have stimulated them to search for the safer natural alternatives that would be useful in organic livestock farming. Moreover, the scarcity of feed resources often imposes a major challenge to the development of animal production in the tropics and subtropics. For years ago, it has been shown that tree leaves containing bioactive compounds are the alternative feed source for ruminant and can help to minimize the wide gap between availability and supply of nutrients, and improve the animal growth and productivity (Wanapat *et al.*, 2012). On the other hand, the increasing cost of feeds and climatic problem associated with low production of cereals, which are key in keeping consistent supply

of concentrate used to feed ruminants, may reduce the applicability of using concentrate as a supplement for ruminant. In many arid and semi arid areas of the tropics, the available alternative would be to supplement the ruminant diet using shrubs and tree leaves as they have been proven to be rich in crude protein as compared to straws and grasses. Tree leaves were found to be rich in protein, soluble carbohydrates, minerals and vitamins, and showed great potential as an alternative feed resource (Bakshi and Wadhwa, 2007). Plants containing bioactive compounds have shown to be more effective in reducing rumen methanogenesis and rumen BH (Jayanegara *et al.*, 2011). Plant extracts containing bioactive compounds such as essential oils, tannins, saponins, and flavonoids have been shown to improve rumen metabolism, to decrease the methanogenesis and protein degradation in the rumen and to increase the microbial protein production (Patra and Saxena 2011; Wallace *et al.*, 2004). Plant extracts have also been used for centuries for various purposes such as traditional medicine and food preservatives due to their antimicrobial properties (Busquet *et al.*, 2006; Davidson and Naidu, 2000). Jayanegara *et al.* (2011) showed that phenols from tropical forages were able to modulate BH and methanogenesis processes simultaneously and both towards a desirable direction. The papaya tree is a tree-like plant, a member of the small family *Caricaceae* and widely cultivated for its edible fruits and can be found in all tropical countries and many sub-tropical regions of the world (Canini *et al.*, 2007). Parts of this plant are used for special purposes. Papaya leaf has been shown as a nutrient source and also as the bioactive compound (presence of phenolic acids) (Ayoola and Adeyeye 2010; Canini *et al.*, 2007). Papaya leaf exhibited the highest phenolic content among other parts of the papaya such as unripe papaya fruit, ripe papaya fruit and seed (Maisareh *et al.*, 2014). Presence of phytochemical contents and antimicrobial activities is also confirmed in PL extracts by ethanol, methanol, Ethyl acetate, Petroleum ether, hexane and water solvents (Baskaran *et al.*, 2012). By considering the negative effect of rumen methanogenesis and rumen BH on the environment and health of consumers, respectively, and necessity of a safe feed alternative, the current study designed to address these issues in ruminant with the following hypotheses and objectives:

1.2 Hypothesis Statements

1. Incorporation of diets with PL can increase blood antioxidant activity in the goat *in vivo*.
2. Incorporation of diets with PL, PL extracts and PL fractions will not have any adverse effects on the rumen fermentation and rumen microbial population in the rumen fluid obtained from goat *in vitro*.
3. Incorporation of diets with PL, PL extracts and PL fractions will reduce the rumen BH in the goat.

REFERENCES

- Adeyemo, G.O., & Akanmu, A.M. (2013). Effects of neem (*Azadirachta indica*) and Pawpaw (*Carica papaya*) leave supplementation on performance and carcass characteristics of broilers. Available from: <http://www.enngormix.com/MA-poultry>.
- Agarwal, N., Kamra, D.N., Chaudhary, L.C., & Patra, A.K. (2011). Effect of *Sapindus mukorossi* Extracts on *in vitro* Methanogenesis and Fermentation Characteristics in Buffalo Rumen Liquor. *Journal of Applied Animal Research*, 30, 1- 4.
- Ahmad, N., Fazal, H., Ayaz, M., Abbasi, B.H., Mohammad, I., & Fazal, L. (2011). Dengue fever treatment with *Carica papaya* leaves extracts. *Asian Pacific Journal of Tropical Biomedical*, 1(4), 330-333.
- Anantasook, N., Wanapat, M., & Cherdthong, A. (2014). Manipulation of ruminal fermentation and methane production by supplementation of rain tree pod meal containing tannins and saponins in growing dairy steers. *Journal of Animal Physiology and Animal Nutrition*, 98, 50–55.
- AOAC, (1990). Official Methods of Analysis. 15th edition. Association of Official Analytical Chemists, Herlick, K. (Ed), Arlington, VA, USA, 1230 pp.
- Ayoola, P.B., & Adeyeye, A. (2010). Phytochemical and nutrient evaluation of *Carica papaya* (pawpaw) leaves. *International Journal of Research and Reviews in Applied Sciences*, 5 (3).
- Bakshi, M.P.S., & Wadhwa, M. (2007). Tree leaves as complete feed for goat kids. *Small Ruminant Research*, 69, 74-78.
- Baskaran, C., Ratha, B.V., Velu, S., & Kumaran, K. (2012). The efficacy of *Carica papaya* leaf extract on some bacterial and a fungal strain by well diffusion method. *Asian Pacific Journal of Tropical Disease*, S658-S662.
- Bhatta, R., Uyeno, Y., Tajima, K., Takenaka, A., Yabumoto, Y., Nonaka, I., Enishi, O., & Kurihara, M. (2009). Difference in the nature of tannins on *in vitro* ruminal methane and volatile fatty acid production, and methanogenic archaea and protozoal populations. *Journal of Dairy Science*, 92, 5512–5522.
- Bhatta, R., Saravanan, M., Baruah, L., & Sampath, K. (2012). Nutrient content, *in vitro* ruminal fermentation characteristics and methane reduction potential of tropical tannin-containing leaves. *Journal of Science and Food Agriculture*, 92, 2929–2935.

- Bhatta, R., Saravanan, M., Baruah, L., Sampath, K.T., & Prasad, C.S. (2013). Effect of plant secondary compounds on *in vitro* methane, ammonia production and ruminal protozoa population. *Journal of Applied Microbiology*, 115, 455-465.
- Beauchemin, K.A., & McGinn, S. (2006). Methane emissions from beef cattle: effects of fumaric acid, essential oil, and canola oil. *Journal of Animal Science*, 84, 1489.
- Benchaar, C., Calsamiglia, S., Chaves, A.V., Fraser, G.R., Colombatto, D., McAllister, T.A., & Beauchemin, K.A. (2008). A review of plant-derived essential oils in ruminant nutrition and production. *Animal Feed Science and Technology*, 145, 209–228.
- Bodas, R., Lopez, S., Fernandez, M., García-González, R., Rodríguez, A.B., Wallace, R.J., & Gonzalez, J.S. (2008). *In vitro* screening of the potential of numerous plant species as antimethanogenic feed additives for ruminants. *Animal Feed Science and Technology*, 145, 245–258.
- Bodas, R., Prieto, N., García-González, R., Andrésa, S., Giráldeza, F.J., & López, S. (2012). Manipulation of rumen fermentation and methane production with plant secondary metabolites. *Animal Feed Science and Technology*, 176, 78 – 93.
- Buccioni, A., Minieri, S., Rapaccini, S., Antongiovanni, M., & Mele, M. (2011). Effect of chestnut and quebracho tannins on fatty acid profile in rumen liquid- and solid-associated bacteria: an *in vitro* study. *Animal*, 5, 1521–1530.
- Buccioni, A., Decandia, M., Minieri, S., Molle, G., & Cabiddu, A. (2012). Lipid metabolism in the rumen: New insights on lipolysis and biohydrogenation with an emphasis on the role of endogenous plant factors. *Animal Feed Science and Technology*, 174, 1– 25.
- Buccioni, A., Pauselli, M., Viti, C., Minieri, S., Pallara, G., Roscini, V., Rapaccini, S., Tralbalz, M., Marinucci, P., Conte, G., & Mele, M. (2015). Milk fatty acid composition, rumen microbial population, and animal performances in response to diets rich in linoleic acid supplemented with chestnut or quebracho tannins in dairy ewes. *Journal of Dairy Science*, 98, 1145–1156.
- Bueno, C., Brandi, R.A., Franzolina, A., Benete, G., Fagundes, G.M., Abdalla, A.L., Louvandini, H., & Muir, J.P. (2015). *In vitro* methane production and tolerance to condensed tannins in five ruminant species. *Animal Feed Science and Technology*, 205, 1–9.
- Busquet, M., Calsamiglia, S., Ferret, A., & Kamel, C. (2006). Plant extracts affect *in vitro* rumen microbial fermentation. *Journal of Dairy Science*, 89, 761–771.
- Cabiddu, A., Salis, L., Tweed, J.K., Molle, G., Decandia, M., & Lee, M. (2010). The influence of plant polyphenols on lipolysis and biohydrogenation in dried

forages at different phenological stages: *in vitro* study. *Journal of Science and Food Agriculture*, 90, 829–835.

- Caldeira, A., Belo, C., Santos, M., Vazques, A., & Portugal, A.V. (2007). The effect of body condition score on blood metabolites and hormonal profiles in ewes. *Small Ruminant Research*, 68(3) 233–241.
- Canini, A., Alesiani, D., Arkangelo, G., & Tagliatesta, P. (2007). Gas chromatography–mass spectrometry analysis of phenolic compounds from *Carica papaya* L. leaf. *Journal of Food Composition and Analysis*, 20, 584–590.
- Cardozo, P.W., Calsamiglia, S., Ferret, A., & Kamel, C. (2004). Effects of natural plant extracts on ruminal protein degradation and fermentation profiles in continuous culture. *Journal of Animal Science*, 82, 3230–3236.
- Carreno, D., Hervás, G., Toral, P.G., Belenguer, A., & Frutos, P. (2015). Ability of different types and doses of tannin extracts to modulate *in vitro* ruminal biohydrogenation in sheep. *Animal Feed Science and Technology*, 202, 42–51.
- Castagnino, P.S., Messanaa, J.D., Fiorentini, G., De Jesusa, R.B., San Vitoa E., Carvalhoa, I.P.C., & Berchielli, T.T. (2015). Glycerol combined with oils did not limit biohydrogenation of unsaturated fatty acid but reduced methane production *in vitro*. *Animal Feed Science and Technology*, 201, 14–24.
- Cherdthong, A., Wanapat, M., Kongmun, P., Pilajun, R., & Khekornsart, P. (2010). Rumen fermentation, microbial protein synthesis and cellulolytic bacterial population of swamp buffaloes as affected by roughage to concentrate ratio. *Journal of Animal Veterinary Advances*, 9, 1667–1675.
- Chilliard, Y., Glasser, F., Ferlay, A., Bernard, L., Rouel, J., & Doreau, M., (2007). Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *European Journal of Lipid Science and Technology*. 109, 828–855.
- Cieslak, A., Szumacher-Strabel, Stochmal., M.A., & Oleszek, W. (2013). Plant components with specific activities against rumen methanogens. *Animal*, 7, 253–265.
- Davidson, P.M., & Naidu, A.S. (2000). Phyto-phenols. Natural Food Antimicrobial Systems. A. S. Naidu, ed. CRC Press, Boca Raton, FL, pp 265–293.
- Durmic, Z., McSweeney, C.S, Kemp, G.W., Hutton, P., Wallace, R.G., & Vercoe, P.E. (2008). Australian plants with potential to inhibit bacteria and processes involved in ruminal biohydrogenation of fatty acids. *Animal Feed Science and Technology*, 145, 271–284.

- Ebrahimi, M., Rajion, M. A., Goh, Y. M., & Sazili, A. Q. (2012). Impact of different inclusion levels of oil palm (*Elaeis guineensis* Jacq.) fronds on fatty acid profiles of goat muscles. *Journal of Animal Physiology and Animal Nutrition*, 96(6), 962–969.
- Ebrahimi, M. (2012). Production of omega-3 polyunsaturated fatty acid- enriched chevon using treated oil palm (*ELAEIS GUINEENSIS JACQ.*) frond silage. PhD thesis, universiti putra Malaysia, Malaysia, pp. 28.
- Edward, E., & Fredy B. (2013). An Overview of Global Papaya Production, Trade, and Consumption. University of Florida, *IFAS extension*, 913.
- EL-Sabagh, M.R., Abd - Eldaim, M.A., Mahboub, D.H., & Abdel-Daim, M. (2014). Effects of spirulina platensis algae on growth performance, antioxidative status and blood metabolites in fattening lambs. *Journal of Agricultural Science*, 6 (3).
- Evans, J.D., & Martin, S.A. (2000). Effects of thymol on ruminal microorganisms. *Current Microbiology*, 41, 336-340.
- Fievez, V., Babayemi, O. J., & Demeyer, D. (2005). Estimation of direct and indirect gas production in syringes: A tool to estimate short chain fatty acid production that requires minimal laboratory facilities. *Animal Feed Science and Technology*, 123(1), 197–210.
- Fievez, V., Vlaeminck, B., Jenkins, T., Enjalbert, F., & Doreau, M. (2007). Assessing rumen biohydrogenation and its manipulation *in vivo*, *in vitro* and *in situ*. *European journal of lipid science and Technology*, 109,740–756.
- Folch, J., Lees, M., & Sloane Stanely, G. H. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry*, 226(1), 497–50.
- Ghaffari, M.H., Tahmasbi, AM., Khorvash, M., Naserian, .AA., Ghaffari, A.H., & Valizadeh, H. (2013). Effects of pistachio by-products in replacement of alfalfa hay on populations of rumen bacteria involved in biohydrogenation and fermentative parameters in the rumen of sheep. *Journal of Animal Physiology and Animal Nutrition*. DOI: 10.1111/jpn.12120.
- Ghasemi, S., Naserian, A.A., Valizadeh, R., Tahmasebi, A.M., Vakili, A.R., Behgar, M., & Ghovvati, S. (2012). Inclusion of pistachio hulls as a replacement for alfalfa hay in the diet of sheep causes a shift in the rumen cellulolytic bacterial population. *Small Ruminant Research*, 104, 94–98.
- Griinari, J. M., & Bauman, D. E. (1999). Biosynthesis of conjugated linoleic acid and its incorporation into meat and milk in ruminants. Yurawecz, M. P. , Mossoba, M., Kramer, J. K. G., Nelson, G., and Pariza, M. W. (Eds), *Advances in Conjugated Linoleic Acid Research, Vol. 1* Champaign, IL, USA: AOCS Press, pp.180–200.

- Gunal, M., Ishlak, A., AbuGhazaleh, A.A., & Khattab, W. (2014). Essential oils effect on rumen fermentation and biohydrogenation under *in vitro* conditions. *Czech Journal of Animal Science*, 10, 450–459.
- Goel, G., Makkar, H.P.S., & Becker, K., (2008). Changes in microbial community structure, methanogenesis and rumen fermentation in response to saponin-rich fractions from different plant materials. *Journal of Applied Microbiology*, 105, 770–777
- Goiri, I., Induraina, G., Insausti, K., Sarries, V., & Garcia-Rodriguez, A. (2010). Ruminant biohydrogenation of unsaturated fatty acids *in vitro* as affected by chitosan. *Animal Feed Science and Technology*, 159, 35–40.
- Gurr, M. I., & Harwood, J. L. (1991). *Lipid Biochemistry: An Introduction*, 4th edition. London: Chapman and Hall, 320 pp.
- Hart, K.J., Yanez-Ruiz, D.R., Duval, S.M., McEwan, N.R., & Newbold, C.J. (2008). Plant extracts to manipulate rumen fermentation. *Animal Feed Science and Technology*, 147, 8–35.
- Hatew, B., Stringano, E., Harvey, M., Hendriks, W.H., Hayot Carboner, C., Smith. & Pellikaan, W.F. (2015). Impact of variation in structure of condensed tannins from sainfoin (*Onobrychis viciifolia*) on *in vitro* ruminal methane production and fermentation characteristics. *Journal of Animal Physiology and Animal Nutrition*. DOI: 10.1111/jpn.12336.
- Hook S.E., Wright, A.D.G., & McBride, B.W. (2010). Methanogens: methane producers of the rumen and mitigation strategies. *Archaea*, 1 <http://dx.doi.org/10.1155/2010/945785>
- Jafari, S., Alizadeh, A., & Imani, A. (2011). Nutritive value of different varieties of almond (*Prunus dulcis*) hulls. *Research Opinion in Animal and Veterinary Sciences*, 2221-1896.
- Jayanegara, A., Togtokhbayar, N., Makkar, H. P. S., & Becker, K. (2009). Tannins determined by various methods as predictors of methane production reduction potential of plants by an *in vitro* rumen fermentation system. *Animal Feed Science and Technology*, 150, 230–237.
- Jayanegara, A., Goel, G., Makkar, H.P.S., & Becker, K. (2010). Reduction in Methane Emissions from Ruminants by Plant Secondary Metabolites: Effects of Polyphenols and Saponins. *Food and Agriculture Organization of the United Nations*, 151-157.
- Jayanegara, A, Kreuzer, M, Wina, E and Leiber, E. (2011). Significance of phenolic compounds in tropical forages for the ruminal bypass of polyunsaturated fatty acids and the appearance of biohydrogenation intermediates as examined *in vitro*. *Animal Production Science*, 51, 1127–1136.

- Jayanegara, A., Kreuzer, M., & Leiber, F. (2012). Ruminant disappearance of polyunsaturated fatty acids and appearance of biohydrogenation products when incubating linseed oil with alpine forage plant species *in vitro*. *Livestock Science*, 147, 104–112.
- Jayanegara, A., Goel, G., Makkar, P. S. H., & Becker, K. (2015). Divergence between purified hydrolysable and condensed tannin effects on methane emission, rumen fermentation and microbial population *in vitro*. *Animal Feed Science and Technology*, 209, 60–68.
- Jenkins, T. C. (2002). Lipid transformations by the rumen microbial ecosystem and their impact on fermentative capacity. pp 103-117 in *Gastrointestinal Microbiology in Animals*, S. A. Martin (Ed.), Research Signpost, Kerala, India.
- Jenkins, T. C., Wallace, R. J., Moate, P. J., & Mosley, E. E. (2008). Board-invited review: Recent advances in biohydrogenation of unsaturated fatty acids within the rumen microbial ecosystem. *Journal of Animal Science*, 86, 397-412.
- Johnson and Johnson, K.A. & Johnson, D.E. (1995). Methane emissions from cattle. *Journal of Animal Science*, 73, 2483–2492.
- Jouany, J., & Morgavi, D.P. (2007). Use of ‘natural’ products as alternatives to antibiotic feed additives in ruminant production. *Animal*, 1, 1443–1466.
- Halim, s.z., Abdollah, N.R., Afzan, A., Abdollah Rashid. B.A., Jantan. A., & Ana Ismaeil, Z. (2011). Acute toxicity study of *Carica papaya* leaf extract in Sprague Dawley rats. *Journal of Medicinal Plants Research*, 5, 1867-1872.
- Harfoot, G., & Hazlewood, G. P. (1997). Lipid metabolism in the rumen. In: Hobson, P. N., (Ed), *The Rumen Microbial Ecosystem*, Elsevier, page (382-426), London, UK.
- Hartmann, T. (2007). From waste products to ecochemicals: Fifty years research of plant secondary metabolism. *Phytochemistry*, 68, 2831–2846.
- Head, W.F. Jr., & Lauter, M. (1956). Phytochemical examination of the leaves of *Carica papaya* L. *Economic Botany*, 258-26.
- Heidarian, M., Tyagi, V., Ebrahimi, A.K., & Mohini, S.H.M., (2013). Plant extracts enhanced ruminal CLA concentration, *in vitro*. *Journal of Animal Feed Science and Technology*, 22, 219–228.
- Heidarian, M., Ebrahimi, S.H., & Tyagi, A.K. (2015). The effect of cumin (*Cuminum cyminum*) seed extract on the inhibition of PUFA biohydrogenation in the rumen of lactating goats via changes in the activity of rumen bacteria and linoleate isomerase enzyme. *Small Ruminant Research*, 125, 56–63.

- Hess, H.D., Kreuzer, M., Diaz, T.E., Lascano, C.E., Carulla, J.E., Soliva, C., & Machmüller, A. (2003). Saponin rich tropical fruits affect fermentation and methanogenesis in faunated and defaunated rumen fluid. *Animal Feed Science and Technology*, 109, 79–94.
- Hristov, A. N., T. A. McAllister, F. H., Van Herk, K. J., Cheng, C. J., & Newbold, P. R. (1999). Effect of *Yucca schidigera* on ruminal fermentation and nutrient digestion in heifers. *Journal of Animal Science*, 77, 2554–2563.
- Hungate, R.E. (1966). The rumen and its microbes. Academic Press, New York. 533 pp.
- Ikram, E.H.K., Stanley, R., Netzel, M., & Fanning, K., (2015). Phytochemicals of papaya and its traditional health and culinary uses – A review. *Journal of Food Composition and Analysis*, 41, 201–211.
- IPCC.(2001). Intergovernmental panel on climate change. (2001).The Scientific Background, vol 94 ed. Houghton, J.T. Cambridge: Cambridge University Press, pp. 1–94.
- Kamra, D.N., Agarwal, N., & Chaudhary, L.C. (2006).Inhibition of ruminal methanogenesis by tropical plants containing secondary compounds. *International Congress Series*, 1293, 156–163.
- Kamra, D.N., Patra, A.K., & Chatterjee, P.N. (2008). Effect of plant extract on methanogenesis and microbial profile of the rumen of buffalo: a brief overview. *Australian Journal of Experimental Agriculture*, 48,175–178.
- Kamra, D.N., Perera, A.N.F., & Xie, Z.Q. (2009).The use of plant secondary metabolites as medicine and feed additives for eco-friendly livestock production. In: Scaife JR, Vercoe PE (eds) Harvesting knowledge – pharming opportunities. *British Society of Animal Science*, pp 8–20.
- Kamra, D.N., Pawar, M., & Singh, B. (2012). Effect of Plant Secondary Metabolites on Rumen Methanogens and Methane Emissions by Ruminants.A.K. Patra (ed.), *Dietary Phytochemicals and Microbes*, chapter 12.
- Keeney, M. (1970). Lipid metabolism in the rumen, In: Phillipson A. T. (Ed), *Physiology of Digestion and Metabolism in the Ruminant*, Proceedings of the 3rd International Symposium, Cambridge, England, August, England, 636 pp.
- Kemp, P. L. D. (1984). Hydrogenation *in vitro* of alpha-linolenic acid to stearic acid by mixed cultures of pure strains of rumen bacteria. *Journal of General Microbiology*, 130, 527–533.
- Kim, J. E., Sharon, A.H., Michael R.F., & Nigel, D.S. (2010).Dietary Transformation of Lipid in the Rumen Microbial Ecosystem. *Asian-Australian Journal of Animal Sciences*, 22(9), 1341-1350.

- Koike, S., & Kobayashi, Y. (2001). Development and use of competitive PCR assays for the rumen cellulolytic bacteria: *Fibrobacter succinogenes*, *Ruminococcus albus* and *Ruminococcus flavefaciens*. *FEMS Microbiology Letters*, 204, 361-366.
- Krishna, K.L., Paridhavi, M., & Patel, J.A (2008). Review on nutritional, medicinal and pharmacological properties of Papaya (*Carica papaya*). *Natural Product Radiance*, 7(4), 364-373.
- Lee, O.H., Lee, B.Y., Lee, J., Lee, H.B., Son, J.Y., Park, C.S., Shetty, K., & Kim, Y.C., (2009). Assessment of phenolics-enriched extract and fractions of olive leaves and their antioxidant activities. *Bioresource Technology*, 100, 6107-6113.
- Lee, Y (2013).Effect of pH on Conjugated Linoleic Acid (CLA) Formation of linolenic acid biohydrogenation by ruminal microorganisms. *Journal of Microbiology*, 51 (4), 471-476.
- Liu, H., Vaddella, V., & Zhou, D. (2012). Effects of chestnut tannins and coconut oil on growth performance, methane emission, ruminal fermentation, and microbial populations in sheep. *Journal of Dairy Science*, 94, 6069-6077.
- Lourenco, M., Ramos-Morales, E., &Wallace, R.J. (2010).The role of microbes in rumen lipolysis and biohydrogenation and their manipulation. *International Journal of Animal Biology science*, 4, 1008-1023.
- Machmuller, A., Soliva, C. R., & Kreuzer, M. (2003).Methane-suppressing effect of myristic acid in sheep as affected by dietary calcium and forage proportion. *British Journal of Nutrition*, 3, 529-540.
- Maia, M. R., Chaudhary, L. C., Figueres, L., & Wallace, R. J. (2007). Metabolism of polyunsaturated fatty acids and their toxicity to the microflora of the rumen. *Antonie van Leeuwenhoek*, 91, 303-314.
- McIntosh, F.M., Williams, P., Losa, R., Wallace, R.J., Beever, D.A., & Newbold, C.J. (2003).Effects of essential oils on ruminal microorganism and their protein metabolism. *Applied and Environmental Microbiology*, 69, 5011-5014.
- Makkar, H.P.S., & Becker, K. (2003).Nutritional value and whole and ethanol antinutritional components of extracted *Moringa oleifera* leaves. *Animal Feed Science and Technology*, 63, 21 I-228.
- Min, B. R., Attwood, G. T., McNabb, W.C., Molan, A. L., & Barry, T. N. (2004).The effect of condensed tannins from *Lotus corniculatus* on the proteolytic

- activities and growth of rumen bacteria. *Animal Feed Science and Technology*, 121, 45–58.
- Monteny, G.J., Bannink, A., & Chadwick, D. (2006). Greenhouse gas abatement strategies for animal husbandry. *Agriculture Ecosystem Environment*, 12, 163–170.
- Ramos-Morales, E., Martínez-Fernández, G., Abecia, L., Martín-García, A.I., Molina-Alcaide, E., & Yáñez-Ruiz, D.R. (2013). Garlic derived compounds modify ruminal fatty acid biohydrogenation and induce shifts in the *Butyrivibrio* community in continuous-culture fermenters. *Animal Feed Science and Technology*, 184, 38 – 48.
- Naumann, H.D., Lambert, B.D., Armstrong, S.A., Fonseca, M. A., Tedeschi, L.O., & Muir, J.P. 2015. Effect of replacing alfalfa with panicle-tick clover or *sericea lespedeza* in corn-alfalfa-based substrates on *in vitro* ruminal methane production. *Journal of Dairy Science*, 98, 3980–3987.
- Navid, S., Sheikhlari, A., & Kaveh, K. (2011). Influence of combination of vitamin D3 and papaya leaf on meat quality and spent layer hen. *Agricultural Journal*, 6(4), 197-200.
- Newbold, C.J., Lassalas, B., & Jouany, J.P. (1995). The importance of methanogens associated with ciliate protozoa in ruminal methane production *in vitro*. *Letters in Applied Microbiology*, 21, 230–234.
- Newbold, C.J., McIntosh, F.M., Williams, P., Losa, R., & Wallace, R.J., (2004). Effects of a specific blend of essential oil compounds on. *Animal Feed Science and Technology*. 114, 105–112.
- NRC.(2007). Nutrient requirements of small ruminant. Sixth edition, National Academy Press, Washington D.C., USA. 384 pp.
- Otsuki, N., Dang, N.H., Kumagai, E., Kondoc, A., Iwataa, S., & Morimoto, C. (2010). Aqueous extract of *Carica papaya* leaves exhibits anti-tumor activity and immunomodulatory effects. *Journal of Ethnopharmacology*, 127, 760–76.
- Patil, T., Patil, S., Patil, A., & Patil, S.H. (2014). *Carica Papaya* Leaf Extracts – An Ethnomedicinal Boon. *International Journal of Pharmaceutical Phytochemistry Research*, 6(2), 260-265.
- Patra, A.K., & Saxena, J. (2009). A new perspective on the use of plant secondary metabolites to inhibit methanogenesis in the rumen. *Phytochemistry*, 71, 1198–1222.

- Patra, M., & Saxena, J. (2011). Exploitation of dietary tannins to improve rumen metabolism and ruminant nutrition. *Journal of Food and Agriculture*, 91, 24–37.
- Patra, A.K., & Yu, G. (2014). Combinations of nitrate, saponin, and sulfate additively reduce methane production by rumen cultures *in vitro* while not adversely affecting feed digestion, fermentation or microbial communities. *Bioresource Technology*, 155, 129–135.
- Pedraza, R. (1988). Use of *in vitro* gas production technique to assess the contribution of both soluble and insoluble fractions on the nutritive value of forages. PhD thesis, Universidad de Camagüey, Cuba.
- Pen, B., Sar, C., Mwenya, B., Kuwaki, K., Morikawa, R., & Takahashi, J. (2006). Effects of *Yucca schidigera* and *Quillaja saponaria* extracts on *in vitro* ruminal fermentation and methane emission. *Animal Feed Science and Technology*, 129, 175–186.
- Pinski, B., Günal, M., & AbuGhazaleh, A.A. (2015). The effects of essential oil and condensed tannin fermentation. *Animal Production Science*, <http://dx.doi.org/10.1071/AN15069>.
- Porter, L.J., & Hrstich, B.G. (1986). The conversion of proanthocyanidins and prodelphinidins to cyanidin and delphinidin. *Phytochemistry*, 25, 223–230.
- Ørskov, E. R. (1998). Feed evaluation with emphasis on fibrous roughages and fluctuating supply of nutrients: a review. *Small Ruminant Research*, 28, pp. 1–8.
- Rajion, M. A., McLean, J. G., Cahill, R. N., N.P., & Cahill, R. (1985). Essential fatty acids in the fetal and newborn lamb. *Australian Journal of Biological Sciences*, 38(1), 33–40.
- Rira, M., Chentli, A., Boufener, S., & Boussebou, H. (2015). Effects of plants containing secondary metabolites on ruminal methanogenesis of sheep *in vitro*. *Energy Procedia*, 74, 15 – 24.
- Riss, J., econd'e, K. D., & Sutra, T. (2007). Phycobiliprotein C-phycoyanin from *Spirulina platensis* is powerfully responsible for reducing oxidative stress and NADPH oxidase expression induced by an atherogenic diet in hamsters. *Journal of Agriculture and Food Chemistry*, 55, 7962–7967.
- Runnie, I., Salleh, M.N., Mohamed, S., Head, R.J., & Abeywardena, M.Y. (2004). Vasorelaxation induced by common edible tropical plant extracts in isolated rat aorta and mesenteric vascular bed. *Journal of Ethnopharmacology*, 92, 311–316.

- Samanta, A.K., Singh, K.K., Das, M.M., Maity, S.B., & Kundu, S.S. (2003). Effect of complete feed block on nutrient utilization and rumen fermentation in Barbari goats. *Small Ruminant Research*, 48, 95–102.
- Santos-Buelga, C., Gonzalez-Manzano, S., Dueñas, M., & Gonzalez-Paramas, A.M. (2012). Extraction and isolation of phenolic compounds. *Methods in Molecular Biology*, 864, 427-464.
- SAS (2003). SAS User's Guide: Statistics, Version 9.1 edition. NC, USA: SAS Inst., Inc., Cary.
- Scalbert, A., Johnson, I.T., & Saltmarsh, M. (2005). Polyphenols: antioxidants and beyond. *American Journal of Clinical Nutrition*, 81, 215S–7S.
- Skenjana, A., (2011). The potential nutritive value of waste products from the subtropical fruit processing industry as livestock feed. MSc Agric. Thesis, University of Pretoria, South Africa.
- Soliva, C.R., Meile, L., Cieslak, A., Kreuzer, M., & Machmuller, A. (2004) Rumen simulation technique study on the interactions of dietary lauric and myristic acid supplementation in suppressing ruminal methanogenesis. *British Journal of Nutrition*, 92, 689–700.
- Solorzano, L. (1969). Determination of ammonia in natural waters by the phenylhypochlorite method. *Limnology and Oceanography*, 14(5), 799–801.
- Spanghero, M., Zanfi, C., Fabbro, E., Scicutella, N., & Camellini, C. (2008). Effects of a blend of essential oils on some end products of *in vitro* rumen fermentation. *Animal Feed Science and Technology*, 145, 364–374.
- Surai, P.F. (2014). Polyphenol compounds in the chicken/animal diet: from the past to the future. *Journal of Animal Physiology and Animal Nutrition*, 98, 19–31
- Sylvester, J.T., Karnati, S.K.R., Yu, Z., Morrison, M., & Firkins, J.L. (2004). Development of an assay to quantify rumen ciliate protozoal biomass in cows using real-time PCR. *The Journal of Nutrition*, 134 (12), 3378-3384.
- Szumacher-Strabel, M., & Cieslak, A. (2010). Potential of phytofactors to mitigate rumen ammonia and methane production. *Journal of Animal Feed Science*, 19, 319–337.
- Tan, H.Y., Sieo, C.C., Abdullah, N., Liang, J.B., Huang, X.D., & Ho, Y.W., (2011). Effects of condensed tannins from *Leucaena* on methane production, rumen fermentation and populations of methanogens and protozoa *in vitro*. *Animal Feed Science and Technology*, 169, 185–193.

- Tavendale, M.H., Meagher, L.P., Pacheco, D., Walker, N., Attwood, G.T., & Sivakumaran, S. (2005) Methane production from *in vitro* rumen incubations with *Lotus pedunculatus* and *Medicago sativa*, and effects of extractable condensed tannin fractions on methanogenesis. *Animal Feed Science and technology*, 123,403–419.
- Tiwari, P., Kumar, B., Kaur, M., Kaur, G., & Kaur, H. (2011). Phytochemical screening and Extraction: A Review. *Internationale Pharmaceutica Scientia*, 1(1).
- Toral, P.G., Hervás, G., Bichi, E., Belenguer, A., & Frutos, P., (2011). Tannins as feed additives to modulate ruminal biohydrogenation: effects on animal performance, milk fatty acid composition and ruminal fermentation in dairy ewes fed a diet containing sunflower oil. *Animal Feed Science and technology*, 164, 199–206.
- Troegeler-Meynadier, A., Bret-Bennis, L., & Enjalbert, F. (2006). Rates and efficiencies of reactions of ruminal biohydrogenation of linoleic acid according to pH and polyunsaturated fatty acids concentrations. *Reproduction Nutrition Development*, 26, 713–724.
- Tug, T., Karatas, F., Terzi, S.M., & Ozdemir, N. (2005). Comparison of serum malondialdehyde levels determined by two different methods in patients with copd: HPLC or TBARS methods. *Lab Medicine*.36 (1).
- Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991).Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74(10), 3583–3597.
- Vasta, V., Makkar, H. P. S., Mele, M., & Priolo, A. (2009). Ruminal biohydrogenation as affected by tannins *in vitro*. *The British Journal of Nutrition*, 102(1), 82–92.
- Vasta, V., Yanez-Ruiz, D.R., Mele, M., Serra, A., Luciano, G., Lanza, M., Biondi, L., Priolo, A. (2010). Bacterial and protozoal communities and fatty acid profile in the rumen of sheep fed a diet containing added tannins. *Applied Environmental Microbiology*, 76, 2549–2555.
- Vasta, V., Pagano, R. I., Luciano, G., Scerra, M., Caparra, P., Foti, F., Cilione, C., Biondi, L., Priolo, A., & Avondo, M. (2012). Effect of morning vs. Afternoon grazing on intramuscular fatty acid composition in lamb. *Meat Science*, 90:93.
- Vuong, Q.V., Hiruna, S., Roach, P.D., Bowyer, M.C., Phillips, P.A., & Scarlett, C.J. (2013). Effect of extraction conditions on total phenolic compounds and

antioxidant activities of *Carica papaya* leaf aqueous extracts. *Journal of the Herbal Medicine*, 3, 104–111.

- Yanez-Ruiz, D.R., Martin Garcia, A.I., Moumen, A., & Molina Alcaide, E. (2004). Ruminant fermentation and degradation patterns, protozoa population and urinary purine derivatives excretion in goats and wethers fed diets based on olive leaves. *Journal of Animal Science*, 82, 3006 – 3014.
- Yongjae, L. (2013). Effect of pH on Conjugated Linoleic Acid (CLA) Formation of Linolenic Acid Biohydrogenation by Ruminant Microorganisms. *Journal of Microbiology*, 54, 471–476.
- Yusha'u, M., Onuorah, F.C., & Murtala, Y. (2009). *In vitro* sensitivity pattern of some urinary tract isolates to *Carica papaya* extracts. *Bayero Journal of Pure and Applied Sciences*, 2(2), 75 - 78.
- Vakili, A.R., Khorrami, B., Danesh Mesgaran, M., & Parand, E. (2013). The effects of thyme and cinnamon essential oils on performance, rumen fermentation and blood metabolites in Holstein calves consuming high concentrate diet. *Asian Australasian Journal of Animal Science*, 26 (7), 935 - 944.
- Varlyakov, I., Radev, V., Slavov, T., & Ganchev, G. (2013). Blood parameters in yearling sheep fed Paulownia (*Paulownia spp.*) leaves. *Agricultural Science and Technology*, 5 (4), 405-409.
- Wallace, R. J., McEwan, N. R., McIntosh, F. M., Teferedegne, B., & Newbold, C. J. (2002). Natural products as manipulators of rumen fermentation. *Asian-Australasian Journal of Animal Science*, 15(10), 1458–1468.
- Wallace, R.J. (2004). Antimicrobial properties of plant secondary metabolites. *Proceeding of Nutrition Society*, 63, 621–629.
- Wanapat, M., Cherdthong, A., Pakdee, P., & Wanapat, S. (2008). Manipulation of rumen ecology by dietary lemongrass (*Cymbopogon citrates Stapf.*) powder supplementation. *Journal of Animal Science*, 86, 3497–3503.
- Wanapat, M., Kongmun, P., Pongchompu, O., Cherdthong, A., Khejornsart, P., Pilajun, R., & Kaenpakdee, S. (2012). Effects of plants containing secondary compounds and plant oils on rumen fermentation and ecology. *Tropical Animal Health and Production*, 44, 399–405.
- Wanapat, M., Kang, S., & Polyorach, S. (2013). Development of feeding systems and strategies of supplementation to enhance rumen fermentation and ruminant production in the tropics. *Journal of Animal Science and Biotechnology*, 4, 32.

- Wanapat, M., Chanthakhoun, V., Phesatcha, K., & Kang, S. (2014). Influence of mangosteen peel powder as a source of plant secondary compounds on rumen microorganisms, volatile fatty acids, methane and microbial protein synthesis in swamp buffaloes. *Livestock Science*, 162, 126–133.
- Wang, X.F., Mao, S.Y., Liu, J.H., Zhang, L.L., Cheng, Y.F., Jin, & Wand Zhu, W.Y. (2011). Effect of the gynosaponin on methane production and microbe numbers in a fungusmethanogen co-culture. *Journal of Animal and Feed Sciences*, 20, 272–284.
- WHO, World Health Organization (2003). Diet, nutrition and the prevention of chronic diseases. Technical Report Series 916. WHO: Geneva.
- Wischer, G., Boguhn, J., Steinga, H., Schollenberger, M., & Rodehutsord, M. (2013). Effects of different tannin-rich extracts and rapeseed tannin monomers on methane formation and microbial protein synthesis *in vitro*. *Animal*, 7(11), 1796–1805.
- Zamiri, M.J., Sharifabadi, H.R., Bagheri, A.R., & Solhjoo, A. (2015). Effects of inclusion of licorice (L.) leaves, a tannin-containing plant, in a low-protein diet on feedlot performance and carcass characteristics of fat-tailed lambs. *Tropical Animal and Health Production*, 47, 597–602.
- Zhang, C.M., Guo, Y.Q., Yuan, Z.P., Wu, Y.M., Wang, J.K., Liu, J.X., & Zhu, W.Y. (2008). Effect of octadeca carbon fatty acids on microbial fermentation, methanogenesis and microbial flora *in vitro*. *Animal Feed Science and Technology*, 146, 259–269.
- Zhou, C.S., Xiao, W.J., Tan, Z.L., Salem, A.Z.M., M.M., Geng, M.M., Tang, S.X., Wang, M., Han, X.F., & Kang, J.H. (2012). Effects of dietary supplementation of tea saponins (*Ilex kudingcha C.J.Tseng*) on ruminal fermentation, digestibility and plasma antioxidant parameters in goats. *Animal Feed Science and Technology*, 176, 163 – 169.
- Zmora, P., Cieslak, A., Pers-Kamczyc, E., Nowak, A., Szczechowiak, J., & Szumacher-Strabel, M. (2013). Effect of *Mentha piperita L.* on *in vitro* rumen methanogenesis and fermentation. *Acta Agriculture*, 62, 46-52.

APPENDIX



Plate 1: Separation of papaya leaf from the stem



Plate 2. Rumen fistulated goat used for *in vitro* and *in vivo* studies.

BIODATA OF STUDENT

Saeid Jafari was born in Tehran, capital of Iran on 7th September 1984. He completed his primary, secondary and high school education in Tehran. He pursued his B.Sc degree at the Faculty of Agriculture, Islamic Azad University of Saveh and graduated in 2007.

The author continued his M.Sc program in the same university focusing his research interests in the areas of agricultural byproducts, namely almond hull. For continuation of his study in PhD program he travelled to Malaysia at February, 2012.



LIST OF PUBLICATIONS

- Saeid Jafari**, Goh Yong Meng, Mohamed Ali Rajion, Mahdi Ebrahimi. Effect of papaya (*Carica papaya*) leaf on gas production *in vitro* gas production technique. Proceedings of the 35th Malaysian Society for Animal Production (MSAP) Annual Conference. 4 - 6 th June, 2014, Sarawak, Malaysia, PP. 69-70.
- Saeid Jafari**, Y.M. Goh, M.A. Rajion, Y. Hammali and M. Ebrahimi. Impact of papaya leaf extract on *in vitro* methane production. Proceedings of the 36th Malaysian Society for Animal Production (MSAP) Annual Conference, 6-8 June, 2015, Negari Sembilan, Malaysia.
- Saeid, J**, Goh, Y.M, Mohamed Ali, R., Yusuf, H., Mahdi, E. and Elaheh, M. Replacing alfalfa with bamboo leaf did not affect *in vitro* rumen gas production in the goat. International Agriculture Congress 2014, Putrajaya, Malaysia, PP. 314-317.
- Said Jafari**, Goh Yang Meng, Mohammed Ali Rajion, Yusuf Hammali and Mahdi Ebrahimi. Impact of Papaya Leaf on *In Vitro* Methane Production. Proceedings of the Asian- Australasian animal production (AAAP) Conference, 2014, Yogyakarta, Indonesia.
- Saeid Jafari**, Goh Yong Meng, Mohamed Ali Rajion, Mohammad Faseleh Jahromi, Mahdi Ebrahimi. Ruminant methanogenesis and biohydrogenation reduction potential of papaya (*Carica papaya*) leaf: an *in vitro* study. (Published)
- Saeid Jafari**, Goh Yong Meng, Mohamed Ali Rajion, Mohammad Faseleh Jahromi, Mahdi Ebrahimi. Papaya (*Carica papaya*) leaf methanolic extract modulates *in vitro* rumen methanogenesis and rumen biohydrogenation. (accepted).
- Saeid Jafari**, Goh Yong Meng, Mohamed Ali Rajion, Mohammad Faseleh Jahromi, Mahdi Ebrahimi. Polyphenol rich solvent fractions from papaya leaf can modulate rumen fermentation characteristics, rumen biohydrogenation and rumen microbial population *in vitro*. (Published).
- Saeid Jafari**, Goh Yong Meng, Mohamed Ali Rajion, Mohammad Faseleh Jahromi, Mahdi Ebrahimi. Papaya (*Carica papaya*) leaf as a possible natural alternative for modulating rumen fermentation characteristics and rumen microbial population in goats: an *in vivo* study (Submitted).



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION: _____

TITLE OF THESIS / PROJECT REPORT:

EVALUATION OF POLYPHENOL RICH PAPAYA (*CARICA PAPAYA* L.) LEAF AND ITS
EXTRACTS ON RUMEN MICROBES, BIOHYDROGENATION AND METHANE PRODUCTION

NAME OF STUDENT: SAEID JAFARI

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentiality or restricted.]