



***GROWTH, CARCASS AND DIGESTIVE SYSTEM CHANGES IN
HYBRID VILLAGE CHICKENS AFTER UROPYGIALECTOMY***

HASAN SAAD ABDULHUSSEIN JAWAD

FPV 2016 10



**GROWTH, CARCASS AND DIGESTIVE SYSTEM CHANGES IN
HYBRID VILLAGE CHICKENS AFTER UROPYGGIALECTOMY**

By

HASAN SAAD ABDULHUSSEIN JAWAD

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

June 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

GROWTH, CARCASS AND DIGESTIVE SYSTEM CHANGES IN HYBRID VILLAGE CHICKENS AFTER UROPYGLIALECTOMY

By

HASAN SAAD ABDULHUSSEIN JAWAD

June 2016

Chairman : Lokman Hakim Bin Idris, PhD
Faculty : Veterinary Medicine

Akar Putra is a hybrid village chicken; the cross breeding process was happened by chance when the wild jungle fowl interred Universiti Putra Malaysia (UPM) ground and mated with their Ayam Kampung (Native chicken). Nonetheless, there is a dearth of information relating to the Akar Putra chicken strain, particularly the morphology of its digestive system.

Uropygial Gland (UG) is the most prominent integument gland in the birds. It is puzzling that little is known about its morphology and function. The suggested functions of this gland can be placed into four groups: 1) feather maintenance; 2) water-proofing; 3) intraspecific communication and 4) defense against predators.

This thesis introduces a new technique to improve the production performance of the chicken through ablation of the uropygial gland which called Uropygialectomy (UP). This application will cause an upset in the poultry industry as well as will significantly contribute towards its development by increasing the economic viability obtained through an increase the chicken product. Previous method of Uropygialectomy included removing the uropygial gland completely and that usually attached by severe bleeding with hard stress exposure of the chicken. While, our modification method includes removing parts from the gland (half lobes, half isthmus and papillae). That modification was taken place in order to make UP operation safer, practical, applicable and shows a significant improvement of production performance.

Therefore, this study was carried out to address the production performance, carcass characteristics, external morphological changes, anatomical changes of digestive system, growth hormone concentration and histology of digestive system investigation of 120 Akar Putra chickens strain following the ablation of the uropygial gland. The experiment comprised five treatments, with 3 replicates for each. The treatments consisted of a control T1; Uropygialectomy was applied with T2, T3, T4 and T5 treatments at 3, 4, 5 and 6 weeks of age respectively.

The results revealed remarkable significant ($P < 0.05$) enhancing for UP treatments than a control group in all of males and females' body weight, weight gain, feed intake and feed conversion ratio measurements. Furthermore, the results indicated that UP treatments caused significantly improvement ($P < 0.05$) concerning body weight, carcass weights and dressing percentage with or without eating giblets. Additionally, significant different at level ($P < 0.05$) was observed in the traits of males' breast and back relative weight. Additionally, there was a significant effect at level ($P < 0.05$) in the females' breast relative weight trait; however, T2 surpasses other treatment groups T1, T3, T4 and T5 with relation to the most carcass traits involved in this experiment.

The external morphological comparison results between UP treatments and control at week 12 shows that the males of UP treatments had higher values in bird length, growth rate, breast diameters and lengths of neck, back, keel bone and extremities than males of the control group. Likewise, females of PU treatments surpassed females of the control group in bird length, growth rate, neck length, breast width and comb length.

The anatomical evaluations revealed that the males of UP treatments had more length ($P < 0.05$) esophagus 9.9-16.2%, proventriculus 11.1-34.4%, gizzard 26.7-220%, pancreas 0-20.4%, jejunum 4.9-26.1 and colon 18.1-60.6 than their control group counterparts. Furthermore, females of UP treatments had ($P < 0.05$) longer esophagus 6.8-22.3%, pancreas 8.3-33.3% and cecum 13-26% compared with females in control.

Histologically, surgical removing of the uropygial gland, especially at week 3 had greater ($P < 0.05$) effect on the total duodenum, jejunum and ilium wall thickness. In addition, effects ($P < 0.05$) were observed on the wall thickness of males' cecum and colon. Moreover, the wall layers of: esophagus, proventriculus, gizzard and rectum were not affected by the treatment. However, removing the uropygial gland showed significant impact ($P < 0.05$) in males' growth hormone concentration level at week 7 and ($P < 0.05$) effect at week 12 in both sexes.

In conclusion, the results of current study demonstrated that partial ablation of the uropygial gland, especially at week 3 of age had a positive effect on the production performance, carcass characteristics, external morphological measurements, anatomy and histology of digestive system organs as well as the growth hormone concentration of the Akar Putra chicken strain. These enhancement in the body performance can be justified that UP will contribute in retention the essential fatty acids in the body and avoid its attraction inside the UG, then secreted out of the body. In another hand, it will support the work of prostaglandins, which drives from Arachidonic fatty acid, resulting in the production of growth hormone. The improvement in the productive performance, carcass characteristics, phenotypic traits, as well as the anatomy and histology of the digestive system were as a consequence to increase of growth hormone secretion by the anterior pituitary gland. Moreover, the improvement in the body morphology was as a consequence to increase of the steroid hormones' levels by stopping the function of the uropygial gland in converting progesterone to testosterone.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PERUBAHAN TUMBESARAN, KARKAS DAN SISTEM PENCERNAAN DI
DALAM AYAM KAMPUNG HYBRID SELEPAS UROPIGIALEKTOMI**

Oleh

HASAN SAAD ABDULHUSSEIN JAWAD

Jun 2016

Pengerusi : Lokman Hakim Bin Idris, PhD
Fakulti : Perubatan Veterinar

Ayam Akar Putra merupakan sejenis baka kacukan yang telah melalui proses pembiakan silang yang berlaku secara kebetulan apabila ayam hutan liar memasuki kawasan Universiti Putra Malaysia (UPM) dan mengawan dengan Ayam Kampung (ayam tempatan). Walau bagaimanapun, terdapat kekurangan maklumat berkaitan dengan ayam Akar Putra, terutamanya morfologi sistem pencernaan.

Kajian ini memperkenalkan kaedah yang baharu untuk mempertingkatkan prestasi pengeluaran melalui penyingkiran kelenjar urophagial. Kaedah ini akan memberi kesan yang ketara kepada nilai ekonomi melalui peningkatan produk ayam. Kaedah yang terdahulu, dimana kelenjar ini dibuang sepenuhnya melalui pembedahan mengakibatkan pendarahan dan tekanan yang teruk kepada ayam. Kaedah yang baharu ini adalah penyingkiran separuh dari kelenjar urophagial (separuh lobes, isthmus dan papillae) melalui pembedahan. Pengubahsuaian kaedah ini adalah untuk menghasilkan pembedahan yang lebih selamat, praktikal, boleh digunakan dan menghasilkan perubahan yang ketara kepada prestasi pengeluaran.

Oleh itu, kajian ini dijalankan bagi mengkaji prestasi pengeluaran, ciri-ciri karkas, perubahan morfologi luaran, perubahan anatomi sistem penghadaman, kepekatan hormon pertumbuhan dan histologi sistem penghadaman pada 120 ayam Akar Putra selepas penyingkiran sebahagian kelenjar uropygial (Uropygiallectomy). Ekperimen ini dibahagikan kepada lima kumpulan dan 3 replicate. Kumpulan ini terdiri dari; T1 untuk kawalan, T2, T3, T4 dan T5 pada minggu ke 3, 4, 5 dan 6.

Hasil kajian menunjukkan penyingkiran sebahagian dari kelenjar uropygial (PU) menunjukkan perbezaa yang ketara ($P < 0.05$) diantara kumpulan kawalan dan kumpulan eksperimen pada ayam jantan dan betina dari segi pertambahan berat badan, pengambilan makanan, kadar penukaran makanan, dan berat reletif otot dada bagi jantan dan betina. Ianya juga menunjukkan perbezaan yang ketara ($P < 0.05$) bagi berat hidup dan berat karkas dengan giblek dan juga tanpa giblek. Kumpulan T2

menunjukkan kesan yang paling ketara diantara kumpulan-kumpulan eksperimen yang terlibat.

Perbandingan morfologi luaran pada minggu ke 12 menunjukkan, ayam jantan menghasilkan nilai yang lebih tinggi untuk berat dan panjang badan, kadar tumbesaran, diameter otot dada dan panjang leher, belakang dan tulang kaki daripada kumpulan kawalan, manakala ayam betina menunjukkan nilai yang lebih tinggi pada berat badan dan panjang badan, kadar tumbesaran, panjang leher dan balung serta lebar otot dada.

Penilaian anatomi menunjukkan PU pada ayam jantan menghasilkan ($P < 0.05$) 9.9-16.2% esophagus lebih panjang, 11.1- 34.4% pada proventrikulus, 26.7 %-220% pada hempedal, 0-20.4% pankreas, 4.9-26.1% jujunum dan 18.1-60.6% kolon daripada kumpulan kawalan. Manakala PU pada ayam betina menunjukkan ($P < 0.05$) 6.8-22.3% esophagus lebih panjang, 8.3-33.3% pankreas dan 13-26% pada sekum berbanding dengan kumpulan kawalan.

Secara histologinya, PU pada umur 3 minggu menunjukkan kesan yang ketara ($P < 0.05$) pada ketebalan duodenum, jujunum dan ilium. Walaubagaimana pun ketebalan esophagus, proventrikulus, hempedal dan rektum tidak menunjukkan perbezaan yang ketara. PU pada ayam jantan menunjukkan perbezaan yang ketara ($P < 0.05$) bagi hormon tumbesaran pada umur 7 minggu dan perbezaan yang ketara ($P < 0.05$) pada umur 12 minggu untuk ayam jantan dan betina.

Kesimpulannya, PU pada umur 3 minggu menunjukkan kesan yang positif kepada prestasi pertumbuhan, kriteria karkas, ukuran morfologi luaran, anatomi dan histologi organ-organ pencernaan termasuk konsentrasi hormon tumbesaran pada ayam Akar Putra.

Penambahbaikan prestasi badan ayam ini disebabkan kerana, kaedah UP ini menyumbang kepada pengekalan asid lemak penting di dalam badan dan menghalang kepada tarikan kepada kelenjar urophagial. Ia juga membantu fungsi hormon prostaglandin yang terhasil dari asid lemak Arachidonic dalam penghasilan hormon tumbesaran. Peningkatan prestasi pengeluaran, kriteria karkas, sifat-sifat fenotip termasuk anatomi dan histologi system pencernaan adalah akibat dari peningkatan penghasilan hormon tumbesaran yang dihasilkan dari kelenjar Pitutari anterior. Tambahan pula peningkatan kepada morfologi badan adalah disebabkan peningkatan paras hormon steroid seterusnya penukaran hormon progesteron kepada testosteron dengan menyekat fungsi kelenjar urophagial.

ACKNOWLEDGEMENTS

In the name of Allah, the most Benevolent and Most Merciful... Alhamdulillah, I am thankful for giving me the strength, which has enabled me to complete this study.

I deeply express my gratitude to my supervisor Dr. Lokman Hakim Bin Idris, for giving me an opportunity to complete my thesis. He devoted his time for invaluable guidance, advice, supervision and support throughout the course of this study.

I wish to express my sincere gratitude to my co-supervisors, Professor Dr. Md Zuki Abu Bakar, and Associate Professor Dr. Azhar bin Kasim for investing their time and knowledge in my study.

I am especially grateful to the Iraqi Ministry of Higher Education and Scientific Research/ Baghdad University for providing the three years the Research Scholarship to perform this study.

It is a pleasure to express my gratitude to Prof. Dr. Saad Abdulhussein Naji who provided advices that improved this study.

My grateful thanks to Mr. Humam Ali Merza for his assistance in poultry management during this experiment.

My grateful thanks extend to all faculty staff and all staff at the Universiti Putra Malaysia for everything they have done for me and whom not mentioned here are deeply appreciated.

I certify that a Thesis Examination Committee has met on 9 June 2016 to conduct the final examination of Hasan Saad Abdulhussein on his thesis entitled "Growth, Carcass and Digestive System Changes in Hybrid Village Chickens After Uropygialectomy" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Goh Yong Meng, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Dahlan bin Ismail, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Noordin bin Mohamed Mustapha, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Abdel-Hamid Kamel Osman, PhD

Professor
Suez Canal University
Egypt
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 23 August 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Lokman Hakim Bin Idris, PhD

Senior Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

MD Zuki Abu Bakar@Zakaria, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Azhar Bin Kasim, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Hasan Saad Abdulhussein Jawad, GS37472

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: Dr. Lokman Hakim Bin Idris

Signature: _____
Name of
Member of
Supervisory
Committee: Professor Dr. Md Zuki Abu Bakar@Zakaria

Signature: _____
Name of
Member of
Supervisory
Committee: Associate Professor Dr. Azhar Bin Kasim

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xix
CHAPTER	
1. INTRODUCTION	1
2. LITERATURE REVIEW	4
2.1 Red jungle fowl	4
2.2 Village chickens	5
2.3 Hybrid village chicken (Akar Putra strain)	5
2.4 Uropygial gland overview	7
2.4.1 Uropygial gland functions	7
2.4.2 Uropygial gland morphology	8
2.4.3 Uropygial gland histology	9
2.5 Hormonal control of growth in chicken	12
2.5.1 Chicken growth hormone	13
2.6 Uropygiaectomy and production	14
2.7 Uropygiaectomy, prostaglandin and growth hormone	16
2.8 Uropygiaectomy, sex hormones and chicken external morphology	20
3. MATERIALS AND METHODS	25
3.1 Animals and housing	25
3.2 Experimental design	26
3.2.1 Uropygiaectomy (UP)	26
3.3 Observations:	28
3.3.1 Production performance	28
3.3.2 Carcass characteristics	28
3.3.3 External morphological characteristics	29
3.3.4 Anatomy of digestive system	31
3.3.5 Growth hormone (GH) concentration and digestive system histometrical aspect.	31
3.4 Data Analysis	31

4.	ABLATION OF UROPYGIAL GLAND EFFECT ON PRODUCTION PERFORMANCE OF HYBRID VILLAGE CHICKEN	32
4.1	Introduction	32
4.2	Materials and methods	32
4.2.1	Animals feeding and management	32
4.2.2	Production parameters	32
4.2.3	Research design and data analysis	33
4.3	Results	33
4.3.1	Body weight	33
4.3.2	Feed intake	34
4.3.3	Weight gain	34
4.3.4	Feed conversion ratio	39
4.4	Discussion	44
5.	ABLATION OF UROPYGIAL GLAND EFFECT ON CARCASS CHARACTERISTICS OF HYBRID VILLAGE CHICKEN	50
5.1	Introduction	50
5.2	Materials and methods	50
5.2.1	Animals feeding and management	50
5.2.2	Carcass characteristics	50
5.2.3	Research design and data analysis	51
5.3	Results	51
5.4	Discussion	54
6.	ABLATION OF UROPYGIAL GLAND EFFECT ON THE EXTERNAL MORPHOLOGICAL CHARACTERISTICS OF HYBRID VILLAGE CHICKEN	56
6.1	Introduction	56
6.2	Materials and methods	56
6.2.1	Animals feeding and management	56
6.2.2	External morphological characteristics	56
6.2.3	Research design and data analysis	58
6.3	Results	58
6.3.1	Zoometrical measurements at week 7 of age:	58
6.3.2	Zoometrical measurements at week 10 of age:	64
6.3.3	Zoometrical measurements at week 12 of age:	64
6.4	Discussion	74
7.	ABLATION OF UROPYGIAL GLAND EFFECT ON THE ANATOMY OF THE DIGESTIVE SYSTEM OF HYBRID VILLAGE CHICKEN	77
7.1	Introduction	77
7.2	Materials and methods	78
7.2.1	Animals feeding and management	78
7.2.2	Anatomy of digestive system	78
7.2.3	Data analysis:	79
7.3	Results	79
7.3.1	GIT organs length	79
7.3.2	GIT organs weight	85

	7.3.3	GIT organs density	93
	7.4	Discussion	96
8.		PARTIAL ABLATION OF UROPYGIAL GLAND EFFECTS ON GROWTH HORMONE CONCENTRATION AND DIGESTIVE SYSTEM HISTOLOGICAL ASPECT OF HYBRID VILLAGE CHICKEN	101
	8.1	Introduction	101
	8.2	Materials and methods	102
		8.2.1 Animals feeding and management	102
		8.2.2 Growth hormone assay	102
		8.2.3 GIT Histometrical assay	102
		8.2.4 Statistical analysis	103
	8.3	Results	103
		8.3.1 Serum GH concentration	103
		8.3.2 Histometrical parameters of digestive system	103
	8.4	Discussion	112
		8.4.1 Effect of PU operation on GH concentration	112
		8.4.2 Effect of UP operation on histological parameters of digestive system	112
9.		GENERAL DISCUSSION	114
	9.1	UP plays a big role in increasing the GH concentration in the blood	114
	9.2	UP plays a big role in supporting the external morphological development of Akar Putra chicken strain	116
	9.3	Conclusion	117
	9.4	Recommendations for futher study	119
		REFERENCES	120
		APPENDICES	134
		BIODATA OF STUDENT	151
		LIST OF PUBLICATIONS	152

LIST OF TABLES

Table		Page
1	Composition of basal diet.	26
2	(Mean \pm S.E.) males' body weight (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	35
3	(Mean \pm S.E.) females body weight (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	36
4	(Mean \pm S.E.) males feed intake (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	37
5	(Mean \pm S.E.) females feed intake (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	38
6	(Mean \pm S.E.) males' weight gain (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	40
7	(Mean \pm S.E.) females weight gain (g/bird/week) for control and uropygialectomy treatments from 1-12 week.	41
8	(Mean \pm S.E.) males feed conversion ratio (g .feed/ g .gain) for control and uropygialectomy treatments from 1-12 week.	42
9	(Mean \pm S.E.) females feed conversion ratio (g .feed/ g .gain) for control and uropygialectomy treatments from 1-12 week.	43
10	(Mean \pm S.E.) males' carcass characteristics of uropygialectomy treatments at week 12 of age (% relative carcass weight).	52
11	(Mean \pm S.E.) females carcass characteristics for uropygialectomy treatments at week 12 of age (% relative weight).	53
11	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 7 of age.	60
12	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 7 of age.	61
13	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 10 of age.	65

14	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 10 of age.	66
15	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 12 of age.	70
16	(Mean \pm S.E.) zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 12 of age.	71
17	Classification of phenotypic traits based on starting appearance of significant effects between treatments.	74
18	(Mean \pm S.E.) males GIT parts length (cm) of uropygialectomy treatments at week 12 of age.	81
19	Relative length (%) of males GIT parts than total GIT length of uropygialectomy treatments at week 12 of age.	81
20	(Mean \pm S.E.) females GIT parts length (cm) of uropygialectomy treatments at week 12 of age.	82
21	Relative length (%) of females GIT parts than total GIT length of uropygialectomy treatments at week 12 of age.	83
22	(Mean \pm S.E.) males GIT parts weight (g) of uropygialectomy treatments at week 12 of age.	86
23	Relative weight (%) of males GIT parts than total GIT weight of uropygialectomy treatments at week 12 of age.	88
24	Relative weight (%) of males GIT parts than total live body weight of uropygialectomy treatments at week 12 of age.	89
25	(Mean \pm S.E.) females GIT parts weight (g) of uropygialectomy treatments at week 12 of age.	90
26	Relative weight (%) of females GIT parts than total GIT weight of uropygialectomy treatments at week 12 of age.	91
27	Relative weight (%) of females GIT parts than total live body weight of uropygialectomy treatments at week 12 of age.	92
28	(Mean \pm S.E.) males GIT parts density (g/cm) of uropygialectomy treatments at week 12 of age.	94
29	(Mean \pm S.E.) females GIT parts density (g/cm) of uropygialectomy treatments at week 12 of age.	95

30	(Mean \pm S.E.) effect of UP operation on serum growth hormone concentration (pg/ml) of Akar Putra chicken in different periods.	104
31	(Mean \pm S.E.) effect of UP on the histometrical parameters (mm) of upper part of the digestive system at week 12 of age.	106
32	(Mean \pm S.E.) effect of UP on the histometrical parameters (mm) of males' small intestinal parts at week 12 of age.	108
33	(Mean \pm S.E.) effect of UP on the histometrical parameters (mm) of females' small intestinal parts at week 12 of age.	109
34	(Mean \pm S.E.) effect of UP on the histometrical parameters (mm) of males' large intestinal parts at week 12 of age.	110
35	(Mean \pm S.E.) effect of UP on the histometrical parameters (mm) of females' large intestinal parts at week 12 of age.	111

LIST OF FIGURES

Figure		Page
1	Akar Putra chicken strain.	6
2	Modified drawing of a UG illustrating anatomical organization.	9
3	Generalized UG structure from follicle through to primary sinus.	11
4	Illustration of a follicle from the UG of a Single Comb White Leghorn Chicken. Scale = 0.01mm.	11
5	Relative importance of hormones in vertebrate growth: B, birth or hatching; and P, puberty.	12
6	Ontogeny of GH, IGF-I, T3, GH binding and GHR expression (mRNA) in the chicken; H, hatching.	13
7	Evolution process of prostaglandins from Arachidonic fatty acid.	18
8	The proposal role of uropygial gland in its effect on the balance of the concentrations of steroid hormones in the avian body.	19
9	Main reproductive organs in chicken.	22
10	Vertical view of follicle and sexual steroid hormones sorter cells.	22
11	Hormonal interaction between the Hypothalamus, Anterior Pituitary and ovary that result in ovulation.	22
12	Hypothalamic-pituitary GH-IGF-I axis. Pituitary GH production and secretion is regulated by the hypothalamic hormones somatostatin and GH releasing hormone (GHRH). GH has direct effect on muscle, bone, fat, lipolysis and reproduction. GH also has indirect effect on those organs by stimulating liver to secrete IGF-I.	23
13	The proposal role of uropygial gland in its effect on the balance of the concentrations of hormones in the avian body.	24
14	Photograph of Uropygialectomy steps.	27
15	Photograph of measuring body weight.	28

16	Measuring process of external morphological characteristics including: Bird Length (BL), Neck length (NL), Back length (BAL), Breast length (BRL) and wide (BRW), Keel bone length (KBL), Wing length (WL), leg length (LL), Comb width (CW), Comb length (CL), Beak width (BEW), Beak length (BEL), Wattles' width (WAW), Wattles' length (WAL).	30
17	Males body weight effectiveness by UP at week 3 (T2); 4 (T3); 5 (T4) and 6 (T5) of age. (Mean \pm Stander error).	47
18	Females body weight effectiveness by UP at week 3 (T2); 4 (T3); 5 (T4) and 6 (T5) of age. (Mean \pm Stander error).	49
18	Zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 7 of age.	62
19	Zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 7 of age.	63
20	Zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 10 of age.	67
21	Zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 10 of age.	68
22	Zoometrical measurements (weight: g; length and wide: cm) of males in control group and UP treatments at week 12 of age.	72
23	Zoometrical measurements (weight: g; length and wide: cm) of females in control group and UP treatments at week 12 of age.	73
24	Males' growth rate gauge variation ratio of UP treatments than the control group.	76
25	Females' growth rate gauge variation ratio of UP treatments than the control group.	76
26	Variation ratio (%) of males GIT parts length of UP treatments than the control group.	98
27	Variation ratio (%) of females GIT parts length of UP treatments than the control group.	98
28	Variation ratio (%) of males GIT parts weight of UP treatments than the control group.	99
29	Variation ratio (%) of females GIT parts weight of UP treatments than the control group.	99

29	Variation ratio (%) of males GIT parts density of UP treatments than the control group.	100
30	Variation ratio (%) of females GIT parts density of UP treatments than the control group.	100



© COPYRIGHT UPM

LIST OF ABBREVIATIONS

UP	Uropygialectomy
UG	Uropygial gland
PI	Production Index
EPEF	European Production Efficiency Factor
ANOVA	Analysis of variance
cm	Centimeter
d	Day
g	Gram
GIT	Gastrointestinal tract
H & E	Haematoxylin and Eosin
ml	Milliliter
NS	Normal Solution
NBF	Neutral Buffered Formalin
nm	Nanometer
PBS	Phosphate Buffered Saline
SE	Standard Error
SPSS	Statistical Package for the Social Sciences
vol	Volume
♂	Male
♀	Female
trt	Treatment
rept	Replicate
RW	Relative Weight
RL	Relative Length
GH	Growth Hormone
GHRH	Growth Hormone Releasing Hormone
IGF-I	Insulin-like Growth Hormone

CHAPTER 1

INTRODUCTION

Akar Putra is a hybrid village chicken; the cross breeding process was happened by chance when the wild jungle fowls entered Universiti Putra Malaysia grounds and mated with their chicken (ayam kampung). Akar Putra chicken has a more robust growth process than its parents because the maturation period is shorter (less 13 weeks). It can lay 120-200 eggs per year, and it has more resistance for diseases (Kasim, 2007).

Uropygial gland is the only subcutaneous gland in birds' body (McLelland, 1990). It has many names like preen gland, based on its function in preening the bird's feather (Lucas and Stettenheim, 1972a; King and McLelland, 1984) and oil gland depending on its oily secretion (Schultz *et al.*, 2002). Furthermore, it called uropygial gland based on its position, which is on the base of the tail, dorsally between the fourth caudal vertebrae and the pygostile (Lucas and Stettenheim, 1972a,b ; Sawad, 2006a). The function of the gland is still a subject of controversy. There are many accepted functions of gland secretions like conferring water-repellent properties on the feather coat and maintaining the suppleness of it. In addition, it has proposed to be associated to pheromone production, control of plumage hygiene, thermal insulation and defense against predators because of its foul smell in some birds (Jacob, 1992; Montalti *et al.*, 2000; Soler *et al.*, 2012; Vincze *et al.*, 2013). Uropygial gland is completely absent in Struthionidae, Rheidae, Casuaridae, Dromidae and in a few species of Columbidae and Psittacidae (Johnston, 1988). Montalti and Salibián (2000) mentioned that the oil of the uropygial gland is not important to the birds who do not have it. While Goodwin (1970) said, the uropygial gland in some of the birds is non-active. Moreover, Moyer *et al.*, (2003a, b) gave an explanation that the birds that do not have uropygial gland use dusts bath to keep and clean their feather.

Modern commercial breeds of meat chicken characterized by super-fast growth and high efficiency of a food conversion ratio as a result of intense genetic selection. Wepruk and Church (2003) observed that the final body weight of broiler in 1976 was 2 kg at the age of 63 days while the same average of body weight was arrived at age 35 days in 2001. This improvement in the growth rate reflected negatively on the disease resistance and immune response of these birds, because of a negative genetic link coefficient was observed between the growth speeds and immune response (Qureshi and Havenstein, 1994). In this context, increasing in a mortality ratio in these strains of birds happened due to increasing their susceptibility to bacterial diseases and metabolic diseases. These occurred as a consequence of irregular metabolic processes, an imbalance in the acid-base balance of body fluids such as ascites disease, sudden-death syndrome (SDS) and increased skeletal disorders like leg abnormalities. It has been scientifically proven that highest rates of those pathological conditions were shown in flocks and individual rapid growth chicken at 3rd and 4th weeks of age (Robinson *et al.*, 1992; Julian, 1997, 1998; Leeson and Summer, 1997; Gonzales *et al.*, 1998, 2000). Based on the limitation of the problem, this research was planned to innovate a safe

technique to raise the level of poultry production performance in general and local Malaysian chicken (Akar Putra strain) particularly without using the genetic improvement methods, which have proven that it has negative impacts on birds' immunity. Furthermore, this study planned to reduce the gap in knowledge in terms of the production performance, carcass characteristics and digestive system morphology of the hybrid village chicken. Regarding on uropygialectomy (UP), earlier studies were described that application as one of the improvement methods to enhance the body performance of chicken. Previous method of UP included removing the uropygial gland completely and that usually attaches with severe bleeding which sometimes case decimation of the experimental birds. Additionally, it has been reported that high numbers of the treated birds could be dying after the operation because of cannibalism, which is sometimes difficult to overcome. The reason behind that may be the remaining blood on the feathers around the incision area after the operation, which is considered as an irritation factor for the chicken and urges them to cannibalism. Furthermore, the previous researchers recommended to apply the uropygialectomy at the earlier ages (1-6) weeks of chicken age in order to give the body sufficient time to grow up (AI-Daraji *et al.*, 2006). In addition, because of their observation that the adult chicken has to be less responsive to the uropygialectomy compared than the chicken in growing stage. The UP in this study has been modified to include removing about half lobes, half isthmus and papillae at the growing stage (3-6 weeks). That modification was taken place in order to make UP operation safer, practical and applicable (quickly and accurately), which encourages to perform it on a large scale in the poultry farms.

The aim of this study is to describe the anatomical and histological changes of the digestive system and the growth patterns of the digestive organs as well as the carcass characteristics after ablation of the uropygial gland, which was at weeks 3, 4, 5 and 6 of age on T2, T3, T4 and T5 treatments respectively. In this study, we also examined the weekly production performance, production index (PI) and European production efficiency factor (EPEF), which are considered the most confidence standards to evaluate poultry farm experiments in the economic term. Furthermore, the external morphology and plasma growth hormone at weeks 7, 10 and 12 of age for males and females separately as an attempt to cover their changes in the period after the last application of uropygialectomy at week six until the end of the experiment (week 12). The hypothesis that the morphology and histology of digestive system organs as well as production performance, carcass characteristics, external morphology and plasma growth hormone are affected by UP. It is worthy to mention that the objectives of this study have been covered for males and females separately in order to be more economic importance in the poultry industry. Therefore, it is known that the new trend of the poultry industry is a single-sex breeding. For example, some companies adopt a male breeding, and others adopt breeding females, in line with the markets need and the desire of consumers. Moreover, the current study has compared each part, organ or tissue between the treatments separately without any interaction in order to provide the maximum amount of information and detail that shows which treatment was more affected by ablation of the uropygial gland. That corresponds with the data analysis new approach in the most professional poultry journals. This study was undertaken with the following objectives:

- To define the production performance of Akar Putra chicken as a new strain of village chicken and its effectiveness by UP at different ages.
- To compare the carcass characteristics between UP treatments and control group.
- To compare the external morphological parameters at 3 different ages (7, 10 and 12 weeks) between UP treatments and control group.
- To investigate the anatomical changes of the digestive system organs after ablation of the uropygial gland.
- To assay the histological changes of the digestive system organs as well as the plasma growth hormone at 3 different ages (7, 10 and 12 weeks) between UP treatments and control group.



REFERENCES

- AI-Daraji, H.J., Abdul-Hassan, I.A., Abmad, A.S. and Razuki, R.H. (2006). Effect of Uropygialectomy at two different ages on semen quality of white leghorn males. *The Iraqi Journal of Agricultural Sciences*. 37: 213 - 218.
- Akester, A.R. (1987). Form and Function in Birds. *Journal of anatomy*, 150, 288.
- AI-Daraji, H.J., Naji, S.A., AI-Tikriti, B.T.O., and AI-Rawi, A.A. (2002). Influence of uropygialectomy (Iraqi Method) on reproductive performance of different flocks of chicken. *The Iraqi Journal of Agricultural Sciences*, 33(2):165-172. Cited by AI-Daraji *et al.*, 2006.
- AI-Mahdawy, R.S. (2003). The effect of uropygialectomy (Iraqi method) on the productive and physiological performance of broilers. M.Sc. thesis, *Department of Animal Production, Collage of Agriculture, University of Baghdad*. Cited by AI-Daraji *et al.*, 2006.
- Arab Agricultural Statistics Yearbook. (2015). Vol 30. <http://www.aoad.org/EAASYXX.htm>.
- Aslan, K., Ozcan, S., and Kürtül, I. (2000). Arterial vasclarization of the uropygial gland (gl. Uropygialis) in geese (*Anser anser*) and ducks (*Anas platyrhynches*). *Anatomy Histology and Embryology*. 29(5), 291_293.
- Al-Hassani, D.H., Abdul-Hussan, I.A., and Naji, S.A. (2008). Effect of uropygialectomy on some semen traits of broiler breeder males. Paper presented in World Poultry Congress conference in Australia. Vol 64. *World Poultry Science Journal*.
- Assan, N. (2013). Bioprediction of body weight and carcass parameters from morphometric measurements in livestock and poultry. *Scientific Journal of Review*, 2(6), 140-150.
- Akanno, E.C., Ole, P.K., Okoli, I.C., and Ogundu, U.E. (2007). Performance characteristics and prediction of body weight of broiler strains using linear body measurements. *In Proceedings of the 22nd Annual Conference of Nigerian Society for Animal Production, Calabar, Cross River State, Nigeria* (pp. 162-164).
- Abdullah, A.Y., Al-Beitawi, N.A., Rjoup, M.M., Qudsieh, R.I., and Ishmais, M.A. (2010). Growth performance, carcass and meat quality characteristics of different commercial crosses of broiler strains of chicken. *The journal of poultry science*, 47(1), 13-21.
- Apuno, A.A., Mbap, S.T., and Ibrahim, T. (2011). Characterization of local chickens (*Gallus gallus domesticus*) in shelleng and song local government areas of Adamawa State, Nigeria. *Agriculture and Biology Journal of North America*, 2(1), 6-14.

- Ajayi, F.O., Ejiofor, O., and Ironkwe, M.O. (2008). Estimation of body weight from linear body measurements in two commercial meat-type chicken. *Global Journal of Agricultural Sciences*, 7(1), 57-59.
- Ann, J. (2004). *Veterinary Histology*. Chapter 32. University of Illinois.
- Apandi, M., and Edwards, H.M. (1964). Studies on the composition of the secretions of the uropygial gland of some avian species. *Poultry Science*, 43(6), 1445-1462.
- Aptekmann, K.P., Artoni, S.M., Stefanini, M.A., and Orsi, M.A. (2001). Morphometric analysis of the intestine of domestic quails (*Coturnix coturnix japonica*) treated with different levels of dietary calcium. *Anatomia, histologia, embryologia*, 30(5), 277-280.
- Al-Tememy, H.S.A., Al-Jaff, F.K., Al-Mashhadani, E.H., and Hamodi, S.J. (2011). Histological effect of inclusion different levels of coriander oil in broiler diet on small intestine. *Diyala Agricultural Sciences Journal*, 3(2), En1-En11.
- Aughey, E., and Frye, F.L. (2001). Comparative veterinary histology with clinical correlates, First edition. *Manson publishing / the veterinary press*.
- Azahan, E.E.A. and Zahari, W.M. (1983). Observation on some characteristics of carcass and meat of Malaysian kampong chickens. *Mardi Research Bulletin*. 11: 225-232.
- Aini, I. (1990). Indigenous Chicken Production in South-East Asia. *Worlds Poultry Science Journal*. 46: 51-57.
- Abdul-Hassan, I. A. 2005. Effect of Iraqi method (uropygialectomy) on some physiological and reproductive traits of broiler breeder males. Dissertation Ph. D, College of Agriculture, University of Baghdad, Iraq.
- Al-Hayani, W. K. A. 2005. The use of Iraqi method represented with surgical removal of the uropygial gland to improve productive, physiological performance and immunological responses to broiler Ross strain. Thesis Msc, College of Agriculture, University of Al-Anbar, Iraq.
- Al-Mahdawy, R. S. R. 2008. The effect of uropygialectomy before the wexual maturity on productive performance and for curing the delay in sexual maturity in commercial laying hens stocks, Dissertation PhD, College of Agriculture, University of Baghdad, Iraq.
- Al-Shamire, J. S. H. 2009. The effect of uropygialectomy and diet supplementation with probiotics on productive and physiological Performanc of Japanese quails. Thesis Msc, College of Agriculture, University of Baghdad, Iraq.
- Brush, A.H. (1993). The evolution of feathers: A novel approach. *Avian Biology*. 9: 121-162.
- Bandyopadhyay, A., and Bhattacharyya, S.P. (1996). Influence of fowl uropygial gland and its secretory lipid components on growth of skin surface bacteria of fowl. *Indian journal of experimental biology*, 34(1), 48-52.

- Bandyopadhyay, A., and Bhattacharyya, S.P. (1999). Influence of fowl uropygial gland and its secretory lipid components on the growth of skin surface fungi of fowl. *Indian Journal of Experimental Biology*, 37(12), 1218-1222.
- Bertness, M.D. (1991). Zonation of *Spartina patens* and *Spartina alterniflora* in New England salt marsh. *Ecology*, 72(1), 138-148.
- Baumel, J.J., King, A.S., Breazile, J.E., Evans, H.E., and Vanden Berge, J.C. (1993). Handbook of avian anatomy: Nomina Anatomica Avium. Massachusetts. *Publications of the Nuttall Ornithological Club, Cambridge*.
- Bhattacharyya, S.P. (1972). A comparative study on the histology and histochemistry of uropygial glands. *La Cellule*, 69(2), 111.
- Baddeley, A.J., Gundersen, H.J.G., and Cruz-Orice, L.M. (1986). Estimation of surface area from vertical sections. *Journal of Microscopy* 142: 259-276.
- Brody, S. (1945). Bioenergetics and growth; with special reference to the efficiency complex in domestic animals.
- Bosch, M. (1996). Sexual Size Dimorphism and Determination of Sex in Yellow-Legged Gulls (Dimorfismo Sexual de Tamaño y Determinación del Sexo en Gaviotas Patiamarillas). *Journal of Field Ornithology*, 534-541.
- Bacon, W.L., Burke, W.H., Anthony, N.B., and Nestor, K.E. (1987). Growth Hormone Status and Growth Characteristics of Japanese quail Divergently Selected for Four-Week Body Weights¹. *Poultry science*, 66(9), 1541-1544.
- Bacon, W.L., Brown, K.I., and Musser, M.A. (1980). Changes in plasma calcium, phosphorus, lipids and estrogens in turkey hens with reproductive state. *Poultry Science*. 59:444-452.
- Bacon, W.L., Vizcarra, J.A., Morgan, J.L.M., Yang, J., Liu, H.K., Long, D.W., and Kirby, J.D. (2002). Changes in plasma concentrations of luteinizing hormone, progesterone, and estradiol-17 β in peripubertal turkey hens under constant or diurnal lighting. *Biology Reprod*. 67:591-598.
- Bluhm, C.K., Phillips, R.E., and Burke W.H. (1983). Serum levels of luteinizing hormone (LH), prolactin, estradiol, and progesterone in laying and nonlaying canvasback ducks (*Aythya valisineria*). *Gen. Comp. Endocrinol*. 52:1-16.
- Beebe, W. (1926). Pheasants: Their lives and homes. Garden City, New York: Doubleday. pp. 240-245.
- Burnside, J. and L. A. Cogburn. 1992. Developmental expression of hepatic growth hormone receptor and insulin-like growth factor-I mRNA in the chicken. *Molecular Cell Endocrinology*. 89:91-96.
- Calislar, T. (1986). Anatomy of domestic animals II. Dissection of Horse and Hen. Istanbul: *Faculty of Veterinary Medicine, Istanbul University*, 281-282.

- Cater, D.B., and Lawrie, N.R. (1950). Some histochemical and biochemical observations on the preen gland. *The Journal of physiology*, 111(3-4), 231-243.
- Chen, Y.H., Kou, M.J., Pan, F.M. and Lu, L.L. (2003). Effects of uropygial gland removal on the growth performance and plasma characteristics in female white roman goslings from 3 to 10 weeks of age. *Tunghai Journal*. vol. 44, p. 7-13.
- Cabarles, J.C., Lambio, A.L., Vega, S.A., Capitan, S.S., and Mendiolo, M.S. (2012). Distinct morphological features of traditional chickens (*Gallus gallus domesticus* L.) in Western Visayas, Philippines. *Animal Genetic Resources/Ressources génétiques animales/Recursos genéticos animales*, 51, 73-87.
- Crawford, R.D. (2003). Origin and history of poultry species. In poultry breeding and genetics. Elsevier, *Amsterdam*.pp1-41.
- Collias, N.E. and Saichuae, P. (1967). Ecology of the red jungle fowl in Thailand and Malaya with reference to the origion of domestication. *Natural History of Bulletin, Siam* 22: 189-209.
- Cabello, G. and C. Wrutniak. 1989. Thyroid hormone and growth: relationships with growth hormone effects and regulation. *Reproductive and Nutrition Development*. 387-402.
- Carter, C., J. Schwartz, and L. S. Smit, 1998. Molecular mechanism of growth hormone action. *Annual Review Physiology*. 58:187-207.
- Chambers, J. R. 1990. Genetics of growth and meat production in chickens. Quantitative Genetics and Selection. R. D. Crawford, ed. *Poultry Breeding and Genetics*. Elsevier, Amsterdam. 599-643.
- Chizzolini, R. E. Zanardi, V. Dorigoni, and S. Ghidini. 1999. Calorific value and cholestrol content of normal and low fat meat and meat products. *Trends Food Science Technology*. 10:119-128.
- Dellman, H.D., and Brown, E.S. 1976. Textbook of Veterinary Histology. Second Ed. Lea and Febiger Philadelphia. Pp. 272_340
- Duncan, D.B. (1955). Multiple range and multiple F tests. *Biometrics*, 11(1), 1-42.
- Dial, K.P. 1992. Avian forelimb muscles and nonsteady flight: can birds fly without using the muscles in their wings? *Auk* 109: 874-885.
- Davidson, M.W., and Abramowitz, M. (2002). www.micro.magnet.fsu.edu/micro/gallery/prostaglandin/prostaglandin.html
- Elder, W., and William, H. (1954). The oil gland of birds. *The Wilson Bulletin*, 6-31.
- Euribrid, B.V., 1994. Technical information for Hybro broilers, pp.22 (Boxmeer, The Netherlands, Euribrid Poultry Breeding Farm).

- El-Ghany, W. A. A., and Madian, K. (2011). Control of experimental colisepticaemia in broiler chickens using sarafloxacin. *Life Science Journal*, 8(3), 318-328.
- Francesch, A., Villalba, I., and Cartana, M. (2011). Methodology for morphological characterization of chicken and its application to compare Penedesenca and Empordanesa breeds. *Animal Genetic Resources/Ressources génétiques animales/Recursos genéticos animales*, 48, 79-84.
- Falconer, D.S. (1989). Introduction to Quantitative Genetics. 3rd Edition, Longman, London. Pp. 438.
- Floch, J.Y., Morfin, R.F., Daniel, J.Y., and Floch, H.H. (1988). Testosterone metabolism and its testosterone-dependent activation in the uropygial gland of quail. *Endocrine Research*. 14 (1): 93-107.
- Fennell, M.J., and Scanes, C.G. (1992). Effects of androgen (testosterone, 5 α -dihydrotestosterone, 19-nortestosterone) administration on growth in turkeys. *Poultry Science*. 71: 539-547.
- Floch, J.Y., Morfin, R., Picart, D., Daniel, J.Y., and Floch, H.H. (1985). Testosterone metabolism in the uropygial gland of the quail. *Steroids* 45 (5): 391-401.
- Flower, R.J., Cheung, H.S., and Cushman, D.W. (1973). Quantitative determination of prostaglandins and malondialdehyde formed by the arachidonate oxygenase (prostaglandin synthetase) system of bovine seminal vesicle. *Prostaglandins*, 4(3), 325-341.
- Fumihito, A., Miyake, T., Sumi, S., Takada, M., Ohno, S. and Kondo, N. (1994). One Subspecies of the Red Jungle fowl (*Gallus gallus gallus*) Suffices as the Matriarchic Ancestor of all Domestic Breeds. *Proceedings of the National Academy of Sciences of the United States of America*. 91: 12505-12509.
- Goodwin, D. (1970). Pigeons and doves of the world (Publication-British Museum) (p. 446). *Trustees of the British Museum*.
- Gonzales, A., Oporta M., Martinez A. and Lopez Y.C. (2000). Feed restriction and salbutamol to control ascites syndrome in broilers: 1- Productive performance and carcass traits. *Publicado Como Articulo en Agrociencia* 34(3), 283-292.
- Gonzales, E., Buyse, J., Loddi, M.M., Takita, T.S., Buys, N. and Decuyper, E. (1998). Performance, incidence of metabolic disturbances and endocrine variables of food-restricted male broiler chickens. *British poultry science*, 39(5), 671-678.
- Gezici, M. (2002). Skin and epithermoidal features. In *Anatomy of Domestic Birds, Turkey: Medisan pub-Lishing Company*, No 49. p. 222. Cited by Aslan, 2000.
- Getty, R. (1975). The Anatomy of Domestic Animals. 5: 255-348. *Saunders Company Philadelphia*. London. Toronto.

- Gaspers, and Vincent. (1991). *Teknik Penarikan Contohuntuk Penelitian Survey*, Tarsito, Bandung.
- Genstat. (2003). Genstat 5.0 Release 4.23DE. Lawes Agric, *Trust, Rothamsted Exp. Stn.*, UK.
- Gibson, J., Gamage, S., Hanotte, O., Iñiguez, L., Maillard, J.C., Rischkowsky, B., Semambo, D., and Toll, J. (2006). Options and strategies for the conservation of farm animal genetic resources: Report of an international workshop (7–10 November 2005, Montpellier, France). CGIAR System-wide Genetic Resources Programme (SGRP)/*Bioversity International*, Rome, Italy, 5.
- Gelis, S. (2006). Evaluating and treating the gastrointestinal system. *Clinical avian medicine*, 1, 411-440.
- Girouard, H., and Savard, R. (1998). The lack of bimodality in the effects of endogenous and exogenous prostaglandins on fat cell lipolysis in rats. *Prostaglandins*. 56: 43-52.
- Humason, G.L. (1972). Animal tissue techniques. 3rd edn. W.H. *Freeman and Company, San Francisco*, pp 180-182.
- Hayes, W.J., and Laws, R.R. (1991). *Handbook of Pesticide Toxicology*, Vol. II. New York: Academic Press, p. 791.
- Hybro. (2004). History of Hybro. Web Site (WWW.hybro.com). E.mail (Hybro@euribrid.com).
- Hassouna, E.M.A. (2001). Some anatomical and morphometrical studies on the intestinal tract of chicken, duck, goose, turkey, pigeon, dove, quail, sparrow, heron, jackdaw, hoopoe, kestrel and owl. *Assiut Veterinary Medical Journal*, 44(88), 47-78.
- Hodges, R.D. (1974). *The histology of the fowl Academic Press*, London, Kap. 2 “The Digestive System” S. 35-89.
- Haddad, E.E., and Mashaly, M.M. (1990). Effect of thyrotropin-releasing hormone, triiodothyronine, and chicken growth hormone on plasma concentrations of thyroxine, triiodothyronine, growth hormone, and growth of lymphoid organs and leukocyte populations in immature male chickens. *Poultry science*, 69(7), 1094-1102.
- Harden, R.L., and Oscar, T.P. (1993). Thyroid hormone and growth hormone regulation of broiler adipocyte lipolysis. *Poultry science*, 72(4), 669-676.
- Harvey S, Johnson CDM & Sanders EJ 2000 Extra-pituitary growth hormone in peripheral tissues of early chick embryos. *Journal of Endocrinology* 166 489–502.

- Harvey, S., R. A. Fraser, and R. W. Lea, 1991. Growth hormone secretion in poultry. *Critical Review in Poultry Biology*. 3:239–282.
- Huybrechts, L. M., E. Decuypere, J. Buyse, E. R. Kuhn, and M. Tixier-Boichard, 1992. Effect of recombinant human insulin-like growth factor-I on weight gain, fat content and hormonal parameters in broiler chickens. *Poultry Science*. 71: 181–187.
- Havenstein, G. B., P. R. Ferket, and M. A. Qureshi. 2003. Carcass composition and yield of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poultry Science* 82:1509–1518.
- Idris, L.H., (2011). Carcass composition, muscle morphology, and meat characteristics of the red jungle fowl (*Gallus gallus spadicius* Linnaeus), village chicken, and commercial broiler. PhD thesis, Veterinary Medicine Faculty, University Putra Malaysia, Malaysia.
- Ige, A.O., Akinlade, J.A., Ojedapo, L.O., Oladunjoye, I.O., Amao, S.R., and Animashaun, A.O. (2006). Effect of sex on interrelationship between body weight and linear body measurements of commercial broilers in a derived savannah environment of Nigeria. In Proc. *11th Annual Conf. of the Animal Science Association of Nigeria* (pp. 18-21).
- Julian, R.J. (1997). Causes and prevention of ascites in broilers. *Zootecnica International*, 20, 52-53.
- Jacob, J. (1992). Systematics and the analysis of integumental lipids: the uropygial gland. Bull. Br. *Ornithol. Club Suppl. A*, 112, 159-167.
- Johnston, D.W. (1988). A morphological atlas of the avian uropygial gland. *Bulletin of the British Museum of Natural History (Zoology)* 54: 199-259.
- Julian, R.J. (1998). Rapid growth problems: ascites and skeletal deformities in broilers. *Poultry Science* 77: 1773-1780.
- Jacob, J., and Ziswiler, V. (1982). The uropygial gland. *Avian Biology*, 6, 199-324.
- Johnston, D.W. (1988). Morphological atlas of the avian uropygial gland. *British Museum (Natural History)*. 54(5): 199_259.
- Jacob, J., Eigener, U., and Hoppe, U. (1997). The structure of preen gland waxes from peleciform birds containing 3, 7-dimethyloctan-1-ol-An active ingredient against dermatophytes. *Zeitschrift für Naturforschung C*, 52(1-2), 114-123.
- Jacob M.C. (1976). Bird waxes. In Kolattukudy, PE. (Ed.). *Chemistry and Biochemistry of Natural Waxes*. Amsterdam: Elsevier. Pp. 93-146.
- Jawad, H.S., Idris, L.H.B., Naji, S.A., Bakar, M.B., and Kasim, A.B. (2015). Partial Ablation of Uropygial Gland Effect on Production Performance of Akar Putra Chicken. *International Journal of Poultry Science*, 14(4), 213-221.

- Jahan, M. S., Asaduzzaman, M., and Sarkar, A. K. (2006). Performance of broiler fed on mash, pellet and crumble. *International Journal of Poultry Science*, 5(3), 265-270.
- Jackson, S. and Dimond, J.M. (1996). Metabolic and digestive response to artificial selection in chickens. *Evolution* 50: 1638-1650.
- Kasim, A., 2007. University Putra Malaysia website
<http://www.upm.edu.my/berita/details/Bakabaruayamkacukanbi?LANG=en>.
- King, A.S. and McLelland, J. (1984). *Birds Their Structure and Function*. Second Ed. Baillie Tindall. London, Pp. 218_275.
- Kennedy, R.J. (1971). Preen gland weights. *Ibis*, 113(3), 369-372.
- Klasing, K.C. (1999). Avian gastrointestinal anatomy and physiology. In *Seminars in Avian and Exotic Pet Medicine* (Vol. 8, No. 2, pp. 42-50). WB Saunders.
- Kermanshahi, H., Nassiri, M.R., and Heravi Moussavi, A. (2010). Effect of feed restriction and different energy and protein levels of the diet on growth performance and growth hormone in broiler chickens. *Journal of Biological Science*, 10.
- King, D. B. and J. D. May. 1984. Thyroidal influence on body growth. *Journal of Experimental Zoology*. 232:453.
- Kuhn, E. R. 1993. Role of growth hormone in thyroid function of vertebrates. *Academiae Analecta* 55:3-13.
- Lucas, A.M., and Stettenheim, P.R. (1972a). Uropygial gland. *Agriculture handbook*, 362, 613-626.
- Leeson, S., and Summers, J.D. (1997). *Commercial poultry Nutrition*, Second Edition University books, P. O. Box 1326, Guelph, Ontario, Canada.
- Law-Brown, J., and Meyers, P.R. (2003). *Enterococcus phoeniculicola* sp. Nov., a novel member of the enterococci isolated from the uropygial gland of the Red-billed Woodhoopoe, *Phoeniculus purpureus*. *International journal of systematic and evolutionary microbiology*, 53(3), 683-685.
- Lucas, A.M., and Stettenheim P.R. (1972b). Avian anatomy. *Integument. Agriculture Handbook* 362, U.S. Dept. Agric., Washington, D.C.
- Liyanage, R.P., Dematawewa, C.M.B., and Silva, G.L.L.P. (2014). Comparative study on morphological and morphometric features of village chicken in Sri Lanka. *Tropical Agricultural Research*, 26(2), 262-274.
- McLelland, J. (1990). *A Color Atlas of Avian Anatomy*. Wolfe Medical Publications Ltd. Pp. 18.

- Montalti, D., Gutierrez, A.M., Reboredo, G., and Salibian, A. (2000). Ablación de la glándula uropigia y sobrevida de *Columba livia*. *Bollettino del Museo Civico di Storia naturale di Venezia* 50: 263-266.
- Moyer, B.R., Rock, A.N., and Clayton, D.H. (2003a). Experimental test of the importance of preen oil in rock doves (*Columba livia*). *Auk*. vol. 120, no. 2, p. 490-496.
- Montalti, D., and Saliban, A. (2000). Uropygial gland size and avian habitat. *Ornitologia Neotropical* 11:297_306.
- Montalti, D., Quiroga, A., Massone, A., Idiart, J.R., and Salibián, A. (2001). Histochemical and lectin histochemical studies on the gland of rock dove *Columba livia*. *Brazilian Journal of Morphological Science* 18, 33 –39.
- Montalti, Ontalti, D., Gutierrez, A.M., and Salibian, A., (1998). Técnica quirúrgica para la ablación de la glándula uropigia en la paloma casera *Columba livia*. *Revista Brasileira de Biología*. vol. 58, no. 2, p. 193-196.
- Moyer, B.R., Pacjka, A.J., and Clayton, D.H. (2003b). How birds combat ectoparasites. *Current Ornithology*. 117.
- Montalti, D., Gutiérrez, A.M., Reboredo, G.R., and Salibián, A. (2006). Removal uropygial gland does not affect serum lipids, cholesterol and calcium levels in the rock pigeon *Columba livia*. *Acta Biologica Hungarica*. 57, 295–300.
- Montalti, Ontalti, D., Gutierrez, A.M., and Salibian, A., (1998). Técnica quirúrgica para la ablación de la glándula uropigia en la paloma casera *Columba livia*. *Revista Brasileira de Biología*, vol. 58, no. 2, p. 193-196.
- Martín-Vivaldi, M., Ruiz-Rodríguez, M., José Soler, J., Manuel Peralta-Sánchez, J., Méndez, M., Valdivia, E., and Martínez-Bueno, M. (2009). Seasonal, sexual and developmental differences in hoopoe *Upupa epops* preen gland morphology and secretions: evidence for a role of bacteria. *Journal of Avian Biology*, 40(2), 191-205.
- Momoh, O.M., and Kershima, D.E. (2008). Linear body measurements as predictors of body weight in Nigerian local chickens. *ASSET: An International Journal (Series A)*, 8(2), 206-212.
- Monsi, A.L.E.X. (1992). Appraisal of interrelationships among live measurements at different ages in meat type chickens. *Nigerian Journal of Animal Production*, 19(1&2), 15-24.
- Muhiuddin, G., (1993). Estimates of genetic and phenotypic parameters of some Performance traits in beef cattle. *Animal Breeding Abstracts*. 66: 495 – 522.

- Mitchell, M.A., and Smith, M.W. (1991). The effects of genetic selection for increased growth rate on mucosal and muscle weights in the different regions of the small intestine of the domestic fowl (*Gallus domesticus*). *Comparative Biochemistry and Physiology Part A: Physiology*, 99(1), 251-258.
- Muelling, C., and Buda, S. (2002). Morphology and function of the digestive system in turkey in: *4th International Symposium on Turkey Diseases - Gießen: DVG*, S. 89-96.
- McMurtry, J. P., Francis, G., Upton, Z., Rosselot, G., Phelps, P., Czerwinski, S., Caperna, T. & Vasilatos-Younken, R. (1995) Aspects of insulin-like growth factors in poultry growth physiology. *Endocrinology*, Abstracts 77th Annual Meeting, p. 43.
- Mao, J. N., L. A. Cogburn and J. Burnside. 1997. Growth hormone down-regulates growth hormone receptor mRNA in chickens but developmental increases in growth hormone receptor mRNA occur independently of growth hormone action. *Molecular and Cellular Endocrinology*. 129:135-143.
- Nickel, R., Schummer, A., and Seiferle, E. (1977). The Skin Anatomy of the Domestic Birds. 2nd Ed., Verlag Paul Parey. *Berlin and Hamburg*.14-15.
- Naji, S.A. (2001). Surgical removal of uropygial gland (Iraqi method) to convert non-laying to laying hens. *The Iraqi Journal of Agricultural Sciences*, 32(5) '203-212.
- Naji, S.A., Al-Daraji, H.1., Al-Tikriti, B.T.O., and Al-Rawi, A.A. (2002). The effect of uropygialectomy for remedy the non-laying hens uncertain productive traits of Iraqi local chickens. *The Iraqi Journal of Agricultural Sciences*, 33(1):123-130. Cited by Al-Daraji *et al.*, 2006.
- Nasrin, M., Siddiqi, M.N.H., Masum, M.A., and Wares, M.A. (2012). Gross and histological studies of digestive tract of broilers during postnatal growth and development. *Journal of the Bangladesh Agricultural University*, 10(1), 69-77.
- Nishida, T., Hayashi, Y., Shotake, T., Maeda, Y., Yamamoto, Y., Kurosawa, Y., Douge, K. and Hongo, A. (1992). Morphological identification and ecology of the red jungle fowl in Nepal. *Animal Science and Technology* 63: 256-269.
- National Research Council (NRC). *Nutrient Requirements of Poultry*. 8th rev. ed. National Academic Press, Washington, DC. 1984.
- Nishimura, T., A. Hattori, and K. Takahashi. 1999. Structural changes in intramuscular connective tissue during the fattening of Japanese Black cattle: effect of marbling on beef tenderization. *Journal of Animal Science*. 77:93–104.
- Oke, U.K., Herbert, U., and Nwachukwu, E.N. (2004). Association between body weight and some egg production traits in the guinea fowl (*Numida meleagris galatea pallas*). *Livestock Research for Rural Development*, 16(9), 6.

- Prum, R.O., Torres, R., Williamson, S., and Dyck, J. (1999). Two-dimensional Fourier analysis of the spongy medullary keratin of structurally colored feather barbs. *Proceedings of the Royal Society of London B: Biological Sciences*, 266(1414), 13-22.
- Philip, J.S. (1970). *Poultry Feeds and Nutrition*. The Avi publishing co. Inc., West Port Connection USA.
- Pettingill, O.S. (1985). *Ornithology in laboratory and field*. Burgess, Minneapolis, Minnesota, USA, *Academic Press Inc.* pp. 378–380.
- Parmer, T.G., Carew, L.B., Alster, F.A., and Scanes, C.G. (1987). Thyroid function, growth hormone, and organ growth in broilers deficient in phosphorus. *Poultry Science*, 66(12), 1995-2004.
- Proudman, J.A., Krishnan, K.A., and Maruyama, K. (1995). Ontogeny of pituitary and serum growth hormone in growing turkeys as measured by radioimmunoassay and radioreceptor assay. *Poultry science*, 74(7), 1201-1208.
- Phornphutkul, C., K. Y. Wu, X. Yang, Q. Chen, and P. A. Gruppuso. 2004. Insulin-like growth factor-I signaling is modified during chondrocyte differentiation. *Journal of Endocrinology*. 183: 477-486.
- Qureshi, M.A., and Havenstein, G.B. (1994). A comparison of the immune performance of a 1991 commercial broiler with a 1957 random bred strain when fed typical 1957 and 1991 broiler diets. *Poultry Science* 73: 312-319.
- Qi, K. K., Chen, J. L., Zhao, G. P., Zheng, M. Q., & Wen, J. (2010). Effect of dietary $\omega 6/\omega 3$ on growth performance, carcass traits, meat quality and fatty acid profiles of Beijing-you chicken. *Journal of animal physiology and animal nutrition*, 94(4), 474-485.
- Robinson, F.E., Classen H.L., Hanson J.A., and Onderka, D.K. (1992). Growth performance, feed efficiency and the incidence of skeletal and metabolic disease in full-fed and feed-restricted broiler and roaster chickens. *Journal of Applied Poultry Research* 1: 33-41.
- Raud, H., and Faure, J.M. (1994). Welfare of ducks in intensive units. *Revue scientifique technique (International Office of Epizootics)*, 13(1), 119.
- Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and Desante, D.F. (1993). *Handbook of field methods for monitoring landbirds*. *USDA Forest Service/UNL Faculty Publications*, 105.
- Rosenberg, L.E. (1941). Microanatomy of the duodenum of the turkey. *Hilgardia*, 13(11), 623-654.
- Robinson, F.E., and Renema, R. A. (2000). *Alberta Poultry Research Centre*, University of Alberta Edmonton, AB, Canada T6G 2P5. www.spottedcowpress.ca/chapters/02FemaleAnatomy.pdf

- Schultz, A., Laschat, S., Morr, M., Diele, S., Dreyer, M., and Bringmann, G. (2002). Highly Branched Alkanoic Acids from the Preen-Gland Wax of the Domestic Goose as Building Blocks for Chiral Triphenylenes. *Helvetica chimica acta*, 85(11), 3909-3918.
- Sawad, A.A. (2006a). Morphological and histological study of uropygial gland in Moorhen (*G. gallinula* C. Choropus). *International journal of poultry science*, 5(10), 938-941.
- Soler, J.J., Peralta J.M., Martín A.M., Martín V.M., Martínez, B.M., and Møller, A.P. (2012). The evolution of size of the uropygial gland: mutualistic feather mites and uropygial secretion reduce bacterial loads of eggshells and hatching failures of European birds. *Journal of evolutionary biology*, 25(9), 1779-1791.
- Scott, M.L., and Dean, W.F. (1991). *Nutrition and management of ducks*. 12.
- Sisson, S., (1975). *The Anatomy of Domestic Animals*, 5th Ed., W. B. Saunders Company. pp. 2099_2093. London.
- Stettenheim, P.R. (2000). The integumentary morphology of modern birds an overview. *American Zoologist*, 40(4), 461-477.
- Sawad. (2006b). Discerning adaptive value of seasonal variation in preen waxes: comparative and experimental approaches. *Acta Zoologica Sinica*, vol. 52, suppl, p. 272-275.
- Shawkey, M.D., Pillai, S.R., and Hill, G.E. (2003). Chemical warfare? Effects of uropygial oil on feather-degrading bacteria. *Journal of Avian Biology*, 34(4), 345-349.
- Salibian, A., and Montalti, D. (2009). Physiological and biochemical aspects of the avian uropygial gland. *Brazilian Journal of Biology*. 69(2), 437-446.
- Scott, B. (1982). Clave del observador de aves. In S.A. Omega, ed., Barcelona. Pp: 13.
- Scanes, C.G., Telfer, S.B., Hackett, A.F., Nightingale, R., and Sharifuddin, B.A.K. (1975). *Effects of growth hormone on tissue metabolism in broiler chicks*.
- Scanes, C.G., Harvey, S., Marsh, J.A., and King, D.B. (1984). Hormones and growth in poultry. *Poultry Science*, 63(10), 2062-2074.
- Sakamoto, K., Hirose, H., Onizuka, A., Hayashi, M., Futamura, N., Kawamura, Y., and Ezaki, T. (2000). Quantitative study of changes in intestinal morphology and mucus gel on total parenteral nutrition in rats. *Journal of Surgical Research*, 94(2), 99-106.
- Starling, M.B., and Elliott, R.B. (1974). The effects of prostaglandins, prostaglandin inhibitors, and oxygen on the closure of the ductus arteriosus, pulmonary arteries and umbilical vessels in vitro. *Prostaglandins*, 8(3), 187-203.

- Sugimoto, Y., Ohta, Y., Morikawa, T., Yamashita, T., Yoshida, M., and Tamaoki, B. I. (1990). In vitro metabolism of testosterone on hepatic tissue of chicken (*Gallus domesticus*). *Journal of steroid biochemistry*, 35(2), 271-279.
- Sturkie, P.D. (1986). *Avian Physiology*. 4th edn. , Springer-Verlag, New York, Berlin, Heidelberg, Tokyo.
- Steimer, T., and Hutchinson, J. B. (1981). Metabolic control of the behavioral action of androgens in the dove brain: testosterone inactivation by 5 β reduction. *Brain Research*: 189:209.
- Sharp, P.J. (1983). Hypothalamic control of gonadotropin secretion in birds. Pages 124–176 in: *Progress in Nonmammalian Brain Research*, Vol. 3. CRC Press, Inc. Boca Raton, FL.
- Sainsbury, D. (1984): Systems of management. Ch.9, P.102. In *Poultry Health and Management*. 2nd Ed. By Sainsbury. Granada Publishing LTD. 8 Grafton Street, London W1X3 LA.
- Taylor, R.D., and Jones, G.P.D. (2004). The incorporation of whole grain into pelleted broiler chicken diets. II. Gastrointestinal and digesta characteristics. *British poultry science*, 45(2), 237-246.
- Tsafiriri, A., Lindner, H.R., Zor, U., and Lamprecht, S.A. (1972). Physiological role of prostaglandins in the induction of ovulation. *Prostaglandins*, 2(1), 1-10.
- Thomas, H.R., Miell, J.P., Taylor, A.M., 1993. Endocrine and cardiac paracrine actions of insulin-like growth factor-I (IGF-I) during thyroid dysfunction in the rat: is IGF-I implicated in the mechanism of heart weight/body weight change during abnormal thyroid function? *Journal of Molecular Endocrinology*. 10, 313–323.
- Udeh, I., Ugwu, S.O.C., and Ogagifo, N.L. (2011). Predicting semen traits of local and exotic cocks using linear body measurements. *Asian Journal of Animal Science*, 5(4), 268-276.
- Vincze, O., Vágási, C.I., Kovács, I., Galván, I., and Pap, P.L. (2013). Sources of variation in uropygial gland size in European birds. *Biological Journal of the Linnean Society*, 110(3), 543-563.
- Verdal, H., Mignon-Grasteau, S., Jeulin, C., Le Bihan-Duval, E., Leconte, M., Mallet, S., and Narcy, A. (2010). Digestive tract measurements and histological adaptation in broiler lines divergently selected for digestive efficiency. *Poultry science*, 89(9), 1955-1961.
- Vasilatos-Younken, R., and Scanes, C.G. (1991). Growth hormone and insulin-like growth factors in poultry growth: required, optimal, or ineffective? *Poultry science*, 70(8), 1764-1780.

- Vasilatos-Younken, R., Wang, X.H., Zhou, Y., Day, J.R., McMurtry, J.P., Rosebrough, R.W., and Tomas, F. (1999). New insights into the mechanism and actions of growth hormone (GH) in poultry. *Domestic animal endocrinology*, 17(2), 181-190.
- Vanderpooten, A., L. M. Huybrechts, P. Rudas, E. Decuypere and E. R. Kuhn. 1991a. Differences in hepatic growth hormone receptor binding during development of normal and dwarf chickens. *Reproduction Nutrition Development*. 31:47-55.
- Vanderpooten, A., V. M. Darras, L. M. Huybrechts, P. Rudas, E. Decuypere and E. R. Kuhn. 1991b. Effect of hypophysectomy and acute administration of growth hormone (GH) and GHreceptor binding in chick liver membranes. *Journal of Endocrinology*. 129:275-281.
- Wepruk, J., and Church, S. (2003). Balancing production and welfare. Complex animal care issues. *Alberta Farm Animal Care (AFAC)*. Association 2-8.
- Watts, G., and Kennett, C. (1995). The broiler industry. *Poultry Tribune, Centennial Edition*. Watt.
- Wang, C.Y., Wang, Y., Li, J., and Leung, F.C. (2006). Expression profiles of growth hormone-releasing hormone and growth hormone-releasing hormone receptor during chicken embryonic pituitary development. *Poultry science*, 85(3), 569-576.
- William, J., and Linda, M. (2000). Color atlas of veterinary histology, second edition, *Awolter Klumer Company*.
- Walzem, R.L. (1996). Lipoproteins and laying hens: Form follows function. *Poultry. Avian Biology Rev.* 7:31-64.
- Wiltbank, M.C., Gallagher, K. P., and Dysko, R.C. (1989). Regulation of blood flow to the rabbit corpus luteum: Effects of estradiol and human chorionic gonadotropin. *Endocrinology* 124:605-611.
- Yamauchi, K.E., and Isshiki, Y. (1991). Scanning electron microscopic observations on the intestinal villi in growing White Leghorn and broiler chickens from 1 to 30 days of age. *British Poultry Science*, 32(1), 67-78.
- Zheng, J.X., Liu, Z.Z., and Yang, N. (2007). Deficiency of growth hormone receptor does not affect male reproduction in dwarf chickens. *Poultry science*, 86(1), 112-117.
- Ziswiler, V., and Farner, D.S. (1972). Digestion and digestive system. *Avian Biology*. Vol 2. pp: 343-430. Academic press. London.
- Zerehdaran, S., Vereijken, A. J., Van Arendonk, J. A. M., & van Der Waaijt, E. H. (2004). Estimation of genetic parameters for fat deposition and carcass traits in broilers. *Poultry Science*, 83(4), 521-525.

APPENDICES

CHAPTER 4

Appendix 4.1




Figure 4.1: Graph of males' body weight effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.2




Figure 4.2: Graph of males' feed intake effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.3




Figure 4.3: Graph of males' weight gain effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.4




Figure 4.4: Graph of males' feed conversion ratio effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.5




Figure 4.5: Graph of females' body weight effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.6




Figure 4.6: Graph of males' feed intake effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.7




Figure 4.7: Graph of females' weight gain effectiveness according to the proceeding time of uropygialectomy.

Appendix 4.8




Figure 4.8: Graph of females' feed conversion ratio effectiveness according to the proceeding time of uropygialectomy.

CHAPTER 6

Appendix 6.1









Figure 6.1: Photographs of external morphological characteristics measuring process. Bird length (BL): 2.1A; Neck length (NL): 2.1B; Back length (BAL): 2.1C; Breast length (BRL): 2.1D; Breast width (BRW): 2.1E; Keel of sternum length (KBL): 2.1F; Fore limbs length (WL): 2.1G; Hand limbs length (legs) (LL): 2.1H; Comb width (CW): 2.1I; Comb length (CL): 2.1J; Beak width (BEW): 2.1K). Beak length (BEL): Length from the tip of the beak until insertion of the beak into the skull (Appendix 2.1L; Wattles' width (WAW): 2.1M; Wattles' length (WAL): 2.1N.

Appendix 6.2






Figure 6.2: Graph of linear development of the morphological characteristics based on the age of measurement (7, 10, 12 weeks). BW: Bird weight; BL: Bird length; NL: Neck length; BAL: Back length; BRD: Breast diameters; KBL: Keel bone length; WL: wing length; LL: Leg length; CD: Comb diameters; BED: Beak diameters; WAD: Wattle diameters; L: Length; W: Wide.

CHAPTER 7

Appendix 7.1




Figure 7.1: Parts of the digestive tract of a chicken (McLelland, 1990).

CHAPTER 8

Appendix 8.1



Figure 8.1. Photographs of blood collection method.

Appendix 8.2

CHICKEN GROWTH HORMONE (GH) ELISA Kit

Catalog Number. CSB-E09866Ch

For the quantitative determination of chicken growth hormone (GH) concentrations in serum, plasma, tissue homogenates.

PRINCIPLE OF THE ASSAY

This assay employs the competitive inhibition enzyme immunoassay technique. The microtiter plate provided in this kit has been pre-coated with an antibody specific to GH. Standards or samples are added to the appropriate microtiter plate wells with Biotin-conjugated GH. A competitive inhibition reaction is launched between GH (Standards or samples) and Biotin-conjugated GH with the pre-coated antibody specific for GH. The more amount of GH in samples, the less antibody bound by Biotin-conjugated GH. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Substrate solution is added to the wells and the color develops in opposite to the amount of GH in the sample. The color development is stopped and the intensity of the color is measured.

DETECTION RANGE

625 pg/ml-10000 pg/ml.

SENSITIVITY

The minimum detectable dose of chicken GH is typically less than 312.5 pg/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest chicken GH concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.

SPECIFICITY

This assay has high sensitivity and excellent specificity for detection of chicken GH. No significant cross-reactivity or interference between chicken GH and analogues was observed.

Note: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between chicken GH and all the analogues, therefore, cross reaction may still exist.

PRECISION

Intra-assay Precision (Precision within an assay): CV % < 15% Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV % < 15% Three samples of known concentration were tested in twenty assays to assess.

LIMITATIONS OF THE PROCEDURE

- FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.
- The kit should not be used beyond the expiration date on the kit label.
- Do not mix or substitute reagents with those from other lots or sources.
- If samples generate values higher than the highest standard, dilute the samples and repeat the assay.
- Any variation in operator, pipetting technique, washing technique, incubation time or temperature, and kit age can cause variation in binding.
- This assay is designed to eliminate interference by soluble receptors, binding proteins, and other factors present in biological samples. Until all factors have been tested in the Immunoassay, the possibility of interference cannot be excluded.

MATERIALS PROVIDED

Reagents	Quantity
Assay plate	1(96 wells)
Standard	5 x 1 ml
Conjugate	1 x 6 ml
HRP-avidin	1 x 6 ml
Wash Buffer (20 x concentrate)	1 x 15 ml
Substrate A	1 x 7 ml
Substrate B	1 x 7 ml
Stop Solution	1 x 7 ml
Adhesive Strip (For 96 wells)	4
Instruction manual	1

STANDARD CONCENTRATION

Standard	S1	S2	S3	S4	S5
Concentration (pg/ml)	625	1250	2500	5000	10000

STORAGE

Unopened kit	Store at 2 - 8°C. Do not use the kit beyond the expiration date.
Opened kit	May be stored for up to 1 month at 2 - 8° C.

OTHER SUPPLIES REQUIRED

- Microplate reader capable of measuring absorbance at 450 nm, with the correction wavelength set at 600 nm - 630 nm.
- An incubator which can provide stable incubation conditions up to 37°C±0.5°C.
- Squirt bottle, manifold dispenser, or automated microplate washer.
- Absorbent paper for blotting the microtiter plate.

- 100 mL and 500 mL graduated cylinders.
- Deionized or distilled water.
- Pipettes and pipette tips.
- Test tubes for dilution.

PRECAUTIONS

The Stop Solution provided with this kit is an acid solution. Wear eye, hand, face, and clothing protection when using this material.

SAMPLE COLLECTION AND STORAGE

- **Serum:** Use a serum separator tube (SST) and allow samples to clot for two hours at room temperature or overnight at 4°C before centrifugation for 15 minutes at 1000 ×g. Remove serum and assay immediately or aliquot and store samples at -20°C or -80°C. Avoid repeated freeze-thaw cycles.
- **Plasma:** Collect plasma using EDTA, or heparin as an anticoagulant. Centrifuge for 15 minutes at 1000 ×g at 2-8°C within 30 minutes of collection. Assay immediately or aliquot and store samples at -20°C or -80°C. Avoid repeated freeze-thaw cycles.
- **Tissue Homogenates:** 100mg tissue was rinsed with 1X PBS, homogenized in 1 ml of 1X PBS and stored overnight at -20°C. After two freeze-thaw cycles were performed to break the cell membranes, the homogenates were centrifuged for 5 minutes at 5000 x g, 2 - 8°C. The supernate was removed and assayed immediately. Alternatively, aliquot and store samples at -20°C or -80°C. Centrifuge the sample again after thawing before the assay. Avoid repeated freeze-thaw cycles.

ASSAY PROCEDURE

Bring all reagents and samples to room temperature before use. Centrifuge the sample again after thawing before the assay. It is recommended that all samples and standards be assayed in duplicate.


1. Prepare all reagents and samples as directed in the previous sections.
2. Determine the number of wells to be used and put any remaining wells and the desiccant back into the pouch and seal the ziploc, store unused wells at 4°C.
3. Set a Blank well without any solution.
4. Add 50µl of Standard or Sample per well. Standard need test in duplicate.
5. Add 50µl of Conjugate to each well (not to Blank well). Mix well and then incubate for 60 minutes at 37°C.
6. Aspirate each well and wash, repeating the process two times for a total of three washes. Wash by filling each well with Wash Buffer (200µl) using a squirt bottle, multi-channel pipette, manifold dispenser, or autowasher, and let it stand for 10 seconds, complete removal of liquid at each step is essential to good performance. After the last wash, remove any remaining Wash Buffer by aspirating or decanting. Invert the plate and blot it against clean paper towels.
7. Add 50µl of HRP-avidin to each well (not to Blank well). Mix well and then incubate for 30 minutes at 37°C.
8. Repeat the aspiration/wash process for three times as in step 6.

9. Add 50µl of Substrate A and 50µl of Substrate B to each well, mix well. Incubate for 15 minutes at 37°C. Keeping the plate away from drafts and other temperature fluctuations in the dark.
10. Add 50µl of Stop Solution to each well, gently tap the plate to ensure thorough mixing.
11. Determine the optical density of each well within 10 minutes, using a microplate reader set to 450 nm.

Note:

1. The final experimental results will be closely related to validity of the products, operation skills of the end users and the experimental environments.
2. Samples or reagents addition: Please carefully add samples to wells and mix gently to avoid foaming. Do not touch the well wall as possible. For each step in the procedure, total dispensing time for addition of reagents or samples to the assay plate should not exceed 10 minutes. This will ensure equal elapsed time for each pipetting step, without interruption. Duplication of all standards and specimens, although not required, is recommended. To avoid cross-contamination, change pipette tips between additions of each standard level, between sample additions, and between reagent additions. Also, use separate reservoirs for each reagent.
3. Incubation: To ensure accurate results, proper adhesion of plate sealers during incubation steps is necessary. Do not allow wells to sit uncovered for extended periods between incubation steps. Once reagents have been added to the well strips, DO NOT let the strips DRY at any time during the assay. Incubation time and temperature must be observed.
4. Washing: The wash procedure is critical. Complete removal of liquid at each step is essential to good performance. After the last wash, remove any remaining Wash Solution by aspirating or decanting and remove any drop of water and fingerprint on the bottom of the plate. Insufficient washing will result in poor precision and falsely elevated absorbance reading. When using an automated plate washer, adding a 30 second soak period following the addition of wash buffer, and/or rotating the plate 180 degrees between wash steps may improve assay precision.
5. Controlling of reaction time: Observe the change of color after adding Substrates (e.g. observation once every 10 minutes). Substrates should change from colorless or light blue to gradations of blue. If the color is too deep, add Stop Solution in advance to avoid excessively strong reaction which will result in inaccurate absorbance reading.
6. Substrates are easily contaminated. Substrates should remain colorless or light blue until added to the plate. Please protect it from light.
7. Stop Solution should be added to the plate in the same order as the Substrates. The color developed in the wells will turn from blue to yellow upon addition of the Stop Solution. Wells that are green in color indicate that the Stop Solution has not mixed thoroughly with the Substrates.

ASSAY PROCEDURE SUMMARY



CALCULATION OF RESULTS

Using the professional soft "Curve Expert 1.3" to make a standard curve is recommended, which can be downloaded from our web.

Average the duplicate readings for each standard and sample and subtract the average optical density of Blank. Create a standard curve by reducing the data using computer software capable of generating a four parameter logistic (4-PL) curve-fit. As an alternative, construct a standard curve by plotting the mean absorbance for each standard on the x-axis against the concentration on the y-axis and draw a best fit curve through the points on the graph. The data may be linearized by plotting the log of the GH concentrations versus the log of the O.D. and the best fit line can be determined by regression analysis. This procedure will produce an adequate but less precise fit of the data. If samples have been diluted, the concentration read from the standard curve must be multiplied by the dilution factor.

Appendix 8.3

Chemical preparation for light microscopy

1- 10% Neutral Buffered Formalin (NBF)

Formalin (37-40% Formaldehyde)	100.00 ml
Sodium Phosphate Monobasic	4.00 g
Sodium Phosphate Dibasic (anhydrous)	6.50 g
Distilled water	900.00 ml

Dissolved the sodium phosphate monobasic and sodium phosphate dibasic in 900 ml distilled water and add 100 ml of 40% formalin.

2- Harris Hematoxylin and Eosin

Hematoxylin	1.0 g	
Potassium alum (KAI (SO ₄) ₂ .12H ₂ O)		20 g
Mercuric oxide		0.5 g
Ethyl alcohol		10 ml
Distilled water	200 ml	

Dissolve hematoxylin in ethyl alcohol. Dissolve potassium alum in water and boil. Add hematoxylin and boil ½ minute. Add mercuric oxide. Cool rapidly. Add a few drops of acetic acid.

Eosin

Eosin Y (C.I. 45380)	1.0 g
Potassium dichromate	0.5 g
Saturated aqueous picric acid	10.0 ml
Absolute ethyl alcohol	10.0 ml
Distilled water	80.0 ml
Acetic acid	1 drop

Procedure

- 1- Xylene: 2-3 minutes (two changes).
- 2- absolute alcohol: 2-3 minutes.
- 3- 95% alcohol: 2-3 minutes.
- 4- 80% alcohol: 2-3 minutes.
- 5- 75% alcohol: 2-3 minutes.
- 6- Running water: 3 minutes.
- 7- Harris hematoxylin: 2-5 minutes. Check after 1 minute.
- 8- Running water: 3-5 minutes.
- 9- Counterstain: eosin or 0.5% aqueous phloxin B.
- 10- Dehydration through series of alcohol 70%, 95%, absolute alcohol.
- 11- Xylene: 2-3 minutes two changes.
- 12- Mounting medium and add cover glass.

BIODATA OF STUDENT

The student was born on March 3, 1983 in Baghdad, Iraq. He attended his primary and secondary school in his hometown, Baghdad. He pursued his study in Veterinary Medicine at University of Baghdad. He completed his DVM degree in 2006, ranking 85th out of 177. At the same year, he started his post-graduate study for MSc in the Department of Anatomy and Histology at the Faculty of Veterinary Medicine, University of Baghdad, and obtained the degree in 2008. After that, he was directly appointed in Department of Animal Production of the Agriculture faculty. Thereafter, he served at Agriculture Faculty as an assistant lecturer until 2012, and then he was promoted to lecturer. His interest in academic and research fields led him to continue his study for a PhD degree in Anatomy and Histology at University Putra Malaysia (UPM) under the supervision of Dr. Lokman Hakim Idris at Faculty of Veterinary Medicine, University Putra Malaysia in 2013.

LIST OF PUBLICATIONS

- Jawad, H.S., Idris, L.H.B., Naji, S.A., Bakar, M.B., and Kasim, A.B. (2015). Partial Ablation of Uropygial Gland Effect on Production Performance of Akar Putra Chicken. *International Journal of Poultry Science*, 14(4), 213-221.
- Jawad, H.S., Idris, L.H.B., Bakar, M.B., and Kasim, A.B. (2015). Anatomical Changes of Akar Putra Chicken Digestive System after Partial Ablation of Uropygial Gland. *American Journal of Animal and Veterinary Sciences*, 10(4), 217-229.
- Jawad, H.S., Idris, L.H.B., Naji, S.A., Bakar, M.B., and Kasim, A.B. (2015). Anatomical Changes of Akar Putra Chicken Digestive System after Partial Ablation of Uropygial Gland. Paper participated as poster in the 2^{ed} World Veterinary Poultry Association and World Poultry Science Association (Malaysia Branch) Scientific Conference, "Enhancing Innovation in Poultry Health and Production", ISBN 978-983-2408-28-4 21-22th September 2015. Kuala Lumpur convention Centre.
- Jawad, H.S., Lokman, I.H., Zuki, A.B.Z., and Kasim, A.B. (2016). Partial ablation of uropygial gland effects on growth hormone concentration and digestive system histometrical aspect of Akar Putra chicken. *Poultry Science*, 95(4): 966-973. doi: 10.3382/ps/pev444.
- Jawad, H.S., Lokman, I.H., Zuki, A.B.Z., and Kasim, A.B. (2016). Partial ablation of uropygial gland effects on carcass characteristics of Akar Putra chicken. *Poultry Science*. doi: 10/3382/ps/pev125.



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

GROWTH, CARCASS AND DIGESTIVE SYSTEM CHANGES IN HYBRID VILLAGE
CHICKENS AFTER UROPYGIALECTOMY

NAME OF STUDENT : HASAN SAAD ABDULHUSSEIN JAWAD

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]