



UNIVERSITI PUTRA MALAYSIA

**CYTOGENETIC AND RANDOM AMPLIFIED POLYMORPHISM DNA
ANALYSIS OF SUBSPECIES OF ASIAN ELEPHANTS (ELEPHAS
MAXIMUS)**

SAMSUL BARIAH SHARUDIN

FPV 2005 9

**CYTOGENETIC AND RANDOM AMPLIFIED POLYMORPHISM DNA
ANALYSIS OF SUBSPECIES OF ASIAN ELEPHANTS (*ELEPHAS MAXIMUS*)**

By

SAMSUL BARIAH SHARUDIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia in
Fulfilment of the Requirements for the Degree of Master of Science**

April 2005



DEDICATION

Dedicated to.....

My beloved husband and daughter

My Parents, My sisters and Brother

And all my beloved friends.....



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

CYTOGENETIC AND RANDOMLY AMPLIFIED POLYMORPHISM DNA ANALYSIS OF SUBSPECIES OF ASIAN ELEPHANTS (*ELEPHAS MAXIMUS*)

By

SAMSUL BARIAH SHARUDIN

April 2005

Chairman : Rosnina Hj. Yusoff, PhD

Faculty : Veterinary Medicine

A study was carried out to investigate the chromosome constitution of 3 subspecies of elephants from cultured lymphocytes; karyotypes were constructed according to standard procedure. Lymphocytes were cultured in RPMI 1640 medium supplemented with a mitogen (phytohemagglutinin-PHA or pokeweed-PWM), penicillin- streptomycin and bovine calf serum. The karyotype showed that the Asian elephant has a diploid number (2n) of 56 and fundamental number (NF) of 66 in both male and female. The autosomes comprised 6 pairs of submetacentric, 10 pairs of large acrocentric and the remaining 11 pairs are characterized as small acrocentric. On the other hand, the sex chromosomes consisted of the largest submetacentric X chromosome while the Y chromosome is a small acrocentric.



G-, C- and NORs bandings were carried out to assist the conventional banding. G-banded karyotypes of the three subspecies were identical as well as for the C- banded karyotypes. NORs banding revealed active nucleolar organizer regions chromosomes 2, 4, 13 and 17. These results were the first banded karyotypes established for the Asian elephants.

Investigations of DNA polymorphisms using RAPD technique provided identical information between the three subspecies. Blood samples from three subspecies of the Asian elephants were employed in the study. Based on band sharing frequency value of pair-wise comparison within subspecies, the genetic relationship between Malayan and Indian elephants was determined to be low and likewise between the Indian and Myanmar elephants. Interestingly, the band sharing frequency value between Malayan elephant and Myanmar elephant are genetically closely similar to the Malayan elephants compared with the Indian elephants. With regards to the similarities in chromosome morphology as well as at the DNA level, they are able to breed amongst themselves without any complication related to parental chromosomal incompatibility.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains.

**ANALISIS SITOGENETIK DAN RAPD SUBSPESIES GAJAH ASIA
(*ELEPHAS MAXIMUS*)**

Oleh

SAMSUL BARIAH SHARUDIN

April 2005

Pengerusi : Rosnina Hj. Yusoff, PhD

Fakulti : Perubatan Veterinar

Pada permulaan perkembangan metodologi, penyelidikan untuk mengkaji kromosom bagi 3 subspecies gajah Asia; Gajah Malaya (*Elephas maximus sumatranus*), Gajah Myanmar (*Elephas maximus indicus*) dan Gajah India (*Elephas maximus maximus*) dengan menggunakan teknik pengkulturan dan kariotip disediakan mengikut prosedur standard. Kombinasi kultura yang terdiri daripada media kultur (80 %) RPMI 1640, phytohemagglutinin (PHA) atau Pokeweed (PWM), streptomycin (20%) dan serum fetus anak lembu.

Hasil daripada analisis yang dijalankan didapati gajah mempunyai kromosom komplemen iaitu $2n = 56$ dan nombor fundanental (NF) bagi gajah jantan dan gajah betina adalah 66. Kariotip tersebut terdiri daripada 6 pasang submetasentrik, 10 pasang

akrosentrik besar dan selebihnya adalah 11 pasang akrosentrik kecil. Kromosom seks terdiri daripada kromosom X adalah submetasentrik yang paling besar dan kromosom Y adalah akrosentrik yang kecil.

Teknik penwarnaan G-, C- dan NORs dijalankan untuk menyokong teknik penwarnaan konvensional. Bagi teknik penwarnaan G- kariotip bagi ketiga-tiga subspesies gajah Asia ini adalah serupa, begitu juga bagi kariotip daripada teknik penwarnaan C-. Daripada teknik penwarnaan NORs, didapati NORs yang aktif adalah pada kromosom nombor 2, 4, 13 dan 17. Keputusan ujikaji ini merupakan kariotip penwarnaan yang pertama dihasilkan bagi gajah Asia.

Hasil penyelidikan polimorfasi DNA dengan menggunakan teknik RAPD menunjukkan maklumat yang sama bagi ketiga-tiga subspesies gajah Asia. Sampel darah daripada ketiga-tiga subspesies gajah Asia digunakan dalam penyelidikan ini. Berasaskan kepada nilai kekerapan perkongsian jalur terhadap perbandingan pasangan jalur yang sama di antara subspesies gajah, perhubungan genetic di antara gajah Malaya dan India telah ditentukan rendah, begitu juga dengan hubungan genetic di antara gajah India dan Myanmar. Apa yang menarik ialah nilai kekerapan perkongsian jalur di antara gajah Malaya dan gajah Myanmar adalah sangat tinggi. Ini menunjukkan bahawa gajah Myanmar adalah amat rapat secara genetic berbanding dengan gajah India.

Dengan merujuk kepada persamaan kromosom dan DNA, dengan demikian mereka boleh diperbaiki antara satu sama lain tanpa kecacatan berkaitan dengan ketidaksepadanan kromosom induk.



ACKNOWLEDGEMENTS

I would like to sincerely acknowledge the chairman of the supervisory committee, Associate Professor Dr. Rosnina Hj. Yusoff for her invaluable guidance and continuous support (not only in academic matters) throughout of this project. I am also profoundly grateful to the members of the supervisory committee, Professor Dr. Mohd Azmi Mohd Lila and Associate Professor Dr. Abd Wahid Haron, for providing prompt and helpful advice and suggestions throughout the course of any research and even after that.

My sincere appreciation also goes to Dr. Zainal Zahari Zainuddin, Dr. Aidi Mohamad and Mrs. Jawahir Jaafar from Zoo Melaka, Dr. Kevin Lazarus from Zoo Taiping and Pusat Pengurusan Gajah Lanchang, Temerloh, Pahang for providing some of the blood samples and for their technical support during the study. I would also like to thank Mr. P. Ganeshamurthi and Miss Norbazlin Md Marham who assisted me during the course of this project.

Last but not least, I would like to thank everyone who has contributed in one way or another in this project, especially to my husband, Mohd Shariman Hj. Omar for his continuous tolerance and constant encouragement gave me the inspiration not to give up at times of frustration. My heart goes to my daughter, Nur Rihhadatulnisa' Aqilah Mohd Shariman. My apology for neglecting both of you, unwittingly at times. I sincerely hope, that in due course, I shall make up for the lost time.



I certify that an Examination Committee met on 12th April 2005 to conduct the final examination of Samsul Bariah Sharudin on her Master of Science thesis entitled “Cytogenetic and Randomly Amplified Polymorphism DNA Analysis of Subspecies of Asian Elephants (*Elephas maximus*)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Kalthum Hashim, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Jalila Abu, PhD


Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Mohamad Hilmi Abdullah, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Mohd Nizam Isa, PhD

Professor
Faculty of Medicine
Universiti Teknologi MARA
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **21 JUL 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows :

Rosnina Hj. Yusoff, PhD
Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Mohd Azmi Mohd Lila, PhD
Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Abd Wahid Haron, PhD
Associate Professor.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



AINI IDERES, PhD
Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : 11 AUG 2005



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SAMSUL BARIAH SHARUDIN

Date : 20 JULY 2005



TABLES OF CONTENTS

| | Page |
|---|------|
| DEDICATION | ii |
| ABSTRACT | iii |
| ABSTRAK | v |
| ACKNOWLEDGEMENTS | viii |
| APPROVAL | ix |
| DECLARATION | xi |
| LIST OF PLATES | xv |
| LIST OF TABLES | xvi |
| LIST OF FIGURES | xvii |
| LIST OF ABBREVIATIONS | xix |
| | |
| CHAPTER | |
| I INTRODUCTION | 1 |
| II LITERATURE REVIEW | 6 |
| Elephant | 6 |
| Taxonomic Classification | |
| Phenotype | |
| Chromosomes | 9 |
| Chromosomes Preparation Technique | 10 |
| Harvesting | |
| Karyotype | 11 |
| Karyotype of Asian Elephant | |
| Chromosome Banding | 13 |
| General View | |
| Centromeric (C) – banding | |
| Giemsa (G) – banding | |
| Nucleolar Organizer Regions (NOR) – banding | |
| DNA Molecule and Structure of DNA | 19 |
| Genetic Polymorphism | 19 |
| DNA Fingerprinting | 20 |
| DNA Fingerprinting Analysis | |
| Polymerase Chain Reaction (PCR) | 22 |
| Genetic Marker | 23 |
| Amplified Fragment Length Polymorphism (AFLP) | |
| Restriction Fragment Length Polymorphism (RFLP) | |



| | | |
|------------|--|-----------|
| | Microsatellites | |
| | Minisatellites | |
| | Random Amplified Polymorphism DNA (RAPD) | |
| | Interpretation of Polymorphic Results | 32 |
| III | MATERIALS AND METHODS | 34 |
| | Materials | 34 |
| | Animals and Samples | |
| | Cytogenetic Procedure | 34 |
| | Blood Culture | |
| | Collection of Samples | |
| | Culture Conditions | |
| | Harvesting The Culture | |
| | Preparation Of Slides For Staining | |
| | Banding Techniques | 40 |
| | Giemsa (G -) Banding Technique | |
| | Centromeric (C -) Banding Technique | |
| | Nucleolar Organizer Regions (NORs -) Banding Technique | |
| | Photography And Construction Of Karyotypes | 41 |
| | Molecular Biology Procedure | 42 |
| | Blood Samples And Isolation Of DNA | |
| | Purity Of DNA | |
| | Gel Preparation And Electrophoresis | |
| | RAPD Procedure | 45 |
| | PCR Amplification And RAPD Procedures | |
| | Detection Of Amplified Products | |
| | Primers For RAPD | |
| | Band Scoring And Statistical Analysis | 49 |
| IV | RESULTS | 50 |
| | Cytogenetic Studies | 50 |
| | Mitogen | |
| | Chromosome Compliments and Karyotypes | |
| | Banded Karyotypes | 58 |
| | Giemsa (G-) Banding | |
| | Centromeric (C-) Banding | |
| | Nucleolar Organizer Regions (NORs -) Banding | 68 |
| | Molecular Biology Studies | 72 |
| | RAPD Analysis | 72 |
| | Different Reactions Condition | |
| | Reproducibility Testing | |
| | Comparison Between Primers | |



| | | |
|----|--|-----|
| V | DISCUSSION | 90 |
| | Chromosome Preparation | 90 |
| | Mitotic Effect Of Pokeweed Mitogen (PWM) | |
| | Chromosome Number and Conventional Karyotype | 91 |
| | Chromosome Banding | 93 |
| | Giemsa (G-) Banding | |
| | Centromeric (C-) Banding | |
| | Nucleolar Organizer Regions (NORs-) Banding | |
| | RAPD Analysis | 100 |
| VI | CONCLUSION | 107 |
| | REFERENCES | 110 |
| | BIODATA OF THE AUTHOR | 124 |



LIST OF PLATES

| | Page |
|--|------|
| Plate 1 : The Asian Elephant | 7 |
| Plate 2 : Inserting the butterfly catheter into the ear vein | 36 |
| Plate 3 : Blood sample collected into a heparin tube | 37 |



LIST OF TABLES

| | Page |
|--|------|
| Table 1 : Composition of 50 primers (Genemed)® used for amplification of genomic DNA | 47 |
| Table 2 : Composition of 20 primers (OPA)® used for the amplification of genomic DNA | 48 |
| Table 3 : Quality of metaphase spreads using 2 different mitogen | 51 |
| Table 4 : A List of primers for successful amplification via RAPD | 75 |
| Table 5 : Similarity index of the 3 subspecies of Asian elephants | 76 |
| Table 6 : Reproducibility of polymorphisms in different subspecies for 10 different primers. | 78 |
| Table 7 : Similarity index of the male and female <i>E. m. sumatranus</i> of Asian elephant | 78 |



LIST OF FIGURES

| | Page |
|--|------|
| Figure 1 : Frequency distribution of chromosomes in each diploid set of three subspecies of Asian elephant | 52 |
| Figure 2 : Conventional karyotype of the male Malayan elephant (<i>E. m. sumatranus</i>) | 54 |
| Figure 3 : Conventional karyotype of the female Malayan elephant (<i>E. m. sumatranus</i>) | 55 |
| Figure 4 : Conventional karyotype of the female Indian elephant (<i>E. m. maximus</i>) | 56 |
| Figure 5 : Conventional karyotype of the female Myanmar elephant (<i>E. m. indicus</i>) | 57 |
| Figure 6 : G- Banded karyotype of the male Malayan elephant (<i>E. m. sumatranus</i>) | 59 |
| Figure 7 : G- Banded karyotype of the female Malayan elephant (<i>E. m. sumatranus</i>) | 60 |
| Figure 8 : G- Banded karyotype of the female Indian elephant (<i>E. m. maximus</i>) | 61 |
| Figure 9 : G- Banded karyotype of the female Myanmar elephant (<i>E. m. indicus</i>) | 62 |
| Figure 10 : C- Banded karyotype of the female Malayan elephant (<i>E. m. sumatranus</i>) | 65 |
| Figure 11 : C- Banded karyotype of the female Indian elephant (<i>E. m. maximus</i>) | 66 |
| Figure 12 : C- Banded karyotype of the female Myanmar elephant (<i>E. m. indicus</i>) | 67 |



| | Page |
|---|------|
| Figure 13 : NORs banded karyotype of the male Malayan elephant (<i>E. m. sumatranus</i>) | 69 |
| Figure 14 : NORs banded karyotype of the male Indian elephant (<i>E. m. maximus</i>) | 70 |
| Figure 15 : NORs banded karyotype of the female Myanmar elephant (<i>E. m. indicus</i>) | 71 |
| Figure 16 : RAPD patterns using Primer 01 (P1) | 80 |
| Figure 17 : RAPD patterns using Primer 06 (P6) | 81 |
| Figure 18 : RAPD patterns using Primer 16 (P16) | 82 |
| Figure 19 : RAPD patterns using Primer 23 (P23) | 83 |
| Figure 20 : RAPD patterns using Primer 25 (P25) | 84 |
| Figure 21 : RAPD patterns using Primer 33 (P33) | 85 |
| Figure 22 : RAPD patterns using Primer 43 (P43) | 86 |
| Figure 23 : RAPD patterns using Primer 46 (P46) | 87 |
| Figure 24 : RAPD patterns using Primer 47 (P47) | 88 |
| Figure 25 : RAPD patterns using Primer 48 (P48) | 89 |



LIST OF ABBREVIATIONS

| | |
|-------------------------------|--|
| AFLP | Amplified Fragment Length Polymorphism |
| AgNO ₃ | Silver/ Argentum nitrate |
| Ba(OH) ₂ | Barium hydroxide |
| Bp | Base pairs |
| dH ₂ O | Distilled water |
| DNA | Deoxybonucleic acid |
| DNTP | Deoxynucleoside triphosphate |
| EDTA | Dissodium ethylene diamine tetraacetate |
| H ₂ O ₂ | Hydrogen peroxide |
| ISCNDA | International System for Cytogenetic Nomenclature of Domestic Animals |
| IUCN | International Union for the Conservation of Nature |
| KCL | Potassium chloride |
| NaCl | Sodium chloride |
| NORs | Nucleolar Organizer Regions |
| mM | milimolar |
| ng | nanogram |
| OD | Optical density |
| PBS | Phosphate buffer saline |
| PCR | Polymerase Chain Reaction |
| PHA | Phytohemagglutinin A |



| | |
|------|--|
| PWM | Pokeweed mitogen |
| RAPD | Random Amplified Polymorphism DNA |
| RE | Restriction enzymes |
| RFLP | Restriction Fragment Length Polymorphism |
| RNA | Ribonucleic acid |
| Rpm | Rotor per minute |
| RPMI | Roswell Park Memorial Institute |
| RRNA | Ribosomal RNA |
| SSC | Standard saline citrate (sodium chloride and nitrate solution) |
| TAE | Tris Acetate EDTA |
| ul | microliter |
| V | volt |



CHAPTER I

INTRODUCTION

Throughout history, the elephant has played a crucial role in both the animal and the plant ecosystems. Elephants modify the habitat by converting woodlands to grasslands, providing water for other species by digging water holes in dry riverbeds, acting as seeds dispersers through fecal matter and creating fire-breakers and rain water conduits through their paths. They also influence the economy, religion and culture of mankind. Therefore, they are considered as keystone species (Wilson, 1993; Shoshani, 1998) whereby the existence of a large number of other species in the ecosystem depends on the elephants such as deer, wild cattle and other small mammals like porcupine and rabbits. If these elephants are extirpated from a system, the species they supported and species that are dependent on the elephants also will disappear. Shrinking off their habitats will lower the biodiversity and affect all 'players' in a food web (Wilson, 1993).

The elephant is classified under the order, Proboscidea, family Elephantidae, subfamilies Stegotetra belontidae and Elephantinae, with a total of six genera and 26 species. However, there are only 2 genera existing in this world today. The two species of elephants that exist today are the African elephant (*Loxodonta africana*) and the Asian elephant (*Elephas maximus*), which are the end results of over fifty million years of evolution. The other 4 genera are extinct and the elephants in these 4 genera are thought to possess a well-developed trunk (Burton, 1980). Criteria used to distinguish between the 2 genera of elephants are their physical appearance, size, morphology, habitat and



also their gestation period or physiology. Although the African and Asian elephants descended from the same prehistoric elephant family, *Mammuthus* (the huge hairy mammoth), the hairy skin of the Asian elephant indicates that it is closely related to the mammoth than the African elephant (Ozawa *et al.*, 1997; Noro *et al.*, 1998; Greenwood *et al.*, 1999). Recently, a study on two species of African elephants, (*Loxodonta africana africana* and *Loxodonta africana cyclotis*) revealed that the phylogenetic distinction between the *L. africana africana* and *L. africana cyclotis* is most likely similar of *L. africana cyclotis* and *E. maximus*. The two species of African elephants have been proven to be two separate species within the family of *Loxodonta* (Roca *et al.*, 2001).

The immense size, strength and stature of this largest living land animal have intrigued people of many cultures for hundreds of years. Elephants have served as beasts of burden, entertainer in circuses and festivals around the world. Considered easier to handle and train, the Asian elephant is widely used as a draught animal in India and Southeast Asia. The range of Asian elephants distribution in the past extended from the Tigris-Euphrates river systems in west Asia, eastward through Persia into the Indian sub-continent, South and Southeast Asia and north into China. Today, it is found in only a small fraction of its past distribution and with continuous fragmentation, will further isolate the existing populations. Currently, the Asian elephants are found in 13 countries: Nepal, Bangladesh, Bhutan, Thailand, India, Sri Lanka, China, Myanmar, Cambodia, Laos, Vietnam, Indonesia and Malaysia. Populations are scattered, isolated and many of them have low probability of viability (Santosh, 1999).



Many subspecies of the Asian elephants have been described in the past but few have been accepted as valid by elephant biologists (Deraniyagala, 1955; McKay, 1973; Shoshani and Eisenberg, 1982 and Shoshani, 1991). There are three subspecies of the *Elephas maximus*, *E. maximus sumatranus*, that is found in Malaysia and Sumatra; *E. indicus*, which inhabits the jungle of Indochina and China and *E. maximus*, which roams the vegetation of Sri Lanka and Southern India (Shoshani 1991; Cavendish 1993). It is quite difficult to differentiate them from each other because the three subspecies vary only slightly in body size, skin color, the size as well as shape of their ears. The physical characteristics among these three forms of the Asian elephants change gradually from one extreme to another, beginning at Sri Lanka in the western part of their distribution and ending with Sumatra at the eastern limit. *E. maximus* are the largest and the darkest of the three elephants having largest ears and patches of depigmentation on the ears, face, trunk and belly. *E. sumatranus* is the smallest and has the lightest skin color with least depigmentation, while the *E. indicus* has the combination of features from the other two subspecies. They are also distinguished by physical traits related to their geographic location. For example, *E. maximus* elephants tend to have larger ears which are useful for regulating body temperature in the hotter climate of Sri Lanka (Cavendish, 1993).

The Asian elephant, *Elephas maximus* is classified as an endangered species by the International Union for the Conservation of Nature (IUCN) Species Survival Commission' s Asian Elephant Specialist Group in the Red Data Book (Baillie and Groombridge, 1996). To date, there are only between 38000 and 51000 wild Asian elephants in comparison to their distant cousin, the African elephants with over 600,000



heads. Fragmentation and loss of existing habitat to agriculture and developmental activities pose as main threats to the survival of this population. For centuries, the elephant's massive tusks have been prized for their ivory (Eltringham, 1991). Poaching also causes long-term damage to the demography as only male Asian elephants have tusks and selective poaching of one sex has resulted in highly skewed sex ratios (Santosh, 1999).

Facing the problem of extinction, the survival of Asian elephants in the future depends on their ability to survive and reproduce. Their genetic background is the most important information that could be used in breeding programs and to identify specific genetic markers as to verify the animals in forensic cases as well as enforcement of law regarding ruthless killing of this endangered species (Deacon and Lah, 1989). In the long term, the conservation program of Asian elephants should be a major importance if the species is to survive in this millennium.

Documentation on Asian elephant's karyotype is very scarce. Available literature on the cytogenetics of Asian elephants described the conventional karyotype only. In our literature search, there have been very few reports on the cytogenetic and molecular aspects of the Asian elephants. Moreover, the molecular studies concentrated only on the demonstration of phylogenetic position among African elephants, Asian elephants and Mammoths.

