



**UNIVERSITI PUTRA MALAYSIA**

***GENETIC VARIATION OF SOUTHEAST ASIAN CROCODILE,  
TOMISTOMA SCHLEGELII MULLER, USING MICROSATELLITE AND  
INTER SIMPLE SEQUENCE REPEAT MARKERS***

**BEHNAM SHAFIEI ASTANI**

**FBSB 2015 22**



**GENETIC VARIATION OF SOUTHEAST ASIAN CROCODILE,  
*Tomistoma schlegelii* MULLER, USING MICROSATELLITE AND INTER  
SIMPLE SEQUENCE REPEAT MARKERS**

**By**

**BEHNAM SHAFIEI ASTANI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfillment of the requirement for the Degree of Master of Science**

**January 2015**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



## DEDICATION

This thesis is dedicated to  
My Beloved Parents,  
My wife  
And  
My True Friends



COPYRIGHT

UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Master of Science

**GENETIC VARIATION OF SOUTHEAST ASIAN CROCODILE,  
*Tomistoma schlegelii* MULLER, USING MICROSATELLITE AND INTER  
SIMPLE SEQUENCE REPEAT MARKERS**

By

**BEHNAM SHAFIEI ASTANI**

**January 2015**

**Chairman: Professor Soon Guan Tan, PhD**

**Faculty: Biotechnology and Biomolecular Sciences**

*Tomistoma schlegelii*, also referred to as the “false gharial”, is one of the most exclusive and least known of the world’s fresh water crocodylians, limited to Indonesia and Malaysia. The species has been recognized for over a century from a museum specimen, and its distribution has been a mystery for a long time. Its lack of economic value and local cultural significance (the skin is not traded locally) has also contributed to the lack of studies on the false gharial.

*Tomistoma schlegelii* within the Southeast Asia region has been found to suffer from population reduction. This species has to live in small environments, because of the current habitat destruction, hence leading to a steady reduction of species number and, this can be as a reason for an increase in the rate of inbreeding. Nevertheless, lack of information on the genetic variability and structure of the species in Malaysia remains a matter of concern. Hence, the objectives of this study were to determine the current genetic diversity and structure of this endangered species in Malaysia. To achieve this, two types of molecular markers, namely microsatellites (40 primer pairs) and Inter Simple Sequence Repeats (ISSRs, 45 primers) were used.

The Department of Wildlife and National Parks (DWNP), Malaysia provided the samples. In this regard, 17 *T. schlegelii* specimens were collected by the department from different parts of Peninsular Malaysia and Sarawak, and were then transferred to the farms and zoos across the country. The mentioned crocodiles were later subjected to the blood sampling process. Consequently, DNA was isolated from the blood samples of *T. schlegelii* by using the conventional phenol-chloroform method. The PCR amplification of *T. schlegelii* DNA was optimized to obtain the most effective annealing temperature and  $Mg^{2+}$  concentration for each individual marker.

Out of the 40 microsatellite primer pairs tested 10 were detected as amplifying polymorphic bands, while 36 ISSR primers were detected as producing polymorphic banding patterns from the 45 tested. The observed heterozygosity ( $H_O$ ) and the expected heterozygosity values ( $H_E$ ) of the polymorphic microsatellite loci ranged from 0.588 to 1.000 and 0.470 to 0.891, respectively. The highest average

heterozygosity was observed for locus Cj109 with a value of 0.945 and the lowest average heterozygosity was observed for locus Cs15 with value of 0.529. Among the 45 ISSR markers, 36 showed polymorphic banding patterns (80%). The highest number (6) of polymorphic bands was produced by primer UBC811 and the lowest (1 each) by primers PT2, UBC820, UBC834 and UBC868.

Interestingly, data from both the microsatellite and ISSR marker systems generated two main clusters. The highest repeat motif similarities to the original species from which the loci were developed for three microsatellite loci, namely Cj16, Ami $\mu$ 16 and Ami $\mu$ 15, which originated from *C. johnstoni*, *A. mississippiensis* and *C. siamensis* respectively. The repeat motifs of the Cj16 locus in *T. schlegelii* were found to be a high match with the repeat motif of the same locus in *C. johnstoni*, and the difference was only limited to one repeat unit of "CA". The observed similarity of the repeat motif type of Ami $\mu$ 16 and Ami $\mu$ 15 was "AC" and the sizes of the amplified loci in the present study were shorter than the ones reported for the original species.

Both microsatellite and ISSR marker systems have high potential for studying the genetic variation of *T. schlegelii*, and these markers are suitable for conservation genetic programme of this endangered species. Overall, assessment of the results showed that the Malaysian *T. schlegelii* individuals studied in this research could have originated from a core population of the crocodiles. The lack of opportunity for outbreeding with other populations of crocodiles led to the occurrence of low genetic variation of the species in Malaysia.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**VARIASI GENETIK BUAYA DI ASIA TENGGARA, *Tomistoma schlegelii*  
MULLER, MENGGUNAKAN PENANDA MIKROSATELIT DAN PENANDA  
INTER SIMPLE SEQUENCE REPEAT**

Oleh

**BEHNAM SHAFIEI ASTANI**

Januari 2015

**Pengerusi: Profesor Tan Soon Guan, PhD**  
**Fakulti: Bioteknologi dan Sains Biomolekul**

*Tomistoma schlegelii*, juga dikenali sebagai “false gharial”, merupakan salah satu spesies buaya yang paling kurang diketahui dan hanya boleh dijumpai di Indonesia dan Malaysia. Kewujudan spesies ini telah lama didokumentasi berdasarkan spesimen yang disimpan di muzium. Walaupun begitu, spesies ini kurang mendapat perhatian dalam kajian kerana ia tidak memberi impak ekonomi yang besar dan kurang kepentingan untuk budaya tempatan.

*Tomistoma schlegelii* di rantau Asia Tenggara didapati telah mengalami pengurangan populasi. Disebabkan pemusnahan habitat, spesies ini terpaksa hidup di dalam kawasan yang kecil dan ini akan mengurangkan lagi bilangan spesies ini, justeru meningkatkan kadar ‘inbreeding’ dalam spesies ini. Kekurangan maklumat mengenai variasi dan struktur genetik spesies ini di Malaysia adalah sesuatu yang membimbangkan. Oleh itu, objektif kajian ini adalah untuk menentukan variasi serta struktur genetik spesies ini di Malaysia. Untuk mencapai matlamat ini, dua jenis penanda molekul, iaitu mikrosatelit (40 pasang ‘primer’) dan ‘Inter Simple Sequence Repeats’ (ISSRs, 45 ‘primers’) telah digunakan.

Jabatan Perlindungan Hidupan Liar dan Taman Negara (PERHILITAN) telah membekalkan sampel untuk kajian ini. Sebanyak 17 spesimen buaya *T. schlegelii* telah dikumpulkan oleh PERHILITAN dari peninsula Malaysia dan Sarawak untuk diagihkan ke ladang-ladang penternakan dan zoo-zoo tempatan. Sampel darah kemudiannya diambil daripada spesimen-spesimen buaya ini. DNA telah dikeluarkan dari sampel darah spesies ini dengan menggunakan teknik ‘phenol-chloroform’. Amplifikasi PCR dengan DNA *T. schlegelii* telah dijalankan untuk mendapatkan suhu ‘annealing’ dan konsentrasi  $Mg^{2+}$  yang paling optimum.

Daripada 40 pasang ‘primer’ mikrosatelit, 10 didapati mengamplifikasikan band yang polimorfik, manakala 36 dari 45 ‘primer’ ISSR didapati mengamplifikasikan band yang polimorfik. Nilai-nilai ‘observed heterozygosity ( $H_o$ )’ dan ‘expected heterozygosity ( $H_E$ )’ penanda-penanda mikrosatelit masing-masing telah didapati berada dalam jangka 0.588 ke 1.000 dan 0.470 ke 0.891. Nilai ‘average heterozygosity’ yang paling tinggi telah diperolehi dengan penanda Cj109 dengan nilai

0.945 manakala nilai yang paling rendah telah diperoleh dengan penanda Cs15 dengan nilai 0.529. Antara 45 'primer' ISSR, 36 'primer' (80%) telah menunjukkan band yang polimorfik. Bilangan band yang paling banyak telah diperoleh dengan menggunakan 'primer' UBC811 (6 band) dan bilangan band yang paling rendah telah diperoleh dengan menggunakan 'primer' PT2, UBC820, UBC834 and UBC868 (masing-masing 1 band). Data daripada penanda mikrosatelit dan ISSR telah menunjukkan pembahagian kepada dua kumpulan. 'Repeat motif' yang paling bersamaan dengan spesies lain telah diperoleh dengan menggunakan penanda-penanda Cj16, Ami $\mu$ 16 dan Ami $\mu$ 15 daripada spesies buaya *C. johnstoni*, *A. mississippiensis* dan *C. siamensis*. 'Repeat motif' untuk Cj16 adalah "CA" manakala 'repeat motif' untuk Ami $\mu$ 16 and Ami $\mu$ 15 adalah "AC".

Kedua-dua penanda mikrosatelit dan ISSR mempunyai potensi yang tinggi bagi kajian variasi genetik *T. schlegelii*, dan amat sesuai untuk digunakan oleh program perlindungan genetik bagi spesies ini. Hasil kajian ini telah mendapati bahawa *T. schlegelii* di Malaysia berkemungkinan berasal daripada 1 populasi spesies buaya ini, dan telah mengakibatkan kekurangan variasi genetik dalam kalangan spesies ini di Malaysia.



## ACKNOWLEDGEMENTS

First and foremost, I would like to thank almighty ALLAH for all He has done in my life, and I praise him for everything.

I especially want to extend my truly kind and devoted supervisor Professor Dr. Soon-Guan Tan. I would never have conducted this research without his continuing guidance and effective suggestions. His ever-ethical devotion to science became the inspiration to my research style.

I wish to express my heartfelt thanks and deepest gratitude to my dear co-supervisor Associate Professor Dr Alan Ong Han Kiat who provided this project for me and giving valuable time to discuss and for suggesting valuable information during the study. My grateful appreciation also goes to my co-supervisor Dr. Christina Yong who made it possible for me to complete this work at the Genetic Lab and for providing research facilities. I would like to thank Associate Professor Dr Noorjahan Banu Mohamed Alitheen who was always supportive and never withheld providing whatever I needed.

This study would not have been possible without the participation of the Department of Wildlife and National Parks, Peninsular Malaysia (Perhilitan), and Forest Department of Sarawak and Sarawak Forestry Corporation (SFC). I would also like to thank Universiti Tunku Abdul Rahman (UTAR), for supporting this study financially.

I am certainly grateful and indebted to my amazing and dearest friends Dr. Alireza Valdiani and Dr Wei Lun Ng who constantly encouraged and helps me during my study. I wish to express my sincere appreciation to Taranjeet Kaur for her experienced guidance during trouble shooting process.

My profound and warmest thanks and love to my mother for her love, patience, and encouragement and to my father for his constant support, love and prayers throughout my study. I am forever indebted to my parents. My heartfelt thanks are extended to my wife, for her care, endless emotional support, love and assistance throughout my course of study. I wish to express my sincere appreciation to my close friends Dr Homayoun Hani, Dr Hamidreza Panjehfouladgaran and Sajad Jamalpour for their support and kind wishes. Last but certainly not least, I wish to express my sincere appreciation to all those who not mentioned here that helped me to ensure the completion of my research.

I certify that a Thesis Examination Committee has met on 16/01/2015 to conduct the final examination of Behnam Shafiei Astani on his thesis entitled “Genetic Variation of Southeast Asian Crocodile, *Tomistoma Schlegelii* Muller, Using Microsatellite and Inter Simple Sequence Repeat Markers” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the degree of Master of Science.

Members of the Thesis Examination Committee were as follows:

**Ho Chai Ling, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Jothi Malar, PhD**

Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

**Patimah Binti Ismail, PhD**

Professor

Faculty of Medicine and Health Sciences

Universiti Putra Malaysia

(Internal Examiner)

**Badrul Munir Md Zain, PhD**

Professor

Faculty Science and Technology

Universiti Kebangsaan Malaysia

(External Examiner)

---

Prof. Dr. Zulkarnain Zainal

Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Tan Soon Guan, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Noorjahan Banu Mohamed Alitheen, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

**Christina Yong Seok Yien, PhD**

Senior Lecturer

Faculty of Science

Universiti Putra Malaysia

(Member)

**Alan Ong Han Kiat**

Associate Professor

Faculty of Medicine and Health Sciences

Universiti Tunku Abdul Rahman

(External)

---

**BUJANG BIN KIM HUAT, PhD**

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- This thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of the thesis are fully-owned by Universiti Putra Malaysia, according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from the supervisor and the office of the Deputy Vice-Chancellor (Research and Innovation) before the thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: \_\_\_\_\_


Date: \_\_\_\_\_


Name and Matric No.: Behnam Shafiei Aastani GS31724


## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature:   
Name of  
Chairman of  
Supervisory  
Committee: **PROF. DR. TAN SOON GUAN**  
Pensyarah  
Jabatan Biologi Sel dan Molekul  
Fakulti Bioteknologi dan Sains Biomolekul  
Universiti Putra Malaysia  
43400 UPM Serdang

Signature:   
Name of  
Member of  
Supervisory  
Committee: **Dr. CHRISTINA YONG**  
Jabatan Biologi  
Fakulti Sains  
Universiti Putra Malaysia  
43400 UPM, Serdang  
Selangor Darul Ehsan

Signature:   
Name of  
Member of  
Supervisory  
Committee: **PROF. MADYA DR. NOORJAHAN BANU MOHD ALITHEEN**  
Pensyarah  
Jabatan Biologi Sel dan Molekul  
Fakulti Bioteknologi dan Sains Biomolekul  
Universiti Putra Malaysia  
43400 UPM SERDANG

Signature: \_\_\_\_\_  
Name of  
Member of  
Supervisory  
Committee: \_\_\_\_\_

## TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	
1.1 Study Background	1
1.2 Problem Statement	2
1.3 General Objectives	2
1.4 Specific Objectives	2
<b>2 LITERATURE REVIEW</b>	
2.1 Taxonomy of <i>Tomistoma schlegelii</i>	3
2.2 Morphology	3
2.3 Diet	5
2.4 Distribution	5
2.4.1 Distribution history	5
2.4.2 Present Distribution	6
2.5 Conservation and Threats	7
2.6 Genetic Markers	8
2.6.1 Microsatellite Markers	8
2.6.2 ISSR Markers	8
2.7 Microsatellite Markers as Useful Tools in Genetic Studies of Crocodile Population	9
2.8 Development Techniques and Complications	10
2.9 The Importance of Crocodylian Protection	10
2.10 The Advantage of ISSR Markers for Population Studies	11
<b>3 MATERIALS AND METHODS</b>	
3.1 Population and Origin	13
3.1.1 Sampling Sites	13
3.2 Blood Sample Collection	13
3.2.1 Manual Restraint	13
3.2.2 Blood Samples Preparation	14
3.3 DNA Extraction	15
3.3.1 Measure of DNA Quantity and Quality	15
3.4 Polymerase Chain Reaction (PCR)	15
3.4.1 Optimisation for Microsatellites Markers	15
3.4.2 PCR Optimisation of ISSR Markers	16
3.5 Polymorphism Screening	17
3.6 DNA Sequencing	17

3.7	Data Analysis	17
3.7.1	Genetic Diversity	17
3.7.2	Data Analysis of Microsatellite Markers	18
3.7.3	Data Analysis of ISSR markers	18
3.8	Phylogenetic Tree	18
3.8.1	Methods for the Formation of Phylogenetic Tree	18
3.8.2	Distance Methods	19
3.9	Bayesian Genotypic Clustering Techniques	19
<b>4</b>	<b>RESULTS</b>	
4.1	DNA Extraction and Quantification	20
4.2	Result of Microsatellite Markers	20
4.2.1	Characteristics of Microsatellite Loci	20
4.2.2	Cross-species Amplification of <i>T. schlegelii</i>	24
4.2.3	Genetic Diversity within <i>T. schlegelii</i> Samples	24
4.2.4	Estimation of Allele Size and Allele Frequency	27
4.2.5	Genetic Diversity Components	30
4.2.6	Genetic Structure Revealed by Microsatellite	32
4.2.7	Nei's Genetic Distance and Cluster Analysis	34
4.3	Result of ISSR Markers	37
4.3.1	ISSR Banding and Genetic Variation	37
4.3.2	ISSR Diversity	39
4.3.3	Genetic Distance and Cluster Analysis	42
4.3.4	Genetic Structure Revealed by ISSR Analysis	45
<b>5</b>	<b>DISCUSSION</b>	
5.1	Genetic Relationship among <i>T. schlegelii</i> Individuals	47
5.2	Assessment of Microsatellite and ISSR Marker Genotype	48
5.3	Technical Aspects and Objectives of Experiment	48
5.4	Conservation of Flanking Sequences	49
5.5	Genetic Relationship and Genetic Structure of <i>T. schlegelii</i>	49
<b>6</b>	<b>CONCLUSION</b>	<b>50</b>
	<b>REFERENCES</b>	<b>51</b>
	<b>APPENDICES</b>	<b>59</b>
	<b>BIODATA OF STUDENT</b>	<b>74</b>
	<b>LIST OF PUBLICATIONS</b>	<b>75</b>

## LIST OF TABLES

Table	Page
3.1 Sampling location and sample types of <i>T. schlegelii</i>	13
4.1 Characterization of 10 polymorphic microsatellite loci in <i>T. schlegelii</i>	22
4.2 Characterization of 1 monomorphic microsatellite loci in <i>T. schlegelii</i>	23
4.3 Comparisons of cross-species amplification of <i>T. schlegelii</i> with other species	24
4.4 Allele size of 10 polymorphic loci across <i>T. schlegelii</i>	27
4.5 Overall Allele Frequency 10 polymorphic loci across <i>T. schlegelii</i>	27
4.6 Number of alleles of 10 polymorphic microsatellite loci in <i>T. schlegelii</i>	30
4.7 Estimation of heterozygosities of 10 polymorphic microsatellite markers in <i>T. schlegelii</i>	31
4.8 Overall estimation of genetic diversity contents of 10 microsatellite markers on <i>T. schlegelii</i>	32
4.9 Nei's (1983) measures of genetic distance of 10 polymorphic microsatellite markers among 17 <i>T. schlegelii</i>	35
4.10 Characteristics of the 36 ISSR primers tested on <i>T. schlegelii</i> samples	38
4.11 The number of ISSR bands generated by 36 primers in 17 individuals of <i>T. schlegelii</i>	41
4.12 Simple Matching similarity measures of 36 ISSR markers among 17 <i>T. schlegelii</i>	43



## LIST OF FIGURES

Figure		Page
2.1	Figure 2.1: A slender snouted adult male <i>T. schlegelii</i> in Zoo Negara, Malaysia	4
2.2	One month old hatchlings of <i>T. schlegelii</i> in Jong's Crocodile Farm, Sarawak (Kaur, 2010)	4
2.3	The sharp teeth of an adult male <i>T. schlegelii</i> in Zoo Negara, Malaysia	5
2.4	The adapted map of current distribution of <i>T. schlegelii</i>	6
3.1	Location of the blood sampling sites	14
4.1	Electrophoretic Characterization Of Genomic DNA Isolated From Blood	20
4.2	SSR Polymorphic Banding Pattern Generated by Locus Cj35 for 17 Individuals of <i>T. schlegelii</i>	25
4.3	SSR Monomorphic Banding Pattern Generated by Locus Amip203 for 17 Individuals of <i>T. schlegelii</i>	26
4.4	Frequency of Allelic Size of 10 Polymorphic SSR Markers Tested for <i>T. schlegelii</i> individuals	28
4.5	Genetic Structure Revealed by Microsatellite Analysis	33
4.6	Consensus of UPGMA Dendrogram of 10 Polymorphic Microsatellite Markers Based on Nei's (1983) Genetic Distance Method among <i>T. Schlegelii</i>	36
4.7	ISSR Banding Pattern of <i>T. schlegelii</i>	40
4.8	Consensus UPGMA dendrogram of genetic relationship based on simple matching coefficient method among <i>T. schlegelii</i>	44
4.9	Genetic Structure Revealed by ISSR Analysis	46

## LIST OF ABBREVIATIONS

1X	One time
A	Adenine
AFLP	Amplified Fragment Length Polymerase
APS	Ammonium Persulfate
bp	Base pairs
C	Cytosine
DNA	Deoxyribonucleic acid
dNTP	Dinucleotide triphosphate
EDTA	Ethylene diamine tera acetic Acid
G	Guanine
H <sub>E</sub>	Expected Heterozygosity
H <sub>O</sub>	Observed Heterozygosity
ISSR	Inter Simple Sequence Repeat
MCMC	Markov Chain Monte Carlo
MgCl <sub>2</sub>	Magnesium chloride
mtDNA	Mitochondrial DNA
Na	Observed number of alleles
NaOH	Sodium Hydroxide
Ne	Effective number of alleles
ng	Nanogram
NJ	Neighbor Joining
PAGE	Polyacrylamide Gel
PCR	Polymerase chain reaction
PIC	Polymorphism Information Content
RAPD	Random amplification of polymorphic DNA
SAHN	Sequential, Agglomerative, Hierarchical and Nested clustering
SSR	Simple sequence repeat
T	Thymine
T <sub>a</sub>	Annealing temperature
TBE	Tris-borate EDTA
UPGMA	Unweighted pair group method with arithmetic mean
UV	Ultraviolet
VNTR	Variable Number Tandem Repeat

# CHAPTER 1

## INTRODUCTION

### 1.1 Study Background

*Tomistoma schlegelii* known as Buaya jenjulong in Malaysia is a freshwater crocodilian species that can only be found in Southeast Asia. *T. schlegelii* or false gharial is one of the most unusual and little known crocodilian species (Bezuijen *et al.*, 2010; Kaur *et al.*, 2013). Presently, there are 23 extant crocodilian species in eight genera of three families coming from Crocodyliforms that had existed on earth 200 million years ago. Given their old ancestral lineage with little morphological changes (Salisbury *et al.*, 2006) and important functions in the ecosystem balance, more attention should be given to this crocodilian species. This species is regarded as a keystone species that ensures the balance and function of the habitat through its activities (Rodriguez, 2007).

Presently, the majority of individuals of the endangered species are found in farms and zoos, and they are obtained illegally. Therefore, with 2,500 estimated remaining individuals, the false gharial native to Southeast Asia illustrates a model of an endangered flagship species resulting from large loss of habitat (Rödder *et al.*, 2010). Almost 77 individuals of the species can be found at the farms and zoos in Malaysia (Stuebing *et al.*, 2004), while 88 other individuals can be found in Singapore, Indonesia, and Thailand. Farms and zoos in the United States of America and the Europe have around 57 individuals (Melino, 2011). *T. schlegelii* is a species found in Peninsular Malaysia, West Borneo, Java, and Sumatra. There is still an argument whether this species belongs to the Gavialidae or the Crocodylidae family (Mathew *et al.*, 2011). This species is an endangered species due to habitat loss. *T. schlegelii* inhabits peat swamps, and riverine forests habitat loss and illegal hunting are believed to contribute to the decline in the species number (Rödder *et al.*, 2010).

Aside from rapid status assessments, little is known about the species ecosystem. However, these rapid assessments have found that, overall, the *T. schlegelii* population in Southeast Asia has declined or fragmented due to habitat degradation, while Tanjung Puting National Park in Central Kalimantan still holds a stable population (Auliya *et al.*, 2009). Southeast Asia's tropical forests are rich with species variety probably because they act as a protection through past climate variations. However, these forests are subjected to modifications by human activities that might affect other living organisms (Gatesy *et al.*, 2003). Several survey studies have been done in Sumatra (Bezuijen *et al.*, 2001), East and Central Kalimantan (Fraser and Bernatchez, 2001), Peninsular Malaysia (Simpson *et al.*, 1998), and Sarawak (Stuebing *et al.*, 2004). The results obtained from these surveys show the distribution, breeding biology, and other aspects of *Tomistoma* environmental science (Bezuijen *et al.*, 2010).

As the genetic variation of this endangered species at the population level is not well studied, this makes conservation efforts difficult to be implemented due to lack of information and sources (Kaur *et al.*, 2013). However, conservation effort still needs to be done to avoid total species extinction. In the early stage, conservation efforts for

*Tomistoma* are carried out by focusing on the field surveys to record the existing populations and to identify conservation priorities (Bezuijen *et al.*, 2010).

To study the genetic variation, microsatellite and ISSR can be useful molecular markers. In conservation biology, microsatellite and ISSR markers can be utilised to identify unexpected changes in the inhabitants and influence of population fragmentation. Besides that, these markers are helpful in recognition of new and original inhabitants. Microsatellites and ISSR are powerful genetic markers that provide a fine resolution for discriminating populations (Tanya *et al.*, 2011).

These markers, which have become more important in recent times, provide precise evaluation of genetic structures of many organisms. Microsatellite markers are polymorphic and are distributed randomly in the genome to help differentiate populations (Avise, 2000).

## 1.2 Problem Statement

*Tomistoma schlegelii* within the Southeast Asia region has been found to suffer from population reduction. Human activities such as illegal fishing, urbanisation, as well as the environmental changes, and consequently habitat destruction as the main threatening factors (Auliya *et al.*, 2009) have led to decrease in the number and density of the endangered freshwater crocodile species "*Tomistoma schlegelii*" (Kaur *et al.*, 2013). Therefore, conservation efforts need to be initiated. In line with this, further extensive surveys to accurately identify the wild population size in Malaysia are required. In another word, conservation plans on habitat protection, population monitoring measurements, and land management should be determined. To achieve these, a clear overview on the genetic variation of the species across the country is unavoidable, while the present knowledge on the genetic diversity of *T. schlegelii* population in Malaysia is not sufficient (Harshman *et al.*, 2003; Janke *et al.*, 2005; Roos *et al.*, 2007). It is hypothesized that molecular markers such as microsatellites and ISSRs will generate useful information on the genetic structure and variation of this species. Such information can be employed for developing any conservation program in the future.

## 1.3 General Objectives

The main objective of this study was to determine the molecular genetic diversity of *T. schlegelii* samples obtained from Peninsular Malaysia and Sarawak. Currently, few microsatellite and ISSR marker studies have been conducted for *T. schlegelii*. The species was only included in these studies as an additional crocodylian species to test cross taxa amplificability of the developed markers.

## 1.4 Specific Objectives

The specific objectives of this study are as follows:

1. To determine the genetic diversity of *T. schlegelii* by using microsatellite and ISSR markers.
2. To estimate the allelic numbers and frequencies of the cross species amplified microsatellite markers.
3. To infer the genetic relationships of the *T. schlegelii* samples.

## REFERENCES

- Adams, S. E., Smith, M. H., and Baccus, R. (1980). Biochemical variation in the American alligator. *Herpetologica*, 36(4), 289-296.
- Auliya, M., Schwedick, B., Sommerlad, R., and Brend, S. Samedi (2009). A short-term assessment of the conservation status of *Tomistoma schlegelii* (Crocodylia: Crocodylidae) in Tanjung Puting National Park (Central Kalimantan, Indonesia). Cooperative survey by the Orangutan Foundation (UK) and the Tomistoma Task Force of the IUCN/SSC Crocodile Specialist Group
- Avise, J. C. (1994). Molecular markers: natural history and evolution. New York, USA: Springer.
- Avise, J. C. (2000). Phylogeography: the history and formation of species. USA: Harvard University Press.
- Baker, A. N., Smith, A. N., and Pichler, F. B. (2002). Geographical variation in Hector's dolphin: Recognition of new subspecies of *Cephalorhynchus hectori*. *Journal of the Royal Society of New Zealand*, 32(4), 713-727.
- Bezuijen, M., Hasudungan, F., Kadarisman, R., Wardoyo, S., and Webb, G. (2002). Samedi and Manolis, False Gharial (*Tomistoma schlegelii*) surveys in southeast Sumatra, Indonesia (1995-2002). *Unpublished report, Wildlife Management International Pty. Limited, Darwin.*
- Bezuijen, M. R. (1997). *Project Tomistoma: Second Report on the Ecology of the False Gharial (Tomistoma Schlegelii) in Sumatera: Wildlife Management International Pty.*
- Bezuijen, M. R., Shwedick, B. M., Sommerlad, R., Stevenson, C., and Steubing, R. B. (2010). *Tomistoma Tomistoma schlegelii. Crocodile: Status Survey and Conservation Action Plan. Darwin: Crocodile Specialist Group, 133-138.*
- Bezuijen, M. R., Suryansyah, Huda, I., Andriyono, S., Pratihno, P., Potess, F. and, and Sommerlad, R. (2004). False gharial (*Tomistoma schlegelii*) surveys in West Kalimantan, Indonesia in 2004. URL: <http://tomistoma.org> Accessed on the 29th March 2009.
- Bezuijen, M. R., Webb, G. J., and Hartoyo, P. (2001). Peat swamp forest and the false gharial *Tomistoma schlegelii* (Crocodylia, Reptilia) in the Merang River, eastern Sumatra, Indonesia. *Oryx*, 35(4), 301-307.
- Bhagyawant, S., and Srivastava, N. (2008). Genetic fingerprinting of chickpea (*Cicer arietinum* L.) germplasm using ISSR markers and their relationships. *African Journal of Biotechnology*, 7(24).

- Botstein, D., White, R. L., Skolnick, M., and Davis, R. W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American journal of human genetics*, 32(3), 314.
- Brochu, C. A. (1997). Morphology, fossils, divergence timing, and the phylogenetic relationships of Gavialis. *Systematic Biology*, 46(3), 479-522.
- Brochu, C. A. (2003). Phylogenetic approaches toward crocodylian history. *Annual Review of Earth and Planetary Sciences*, 31(1), 357-397.
- Casu, M., Casu, D., Lai, T., Cossu, P., and Curini-Galletti, M. (2008). A molecular tool for genetic surveys in the red coral (*Corallium rubrum*): an Inter-Simple Sequence Repeats (ISSRs) perspective. *Biochemical Systematics and Ecology*, 36(2), 77-83.
- Chaeychomsri, W., Tabthipwon, P., Noparatnaraporn, N., and Siripholvat, V. (2008). Development of microsatellite markers for siamese crocodile (*Crocodylus siamensis*). *Kasetsart Journal (Natural Sciences)(Thailand)*, 42, 256-262.
- Chennaoui-Kourda, H., Marghali, S., Marrakchi, M., and Trifi-Farah, N. (2007). Genetic diversity of *Sulla* genus (*Hedysarea*) and related species using Inter-simple Sequence Repeat (ISSR) markers. *Biochemical systematics and ecology*, 35(10), 682-688.
- Chong, L., Tan, S., Yusoff, K., and Siraj, S. (2000). Identification and characterization of Malaysian river catfish, *Mystus nemurus* (C&V): RAPD and AFLP analysis. *Biochemical genetics*, 38(3-4), 63-76.
- Cooper, G. M., Stone, E. A., Asimenos, G., Green, E. D., Batzoglou, S., and Sidow, A. (2005). Distribution and intensity of constraint in mammalian genomic sequence. *Genome Research*, 15(7), 901-913.
- Davis, L., Glenn, T., Elsey, R., Brisbin Jr, I., Rhodes, W., Dessauer, H., Sawyer, R., Grigg, G., Seebacher, F., and Franklin, C. (2001). Genetic structure of six populations of American alligators: A microsatellite analysis. *Crocodylian biology and evolution*, 38-50.
- Davis, L. M., Glenn, T. C., Strickland, D. C., Guillette, L. J., Elsey, R. M., Rhodes, W. E., Dessauer, H. C., and Sawyer, R. H. (2002). Microsatellite DNA analyses support an east-west phylogeographic split of American alligator populations. *Journal of Experimental Zoology*, 294(4), 352-372.
- Densmore III, L. D., and Dessauer, H. C. (1984). Low levels of protein divergence detected between *Gavialis* and *Tomistoma*: Evidence for crocodylian monophyly? *Comparative Biochemistry and Physiology. Part B: Comparative Biochemistry*, 77(4), 715-720.
- Dever, J. A., and Densmore, L. D. (2001). Microsatellites in Morelet's crocodile (*Crocodylus moreletii*) and their utility in addressing crocodylian population genetics questions. *Journal of Herpetology*, 35(3), 541-544.

- Dever, J. A., Strauss, R. E., Rainwater, T. R., McMurry, S. T., Densmore III, L. D., and Wood, R. (2002). Genetic diversity, population subdivision, and gene flow in Morelet's crocodile (*Crocodylus moreletii*) from Belize, Central America. *Copeia*, 2002(4), 1078-1091.
- Esa, Y., and Abdul Rahim, K. A. (2013). Genetic Structure and Preliminary Findings of Cryptic Diversity of the Malaysian Mahseer (*Tor tambroides Valenciennes*: Cyprinidae) Inferred from Mitochondrial DNA and Microsatellite Analyses. *BioMed research international*, 2013.
- Evanno, G., Regnaut, S., and Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, 14(8), 2611-2620.
- FitzSimmons, N., Tanksley, S., Forstner, M., Louis, E., Daghli, R., Gratten, J., and Davis, S. (2001). Microsatellite markers for *Crocodylus*: new genetic tools for population genetics, mating system studies and forensics. *Crocodylian biology and evolution*, 1(1), 51-57.
- Fitzsimmons, N. N., Buchan, J. C., Lam, P. V., Polet, G., Hung, T. T., Thang, N. Q., and Gratten, J. (2002). Identification of purebred *Crocodylus siamensis* for reintroduction in Vietnam. *Journal of Experimental Zoology*, 294(4), 373-381.
- FitzSimmons, N. N., Moritz, C., and Moore, S. S. (1995). Conservation and dynamics of microsatellite loci over 300 million years of marine turtle evolution. *Molecular biology and evolution*, 12(3), 432-440.
- Frankham, R. (2005). Genetics and extinction. *Biological Conservation*, 126(2), 131-140.
- Fraser, D. J., and Bernatchez, L. (2001). Adaptive evolutionary conservation: towards a unified concept for defining conservation units. *Molecular Ecology*, 10(12), 2741-2752.
- Gatesy, J., Amato, G., Norell, M., DeSalle, R., and Hayashi, C. (2003). Combined support for wholesale taxic atavism in gavialine crocodylians. *Systematic Biology*, 52(3), 403-422.
- Glenn, T. C., Dessauer, H. C., and Braun, M. J. (1998). Characterization of microsatellite DNA loci in American alligators. *Copeia*, 1998(3), 591-601.
- Glenn, T. C., Stephan, W., Dessauer, H. C., and Braun, M. J. (1996). Allelic diversity in alligator microsatellite loci is negatively correlated with GC content of flanking sequences and evolutionary conservation of PCR amplifiability. *Molecular biology and evolution*, 13(8), 1151-1154.
- Gross, S. S., and Brent, M. R. (2006). Using multiple alignments to improve gene prediction. *Journal of Computational Biology*, 13(2), 379-393.

- Harshman, J., Huddleston, C. J., Bollback, J. P., Parsons, T. J., and Braun, M. J. (2003). True and false gharials: a nuclear gene phylogeny of Crocodylia. *Systematic Biology*, 52(3), 386-402.
- Hashizume, T., Shimamoto, I., and Hirai, M. (2003). Construction of a linkage map and QTL analysis of horticultural traits for watermelon [*Citrullus lanatus* (THUNB.) MATSUM & NAKAI] using RAPD, RFLP and ISSR markers. *Theoretical and Applied Genetics*, 106(5), 779-785.
- Isberg, S., Chen, Y., Barker, S., and Moran, C. (2004). Analysis of microsatellites and parentage testing in saltwater crocodiles. *Journal of Heredity*, 95(5), 445-449.
- Janke, A., Gullberg, A., Hughes, S., Aggarwal, R. K., and Arnason, U. (2005). Mitogenomic analyses place the gharial (*Gavialis gangeticus*) on the crocodile tree and provide pre-K/T divergence times for most crocodylians. *Journal of molecular evolution*, 61(5), 620-626.
- Joshi-Saha, A., and Gopalakrishna, T. (2007). Inheritance and tagging of gene regulating flowering time in the green manure crop *Sesbania rostrata* (Bremek. and Obrem.). *Molecular Breeding*, 20(4), 389-399.
- Kaur, T. (2010). Phylogenetic analyses and the repetitive patterns of *Tomistoma schlegelii* based on mitochondria dna gene regions. UTAR.
- Kaur, T., Japning, J. R. R., Sabki, M. S., Sidik, I., Chong, L. K., and Ong, A. H. (2013). Genetic Diversity of *Tomistoma schlegelii* Inferred from mtDNA Markers. *Biochemical genetics*, 51(3-4), 275-295.
- Kaur, T., and Ong, A. H. (2011). Heteroplasmy, length, and sequence characterization of the mitochondrial control region in *Tomistoma schlegelii*. *Biochemical genetics*, 49(9-10), 562-575.
- Korpelainen, H., Kostamo, K., and Virtanen, V. (2007). Microsatellite marker identification using genome screening and restriction-ligation. *Biotechniques*, 42(4), 479.
- Lawson, R., Kofron, C. P., and Dessauer, H. C. (1989). Allozyme variation in a natural population of the Nile crocodile. *American Zoologist*, 29(3), 863-871.
- Li, Y., Wu, X., Ji, X., Yan, P., and Amato, G. (2007). The Complete Mitochondrial Genome of Salt-water Crocodile (*Crocodylus porosus*) and Phylogeny of Crocodylians. *Journal of Genetics and genomics*, 34(2), 119-128.
- Liu, K., and Muse, S. V. (2005). PowerMarker: an integrated analysis environment for genetic marker analysis. *Bioinformatics*, 21(9), 2128-2129.
- Machkour-M'Rabet, S., Hénaut, Y., Charruau, P., Gevrey, M., Winterton, P., and Legal, L. (2009). Between introgression events and fragmentation, islands are the last refuge for the American crocodile in Caribbean Mexico. *Marine biology*, 156(6), 1321-1333.



- Martins-Lopes, P., Lima-Brito, J., Gomes, S., Meirinhos, J., Santos, L., and Guedes-Pinto, H. (2007). RAPD and ISSR molecular markers in *Olea europaea* L.: Genetic variability and molecular cultivar identification. *Genetic Resources and Crop Evolution*, 54(1), 117-128.
- Mathew, A., Ganesan, M., Majid, R. A., and Beastall, C. (2011). Breeding of false gharial, *Tomistoma schlegelii*, at Zoo Negara, Malaysia. *On-line pdf*. *Zoo Negara*. Accessed September, 25, 2012.
- McAliley, L. R., Willis, R. E., Ray, D. A., White, P. S., Brochu, C. A., and Densmore III, L. D. (2006). Are crocodiles really monophyletic? Evidence for subdivisions from sequence and morphological data. *Molecular phylogenetics and evolution*, 39(1), 16-32.
- Melino, G. (2011). p63 is a suppressor of tumorigenesis and metastasis interacting with mutant p53. *Cell Death & Differentiation*, 18(9), 1487-1499.
- Milián-García, Y., Venegas-Anaya, M., Frias-Soler, R., Crawford, A. J., Ramos-Targarona, R., Rodríguez-Soberón, R., Alonso-Tabet, M., Thorbjarnarson, J., Sanjur, O. I., and Espinosa-López, G. (2011). Evolutionary history of Cuban crocodiles *Crocodylus rhombifer* and *Crocodylus acutus* inferred from multilocus markers. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 315(6), 358-375.
- Naga Jogayya, K., Meganathan, P., Dubey, B., and Haque, I. (2013). Mitochondrial 16S ribosomal RNA gene for forensic identification of crocodile species. *Journal of forensic and legal medicine*, 20(4), 334-338.
- Nei, M. (1996). Phylogenetic analysis in molecular evolutionary genetics. *Annual review of genetics*, 30(1), 371-403.
- Nei, M., Tajima, F., and Tateno, Y. (1983). Accuracy of estimated phylogenetic trees from molecular data. *Journal of molecular evolution*, 19(2), 153-170.
- Nielsen, R., Bustamante, C., Clark, A. G., Glanowski, S., Sackton, T. B., Hubisz, M. J., Fledel-Alon, A., Tanenbaum, D. M., Civello, D., and White, T. J. (2005). A scan for positively selected genes in the genomes of humans and chimpanzees. *PLoS biology*, 3(6), e170.
- Pivorienė, O., and Pašakinskienė, I. (2008). Genetic diversity assessment in perennial ryegrass and *Festulolium* by ISSR fingerprinting. 95(2), 125-133.
- Pollard, K. S., Salama, S. R., Lambert, N., Lambot, M.-A., Coppens, S., Pedersen, J. S., Katzman, S., King, B., Onodera, C., and Siepel, A. (2006). An RNA gene expressed during cortical development evolved rapidly in humans. *Nature*, 443(7108), 167-172.
- Pritchard, J. K., Stephens, M., and Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155(2), 945-959.

- Raveendran, M., Harris, R. A., Milosavljevic, A., Johnson, Z., Shelledy, W., Cameron, J., and Rogers, J. (2006). Designing new microsatellite markers for linkage and population genetic analyses in rhesus macaques and other nonhuman primates. *Genomics*, 88(6), 706-710.
- Rödder, D., Engler, J. O., Bonke, R., Weinsheimer, F., and Pertel, W. (2010). Fading of the last giants: an assessment of habitat availability of the Sunda gharial *Tomistoma schlegelii* and coverage with protected areas. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 20(6), 678-684.
- Rodriguez, D. (2007). Crocodylian evolution, systematics and population genetics: recovery and ecological interactions of the American crocodile (*Crocodylus acutus*).
- Rohlf, F. (2000). NTSYS-PC, numerical taxonomy system for the PC ExeterSoftware, Version 2.1. *Applied Biostatistics Inc Setauket, USA*.
- Roos, J., Aggarwal, R. K., and Janke, A. (2007). Extended mitogenomic phylogenetic analyses yield new insight into crocodylian evolution and their survival of the Cretaceous–Tertiary boundary. *Molecular phylogenetics and evolution*, 45(2), 663-673.
- Ruas, P. M., Ruas, C. F., Rampim, L., Carvalho, V. P., Ruas, E. A., and Sera, T. (2003). Genetic relationship in *Coffea* species and parentage determination of interspecific hybrids using ISSR (Inter-Simple Sequence Repeat) markers. *Genetics and Molecular Biology*, 26(3), 319-327.
- Ryberg, W. A., Fitzgerald, L. A., Honeycutt, R. L., and Cathey, J. C. (2002). Genetic relationships of American alligator populations distributed across different ecological and geographic scales. *Journal of Experimental Zoology*, 294(4), 325-333.
- Salisbury, S. W., Molnar, R. E., Frey, E., and Willis, P. M. (2006). The origin of modern crocodyliforms: new evidence from the Cretaceous of Australia. *Proceedings of the Royal Society B: Biological Sciences*, 273(1600), 2439-2448.
- Schlötterer, C. (2000). Evolutionary dynamics of microsatellite DNA. *Chromosoma*, 109(6), 365-371.
- Sehgal, D., Rajpal, V. R., Raina, S. N., Sasanuma, T., and Sasakuma, T. (2009). Assaying polymorphism at DNA level for genetic diversity diagnostics of the safflower (*Carthamus tinctorius* L.) world germplasm resources. *Genetica*, 135(3), 457-470.
- Selkoe, K. A., and Toonen, R. J. (2006). Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecology letters*, 9(5), 615-629.

- Semagn, K., Bjørnstad, Å., and Ndjiondjop, M. (2006). An overview of molecular marker methods for plants. *African Journal of Biotechnology*, 5(25).
- Siepel, A. (2009). Phylogenomics of primates and their ancestral populations. *Genome Research*, 19(11), 1929-1941.
- Siepel, A., and Haussler, D. (2004). Computational identification of evolutionarily conserved exons. Paper presented at the Proceedings of the eighth annual international conference on Research in computational molecular biology.
- Silvius, M. J., and Suryadiputra, N. (2005). Review of policies and practices in tropical peat swamp forest management in Indonesia. *Wetlands International*.
- Simpson, B. (2004). False Gharials (*Tomistoma schlegelii*) in Tanjung Puting National Park, Kalimantan, Indonesia. Paper presented at the Proceedings of the 16th Working Meeting of the Crocodile Specialist Group, Darwin, NT, Australia.
- Sneath, P. H., and Sokal, R. R. (1973). Numerical taxonomy. The principles and practice of numerical classification (pp; 230-234). W.H. Freeman and Company, San Francisco, California, USA.
- Sokal, R. R., and Micheners, C. D. (1958). A statistical method for evaluating systematic relationships. *Univ Kans Sci Bull*, 38, 1409-1438.
- Strassmann, J. E., Solis, C., Peters, J. M., and Queller, D. (1996). Strategies for finding and using highly polymorphic DNA microsatellite loci for studies of genetic relatedness and pedigrees. *Molecular zoology: advances, strategies and protocols*, 163-180.
- Stuebing, R., Sah, M., Anuar, S., Lading, E., and Jong, J. (2004). The Status Of *Tomistoma schlegelii* (Mueller) In Malaysia.
- Stuebing, R. B., Bezuijen, M. R., Auliya, M., and Voris, H. K. (2006). The current and historic distribution of *Tomistoma schlegelii* the false gharial Mueller, (1838) Crocodylia, Reptilia. *The Raffles Bulletin of Zoology*, 54(1), 181-197.
- Tanya, P., Taepayoon, P., Hadkam, Y., and Srinives, P. (2011). Genetic diversity among *Jatropha* and *Jatropha*-related species based on ISSR markers. *Plant molecular biology reporter*, 29(1), 252-264.
- Tautz, D., and Renz, M. (1984). Simple sequences are ubiquitous repetitive components of eukaryotic genomes. *Nucleic acids research*, 12(10), 4127-4138.
- Venkateswarlu, M., Urs, S. R., Nath, B. S., Shashidhar, H., Maheswaran, M., Veeraiah, T., and Sabitha, M. (2006). A first genetic linkage map of mulberry (*Morus* spp.) using RAPD, ISSR, and SSR markers and pseudotestcross mapping strategy. *Tree Genetics & Genomes*, 3(1), 15-24.

- Vijayan, K., Srivatsava, P., Nair, C., Awasthi, A., Tikader, A., Sreenivasa, B., and Urs, S. R. (2006). Molecular characterization and identification of markers associated with yield traits in mulberry using ISSR markers. *Plant Breeding*, 125(3), 298-301.
- Willis, R. E., McAliley, L. R., Neeley, E. D., and Densmore III, L. D. (2007). Evidence for placing the false gharial (*Tomistoma schlegelii*) into the family Gavialidae: Inferences from nuclear gene sequences. *Molecular phylogenetics and evolution*, 43(3), 787-794.
- Yeh, F., Yang, R., and Boyle, T. (1999). POPGENE ver. 1.32. Microsoft Windows-based freeware for population genetic analysis. Quick user guide. Center Int. For. Res, Univ. of Alberta.
- Zheng, S., Cao, Z., Wang, K., Zhao, M., Mi, J., Sui, X., and Feng, Y. (2009). Effect of habitat fragmentation on the genetic diversity of *Stipa krylovii* Reshov. in an agro-pastoral ecotone in northern China. *Canadian Journal of Plant Science*, 89(5), 875-882.

## BIODATA OF STUDENT

Behnam Shafiei Astani was born on a Lion month of Dragon Year, 8<sup>th</sup> of August 1988, in Sari, a magnificent historical city located in the north of Iran and southern coast of the Caspian Sea. He grew up in an academic atmosphere with lots of love and care and he was always told to pursue his dream no matter what the tradition and the society demand. Due to the advanced, profound and yet fascinating concepts of Genetics, he has always been carrying a thirst to explore more and more about this field. He finished his diploma at Shahed High school in Neka City. In 2005, he enrolled as a bachelor student and accepted in Tehran Central Branch University. He completed his degree at Faculty of Science in Cellular and Molecular of Biology in 2009. Through suggestions of his friends, he chose Malaysia to carry on his education. After admitting in UPM and a vast area of research opportunity, he accepted to Genetic Engineering and Molecular Science and got steps closer to what he constantly would wish for. But by going through the experiences of learning techniques during his M.Sc, he realized that was just a beginning of an abysmal knowledge. The more he learned the more he felt unaware and hence, only keeping on Phd and further can bring a slight satisfaction to his desire of being an expert in Genetics and Biotechnological breakthroughs.

## LIST OF PUBLICATIONS

**Behnam Shafiei-Astani;** Ong H. A, Valdiani A, Tan S. G, Yong C. S. Y, Ahmady F, Alitheen N. B, Ng W. L, and Kuar T. (2015). Molecular genetic variation and structure of Southeast Asian crocodile (*Tomistoma schlegelii*): comparative potentials of SSRs versus ISSRs. *Submitted at Journal of GENE*.

**Behnam Shafiei-Astani,** Tan S. G., Yong S. Y. C., Ong H. and A., Kaur T., (2013). Allelic diversity and microsatellite DNA marker analysis for Southeast Asian crocodile, *Tomistoma schlegelii*. *Submitted at Journal of Animal Biotechnology*.

