



UNIVERSITI PUTRA MALAYSIA

***TOENAIL METAL CONCENTRATION AS BIOMARKER OF EXPOSURE
TO HEAVY METAL IN DRINKING WATER OF THE PASIR MAS
DISTRICT, MALAYSIA***

NURUL HAFIZA BINTI AB RAZAK

FPSK(p) 2016 36



**TOENAIL METAL CONCENTRATION AS BIOMARKER OF EXPOSURE TO HEAVY
METAL IN DRINKING WATER OF THE PASIR MAS DISTRICT, MALAYSIA**

By

NURUL HAFIZA BINTI AB RAZAK

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of
the Requirements for the Doctor of Philosophy**

August 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the Doctor of Philosophy

TOENAIL METAL CONCENTRATION AS BIOMARKER OF EXPOSURE TO HEAVY METAL IN DRINKING WATER OF THE PASIR MAS DISTRICT, MALAYSIA

By

NURUL HAFIZA BINTI AB RAZAK

August 2016

Chairman: Sarva Mangala Praveena, PhD
Faculty: Medicine and Health Sciences

Drinking water consumed by population is generally from piped water to the house, public pipe or standpipes, boreholes, protected dug wells, protected springs and rainwater collection. Population in Pasir Mas use tap water as the main source of drinking water. Even the water sources has been treated before distributed to consumers, drinking water is still contaminated by heavy metal due to less efficient treatment system or leach out from plumbing systems. The first objective of this study was to ascertain the level of knowledge, attitude and practice (KAP) regarding heavy metal contamination of Pasir Mas drinking water. The second objective was to determine the concentration of heavy metals (Al, Cr, Cu, Fe, Ni, Pb, Zn, and Cd) in drinking water and toenail from Pasir Mas. The third objective was to identify potential sources of heavy metal in drinking water from Pasir Mas. The next objective was to investigate correlation between heavy metal (Fe, Cu and Pb) concentration in drinking water and toenail. This study also identified the associations of confounders with heavy metal in toenail. Finally, this study evaluated health risks for heavy metal exposure via drinking water among respondents from Pasir Mas District. Stratified random sampling was used to select 214 respondents in Pasir Mas. A set of modified questionnaire were administered to the respondents to achieve the first objective. Then, analysis of heavy metal in drinking water were carried out using Graphite Furnace Atomic Absorption Spectrophotometry (GF-AAS) and heavy metal in toenail were analysed using Inductively Coupled Plasma Mass Spectrometry (ICP-MS). Statistical analysis involved in this study were cluster analysis and Mann-Whitney U test for the third objective. Spearman correlation analysis was carried out to investigate correlation between heavy metal (Fe, Cu and Pb) concentration in drinking water and toenail. Lastly, Multiple Linear Regression (MLR) was carried out to identify the associations of confounders with heavy metal in toenail. Pasir Mas population has good knowledge (80%), less positive attitude (93%), and good practice (81%) towards heavy metal contamination in drinking water. Heavy metal (Al, Cr, Cu, Fe, Ni, Pb, Zn, and Cd) analysed in this study was found to be under the permissible limit of drinking

water quality set by the Malaysian Ministry of Health. Concentrations of heavy metal in toenail from Pasir Mas population were $35.29 \pm 39.22 \mu\text{g/g}$ (Fe), $3.89 \pm 2.08 \mu\text{g/g}$ (Cu) and $0.30 \pm 0.34 \mu\text{g/g}$ (Pb). Cluster analysis (CA) was performed to classify the heavy metal on the basis of the similarities of their physical and chemical properties into clusters for purposes of source apportionment. Cluster analysis of heavy metal in drinking water showed three clusters which were Ni, Pb and Cr (C1), Cu, Zn and Fe (C2) and Al (C3). Thus, the cluster analysis indicated three main possible sources of heavy metal in drinking water. Cluster 1 (Ni, Pb and Cr) might originate from heavy metal leaching from galvanised iron and mild steel pipeline. Cluster 2 (Cu, Zn and Fe) might originate from corrosion of pipeline and occurrence of the heavy metal in Kelantan river water which were affected by agricultural run-off. Cluster 3 (Al) was likely to originate from usage of alum in conventional water treatment system. Mann-Whitney U test has shown that heavy metal concentration of Pb, Fe, Cu and Zn were significantly ($p < 0.05$) different at house and distribution tanks which indicated that plumbing material were the source of the heavy metal in drinking water sample from Pasir Mas District. This study also showed no significant correlation between heavy metal (Fe, Cu and Pb) in drinking water and toenail. This study found confounder associated with heavy metal in toenail is dietary intake (cockles, sandwich bread and marine fish). A total of 17.5% variation in Fe concentration in toenail was explained by Cu in toenail and dietary intakes of cockles and sandwich bread. Model for Cu showed Fe concentration in toenail and cockles intake had accounted to 13.5% variation in Cu concentration in toenail. Dietary intakes of sandwich bread and marine fishes had 14.2% of variance for Pb concentration in toenail. Health Risk Assessment conducted in this study showed no potential non-carcinogenic and carcinogenic risks from the intake of heavy metal through drinking water. In conclusion, this study provided information on KAP of the Pasir Mas population towards heavy metal in drinking water. Output of this study provided information that conventional treatment system is adequate to treat the Kelantan river water to be used as drinking water. This study also showed that pipeline material of galvanised pipes, mild steel and PVC could be the source of heavy metal in treated drinking water supplied to population in Pasir Mas. Output of this study provided evidence that toenail was a less efficient biomarker to reflect heavy metal (Fe, Cu and Pb) exposure through drinking water ingestion pathway. Lastly, this study provided information to fulfil the knowledge gap on health risk assessment (HRA) regarding heavy metal exposure through ingestion of drinking water. This information could be beneficial for decision makers and authorities to improve drinking water quality and management in the future.

Keywords: Drinking water, Heavy metal, Toenail, Health Risk Assessment, Confounder

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KEPEKATAN LOGAM DALAM KUKU KAKI SEBAGAI PENANDA BIOLOGI TERHADAP PENDEDAHAN KEPADA LOGAM BERAT DALAM AIR MINUMAN DI DAERAH PASIR MAS, MALAYSIA.

Oleh

NURUL HAFIZA BINTI AB RAZAK

Ogos 2016

Pengerusi: Sarva Mangala Praveena, PhD
Fakulti: Perubatan dan Sains Kesihatan

Air minuman yang digunakan oleh populasi umumnya adalah dari air paip rumah, paip awam, lubang digerudi, telaga tertutup, mata air yang dilindungi dan air hujan yang dikumpulkan. Populasi di Pasir Mas menggunakan air paip sebagai sumber utama air minuman. Walaupun sumber air telah dirawat sebelum diedarkan kepada pengguna, air minuman masih boleh tercemar oleh logam berat disebabkan oleh sistem rawatan yang kurang efisien atau kebocoran dari sistem paip. Objektif pertama kajian ini adalah untuk mengetahui tahap pengetahuan, sikap dan amalan (KAP) mengenai pencemaran logam berat di dalam air minuman di Pasir Mas. Objektif kedua adalah untuk menentukan kepekatan logam berat (Al, Cr, Cu, Fe, Ni, Pb, Zn dan Cd) di dalam air minuman dan kuku jari kaki dari Pasir Mas. Objektif ketiga ialah untuk mengenal pasti sumber logam berat dalam air minum dari Pasir Mas. Objektif seterusnya adalah untuk menyiasat hubungan antara kepekatan logam berat (Fe, Cu dan Pb) dalam air minuman dan kuku jari kaki. Kajian ini juga mengenal pasti hubung kait antara faktor perancu dengan logam berat dalam kuku jari kaki. Akhir sekali, kajian ini menilai risiko kesihatan oleh pendedahan logam berat melalui air minuman di kalangan responden dari Daerah Pasir Mas. Seramai 214 responden dipilih secara rawak berstrata. Satu set soal selidik yang telah diubahsuai dilengkapkan oleh responden untuk mencapai objektif pertama. Kemudian, analisis logam berat dalam air minum telah dijalankan menggunakan teknik Spektrometri Penyerapan Atom Relau Grafit (GF-AAS) dan logam berat di kuku jari kaki telah dianalisis menggunakan Spektrometer Jisim Plasma Gandingan Teraruh (ICP-MS). Analisis statistik yang terlibat dalam kajian ini ialah analisis kelompok dan ujian Mann-Whitney U untuk objektif ketiga. Analisis korelasi telah dijalankan untuk menyiasat hubungan antara logam berat (Fe, Cu dan Pb) kepekatan dalam air dan kuku jari kaki minum. Akhir sekali, Regresi Linear Berganda (MLR) telah dijalankan untuk mengenal pasti hubungan di antara faktor perancu dengan logam berat dalam kuku jari kaki. Populasi dari Pasir Mas mempunyai pengetahuan yang baik (80%), sikap kurang positif (93%), dan amalan yang baik (81%) terhadap pencemaran logam berat dalam air minuman. Logam berat (Al, Cr,

Cu, Fe, Ni, Pb, Zn dan Cd) yang dianalisis dalam kajian ini berada dalam julat piawaian kualiti air minuman yang telah ditetapkan oleh Kementerian Kesihatan Malaysia. Kepekatan logam berat dalam kuku jari kaki daripada penduduk Pasir Mas adalah $35.29 \pm 39.22 \mu\text{g/g}$ (Fe), $3.89 \pm 2.08 \mu\text{g/g}$ (Cu) dan $0.30 \pm 0.34 \mu\text{g/g}$ (Pb). Analisis Kelompok (CA) digunakan untuk mengelaskan logam berat berdasarkan persamaan ciri-ciri fizikal dan kimia ke dalam kelompok untuk tujuan pembahagian sumber. Analisis Kelompok (CA) terhadap logam berat dalam air minuman menunjukkan tiga kelompok iaitu Ni, Pb dan Cr (K1), Cu, Zn dan Fe (K2) dan Al (K3). Oleh itu, Analisis Kelompok menunjukkan tiga sumber utama logam berat dalam air minuman. Kelompok 1 (Ni, Pb dan Cr) mungkin berasal dari larut lesap logam berat daripada saluran paip disadur besi dan keluli lembut. Kelompok 2 (Cu, Zn dan Fe) mungkin berasal dari hakisan saluran paip dan kehadiran logam berat dalam air Sungai Kelantan yang terjejas akibat dari aktiviti pertanian. Kelompok 3 (Al) mungkin berasal dari penggunaan alum dalam sistem rawatan air konvensional. Kajian ini juga menunjukkan tiada hubungan yang signifikan antara logam berat (Fe, Cu dan Pb) di dalam air minuman dan kuku jari kaki. Selain itu, kajian ini mendapati faktor perancu yang berkaitan dengan logam berat di dalam kuku jari kaki adalah pengambilan pemakanan (kerang, roti sandwic dan ikan laut). Sebanyak 17.5% variasi dalam kepekatan Fe dalam kuku jari kaki adalah di sumbang oleh Cu dalam kuku jari kaki dan pengambilan pemakanan (kerang dan roti sandwic). Model Cu menunjukkan kepekatan Fe dalam kuku jari kaki dan pengambilan kerang telah menyumbang kepada 13.5% variasi dalam kepekatan Cu dalam kuku jari kaki. Pengambilan pemakanan roti sandwic dan ikan laut menyumbang kepada 14.2% variasi dalam kepekatan Pb dalam kuku jari kaki. Penilaian Risiko Kesihatan (HRA) yang dijalankan dalam kajian ini pula menunjukkan tiada potensi risiko bukan kanser dan kanser melalui pengambilan logam berat dalam air minuman. Kesimpulannya, kajian ini menyediakan maklumat tentang pengetahuan, sikap dan amalan (KAP) penduduk Pasir Mas terhadap pencemaran logam berat dalam air minuman. Output kajian ini memberi informasi berkenaan sistem rawatan konvensional mampu untuk merawat air Sungai Kelantan dengan optimum sebelum digunakan sebagai air minuman. Kajian ini juga menunjukkan bahawa bahan paip seperti besi, keluli lembut dan plastik (PVC) boleh menjadi sumber kepada pencemaran logam berat dalam air minuman dirawat yang dibekalkan kepada penduduk di Pasir Mas. Output kajian ini membuktikan bahawa kuku jari kaki adalah penunjuk biologi yang kurang efisien untuk mencerminkan pendedahan terhadap logam berat (Fe, Cu dan Pb) melalui pengambilan air minuman. Akhir sekali, kajian ini menyediakan maklumat untuk mengisi jurang pengetahuan mengenai penilaian risiko kesihatan (HRA) berkaitan dengan pendedahan terhadap logam berat melalui pengambilan air minuman. Maklumat ini boleh memberi manfaat kepada pembuat keputusan dan pihak berkuasa untuk meningkatkan kualiti air minuman dan pengurusan pada masa hadapan.

Kata kunci: Air minuman, logam berat, kuku kaki, Penilaian Risiko Kesihatan, faktor perancu

ACKNOWLEDGEMENTS

In the name of Allah, the Most Gracious and the Most Merciful.

Firstly, I would like to express my sincere gratitude to my advisor Dr. Sarva Mangala Praveena for the continuous support of my Ph.D study, for her patience, motivation, and immense knowledge. Her guidance helped me in all the time of research and writing of this thesis. I could not have imagined having a better advisor and mentor for my Ph.D study.

Besides my advisor, I would like to thank the rest of my thesis committee: Prof. Dr. Zailina Binti Hashim, Assoc. Prof. Dr. Ahmad Zaharin Bin Aris, and Assoc. Prof. Dr. Hafizan Bin Juahir, for their insightful comments and encouragement.

My sincere thanks also goes to Mrs. Norijah Binti Kassim and Mrs. Ummi Kalthum Binti Abdullah from Environmental Health Laboratory and Institute of Tropical Agriculture. Without their precious support it would not be possible to complete this research.

Also I thank my friends (Aida Soraya Shamsuddin, Nurul Jannah Khairuddin, Fadhilah Othman, Noreen Adila Omar, Nur Aqma Madiha Mazelan, Nurul Ainni Abdul Latiff, Nurul Ain Mustafa Udin, Ayuni Nabilah Alias, Nurul Syazani Yuswir, Maisarah Mutalib and Nur Maziah Hanum Osman) from Faculty of Medical and Health Sciences and Faculty of Environmental Studies for the stimulating discussions, for the sleepless nights we were working together before deadlines, and for all the fun we have had in the last four years.

I would like to thank the staffs from Water Treatment Plant at Kelar, Pasir Mas and the respondents from Pasir Mas District and Port Dickson District. Without their cooperation it would not be possible to conduct this research.

Last but not the least, I would like to thank my family (Ab Razak Mat Piah, Fatimah Yaacob, NurulHajar Ab Razak and Muhammad Hasyim Ab Razak) for supporting me spiritually throughout writing this thesis and my life in general.

I certify that a Thesis Examination Committee has met on 26 August 2016 to conduct the final examination of Nurul Hafiza binti Ab Razak on her thesis entitled "Toenail Metal Concentration as Biomarker of Exposure to Heavy Metal in Drinking Water of the Pasir Mas District, Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

Juliana binti Jalaludin, PhD

Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Chairman)

Azrina binti Azlan, PhD


Associate Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Azhar Abdul Halim, PhD

Associate Professor
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)

Claire De Burbure, PhD

Professor
Université catholique de Louvain
Belgium
(External Examiner)



NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 22 November 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Sarva Mangala Praveena, PhD

Senior Lecturer
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Zailina Binti Hashim, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Ahmad Zaharin Bin Aris, PhD

Associate Professor
Faculty of Environmental Studies
Universiti Putra Malaysia
(Member)

Hafizan Bin Juahir, PhD

Associate Professor
East Coast Environmental Research Institute
Universiti Sultan Zainal Abidin
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Nurul Hafiza binti Ab Razak, GS36103

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of
Chairman of
Supervisory
Committee: _____

Signature: _____

Name of
Member of
Supervisory
Committee: _____

Signature: _____

Name of
Member of
Supervisory
Committee: _____

Signature: _____

Name of
Member of
Supervisory
Committee: _____



TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
CHAPTER	
1	INTRODUCTION
1.1	Background of study 1
1.2	Problem statement 4
1.3	Study justification 6
1.4	Significant of study 7
1.5	Study limitation 8
1.6	Conceptual framework 10
1.7	Research objective and hypotheses 12
1.7.1	General objective 12
1.7.2	Specific objectives 12
1.7.3	Research hypotheses 12
1.8	Conceptual Definition 13
1.8.1	Heavy metal 13
1.8.2	Health Risk Assessment (HRA) 13
1.8.3	Operational Definition 14
2	LITERATURE REVIEW
2.1	Knowledge, attitude and practice (KAP) 16
2.2	Heavy metal contamination in drinking water 17
2.2.1	Heavy metal in Malaysian drinking water 19
2.3	Biomarkers used to monitor human exposure of heavy metal via drinking water 21
2.3.1	Blood 23
2.3.2	Urine 24
2.3.3	Hair 25
2.3.4	Toenail 26
2.4	Confounders 27
2.4.1	Demographic Characteristics 27
2.4.2	Food intake 28
2.4.3	Smoking Habit 28
2.5	Health Risk Assessment application in Malaysian drinking water studies 29
3	METHODOLOGY
3.1	Study Design 31
3.2	Study Location 33
3.3	Study population and sampling technique 35
3.4	Sample size 36

3.5	Sampling Frame	36
3.6	Data and sample collection	37
	3.6.1 Questionnaire	37
	3.6.2 Drinking water collection and preparation	38
	3.6.3 Toenail sampling and preparation	39
3.7	Sample Analysis	40
	3.7.1 Drinking water	40
	3.7.2 Toenail	41
3.8	Quality Control	42
	3.8.1 Apparatus	42
	3.8.2 Drinking water samples	42
	3.8.3 Toenail samples	42
	3.8.4 Standard operating procedure	43
3.9	Data Analysis	43
	3.9.1 Statistical analysis	43
	3.9.2 Health risk assessment	45
3.10	Ethical consideration	46
4	RESULTS AND DISCUSSION	
4.1	Knowledge, attitude and practice (KAP)	47
	4.1.1 Knowledge level	47
	4.1.2 Attitude level	49
	4.1.3 Practice level	51
4.2	Heavy metal distributions	53
	4.2.1 Heavy metal in drinking water	53
	4.2.2 Heavy metal distribution in toenail	55
4.3	Possible sources of heavy metal concentration in drinking water	56
4.4	Correlation between toenail and drinking water	60
4.5	Association between confounders and heavy metal in toenail	62
4.6	Health Risk Assessment	66
	4.6.1 Non-carcinogenic risk	66
	4.6.2 Carcinogenic risk	67
5	CONCLUSION AND RECOMMENDATION	68
5.1	Conclusion	68
5.2	Recommendation	69
	REFERENCES	70
	APPENDICES	93
	BIODATA OF STUDENT	122
	LIST OF PUBLICATIONS	123

LIST OF TABLES

Table	Page
1.1. Drinking water contaminants	3
2.1. Heavy metal concentration in drinking water samples	20
2.2. Studies investigated heavy metal in drinking water and biomarker.	22
3.1. Population for Pasir Mas District (Department of Statistics, 2010) and sample collected in each sub-district.	35
3.2. Proportion for drinking water monitoring (WHO, 1997)	36
3.3. Sample size calculation	36
3.4. List of main distribution tank for each sub-districts.	38
3.5. Limit of detection (LOD), Limit of Quantification (LOQ), Calibration coefficient (r^2) and recovery rates of control samples by GF-AAS (600 AAnalyst, Perkin Elmer, USA)	40
3.6. Limit of detection (LOD), Limit of Quantification (LOQ), Calibration coefficient (r^2) and recovery rates of CRM of human hair (GBW07601) by ICP-MS (ELAN DRC-e, Perkin Elmer, USA)	41
4.1. Knowledge scores for heavy metal contamination in drinking water	48
4.2. Attitude scores towards heavy metal contamination in drinking water	50
4.3. Practice scores towards heavy metal contamination in drinking water	52
4.4. Heavy metal concentration ($\mu\text{g/L}$) in drinking water from Pasir Mas (n=214)	54
4.5. Heavy metal concentration ($\mu\text{g/g}$) in toenail (n=150)	55
4.6. Mann-Whitney U test between heavy metal concentration at houses and at distribution tanks.	58
4.7. Correlation between heavy metal in toenail and drinking water (n=150)	61
4.8. Models for heavy metal using stepwise multiple linear regression on factors influencing toenail metals concentrations	63

LIST OF FIGURES

Figure	Page
1.1. Conceptual framework of this study	11
2.1. Conventional water treatment system schematic diagram	18
2.2. Biomarkers have been used in drinking water studies	21
2.3. Blood components	23
2.4. Hair Diagram	25
2.5. Toenail diagram	26
2.6. Steps in HRA process	29
2.7. Location of previous HRA studies in Peninsular Malaysia	30
3.1. Flow diagram of methodology used in this study	32
3.2. Map of Pasir Mas District	33
3.3. Location of Kelar Water Treatments Plant, distribution tanks and sampling points.	34
4.1. Dendogram for cluster analysis	57
4.2. Hazard Quotient (HQ) for non-carcinogenic health risk	66
4.3. Lifetime Cancer Risks (LCR) for carcinogenic health risk	67

LIST OF ABBREVIATIONS

<	Less than
>	More than
AKSB	Air Kelantan Sdn. Bhd.
Al	Aluminium
APHA	American Public Health Association
As	Arsenic
AT	Averaging time
BMI	Body mass index
BW	Body weight
CA	Cluster analysis
Cd	Cadmium
CDC	Centre of Disease Control and Prevention
CDI	Chronic Daily Intake
C _{dw}	Heavy metal concentration in drinking water
Cr	Chromium
Cu	Copper
Cu _{drinking water}	Copper concentration in drinking water
Cu _{toenail}	Copper concentration in toenail
DID	Department of Irrigation and Drainage
DW	Drinking water
ED	Exposure duration
EF	Exposure frequency
FAO	Food and Agricultural Organisation
Fe	Iron
Fe _{drinking water}	Iron concentration in drinking water
Fe _{toenail}	Iron concentration in toenail
GF-AAS	Graphite Furnace Atomic Absorption Spectrophotometry
H	Hair
HDPE	High density polyethylene
HQ	Hazard Quotient
HRA	Health Risk Assessment
ICP-MS	Inductively Coupled Plasma - Mass Spectrometry
IR	Drinking water ingestion rate
IRIS	Integrated Risk Information System
JECFA	The Joint FAO/WHO Expert Committee in Food Additives
KAP	Knowledge, Attitude and Practice
MLR	Multiple Linear Regression
MMOH	Malaysian Ministry of Health
Mn	Manganese
Ni	Nickel
NWSC	National Water Services Commission
Pb	Lead
Pb _{drinking water}	Lead concentration in drinking water
Pb _{toenail}	Lead concentration in toenail
PVC	Polyvinyl chloride
RfD	Oral reference dose of heavy metal
SAJH	Syarikat Air Johor Holdings
Sf	Slope Factor

SYABAS	Syarikat Bekalan Air Selangor
TN	Toenail
U	Urine
UNESCO	United Nations Educational, Scientific and Cultural Organisation
USEPA	United States Environmental Protection Agency
WHO	World Health Organisation
Zn	Zinc



© COPYRIGHT UPM

CHAPTER 1

INTRODUCTION

1.1 Background of study

Sources of drinking water supply for global population are surface water (rivers, springs or lakes in a watershed) and groundwater. Furthermore, availability of drinking water supply depends on geography, climate, engineering, regulation, and competition for resources. Approximately 11% of global population do not have access to water (UNESCO, 2014; WHO, 2013). Another 89% of global population have access to improved drinking water source (UNESCO, 2014; WHO, 2013). World Health Organisation (2013) has stated that improved drinking water sources included piped water to the house, public pipe or standpipes, boreholes, protected dug wells, protected springs and rainwater collection. According to Department of Statistics Malaysia (2010), 93% of housing units in Malaysia are equipped with treated drinking water supply. For instance, drinking water supply in Malaysia is from rivers and groundwater. Malaysia used 99% drinking water supply from rivers (Azrina et al., 2011). River water in Malaysia is readily available and treated for domestic uses (FAO 2010; Lim et al., 2012). However, rapid development and industrial activity in Malaysia have deteriorated river water quality (Fulazzaky et al., 2010). Major pollution water sources in Malaysia are manufacturing industries, livestock farming and agro-industries (Aini et al., 2007). Furthermore, most of the supplied drinking water in Malaysia is treated using conventional treatment system. Conventional treatment system using alum salt as coagulant is proven to produce aluminium residue in the treated drinking water (Muyibi et al., 2004).

Drinking water can be defined as the water supplied to the consumer that can be safely used for drinking, cooking, and washing (DeZuane, 1997). A survey by Azrina et al., (2012) in a Malaysian urban area (Klang Valley) showed that majority of the respondents consumed tap water. Although drinking water is treated before being supplied, consumers commonly complaint on drinking water quality related to air in water, high turbidity, high colour and presence of foreign particles, taste and odour (Chan 2007). Meanwhile, a study by Aini et al., (2007) showed that Malaysian consumers are dissatisfied with drinking water quality only based on the appearance without realising that drinking water could be contaminated by toxic contaminants, even when the appearance of drinking water seems to be in good quality. Besides, treated drinking water is supplied to consumers through the pipeline system. Pipeline system has been used in Malaysia which varies from mild steel, galvanised iron, stainless steel and plastic (HDPE and PVC) (NWSC, 2015). For instance, galvanised iron pipe has been replaced by HDPE, PVC and stainless steel pipe in West Coast Peninsular Malaysia and South Malaysia such as states of Selangor and Johor (SYABAS, 2013; SAJH, 2012). However, states in East Coast Peninsular Malaysia such as Kelantan has replaced galvanised pipe with mild steel and HDPE pipes (AKSB, 2013). Due to the fact that mild steel pipe is prone to corrosion which is susceptible to heavy metal (Fe, Cu, Cr, Ni and Pb) release (Jung et al., 2009; Pearce,

2013). Thus, corrosion of pipeline can be a problem to drinking water quality in Kelantan.

The United States Environmental Protection Agency (USEPA) (2016) have listed six group of drinking water contaminants including microorganisms, disinfectants, disinfection by-products, inorganic chemicals, organic chemicals and radionuclides (Table 1.1). Inorganic chemical through heavy metal contamination in drinking water is a major concern due to heavy metal exposure, which will lead to various health effect such as liver and kidney damage, neurology impairment and increased risk of cancer (USEPA, 2016; Järup, 2003). Even though the drinking water is treated before being supplied, heavy metal still can present in treated drinking water. Heavy metal contamination in drinking water may be due to contaminated water source, usage of chemical in water treatment, inefficiency of water treatment system, and leaching out from corroded pipeline system (Tamasi and Cini, 2004; Farizwana et al., 2010). Furthermore, due to the unpleasant appearance of supplied drinking water, most consumers have installed private water filter at their home (Aini et al., 2007). This practice is a good alternative to ensure the quality of drinking water, however poor maintenance of the filter unit could also contribute to heavy metal contamination in drinking water. On top of that, household storage practice also could contribute to heavy metal contamination in drinking water (Brick et al., 2004). Besides, heavy metal present in drinking water is in soluble form and readily absorbable (Chiron et al., 2003). Thus, heavy metal can be easily absorbed and accumulated in human body (Chiron et al., 2003).

Table 1.1: Drinking water contaminants

Contaminants	Potential Health Effects from Long-Term Exposure	Sources of Contamination
Microorganisms		
Total Coliforms (including fecal coliform and <i>E. Coli</i>)	Not a health threat in itself but it is used to indicate whether other potentially harmful bacteria may be present.	Human and animal fecal waste.
<i>Cryptosporidium</i>	Gastrointestinal illness (such as diarrhea, vomiting, and cramps)	
Disinfection byproduct		
Chlorite	Anemia; infants and young children: nervous system effects	Byproduct of drinking water disinfection
Disinfectants		
Chlorine	Eye/nose irritation; stomach discomfort	Water additive used to control microbes
Inorganic pollutants		
Arsenic	Skin damage or problems with circulatory systems, and may have increased risk of getting cancer	Erosion of natural deposits, runoff from glass and electronics production wastes
Cadmium	Kidney damage	Corrosion of galvanised pipes, erosion of natural deposits, discharge from metal refineries
Copper	Liver or kidney damage	
Lead	Kidney problems; high blood pressure	Corrosion of household plumbing systems, erosion of natural deposits
Chromium	Allergic dermatitis	Discharge from steel and pulp mills, erosion of natural deposits
Organic pollutants		
Dichloromethane	Liver problems; increased risk of cancer	Discharge from drug and chemical factories
Radionuclides		
Uranium	Increased risk of cancer, kidney toxicity	Erosion of natural deposits

(Source: USEPA, 2009)

1.2 Problem statement

Kelantan river water is treated at Kelar Water Treatment Plant before being used as drinking water in Pasir Mas District (AKSB, 2014). However, Kelantan river water quality was deteriorating due to anthropogenic activities such as logging activity near the upper stream of the river, sand mining and agricultural activity along the riverside (Rahim and Shahwahid, 2011; Yen and Rohasliney, 2013; Yusoff et al., 2015; Asyirah et al., 2016). The increase of land use at the upper river for logging and agricultural activity has resulted changes of river water quality from class I-II at the upper stream to class III at the downstream of Kelantan River (Rahim and Shahwahid, 2011; Yusoff et al., 2015; Asyirah et al., 2016). Furthermore, uncontrolled sand mining activity along the Kelantan River was also contributed to the river water quality deterioration (Yen and Rohasliney, 2013). Besides, land clearing due to logging and agricultural activities and fertilizers run-off from agricultural area were the main source of heavy metal (Fe, Pb, Zn, Cu and Cd) pollution to Kelantan River (Ahmad, 2009; Yen and Rohasliney, 2013; Yusoff et al., 2015). A study done by Yusoff et al., (2015) have found that concentration of Fe in Kelantan river water was exceeded the permissible limit of 0.3 mg/L set by MMOH for raw water guideline with the highest concentration up to 4.37 mg/L.

Water supply categorised as class III such as Kelantan river water requires advanced water purification treatment to remove organic matter and other pollutants before being used as water supply (Kubota and Tsuchiya, 2010; Halim and Khan, 2012). Yet, water treatment plant at Kelar Water Treatment Plant is still practicing conventional treatment system (AKSB, 2014). Usage of alum in conventional water treatment system at coagulation stage will contribute to heavy metal residue in drinking water (Wang and Peng, 2010; Fu and Wang 2011). Moreover, previous studies in Malaysia have found that high concentration of Al in drinking water is due to the excess use of alum in water treatment process (Qaiyum et al. 2011; Farizwana et al., 2010; Dzulfakar et al, 2010). According to Wang and Peng (2010), pH adjustment during coagulation process is crucial to avoid heavy metal residues in treated drinking water. Additionally, higher alum dosage is needed to remove higher turbidity and dissolved organic carbon in raw water such as water from Kelantan river (Srinivasan and Viraraghavan, 2002; Halim and Khan, 2012).

On the other hand, there are possibilities of heavy metal being leached out into drinking water supply during transfer and storage stage (Lim et al., 2013). Treated water is distributed to the Pasir Mas population through pipeline system which consists of mild steel and HDPE pipes (AKSB, 2014). Moreover, majority of pipeline systems in Kelantan are using mild-steel pipe which has low resistance towards corrosion (AKSB, 2013; Pearce, 2013). In this regard, Mustapha and Anuar (2011) has reported that concentration of Fe in drinking water supplied to Pasir Mas District was higher than the drinking water guideline (0.3 mg/L) with concentration of 4.89 mg/L. Besides pipeline, fittings material such as brass and copper also contributed to leach out of heavy metal such as Pb and Cu into drinking water (Tam and Elefsiniotis, 2009). Apart from that, corrosion of pipeline and fittings depends on drinking water characteristic such as pH, high dissolved oxygen and temperature (Turek et al., 2011; Tam and

Elefsiniotis, 2009). Meanwhile, Lim et al. (2013) has found that Pb in first flush was higher than in the fully flush drinking water. The finding shows that pipeline corrosion and stagnation of water in the pipeline will allow contacts between water with leached Pb (Kim et al. 2011; Lim et al. 2013).

Drinking water studies in Malaysia have focused on concentration of heavy metal without further study on the risk and accumulation of heavy metal in population. Most of the heavy metal concentration in the Malaysian drinking water complied with standard from Malaysian Drinking Water Quality Standard, except for Pb, Al and Fe (Ong et al., 2007; Nalatambi, 2009; Azrina et al., 2011; Lim et al., 2013; Qaiyum et al. 2011; Farizwana et al., 2010; Dzulfakar et al, 2010). Although the heavy metal concentration were under permissible limit, there is lack of investigation on the possible sources of heavy metal in the treated drinking water at the point of use by population (Lim et al., 2013). Furthermore, literatures on knowledge, attitude and practice (KAP) related to heavy metal contamination in drinking water is limited. A study by Aini et al., (2007) found that the urban population (Seremban, Negeri Sembilan) practices good drinking water quality improving techniques, by using a filtration unit at home and boiling drinking water. Another study by Azlina et al., (2013) has found lack of knowledge regarding safe drinking water among the rural population (Kelantan). Apart from KAP, there have been few studies investigating health risk of heavy metal to the Malaysia population via drinking water ingestion using Health Risk Assessment (HRA) (Qaiyum et al., 2011; Dzulfakar et al., 2011). However, there is still limited quantitative data on KAP, HRA and biomarker pertaining to the exposure of heavy metal to the Malaysia population through drinking water ingestion pathway, particularly in areas which use treated Kelantan River water.

1.3 Study justification

Drinking water from Pasir Mas District has been reported to contain high Fe concentration ($489\mu\text{g/L}$) by Mustapha and Anuar (2011). Although heavy metal removal using an efficient conventional water treatment system could achieve heavy metal removal of 99% as reported by Chang and Wang (2007), other factors such as heavy metal leaching out during transfer and storage stage could also contaminate drinking water supply (Buchet and Lison, 2000; Lim et al., 2013). Thus, usage of mild steel pipe and fittings of brass and copper might be the factor of heavy metal contamination in drinking water in Pasir Mas District. Previous studies have found that corrosion of pipeline system is a factor of heavy metal contamination in drinking water (Buchet and Lison, 2000; Lim et al., 2013). Therefore, further research is needed to provide a better understanding that conventional treatment system is suitable to treat Kelantan river water which is classified as class III to be used as drinking water. This study is also required to provide knowledge of pipeline system material, which could be the source of heavy metal in treated drinking water supplied in Pasir Mas population.

Heavy metal contamination in Malaysian drinking water has been well studied (Ong et al., 2007; Nalatambi, 2009; Azrina et al., 2011; Lim et al., 2013; Qaiyum et al. 2011; Farizwana et al., 2010; Dzulfakar et al, 2010). However, there is lack of evidence that heavy metal in the drinking water could give adverse health effects to the Malaysia population. Furthermore, there is also very limited information regarding heavy metal accumulation in the Malaysia population due to intake of heavy metal through drinking water. Hence, this study will provide information to fulfil the knowledge gap on health risk assessment (HRA) and biomarker related to the exposure of heavy metal to population through drinking water ingestion pathway.

1.4 Significant of study

Outputs from this study included the level of knowledge, attitude and practice (KAP) of the Pasir Mas population regarding heavy metal contamination in drinking water. Detailed discussion of KAP level regarding heavy metal contamination in drinking water was provided to open wide field of information that will be very useful for authorities to improve drinking water management in the future.

This has study also shown variation, level and distribution of heavy metal concentration in drinking water and toenail from Pasir Mas District. The discussion presented were including the possible sources of heavy metal in drinking water. This study also provided evidence of incorporation of heavy metal investigation in drinking water and biomarker to give a clear picture on heavy metal accumulation in the Pasir Mas population via drinking water. This information provide a clear picture of biomarker as a useful tool to understand the end point of heavy metal once the elements ingested into the human body. This study also contributed an understanding of significant influential parameters or confounders to heavy metal accumulation in toenail, which may be applied for other exposure pathways in future studies.

Lastly, this study also provided evidence of application from Health Risk Assessment (HRA) are a useful tool to estimate more accurate health risk of a population. This will help decision makers to maintain or improve drinking water quality for the sake of public health.

1.5 Study limitation

Design of this study is cross-sectional study, which only provides a snapshot of the outcome and the characteristics associated with it, at a specific point in time (Levin, 2006). The heavy metal in drinking water data presented in this study represent at the point of sampling without consideration of seasonal factor which could affect the heavy metal distribution (Yusoff et al., 2015). However, Mann (2003) has stated that cross-sectional study is quick, less expensive and produces multiple outcomes. Thus, a broad base of knowledge from this study will be obtained to provide more information for future research (Levin, 2006).

Confounder is a contributing factor that may be resulting in underestimation and overestimation of the exposure (McNamee, 2003). The most important step to control confounders is to identify the possible confounders. In this study, possible confounders are demographic, dietary pattern and smoking habit which were obtained using questionnaire as these factors can increase or decrease heavy metal concentration in toenail. Confounders such as demographic, dietary pattern and smoking can be controlled by statistical method. Thus, statistical method using multiple linear regression can be applied to determine the role of confounders are associated with heavy metal concentration in toenail (Slotnick et al., 2008). Sample population chosen in this study is based on available respondents during the day of sampling. Sampling was conducted on weekdays and weekend, which resulting higher response rate among housewives compared to working men. However, previous studies had shown that gender were not a contributing factor to heavy metal concentration in toenail (Coelho et al., 2012; Barbosa et al., 2005).

Furthermore, this study only focused on heavy metal exposure through drinking water intake. This is due to the fact that heavy metal present in drinking water is in soluble form which is can be easily absorbed and accumulated in human body (Chiron et al., 2003). Other routes of exposure such as dietary intake, dermal exposure, inhalation exposure and occupational exposure were not considered in quantitative manner. This study is intended to study the heavy metal exposure on general adult population which does not requires population classified according to their occupational exposure. Similar studies has been conducted by Kavcar et al., (2009) which has reported exposure of heavy metal on general population in Italy. Furthermore, a study done by Saat et al., (2013) have found that toenail heavy metal concentration in farmers from Kelantan is not associated with their occupational exposure.

The order of important heavy metal exposure routes can be rank as absorption < inhalation < ingestion (Beckett et al., 2007; Cornelis and Nordberg, 2007). Ingestion route is indirect exposure by intake of food and drinking water into gastrointestinal tract (Beckett et al., 2007; Cornelis and Nordberg, 2007). Beckett et al., (2007) stated that exposure through the skin is usually less important than inhalation and ingestion pathways. Cornelis and Nordberg (2007) stated that intake of heavy metal through inhalation pathway is usually small compared to ingestion pathway. Thus, ingestion of food and drinking water is the major source of general population exposure (Beckett

et al., 2007; Xu et al., 2006). Analysis of heavy metal in food is a complex process due to various pattern of a population diet and the analysis should represent cooking method (Cornelis and Nordberg, 2007; Żukowska and Biziuk, 2008). Food analysis for heavy metal also could contribute to more confounder factors such as source of the food item, cooking method and utensils used (Żukowska and Biziuk, 2008). Furthermore, heavy metal in drinking water also could be a factor of heavy metal in food item during washing and cooking process such as reported by Ndilila et al., (2013). Thus, this study only focused on drinking water to reduce the possible confounders and the confounders considered can be controlled using qualitative method and statistical analysis.



1.6 Conceptual framework

Figure 1.2 shows the conceptual framework of this study by estimating carcinogenic and non-carcinogenic health risk of heavy metal exposure through drinking water ingestion pathway, along with examining biomarker of exposure (toenail) with consideration of confounders.

Heavy metal occur in environmental media (water, soil and air) due to both natural occurrences and anthropogenic sources (Kavcar et al., 2009; Ahmad et al., 2009). Pollution of heavy metal in water bodies such as groundwater and surface water will affect the quality of drinking water sources, even the drinking water was treated before distributed to the population (Bobaker et al., 2014).

Heavy metal can be absorbed into human through three main routes which are ingestion, inhalation, and dermal contact (USEPA, 2011). However, ingestion of drinking water has been considered as the main pathway of heavy metal exposure due to the solubility of heavy metal in water, which are readily absorbable into the human body (Chiron et al., 2003). Carcinogenic and non-carcinogenic health risks through ingestion pathway were estimated using Health Risk Assessment (HRA).

There are a few biomarkers has been used to monitor exposure of heavy metal through drinking water, such as blood, urine, toenail, fingernail and hair (Esteban and Castaño, 2009). However, previous studies had suggested that toenail is the most suitable biomarker compared to blood, urine, fingernail and hair due to its rapid growth, less external contamination, adequate sample availability and incorporation of elements in the tissue (Adair et al., 2006; Sukumar, 2006; Slotnick et al., 2007).




Figure 1.1: Conceptual framework of this study

1.7 Research objective and hypotheses

1.7.1 General objective

To estimate exposure and health risks of heavy metal (Al, Pb, Fe, Cu, Cd, Cr, Ni & Zn) through drinking water in the population of Pasir Mas District.

1.7.2 Specific objectives

1. To ascertain the level of knowledge, attitude and practice (KAP) on heavy metal contamination in drinking water in Pasir Mas population.
2. To determine heavy metal concentration in drinking water and toenail from Pasir Mas District.
3. To identify possible sources of heavy metal in drinking water from Pasir Mas.
4. To investigate correlations between heavy metal (Fe, Cu and Pb) concentration in drinking water and toenail.
5. To identify the associations of confounders with heavy metal in toenail.
6. To evaluate health risks for heavy metal exposure via drinking water among the population of Pasir Mas District.

1.7.3 Research hypotheses

1. There is significant sources of heavy metal in drinking water from Pasir Mas.
2. There is significant correlation between heavy metal concentration in drinking water and toenail.
3. There is significant associations between confounders and heavy metal in toenail.
4. There are health risks presents for heavy metal exposure via drinking water among the population of Pasir Mas District.

1.8 Conceptual Definition

1.8.1 Heavy metal

Heavy metal is defined as an element with densities greater than 5g/cm^3 (Jaishankar et al., 2014; Tchounwou et al., 2014). Heavy metal is also considered as trace elements or trace metal because of their presence in trace concentrations (ppb range to less than 10ppm) in environmental matrices (Tchounwou et al., 2014). United State Environmental Protection Agency (USEPA, 2014) has listed Al, As, Pb, Fe, Cu, Cd, Cr, Ni & Zn as the target heavy metal analyte to be analysed in the environmental assessment programme under USEPA. Physical properties of heavy metal are listed as below (Cornelis and Nordberg, 2007):

1. High reflectivity that is responsible for the characteristic metallic cluster
2. High electrical conductivity, decreasing with increasing temperature
3. High thermal conductivity
4. Mechanical properties such as strength and ductility

1.8.2 Health Risk Assessment (HRA)

HRA is the process to estimate the nature and probability of adverse health effects in humans who may be exposed to chemicals in contaminated environmental media, now or in the future (USEPA, 2015).

1.8.3 Operational Definition

1.8.3.1 Health Risk Assessment

Health risk assessment was directly calculated. Equation (1) shows calculation of chronic daily intake (CDI) (mg/kg/day) through ingestion route (USEPA, 2011; Lim et al., 2013). This equation was used to estimate exposure to each heavy metal.

$$CDI (mg/kg/day) = \frac{C_{dw} \times IR \times EF \times ED}{BW \times AT} \quad (1)$$

Where;

CDI = Chronic Daily Intake (mg/kg/day)
C_{dw} = Heavy metal concentration in drinking water (mg/L)
IR = Drinking water ingestion rate (L/day)
ED = Exposure duration (years)
EF = Exposure frequency (day/year)
BW = Body weight (kg)
AT = Averaging time (ED x 365 days)

Hazard quotient (HQ) was calculated to estimate non-carcinogenic risk using equation (2) (USEPA, 2011; Kavcar et al., 2009; Lim et al., 2013). RfD were obtained from the USEPA (IRIS, 2012).

$$HQ = \frac{CDI}{RfD} \quad (2)$$

Where;

HQ= hazard quotient,
CDI= chronic daily intake (mg/kg/day)
RfD= oral reference dose of heavy metal (mg/kg/day).

HQ was compared with the values of risk acceptability for non-carcinogenic health risks. It is assumed that no chronic non-carcinogenic health risks faced by population when HQ < 1.

Cancer risk associated with ingestion exposure was calculated using both equation (3) and equation (4) as shown below (USEPA, 2011).

$$CDI (mg/kg/day) = \frac{C_{dw} \times IR \times ED \times EF}{BW \times AT} \quad (3)$$

Where;

CDI = Chronic daily intake (mg/kg/day)
 C_{dw} = Heavy metal concentration in drinking water (mg/L)
IR = Drinking water ingestion rate (L/day)
ED = Exposure duration (years)
EF = Exposure frequency (day/year)
BW = Body weight (kg)
AT = Averaging time (25550 days)

$$\text{Lifetime cancer risk (LCR)} = CDI \times SF \quad (4)$$

Where;

CDI = Chronic daily intake (mg/kg/day)
SF = slope factor (mg/kg/day)

Slope factor was stated on USEPA's (2012) Integrated Risk Information System (IRIS). The acceptable LCR ranges from one in ten thousand to one in a million ($10^{-6} < LCR < 10^{-4}$).

REFERENCES

- Abdulrahman, F.I., Akan, J.C., Chellube, Z.M. and Waziri, M. (2012). Levels of heavy metals in human hair and nail samples from Maiduguri Metropolis, Borno State, Nigeria. *World Environment* 2(4):81-89. doi: 10.5923/j.env.20120204.05.
- Adair, B.M., Hudgens, E.E., Schmitt, M., Calderon, R.L. and Thomas, D.J. (2006). Total arsenic concentrations in toenails quantified by two techniques provide a useful biomarker of chronic arsenic exposure in drinking water. *Environmental Research* 101(2):213-220.
- Agusa, T., Kunito, T., Yasunaga, G., Iwata, H., Subramanian, A., Ismail, A. and Tanabe, S. (2005). Concentrations of trace elements in marine fish and its risk assessment in Malaysia. *Marine pollution bulletin* 51(8), 896-911.
- Ahmad, A.K., Mushrifah, I. and Shuhaimi-Othman, M. (2009). Water quality and heavy metal concentrations in sediment of Sungai Kelantan, Kelantan, Malaysia: A baseline study. *Sains Malaysiana* 38(4):435-442.
- Ahmad, M.K., Islam, S., Rahman, M.S., Haque, M.R. and Islam, M.M. (2010). Heavy metals in water, sediment and some fishes of Buriganga River, Bangladesh. *International Journal of Environmental Research* 4(2), 321-332.
- Aini, M.S., Fakhurul-Razi, A., Mumtazah, O. and Chen, J.M. (2007). Malaysian households' drinking water practices: A case study. *The International Journal of Sustainable Development & World Ecology* 14(5), 503-510.
- Air Kelantan Sdn Bhd (AKSB) (2013). Information on water treatment plant and treatment system until December 2013. AKSB, Kota Bharu, Kelantan. (Unpublished material).
- Air Kelantan Sdn. Bhd. (AKSB) (2014). List of Drinking Water Treatment Plants. AKSB, Kota Bharu. Retrieved 10th June 2015 from <http://airkelantan.com.my/maklumat-pengguna/senarai-loji-peta/>
- Aitio, A., Bernard, A., Fowler, B.A. and Nordberg, G.F. (2007). Biological Monitoring and Biomarkers. In: Fowler, B.A., Nordberg, G.F., Nordberg, M., and Friberg, L. (Ed.), *Handbook on the Toxicology of Metals, 3rd Edn.* (pp. 65-78). Academic Press.

American Public Health Association (APHA) (2005). *Standard Methods for the Examination of Water and Wastewater*. American Water Works Association, Water Environment Federation, Washington, DC, USA, 21st edition.

Aminah, A.H. (2012). *Health risk assessment of aluminium residue exposure in drinking water among residents in two villages in Kuala Terengganu, Terengganu*. (Unpublished degree dissertation). Universiti Putra Malaysia.

Aris, A.Z., Abdullah, M.H., Ahmed, A., Woong, K.K. (2007). Controlling factors of groundwater hydrochemistry in a small island's aquifer. *International Journal of Environmental Science and Technology* 4(4):441-450.

Aris, A.Z., Kam, R.C.Y., Lim, A.P., Praveena, S.M. (2013), Concentration of ions in selected bottled water samples sold in Malaysia. *Applied Water Science* 3(1):67-75.

Asaduzzaman, K., Khandaker, M.U., Amin, Y.M., and Mahat, R. (2015). Uptake and distribution of natural radioactivity in rice from soil in north and west part of peninsular Malaysia for the estimation of ingestion dose to man. *Annals of Nuclear Energy*, 76:85-93.

Asyirah, A.R., Ismail, W.R., Narimah, S., Rahaman, Z.A., Mardiana, A., Widad, F. *Catchment Health and Sub-Catchment Land-use Activities as Risk Indicator for Disaster Risk Reduction: Case Study of Galas River, Kelantan*. Paper presented in Persidangan Kajian Bencana Banjir 2016, April 4-6, 2016 Putrajaya (Malaysia).

Azlan, A., Khoo, H.E., Idris, M.A., Amin, I., Razman, M.R. (2012), Consumption patterns and perception on intake of drinking water in Klang Valley, Malaysia. *Pakistan Journal of Nutrition* 11(6):584-590.

Azlina, A., Nik Rosmawati, N.H., Zaliha, I. (2013). Factors associated with preferences for safe drinking water among outpatients at Hospital Universiti Sains Malaysia (HUSM), Kelantan, Malaysia. *Health and the Environment Journal* 4 (1):31-41.

Azmir, A. (2003). *Kepekatan aluminium dalam air minum dan penilaian risiko kesihatan di dua buah petempatan di Selangor*. (Unpublished degree dissertation). Universiti Putra Malaysia.

- Azrina, A., Khoo, H.E., Idris, M.A., Amin, I., Razman, M.R. (2011). Major inorganic elements in tap water samples in Peninsular Malaysia. *Malaysian Journal of Nutrition* 17(2):271-276.
- Azrina, A., Khoo, H.E., Idris, M.A., Amin, I., Razman, M.R. (2012). Evaluation of minerals content of drinking water in Malaysia. *The Scientific World Journal* 1-10.
- Baig, J.A., Kazi, T.G., Shah, A.Q., Afridi, H.I., Khan, S., Kolachi, N.F., Kandhro, G.A., Wadhwa, S.K., Shah, F. (2011). Evaluation of toxic risk assessment of arsenic in male subjects through drinking water in southern Sindh Pakistan. *Biological Trace Element Research* 143:772-786.
- Barbosa, Jr.F., Tanus-Santos, J.E., Gerlach, R.F., Parsons, P.J. (2005) A critical review of biomarkers used for monitoring human exposure to lead: advantages, limitations, and future needs. *Environmental Health Perspectives* 113(12):1669-1674.
- Basamba, T.A., Sekabira, K., Kayombo, C.M., and Ssegawa, P. (2013). Application of factor and cluster analyses in the assessment of sources of contaminants in borehole water in Tanzania. *Polish Journal of Environmental Studies* 22(2).
- Bashir, F.H., Othman, M.S., Mazlan, A.G., Rahim, S.M., and Simon, K.D. (2013). Heavy metal concentration in fishes from the coastal waters of Kapar and Mersing, Malaysia. *Turkish Journal of Fisheries and Aquatic Sciences*, 13: 375-382.
- Beckett, W.S., Nordberg, G.F., and Clarkson, T.W. (2007) Routes of exposure, dose, and metabolism of metals. In: Fowler, B.A., Nordberg, G.F., Nordberg, M., and Friberg, L. (Ed.), *Handbook on the Toxicology of Metals, 3rd Edn.* (pp. 39-64). Academic Press.
- Bergomi, M., Pellacani, G., Vinceti, M., Bassissi, S., Malagoli, C., Alber, D. and Vivoli, R. (2005). Trace elements and melanoma. *Journal of Trace Elements in Medicine and Biology* 19(1): 69-73.
- Bobaker, A.M., Elkhidir, E.E., Sarmani, S.B. *Modeling and prediction of the total manganese concentration levels in the finished water of the three water treatment plants in the Linggi River Basin of Negeri Sembilan, Malaysia.* Paper presented at the International Conference on Agricultural, Ecological and Medical Sciences (AEMS-2014) Feb. 6-7, 2014 Bali (Indonesia).

- Boogaard, P.J., and Money, C.D. (2008). A proposed framework for the interpretation of biomonitoring data. *Environmental Health* 7(1): 1-6.
- Brick, T., Primrose, B., Chandrasekhar, R., Roy, S., Muliyl, J., and Kang, G. (2004). Water contamination in urban south India: household storage practices and their implications for water safety and enteric infections. *International Journal of Hygiene and Environmental Health* 207(5): 473-480.
- Brima, E.I., Haris, P.I., Jenkins, R.O., Polya, D.A., Gault, A.G., Harrington, C.F. (2006). Understanding arsenic metabolism through a comparative study of arsenic levels in the urine, hair and fingernails of healthy volunteers from three unexposed ethnic groups in the United Kingdom. *Toxicology and Applied Pharmacology* 216(1):122-30.
- Buchet, J.P., Lison, D. (2000). Clues and uncertainties in the risk assessment of arsenic in drinking water. *Food and Chemical Toxicology* 38(1):81-85.
- Button, M., Jenkin, G.R.T., Harrington, C.F., and Watts, M.J., (2009). Human toenails as a biomarker of exposure to elevated environmental arsenic. *Journal of Environmental Monitoring* 11(3): 610–617.
- Caceres, D.D., Pinoa, P., Montesinos, N., Atalah, E., Amigod, H., and Loomis, D. (2005). Exposure to inorganic arsenic in drinking water and total urinary arsenic concentration in a Chilean population. *Environmental Research* 98(2):151-159.
- Carneiro, M.F.H., Grotto, D., Batista, B.L., Rhoden, C.R., and Barbosa, F. (2011). Background values for essential and toxic elements in children's nails and correlation with hair levels. *Biological Trace Element Research* 144: 339-350.
- Centre for Disease Control and Prevention (CDC) (2008). A guide to conducting household surveys for Water Safety Plans. Atlanta: U. S. Department of Health and Human Services. Retrieved 10th March 2013 from http://www.cdc.gov/nceh/ehs/gwash/Publications/Guide_Conducting_Household_Surveys_for_Water_Safety_Plans.pdf.
- Chan, C.L., Zalifah, M.K., and Norrakiah, A.S. (2007). Microbiological and physicochemical quality of drinking water. *Malaysian Journal of Analytical Sciences* 11(2): 414-420.
- Chang, Q., Wang, G., (2007). Study on the macromolecular coagulant PEX which traps heavy metals. *Chemical Engineering Science* 62(17): 4636-4643.

Chen, L., Jia, R. B., and Li, L. (2013). Bacterial community of iron tubercles from a drinking water distribution system and its occurrence in stagnant tap water. *Environmental Science: Processes & Impacts* 15(7): 1332-1340.

Chiba, M., Shinohara, A., Sekine, M., and Hiraishi, S. (2006). Drinking water quality from the aspect of element concentrations. *Journal of Radioanalytical and Nuclear Chemistry* 269(3): 519–526.

Chiron, N., Guilet, R., and Deydier, E. (2003). Adsorption of Cu (II) and Pb (II) onto a grafted silica: isotherms and kinetic models. *Water Research* 37(13):3079-3086.

Coelho, P., Costa, S., Susana, S., Walter, A., Ranville, J., Sousa, A.C.A., Costa, C., Coelho, M., García-Lestón, J., Pastorinhoc, M.R., Laffon, B., Pásaro, E., Harrington, C., Taylor, A., Teixeira, J.P. (2012). Metal(loid) levels in biological matrices from human populations exposed to mining contamination-Panasqueira Mine (Portugal). *Journal of Toxicology and Environmental Health, Part A* 75:893-908.

Cornelis, R., and Nordberg, M. (2007). General chemistry, sampling, analytical methods, and speciation In: Fowler, B.A., Nordberg, G.F., Nordberg, M., and Friberg, L. (Ed.), *Handbook on the Toxicology of Metals, 3rd Edn.* (pp. 38-11). Academic Press.

Daneshmand, S., Huat, B.B., Moayedi, H., Ali, T.A.M. (2011). Study on water quality parameters of Linggi and Melaka rivers catchments in Malaysia. *Engineering Journal* 15(4): 41-52.

Del Razo, L.M., Styblo, M., Cullen, W.R., Thomas, D.J. (2001). Determination of trivalent methylated arsenicals in biological matrices. *Toxicology and Applied Pharmacology* 174(3):282-93.

Department of Environment (DOE) Malaysia (2010). Malaysia environmental quality report 2010. Retrieved 30th April 2013 from http://appsdoegovmy/portal_efiles/e_publication/indexphp?dir=Laporan+-+Reports/Environmental+Quality+Report+%28EQR%29/Environmental+Quality+Report+%28EQR%29+2011,2010.

Department of Environment (DOE) Malaysia (2011). Malaysia environmental quality report 2011. Retrieved 30th April 2013 from http://appsdoegovmy/portal_efiles/e_publication/indexphp?dir=Laporan+-

+Reports/Environmental+Quality+Report+%28EQR%29/Environmental+Quality+Report+%28EQR%29+2011, 2011.

Department of Irrigation and Drainage (DID) of Malaysia. 2009. Study on the River Water Quality Trends and Indexes in Peninsular Malaysia. Water Resources Publication No. 21. Retrieved 15th January 2016 from http://h2o.water.gov.my/man_water/WRP21.pdf

Department of Statistic Malaysia (2010). Population and Housing Census 2010. Retrieved 20th February 2013 from http://www.statistics.gov.my/portal/download_Population/files/census2010/Ta_buran_Penduduk_dan_Ciri-ciri_Asas_Demografi.pdf; 2010

Desai, C.K., Iyer, G., Panchal, J., Shah, S., and Dikshit, R.K. (2011). An evaluation of knowledge, attitude, and practice of adverse drug reaction reporting among prescribers at a tertiary care hospital. *Perspectives in Clinical Research* 2(4): 129.

Deshommes, E., Prévost, M., Levallois, P., Lemieux, F., and Nour, S. (2013). Application of lead monitoring results to predict 0–7 year old children's exposure at the tap. *Water Research* 47: 2409-2420.

DeZuane, J. (1997). *Handbook of drinking water quality*. John Wiley & Sons.

Diaconu, D., Nănău, M.M., Nechifor, E., Nechifor, O. (2009). Aluminium concentration in drinking water from Moldova territory, Romania. *Ovidius University Annals of Chemistry* 20(1): 115–118.

Dzulfakar, M.A., Shaharuddin, M.S., Muhaimin, A.A., Syazwan, A.I. (2011). Risk assessment of aluminum in drinking water between two residential areas. *Water* 3(3):882-893.

Egeghy, P.P., Quackenboss, J.J., Catlin, S., and Ryan, P.B. (2005). Determinants of temporal variability in NHEXAS-Maryland environmental concentrations, exposures, and biomarkers. *Journal of Exposure Science and Environmental Epidemiology* 15(5):388-397.

Esteban, M., and Castaño, A. (2009). Non-invasive matrices in human biomonitoring: A review. *Environment International* 35(2):438-449.

- Falk-Filipsson, A., Hanberg, A., Victorin, K., Warholm, M., and Wallén, M. (2007). Assessment factors-applications in health risk assessment of chemicals. *Environmental Research* 104(1):108-27.
- Farizwana, M.R.S., Mazrura, S., Zurahanim, F.A., and AhmadRohi, G. (2010). Determination of aluminium and physicochemical parameters in the palm oil estates water supply at Johor, Malaysia. *Journal of Environmental and Public Health* 2010(2010): 1-7.
- Field, A., (2013). *Discovering Statistics Using IBM SPSS Statistics, fourth ed.* Sage, Singapore.
- Food and Agricultural Organization of the United Nations (FAO) (2010). FAO's Information System on Water and Agriculture Malaysia. Retrieved 20th June 2013 from http://www.fao.org/nr/water/aquastat/countries_regions/malaysia/index.stm.
- Fu, F., and Wang, Q. (2011). Removal of heavy metal ions from wastewaters: a review. *Journal of Environmental Management* 92(3): 407–18.
- Fulazzaky, M.A., Seong, T.W., & Masirin, M.I.M. (2010). Assessment of water quality status for the Selangor River in Malaysia. *Water, Air, and Soil Pollution* 205(1-4): 63-77.
- Gamble, M.V., Liu, X., Ahsan, H., Pilsner, J.R., Ilievski, V., and Slavkovich, V. (2005). Folate, homocysteine and arsenic metabolism in arsenic-exposed individuals in Bangladesh. *Environmental Health Perspectives* 113(12):1683-1688.
- Gault, A.G., Rowland, H.A.L., Charnock, J.M., Wogelius, R.A., Gomez-Morilla, I., and Vong, S. (2008). Arsenic in hair and nails of individuals exposed to arsenic-rich groundwaters in Kandal province, Cambodia. *Science of the Total Environment* 393(1):168-176.
- Gaur, V.K., Gupta, S.K., Pandey, S.D., Gopal, K., and Misra, V. (2005). Distribution of heavy metals in sediment and water of river Gomti. *Environmental Monitoring and Assessment* 102(1-3): 419-433.
- Georgopoulos, P.G., Sasso, A.F., Isukapalli, S.S., Liroy, P.J., Vallero, D.A., Okino, M., and Reiter, L. (2009). Reconstructing population exposures to environmental chemicals from biomarkers: challenges and opportunities. *Journal of Exposure Science and Environmental Epidemiology* 19(2): 149-171.

- Ghasemi, A., and Zahediasl, S. (2012). Normality tests for statistical analysis: a guide for non-statisticians. *International journal of endocrinology and metabolism* 10(2): 486-489.
- Ghazali, A.R., Abdul Razak, N.E., Othman, M.S., Othman, H., Ishak, I., Lubis, S.H., and Abdullah, R. (2012). Study of heavy metal levels among farmers of Muda Agricultural Development Authority, Malaysia. *Journal of Environmental and Public Health*, 2012.
- Gliem, J.A. and Gliem, R.R. (2003). Proceedings from 2003 Midwest Research to Practice Conference in Adult, Continuing, and Community Education: *Calculating, interpreting, and reporting Cronbach's alpha reliability coefficient for Likert-type scales*.
- Gonzalez, A.P., Ulrike, L., Satia, J.W., Jessie, A., and White, E. (2008). Correlates of toenail zinc in a free-living U.S. population. *Annals of epidemiology* 18(1):74-77.
- Goullé, J.P., Saussereau, E., Mahieu, L., Bouige, D., Groenwont, S., Guerbet M., and Lacroix, C. (2009). Application of inductively coupled plasma mass spectrometry multielement analysis in fingernail and toenail as a biomarker of metal exposure. *Journal of Analytical Toxicology* 33(2): 9-28.
- Guidotti, T.,L., Moses, M.,S., Goldsmith, D.,F., and Ragain, L. 2015. DC Water and sewer authority and lead in drinking water: a case study in environmental health risk management. *Journal of Public Health Management & Practice* 14(1): 33–41.
- Hair, J.F., Black, W.C., Babin, B.J., Anderson, R.E. (2010). *Multivariate Data Analysis: A Global Perspective, seventh ed.* Upper Saddle River: Prentice Hall.
- Halim, S.A. and Khan, M.M.A. (2012). Proceedings from UMT 11th International Annual Symposium on Sustainability Science and Management: *Generation of Landuse Map and Statistical Analysis of Water Samples*.
- Hall, M., Chen, Y., Ahsan, H., Slavkovich, V., van Geen, A., and Parvez, F. (2006). Blood arsenic as a biomarker of arsenic exposure: Results from a prospective study. *Toxicology* 225(2-3):225-233.
- Harkins, D.K., and Susten, A.S. (2003). Hair analysis: Exploring the state of the science. *Environmental Health Perspectives* 111(4):576–578.

- Hinwood, A.L., Sim, M.R., Jolley, D., de Klerk, N., Bastone, E.B., Gerostamoulos, J., and Drummer, O.H. (2003). Hair and toenail arsenic concentrations of residents living in areas with high environmental arsenic concentrations. *Environmental Health Perspectives* 111(2):187–193.
- Housecroft, C.E., and Sharpe, A.G. (2005). The trace metal of life. In: Housecroft, C.E., and Sharpe, A.G., (Ed.) *Inorganic Chemistry, 2nd edn.* (pp 830-833). Madrid: Pearson.
- Hughes, M. (2006) Biomarkers of exposure: a case study with inorganic arsenic. *Environmental Health Perspectives* 1790(11):1790-1797.
- Iannotti, L.L., Tielsch, J.M., Black, M.M., Black, R.E. (2006). Iron supplementation in early childhood: health benefits and risks. *American Journal of Clinical Nutrition* 84:1261–76.
- Idriss, A.A., and Ahmad, A.K. (2012). Heavy metal concentrations (Cu, Cd and Pb) in sediments in the Juru River, Penang, Malaysia. *Journal of Biological Sciences*, 12(7), 376-384.
- Isa, N.M., Aris, A.Z., Lim, W.Y., Sulaiman, W.N.A.W., and Praveena, S.M. (2013). Evaluation of heavy metal contamination in groundwater samples from Kapas Island, Terengganu, Malaysia. *Arabian Journal of Geosciences* 1-14.
- Islami, N., Taib, S.H., Yusoff, I., and Ghani, A.A. (2012). Integrated geoelectrical resistivity, hydrochemical and soil property analysis methods to study shallow groundwater in the agriculture area, Machang, Malaysia. *Environmental Earth Sciences* 65(3):699-712.
- Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B.B., and Beeregowda, K.N. (2014). Toxicity, mechanism and health effects of some heavy metals. *Interdisciplinary toxicology* 7(2): 60-72.
- Järup, L. (2003). Hazards of heavy metal contamination. *British medical bulletin* 68(1): 167-182.
- Jung, H., Kim, U., Seo, G., Lee, H., and Lee, C. (2009). Effect of dissolved oxygen (DO) on internal corrosion of water pipes. *Environmental Engineering Research* 14(3): 195-199.

- Kakkar, P., and Jaffery, F.N. (2005). Biological markers for metal toxicity. *Environmental Toxicology and Pharmacology* 19(2):335-349.
- Karagas, M.R., Tosteson, T.D., Blum, J., Klaue, B., Weiss, J.E., Stannard, V., Spate, V., and Morris, J.S. (2000). Measurement of low levels of arsenic exposure: A comparison of water and toenail concentrations. *American Journal of Epidemiology* 152(1):84-90.
- Kato, M., Onuma, S., Katoa, Y., Thangac, N.D., Yajima, I., Hoque, M.Z., and Shekhar, H.U. (2010). Toxic elements in well water from Malaysia. *Toxicological & Environmental Chemistry* 92(9):1609–1612.
- Kavcar, P., Sofuoglu, A., and Sofuoglu, S.C. (2009). A health risk assessment for exposure to heavy metal via drinking water ingestion pathway. *International Journal of Hygiene and Environmental Health* 212:216-227.
- Khan, N.I., Owens, G., Bruce, D., and Naidu, R. (2009). Human arsenic exposure and risk assessment at the landscape level: a review. *Environmental Geochemistry and Health* 31(1): 143-166.
- Khan, Y.H., Sarriff, A., Khan, A.H., and Mallhi, T.H. (2014). Knowledge, attitude and practice (KAP) survey of osteoporosis among students of a tertiary institution in Malaysia. Benin City: Pharmacotherapy Group, Faculty of Pharmacy, University of Benin. Retrieved 10th March 2015 from <https://tspace.library.utoronto.ca/handle/1807/61291>.
- Khoo, H.E., Azlan, A., Idris, M.A., Ismail, A., and Razman, M.R. (2011). Evaluation of selected metal elements in commercial drinking water and tap water in Peninsular Malaysia. *Malaysian Journal of Health Sciences* 9(1):5-11.
- Kile, M.L., Houseman, E.A., Breton, C.V., Smith, T., Quamruzzaman, Q., Rahman, M., Mahiuddin, G. and Christiani, D.C. (2007). Dietary Arsenic Exposure in Bangladesh. *Environmental Health Perspectives* 115(6): 889-93.
- Kim, E.J., Herrera, J.E., Huggins, D., Braam, J., and Koshowski, S. (2011). Effect of pH on the concentrations of lead and trace contaminants in drinking water: A combined batch, pipe loop and sentinel home study. *Water Research* 45:2763-2774.
- King, M.W. (2016). Copper and Iron Homeostasis. Retrieved 20th October 2016 from <http://themedicalbiochemistrypage.org/iron-copper.php>.

- Kioko, K.J., and Obiri, J.F. (2012). Household attitudes and knowledge on drinking water enhance water hazards in peri-urban communities in Western Kenya. *Jàmbá: Journal of Disaster Risk Studies* 4(1):1-5.
- Kubota, S., and Tsuchiya, Y. (2010). *Water Quality and Standards - Volume I*. Oxford: Encyclopedia of Water Sciences, Engineering and Technology Resources in the global Encyclopedia of Life Support Systems (EOLSS) Publications Co. Ltd.
- LaKind, J.S., Barraj, L., Tran, N., and Aylward, L. (2008). Environmental chemicals in people: Challenges in interpreting biomonitoring information. *Journal of Environmental Health* 70(9):61-4.
- Lasheen, M.R., Sharaby, C.M., El-Kholy, N.G., Elsherif, I.Y., and El-Wakeel, S.T. (2008). Factors influencing lead and iron release from some Egyptian drinking water pipes. *Journal of Hazardous Materials* 160: 675-680.
- Levin, K.A. (2006). Study design III: Cross-sectional studies. *Evidence-Based Dentistry*. 7, 24–25.
- Lim, C.S., Shaharuddin, M.S., and Sam, W.Y. (2013). Risk assessment of exposure to lead in tap water among residents of Seri Kembangan, Selangor State, Malaysia. *Global Journal of Health Science* 5(2):1-12.
- Lim, W.Y., Aris, A.Z., and Zakaria, M.P. (2012). Spatial variability of metals in surface water and sediment in the Langat River and geochemical factors that influence their water-sediment interactions. *The Scientific World Journal* 1:1-14.
- Lim, W.Y., Aris, A.Z., Ismail, T.H.T., and Zakaria, M.P. (2013). Elemental hydrochemistry assessment on its variation and quality status in Langat River, Western Peninsular Malaysia. *Environmental Earth Sciences* 1-12.
- Lin, C.Y., Abdullah, M.H., Musta, B., Praveena, S.M., and Aris, A.Z. (2011). Stability behavior and thermodynamic states of iron and manganese in sandy soil aquifer, Manukan Island, Malaysia. *Natural Resources Research* 20(1):45-56.
- Lin, J., Ellaway, M., and Adrien, R., (2001). Study of corrosion material accumulated on the inner wall of steel water pipe. *Corrosion Science* 43: 2065-2081.

- Ma, H.W., Hung, M.L., and Chen, P.C. (2007). A systemic health risk assessment for the chromium cycle in Taiwan. *Environment International* 33(2):206-18.
- Maas, R.P., Patch, S.C., Christian, A.M., and Coplan, M.J. (2007). Effects of fluoridation and disinfection agent combinations on lead leaching from leaded-brass parts. *Neurotoxicology* 28: 1023-1031.
- Magomya, A.M., Yebpella, G.G., Udiba, U.U., Amos, H.S., and Latayo, M.S. (2013). Potassium Brómate and Heavy Metal Content of Selected Bread Samples Produced in Zaria. *Nigeria International Journal of Science and Technology* 2(2): 232-237.
- Mahata, J., Basu, A., Ghoshal, S., Sarkar, J.N., Roy, A.K., Poddar, G., Nandy, A.K., Banerjee, A., Ray, K., Natarajan, A.T. and Nilsson, R. (2003). Chromosomal aberrations and sister chromatid exchanges in individuals exposed to arsenic through drinking water in West Bengal, India. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 534(1): 133-143.
- Mahmood, Q., Baig, S.A., Nawab, B., Shafqat, M.N., Pervez, A., and Zeb, B.S. (2011). Development of low cost household drinking water treatment system for the earthquake affected communities in Northern Pakistan. *Desalination* 273:316-320.
- Maisarah, Z. (2012). *Health risk assessment of lead exposure in drinking water in two villages in Kuala Terengganu, Terengganu*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Makokha, A.O., Mghweno, L.R., Magoha, H.S., Nakajugo, A., and Wekesa, J.M. (2008). Environmental lead pollution and contamination in food around Lake Victoria, Kisumu, Kenya. *African Journal of Environmental Science and Technology* 2(10): 349-353.
- Malaysian Food and Regulations, (1985). In Hamid, I.N., Huat Y.T., *Malaysian Law on Food and Drugs*. Kuala Lumpur, Malaysia Law Publisher.
- Malaysian Ministry of Health (MMOH) (2009). Drinking water quality standard, Engineering Services Division, Ministry of Health, Malaysia, 2009. Retrieved 17th May 2013 from <http://kmam.moh.gov.my/public-user/drinking-water-quality-standard.html>.

- Mandal BK, Ogra Y, Suzuki KT (2003) Speciation of arsenic in human nail and hair from arsenic-affected area by HPLC-inductively coupled argon plasma mass spectrometry. *Toxicology and Applied Pharmacology* 189(2):73-83.
- Mann, C.J. (2003). Observational research methods. Research design II: cohort, cross sectional, and case-control studies. *Emergency Medicine Journal* 20(1): 54-60.
- Marchiset-Ferlay, N., Savanovitch, C., and Sauvart-Rochat, M.P. (2012). What is the best biomarker to assess arsenic exposure via drinking water? *Environment International* 39(1):150-171.
- McCluskey, A., and Lalkhen, A.G. (2007). Statistics II: Central tendency and spread of data. *Continuing Education in Anaesthesia, Critical Care & Pain* 7(4): 127-130.
- McNamee, R. (2003). Confounding and confounders. *Occupational and environmental medicine* 60(3): 227-234.
- Mohan, D., and Pittman, C.U. (2007). Arsenic removal from water/wastewater using adsorbents—a critical review. *Journal of Hazardous Materials* 142:1-53.
- Mohd Aizuddin, A. (2012). *Health risk assesment of exposure to residual aluminium (Al) in drinking water of residents in Kamunting Perak*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Mok, W.J., Senoo, S., Itoh, T., Tsukamasa, Y., Kawasaki, K.I., and Ando, M. (2012). Assessment of concentrations of toxic elements in aquaculture food products in Malaysia. *Food Chemistry* 133(4): 1326-1332.
- Mooi, E., and Sarstedt, M.A. (2011). *Concise Guide to Market Research*. Springer-Verlag Berlin Heidelberg. Retrieved 3rd October 2015 from <http://www.springer.com/978-3-642-12540-9>.
- Mordukhovich, I., Wright, R.O., Hu, H., Amarasiriwardena, C., Baccarelli, A., Litonjua, A., Sparrow, D., Vokonas, P. and Schwartz, J. (2012) Associations of toenail arsenic, cadmium, mercury, manganese, and lead with blood pressure in the normative aging study. *Environmental Health Perspectives* 120(1): 98-104.
- Muhammad, S., Shah, M. T. and Khan, S. (2011). Health risk assessment of heavy metals and their source apportionment in drinking water of Kohistan region, northern Pakistan. *Microchemical Journal* 98(2): 334-343.

- Muslim, N.Z.M. and Saim, H. (2012). Proceedings from 2nd International Conference on Ecological, Environmental and Biological Sciences 2012 Bali (Indonesia) (EEBS'2012): *Determination of Heavy Metal in Pelagic Fish, Demersal Fish and Shellfish*.
- Mustapha, M.A. and Anuar, T.C., (2011, November 20). Kesehatan rakyat Kelantan terancam. *Utusan Malaysia*. Retrieved 28th March 2014 from http://ww1.utusan.com.my/utusan/info.asp?y=2011&dt=1120&pub=Utusan_Malaysia&sec=Muka_Hadapan&pg=mh_01.htm.
- Muyibi, S.A., Birima, A.H.M., Mohammed, T.A., and Noor, M.J.M. (2004). Conventional Treatment of Surface Water Using Moringa oleifera Seeds Extract as a Primary Coagulant. *IIUM Engineering Journal* 5(1): 25-35.
- Nalatambi, S. (2009) Determination of metals in tap water using atomic absorption spectrometry: a case study in Bandar Sunway residential area. *Sunway Academic Journal* 6:33-46.
- National Cancer Institute (2015). Risk factors for cancer. Retrieved 10th January from <http://www.cancer.gov/about-cancer/causes-prevention/risk>.
- National Water Services Commission (NWSC) (2015). List Of Product For Water Supply - Category A. Retrieved 25th August 2015 from <http://www.span.gov.my/files/guideline/water/Product-Grouping-Under-Category-A-for-Water-Supply-Product.pdf>.
- Ndilila, W., Callan, A.C., McGregor, L.A., Kalin, R.M., Hinwood, A.L. (2013). Environmental and toenail metals concentrations in copper mining and non-mining communities in Zambia. *International Journal of Hygiene and Environmental Health* 217(1): 62-69.
- Nermell, B., Lindberg, A.L., Rahman, M., Berglund, M., Persson, L.A., Arifeen, S.E. and Vahter, M. (2008). Urinary arsenic concentration adjustment factors and malnutrition. *Environmental Research* 106(2):212-8.
- Nor Hasyimah, A.K., Noik, V.J., Teh, Y.Y., Lee, C.Y., and Ng, H.P. (2011). Assessment of cadmium (Cd) and lead (Pb) levels in commercial marine fish organs between wet markets and supermarkets in Klang Valley, Malaysia. *International Food Research Journal* 18(2): 792-802.

- Nor, Z.Z. (2012). *Health risk assessment of lead exposure in drinking water among residents of Felda Palong, Negeri Sembilan*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Nora, A. (2003). *Kepekatan aluminium dalam air minum dan penilaian risiko kesihatan di dua buah petempatan di Johor*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Norimah, A.K., Safiah, M., Jamal, K., Haslinda, S., Zuhaida, H., Rohida, S., Fatimah, S., Siti Norazlin, Poh, B.K., Kandiah, M., Zalilah, M.S., Wan Manan, W.M., Fatimah, S. and Azmy, M.Y. (2008). Food consumption patterns: Findings from the Malaysian adult nutrition survey (MANS). *Malaysian Journal of Nutrition* 14(1): 25-39.
- Nowak, B., and Chmielnicka, J. (2000). Relationship of lead and cadmium to essential elements in hair, teeth, and nails of environmentally exposed people. *Ecotoxicology and Environmental Safety* 46(3):265-274.
- Nur, A.M.M. (2012). *Risk assessment of aluminium residue concentration in daily intake by human from drinking tap water, Felda Jenderak Selatan, Temerloh, Pahang*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Omar, N.A., Praveena, S.M., Aris, A.Z., and Hashim, Z. (2015). Health Risk Assessment using in vitro digestion model in assessing bioavailability of heavy metal in rice: A preliminary study. *Food chemistry* 188: 46-50.
- Onabolu, B., Jimoh, O.D., Igboro, S.B., Sridhar, M.K.C., Onyilo, G., Gege, A., and Ilya, R. (2011). Source to point of use drinking water changes and knowledge, attitude and practices in Katsina State, Northern Nigeria. *Physics and Chemistry of the Earth, Parts A/B/C* 36(14-15):1189-1196.
- Ong, C., Ibrahim, S., and Gupta, B.S. (2007). A survey of tap water quality in Kuala Lumpur. *Urban Water Journal* 4(1):29-41.
- Ouyang Y, Higman J, Thompson J, O'Toole T, Campbell D (2002) Characterization and spatial distribution of heavy metal in sediment from Cedar and Ortega Rivers subbasin. *J Contam Hydrol* 54(1):19-35.
- Özdemir, S., Elliott, M., Brown, J., Nam, P.K., Hien, V.T., and Sobsey, M.D. (2011). Rainwater harvesting practices and attitudes in the Mekong Delta of

Vietnam. *Journal of Water Sanitation and Hygiene for Development* 1(3): 171-177.

Palmeri, A., Pichini, S., Pacifici, R., Zuccaro, P., and Lopez, A. (2000). Drugs in nails. *Clinical pharmacokinetics* 38(2): 95-110.

Pasir Mas District Council, (2011). Kajian Rancangan Tempatan Jajahan Pasir Mas, 2008-2020. Retrieved 30th September 2016 from <http://www.ugisp.com/lipis/pdf/Pengenalan/Profil%20KRT.pdf>.

Pasir Mas Land and District Office, (2015). Pasir Mas District Profile. Retrieved 30th September 2016 from <http://www.ptjpm.kelantan.gov.my/index.php/en/about-us/district-in-pasir-mas>.

Pearce, M.S. (2013). *U.S. Patent No. 8,574,441*. Washington, DC: U.S. Patent and Trademark Office.

Peng, C.Y., Korshin, G.V., Valentine, R.L., Hill, A.S., Friedman, M.J., and Reiber, S.H. (2010). Characterization of elemental and structural composition of corrosion scales and deposits formed in drinking water distribution systems. *Water research* 44(15): 4570-4580.

Phan, K., Sthiannopkao, S., and Kim, K.W. (2011). Surveillance on chronic arsenic exposure in the Mekong River basin of Cambodia using different biomarkers. *International Journal of Hygiene and Environmental Health* 215(1):51-8.

Qaiyum, M.S., Shaharudin, M.S., Syazwan, A.I., and Muhaimin, A. (2011). Health risk assessment after exposure to aluminium in drinking water between two different villages. *Journal of Water Resource and Protection* 3(4):268-274.

Rahim, A.S.A., and Shahwahid, S.O.M. (2011). A panel data analysis of timber harvesting operations and its impact on the cost of water treatment. *Australian Journal of Basic and Applied Sciences* 5(12): 598-601.

Rahim, B.E.E., Yusoff, I., Samsudin, A.R., Yaacob, W.Z.W., Rafek, A.G.M. (2010). Deterioration of groundwater quality in the vicinity of an active open-tipping site in West Malaysia. *Hydrogeology Journal* 18(4):997-1006.

- Rahimah, S. (2012). *Health risk assessment of residual aluminium in drinking water among settlers of Felda Palong 4,5,6,7 and 8, Negeri Sembilan*. (Unpublished degree dissertation). Universiti Putra Malaysia.
- Rajan, S., Nurul, N.M.F., Appukutty, M., and Ramasamy, K. (2012). Effects of climate changes on dissolved heavy metal concentrations among recreational park tributaries in Pahang, Malaysia. *Biomedical Research* 23(1):23-30.
- Rivera-Núñez, Z., Meliker, J.R., and Meeker, J.D. (2012). Urinary arsenic species, toenail arsenic, and arsenic intake estimates in a Michigan population with low levels of arsenic in drinking water. *Journal of Exposure Science and Environmental Epidemiology* 22 (2): 182-90
- Robert, B., and Richard, B., (2008). *Business research methods and statistics using SPSS*. London: Sage Publications LTD.
- Rodushkin, I., and Axelsson, M.D., (2000). Application of double focusing sector field ICP-MS for multielemental characterization of human hair and nails. Part II. A study of the inhabitants of northern Sweden. *Science of the Total Environment* 262: 21-36.
- Saat, N.Z.M., Mohandas, B., Ghazali, A.R., Ishak, I., Lubis, S.H., Mohamed, N., Hamid, A., Hamid, Z.A. and Othman, H. (2013). Study of heavy metals in hair and nails of among farmers in Kelantan, Malaysia. *Research Journal of Applied Sciences* 8(3): 225-229.
- Sabo, A., Gani, A.M., and Ibrahim, A.Q. (2014). Geochemical Partitioning of Metal Pollutants in Agricultural Soil as Index of Human Health Risk Assessment. *Journal of Environment Pollution and Human Health* 2(5): 91-99.
- Saeki, H., and Sugimachi, K. (2001). Carcinogenic Risk Factors. *Japan Medical Association Journal* 44(6): 245-249.
- Salati, S., and Moore, F., (2010). Assessment of heavy metal concentration in the Khoshk River water and sediment, Shiraz, Southwest Iran. *Environmental Monitoring and Assessment* 164: 677-689.
- Samanta, G., Sharma, R., Roychowdhury, T., and Chakraborti, D. (2004). Arsenic and other elements in hair, nails, and skin-scales of arsenic victims in West Bengal, India. *Science of the Total Environment* 326(1-3):33-47.

- Şeker, S., and Kutlu, B., (2014). Determination of Copper (Cu) Levels for Rivers in Tunceli, Turkey. *World Environment* 4(4): 168-171.
- Simeonova, P., Papazova, P., and Lovchinov, V. (2013). Application of Cluster Analysis in Interpretation of Toxic Metals Accumulation in Biomonitoring Marine Organisms. *Bulg. J. Phys* 40: 269-273.
- Slotnick, M.J., and Nriagu, J.O. (2006). Validity of human nails as a biomarker of arsenic and selenium exposure: A review. *Environmental Research* 102(1):125-39.
- Slotnick, M.J., Meliker, J.R., and Nriagu, J.O., (2008b). Intra-individual variability in toenail arsenic concentrations in a Michigan population, USA. *Journal of Exposure Science and Environmental Epidemiology* 18, 149-157.
- Slotnick, M.J., Meliker, J.R., AvRuskin, G.A., Ghosh, D., and Nriagu, J.O. (2007). Toenails as a biomarker of inorganic arsenic intake from drinking water and foods. *Journal of Toxicology and Environmental Health, Part A* 70(2):148-58.
- Slotnick, M.J., Meliker, J.R., Kannan, S., and Nriagu, J.O. (2008a). Effects of nutritional measures on toenail arsenic concentration as a biomarker of arsenic exposure. *Biomarkers* 13(5): 451-466.
- Slotnick, M.J., Nriagu, J.O., Johnson, M.M., Linder, A.M., Savoie, K.L., Jamil, H.J., and Hammad, A.S. (2005). Profiles of trace elements in toenails of Arab-Americans in the Detroit area, Michigan. *Biological Trace Element Research* 107(2):113-126.
- Sobus, J.R., Tan, Y.M., Pleil, J.D., and Sheldon, L.S. (2011). A biomonitoring framework to support exposure and risk assessments. *Science of the Total Environment* 409(22):4875-4884.
- Spickett, J., Katscherian, D., and Goh, Y.M. (2012). A new approach to criteria for health risk assessment. *Environmental Impact Assessment Review* 32(1):118-122.
- Srinivasan, P.T., and Viraraghavan, T. (2002). Characterisation and concentration profile of aluminium during drinking-water treatment. *Water SA* 28(1): 99-106.

- Srogi, K. (2006). Hair analysis for monitoring environmental pollution and the resulting human exposure to trace metals: An overview. *Environnement, Risques & Santé* 5(5):391-405.
- Sukumar, A. (2006). Human nails as a biomarker of element exposure. In: Ware, G.W., Nigg, H.N., and Doerge, D.R. (Ed.), *Reviews of environmental contamination and toxicology* (pp. 141-177). New York: Springer.
- Sukumar, A., and Subramanian, R. (2007). Relative element levels in the paired samples of scalp hair and fingernails of patients from New Delhi. *Science of the Total Environment* 372(2-3):474-479.
- Sun, G., Xu, Y., Li, X., Jin, Y., Li, B., and Sun, X. (2007). Urinary arsenic metabolites in children and adults exposed to arsenic in drinking water in Inner Mongolia, China. *Environmental Health Perspectives* 115(4):648-52.
- Syarikat Air Johor Holdings (SAJH) (2012). List of specification: pipe & specials. Retrieved 21st July 2014 from http://www.saj.com.my/V2/Material_Spec.php?s_mat=2.
- Syarikat Bekalan Air Selangor (SYABAS) (2013). Pipe length and material. (Unpublished material)
- Tabachnick, B.G., and Fidell, L.S. (2014). *Using multivariate statistics fifth ed.* Boston: Pearson Education.
- Tam, Y.S., and Elefsiniotis, P., (2009). Corrosion control in water supply systems: effect of pH, alkalinity, and orthophosphate on lead and copper leaching from brass plumbing. *Journal of Environmental Science and Health Part A* 44(12): 1251–1260.
- Tamasi, G., and Cini, R. (2004). Heavy metals in drinking waters from Mount Amiata (Tuscany, Italy). Possible risks from arsenic for public health in the Province of Siena. *Science of the Total Environment* 327(1): 41-51.
- Taweel, A., Shuhaimi-Othman, M., and Ahmad, A.K. (2013). Assessment of heavy metals in tilapia fish (*Oreochromis niloticus*) from the Langat River and Engineering Lake in Bangi, Malaysia, and evaluation of the health risk from tilapia consumption. *Ecotoxicology and environmental safety* 93: 45-51.

Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K., and Sutton, D.J. (2012). Heavy metal toxicity and the environment. *Molecular, clinical and environmental toxicology* 101: 133–164.

The Joint FAO/WHO Expert Committee on Food Additives (JECFA), (2004). Evaluation of certain food additives and contaminants. WHO Technical Report Series. Geneva, Switzerland. 922, 1-176. Retrieved 12th January 2016 from http://apps.who.int/iris/bitstream/10665/42849/1/WHO_TRS_922.pdf

Thomas KW, Dosemeci M, Coble JB, Hoppin JA, Sheldon LS, Chapa G (2009) Assessment of a pesticide exposure intensity algorithm in the Agricultural Health Study. *Journal of Exposure Science and Environmental Epidemiology* 20(6):559-569.

Turek, N.F., Kasten, L., Lytle, D.A., and Goltz, M.N. (2011). Impact of plumbing age on copper levels in drinking water. *Journal of Water Supply: Research and Technology-Aqua* 60(1): 1-15.

U.S. Geological Survey. (2006). Chapter A4. Collection of Water Samples. Techniques of Water-Resources Investigations. Book 9 Handbooks for Water-Resources Investigations. National Field Manual for the Collection of Water-Quality Data. Retrieved 20th May 2013 from https://water.usgs.gov/owq/FieldManual/chapter4/pdf/Chap4_v2.pdf.

United Nations Educational, Scientific and Cultural Organization (UNESCO), 2014. WWAP (United Nations World Water Assessment Programme). The United Nations World Water Development Report 2014: Water and Energy. Paris, UNESCO. Retrieved 13th March 2016 from <http://unesdoc.unesco.org/images/0022/002257/225741E.pdf>.

United States Environmental Protection Agency (USEPA), (2016). Table of Regulated Drinking Water Contaminants. Retrieved 15th July 2015 from <https://www.epa.gov/your-drinking-water/table-regulated-drinking-water-contaminants>.

USEPA, Integrated Risk Information System (IRIS) (2012). A-Z list of substances. Retrieved 10th June 2014 from <http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction¼iris.showSubstanceList>.

USEPA. (2011). Exposure Factors Handbook 2011 Edition (Final). U.S. Environmental Protection Agency, Washington, DC, EPA/600/ R-09/052F.

Retrieved 6th May 2014 from <http://www.epa.gov/ncea/efh/pdfs/efh-complete.pdf>.

USEPA. (2015). Human Health Risk Assessment. Retrieved 11th November 2013 from <https://www.epa.gov/risk/human-health-risk-assessment>.

Vasile, G., Ciupe, A., Dinu, C., Gheorghe, D., Chiru, E., and Cruceru, L. (2012). *Evaluation of Drinking Water Quality in Three Municipalities of Romania: The Influence of Municipal and Customer's Distribution Systems Concerning Trace Metals*. INTECH Open Access Publisher.

Wang, S., and Peng, Y., (2010). Natural zeolites as effective adsorbents in water and wastewater treatment. *Chemical Engineering Journal* 156(1): 11-24.

Water Research Foundation. (2013). Electricity Use and Management in the Municipal Water Supply and Wastewater Industries. Retrieved 20th November 2015 from <http://www.waterrf.org/PublicReportLibrary/4454.pdf>.

Were, F.H., Njue, W., Murungi, J., and Wanjau, R. (2008). Use of human nails as bio-indicators of heavy metals environmental exposure among school age children in Kenya. *Science of the Total Environment* 393(2-3):376-84.

Wilhelm, M., Pesch, B., Wittsiepe, J., Jakubis, P., Miskovic, P., Keegan, T., Nieuwenhuijsen, M.J., and Ranft, U. (2005). Comparison of arsenic levels in fingernails with urinary As species as biomarkers of arsenic exposure in residents living close to a coal-burning power plant in Prievidza District, Slovakia. *Journal of Exposure Science and Environmental Epidemiology* 15(1): 89-98.

World Health Organization (WHO), (1997). Guidelines for drinking-water quality. WHO, Geneva. Retrieved 18th April 2013 from http://www.who.int/water_sanitation_health/dwq/gdwqvol32ed.pdf; 1997.

World Health Organization (WHO), (2008) Geneva: Guidelines for drinking-water quality Third edition incorporating the first and second addenda, Volume 1 World Health Organization (WHO), 2008. Retrieved 19th May 2013 from http://www.who.int/water_sanitation_health/dwq/gdwq0506.pdf.

World Health Organization (WHO), (2008). Advocacy, communication and social mobilization for TB control: a guide to developing knowledge, attitude and

practice surveys. WHO/HTM/STB/2008.46. Retrieved 17th January 2014 from http://apps.who.int/iris/bitstream/10665/43790/1/9789241596176_eng.pdf.

World Health Organization (WHO), (2013). United Nations children's fund. Progress on sanitation and drinking-water: 2013 update WHO&UNICEF, Geneva. Retrieved 13th March 2016 from http://apps.who.int/iris/bitstream/10665/81245/1/9789241505390_eng.pdf

Xu, P., Huang, S., Wang, Za., and Lagos, G. (2006). Daily Intakes of Copper, Zinc and Arsenic in Drinking Water by Population of Shanghai, China. *Science of the Total Environment* 362: 50– 55.

Yalcin, M.G. (2009). Cluster analysis approach to identify metal sources in bottom sediments of Sultan Marsh canals, Turkey. *International Journal of Environment and Health* 3(1): 106-125.

Yap, C.K., Yusoff, H., Edward, F.B., and Tan, S.G. (2008). Comparison of heavy metal concentrations (Cd, Cu, Fe, Ni and Zn) in the shells and different soft tissues of *Anadara granosa* collected from Jeram, Kuala Juru and Kuala Kurau, Peninsular Malaysia. *Pertanika Journal of Tropical Agricultural Science* 31(2): 205-215.

Yen, T.P., and Rohasliney, H. (2013). Status of water quality subject to sand mining in the Kelantan River, Kelantan. *Tropical life sciences research* 24(1): 19–34.

Yi, Y., Yang, Z., and Zhang, S. (2011). Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze River basin. *Environmental Pollution* 159(10): 2575-2585.

Yunus, K., Suhaimi, M., Zahir, M., John, A., Khan Chowdhury, A. J., Saad, S., Al-Barwani, S.M. and Goddard, J. S. (2011). Bioaccumulation of some metals by Green mussel *Perna viridis* (Linnaeus 1758) from Pekan, Pahang, Malaysia. *International Journal of Biological Chemistry* 5(1): 54-60.

Yusoff, W.A., Jaafar, M., Kamarudin, M.K.A., and Toriman, M.E. (2015). Land Exploration Study and Water Quality Changes in Tanah Tinggi Lojing, Kelantan, Malaysia. *Malaysian Journal of Analytical Sciences* 19(5): 951-959.

Zafar, M. (2014). Knowledge and attitude towards and preventive practices relating to diarrhea among mothers under five years of children: findings of a cross-sectional study in Karachi. *Pakistan Journal Infectious Disease Therapy* 2:2-7

Zamani, A.A., Yaftian, M.R., and Parizanganeh, A. (2012). Multivariate statistical assessment of heavy metal pollution sources of groundwater around a lead and zinc plant. *Iranian journal of environmental health science & engineering*, 9(1), 1.

Zietz, B.P., Dieter, H.H., Lakomek, M., Schneider, H., Keßler-Gaedtke, B., and Dunkelberg, H. (2003). Epidemiological investigation on chronic copper toxicity to children exposed via the public drinking water supply. *Science of the Total Environment* 302(1): 127-144.

Żukowska, J., and Biziuk, M. (2008). Methodological evaluation of method for dietary heavy metal intake. *Journal of food science* 73(2): 21-29.

APPENDIX G: Descriptive statistics of dietary intakes (g)

Food	Minimum	Maximum	Mean	Std. Deviation
rice (g)	68.00	612.00	342.0400	135.67044
sandwich bread (g)	.00	136.00	32.5421	19.93445
eggs (g)	.00	64.00	21.2345	13.61215
fish (g)	.00	128.00	49.2485	26.41287
freshwater fish (g)	.00	84.00	7.2080	10.66382
cockles (g)	.00	28.57	10.5003	10.29016
squid (g)	.00	59.43	18.7089	15.50744
prawn (g)	.00	171.43	45.1552	38.16068
green leafy vegetables (g)	.00	240.00	169.4476	64.40182
tuber vegetables (g)	.00	273.00	19.8233	38.52695
manggo (g)	.00	140.00	24.4370	18.71771
guava (g)	.00	320.00	49.8936	44.00883
plain water (ml)	750.000	4500.000	1988.33333	591.704870
tea (ml)	.000	1000.000	167.14193	218.972757
coffee (ml)	.000	400.000	89.28600	85.248785
chocolate drink (ml)	.000	200.000	64.76093	70.315665
fruit juice (ml)	.000	71.430	9.03773	17.658475

BIODATA OF STUDENT

Nurul Hafiza Binti Ab Razak was born on 30th December 1988 in Tanah Merah, Kelantan. She obtained her primary education at SK Nik Daud and followed her secondary education at SMK Zainab 1, Kota Bharu and MRSM Kuala Berang. She furthered her study at Kolej Matrikulasi Kedah from 2006 until 2007. She joined Universiti Malaysia Terengganu in the year 2007, as a Bachelor Science of Analytical and Environmental Chemistry student and graduated in the year of 2010. In December 2010, she pursued her study at the Master level in Environmental Science area in Faculty of Environmental Studies, Universiti Putra Malaysia and graduated in the year 2012. Then, she further her study at PhD level in February 2013 under supervision of Dr Sarva Mangala Praveena. During her Master Degree, she has been doing a research on the environmental science area for her final year project entitled—Temporal flood incidence forecasting for Segamat River (Malaysia) using auto-regressive integrated moving average (ARIMA) modelling. In contrast, currently, she was doing research in environmental health area entitled — Toenail Metal Concentration as Biomarker of Exposure to Heavy Metal in Drinking Water of the Pasir Mas District, Malaysia. Throughout her postgraduate life, she has become a committee member in International Conference of Environmental and Occupational Health (ICEOH) 2014, a teaching assistant and demonstrator for Environmental Health Principle (EOH 4101) subject and a demonstrator for student mobility programme onbond Walailak University, Thailand. In addition, she also has attended few seminars namely Springer SEA eBooks Publishing Seminar 2013 and Thomson Reuters Seminar 2013 that was held in UPM. In presenting her research in postgraduate life, she also has been presented an oral presentation in International Conference on Environmental Forensics (iENFORCE) 2013, International Conference of Environmental and Occupational Health (ICEOH) 2014 and ICEOH 2016. During the iENFORCE conference, she was presenting a research entitled — Health Risk Assessment of Heavy Metal Exposure to Classroom Dust in Primary School, Serdang (Malaysia) on 12th to 13th Nov 2013 at Marriot Hotel, Putrajaya. During the ICEOH conferences, she was presenting apart of findings of her postgraduate research entitled — Toenail as a Biomarker of Heavy Metal Exposure Via Drinking Water: A Systematic Review on 8th to 9th April 2014 and — Possible Sources of Heavy Metal in Drinking Water and Exposure of Heavy Metal through Drinking Water Intake Among Pasir Mas Population on 11th to 13th April 2016 at Marriot Hotel, Putrajaya. Nevertheless, she has published two review papers that related to her research entitled — Toenail as a biomarker of heavy metal exposure via drinking water: a systematic review and — Drinking water studies: A review on heavy metal, application of biomarker and health risk assessment (a special focus in Malaysia). Besides, she has also published an article entitled — Quality of Kelantan drinking water and knowledge, attitude and practice among the population of Pasir Mas, Malaysia. Currently, she work at Victoria International College as Head of Programme Diploma in Environmental Health.

LIST OF PUBLICATIONS

Published

Ab Razak, N.H., Praveena, S. M., & Hashim, Z. (2014). Toenail as a biomarker of heavy metal exposure via drinking water: a systematic review. *Reviews on environmental health*. 30(1): 1-7. (Scopus)

Ab Razak, N. H., Praveena, S. M., Aris, A. Z., & Hashim, Z. (2015). Drinking water studies: A review on heavy metal, application of biomarker and health risk assessment (a special focus in Malaysia). *Journal of epidemiology and global health*. 5(4): 297-310. (Scopus)

Ab Razak, N. H., Praveena, S. M., Aris, A. Z., & Hashim, Z. (2015). Quality of Kelantan drinking water and knowledge, attitude and practice among the population of Pasir Mas, Malaysia. *Public health*. 131: 103-111. (Scopus, ISI-WoS, 5- year IF: 1.623, Q2 in Public, Environmental and Occupational Health)

Under review

Investigation of the Possible Sources of Heavy Metal in Drinking Water and Biomonitoring of Heavy Metal Exposure through Drinking Water among Pasir Mas District Population. Submitted to *Chemosphere*.

Proceedings

International Conference on Environmental Forensics (iENFORCE 2013). 12-13 Nov 2013. Putrajaya Marriott Hotel, Malaysia. Praveena, S. M., Mutalib, S. A., **Razak, N. H.**, Abidin, E. Z., & Aris, A. Z. (2014). Health Risk Assessment of Heavy Metal Exposure to Classroom Dust in Primary School, Serdang (Malaysia). In *From Sources to Solution* (pp. 83-87). Springer Singapore. **(Oral presentation)**.

International Conference on Environmental and Occupational Health (ICEOH 2014). 8-9 Apr 2014. Putrajaya Marriott Hotel, Malaysia. **Ab Razak, N. H.**, Praveena, S. M., Aris, A. Z., & Hashim, Z. (2015). Toenail as a biomarker of heavy metal exposure via drinking water: a systematic review. **(Oral presentation)**.

Poster Symposium Faculty Excellence Month 2015. 26 Aug 2015. Faculty of Medicine & Health Sciences, UPM. **Ab Razak, N. H.**, Praveena, S. M., Aris, A. Z., & Hashim, Z. (2015). Quality of Kelantan drinking water and knowledge, attitude and practice among the population of Pasir Mas, Malaysia. **(Poster Presentation)**.

International Conference on Environmental and Occupational Health (ICEOH 2016). 11-13 Apr 2016. Putrajaya Marriott Hotel, Malaysia. **Ab Razak, N. H.**, Praveena, S. M., Aris, A. Z., & Hashim, Z. (2015). Possible Sources of Heavy Metal in Drinking Water and Exposure of Heavy Metal through Drinking Water Intake Among Pasir Mas Population. **(Oral presentation)**.





UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

TOENAIL METAL CONCENTRATION AS BIOMARKER OF EXPOSURE TO HEAVY METAL IN DRINKING WATER OF THE PASIR MAS DISTRICT, MALAYSIA

NAME OF STUDENT : NURUL HAFIZA BINTI AB RAZAK

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (v)

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]