



**UNIVERSITI PUTRA MALAYSIA**

***EFFECTIVENESS OF HEALTH EDUCATION IN IMPROVING  
KNOWLEDGE AND ATTITUDE TOWARDS TOXOPLASMOSIS AMONG  
PREGNANT WOMEN IN AL NAJAF, IRAQ***

**ATHEER KADHIM IBADI**

**FPSK(p) 2016 27**



**EFFECTIVENESS OF HEALTH EDUCATION IN IMPROVING  
KNOWLEDGE AND ATTITUDE TOWARDS TOXOPLASMOSIS AMONG  
PREGNANT WOMEN IN AL NAJAF, IRAQ**

**By**

**ATHEER KADHIM IBADI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**October 2016**

## **COPYRIGHT**

All material contained within the thesis, including without limitation text, logos, icons, photographs, and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright© Universiti Putra Malaysia



## **DEDICATION**

TO

Dedicated especially to my parents, my love, and partner of my life (my wife), To my dear daughter Fatemah the sense who make me possible to complete my study successfully



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Doctor of Philosophy

**EFFECTIVENESS OF HEALTH EDUCATION IN IMPROVING  
KNOWLEDGE AND ATTITUDE TOWARDS TOXOPLASMOSIS AMONG  
PREGNANT WOMEN IN AL NAJAF, IRAQ**

By

**ATHEER KADHIM IBADI**

**October 2016**

**Chairman : Titi Rahmawati Binti Hamedon, PhD**  
**Faculty : Medicine and Health Sciences, Department of Community Health**

The aim of this study is to determine the effect of intervention by health education on the knowledge and attitude on toxoplasmosis among pregnant women with toxoplasmosis in Al- Najaf Al- Ashraf – Iraq-2014. The study compared the scores of knowledge and attitudes between the baseline against the first and second posttests and within three stages and groups, and determine the association of these factors with their sociodemographic characteristics, both Experiment and Control groups. Intervention study design, and a simple random sampling technique was used to select the 340 respondents, who were patients from gynecological clinic from three hospitals. Data was collected from 1<sup>st</sup> June to 31<sup>st</sup> October 2015 using a structured pre-tested questionnaire in Arabic language and the response rate was 100 %. The results of this study showed that most of the respondents were housewife, young, had low level of education and live in urban area. At base line there was no significant statistical differences of score between both groups in terms of the overall knowledge on toxoplasmosis. However, the knowledge score of both groups became different statistically at first and second posttests. Findings on the attitude showed that both groups had positive attitude towards toxoplasmosis at baseline and first posttest. However, at second posttest the attitude of the Control Group became negative, whereas the attitude of the Experimental Group remained positive. There was statistically significant difference of the mean score of knowledge between both groups in all different stages of data collections. Repeated measurement using ANOVA with a Greenhouse-Geisser correction showed that the mean score according to all items of knowledge on toxoplasmosis infection were differed significantly within time and also differed significantly in the interaction between groups. The same test also showed that the mean score of attitudes on toxoplasmosis were significantly different within baseline, first posttest, and second posttest and also differed significantly within the time between groups. In conclusion, this study showed that the level of knowledge and attitude related to toxoplasmosis among the pregnant women infected with toxoplasmosis in Al-Najaf province is unsatisfactory at the baseline, but it became better after they were given health education on toxoplasmosis.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KEBERKESANAN PENDIDIKAN KESIHATAN DALAM MEMPERBAIKI  
PENGETAHUAN DAN SIKAP TENTANG TOXOPLASMOSIS DI  
KALANGAN WANITA MENGANDUNG DI AL NAJAF, IRAQ**

Oleh:

**ATHEER KADHIM IBADI**

**Oktober 2016**

**Pengerusi : Titi Rahmawati binti Hamedon, PhD**  
**Fakulti : Perubatan dan Sains Kesihatan**

Tujuan kajian ini adalah untuk menentukan kesan intervensi melalui pendidikan kesihatan ke atas pengetahuan dan sikap terhadap toxoplasmosis di kalangan wanita mengandung yang menghidap penyakit toxoplasmosis di Al-Najaf al-Ashraf - Iraq-2014. Kajian membandingkan skor pengetahuan dan sikap di peringkat awal, ujian pertama dan kedua pada dua kumpulan berbeza (Ujian dan Kawalan) dan untuk menentukan hubungan antara faktor sosiodemografi dengan pengetahuan dan sikap mereka terhadap toxoplasmosis. Dengan menggunakan rekabentuk kajian intervensi, seramai 340 orang peserta kajian telah dipilih secara rambang di klinik sakit puan dari tiga hospital. Data dikumpulkan mulai 1 Jun hingga 30 Oktober 2015 menggunakan borang soal selidik berstruktur yang telah di pra uji, menggunakan Bahasa Arab. Kadar respon adalah 100%. Hasil kajian menunjukkan kebanyakan peserta kajian adalah suri rumahtangga, berusia muda, berpendidikan rendah, serta tinggal di bandar. Di peringkat awal diadapati tiada perbezaan signifikan skor pengetahuan antara dua kumpulan tersebut. Skor tersebut menjadi berbeza secara signifikan pada ujian pertama dan ke dua. Walaubagaimanapun sikap peserta kajian pada ke dua kumpulan adalah positive di peringkat awal. Pada ujian ke dua sikap Kumpulan Kawalan menjadi negative sedangkan sikap kumpulan Kajian tetap positif. Didapati perbezaan statistik yang signifikan untuk purata skor pengetahuan pada ke dua kumpulan di dalam semua peringkat pengumpulan data. Ujian berulang menggunakan kaedah ANOVA dengan pembetulan Greenhouse-Geisser menunjukkan perbezaan skor tersebut masih berbeza secara signifikan mengikut waktu dan interaksi antara kumpulan. Untuk sikap, hasil yang sama juga telah diadapati dan ujian menggunakan kaedah yang sama menunjukkan purata skor sikap berbeza secara signifikan mengikut waktu dan interaksi antara kumpulan. Secara kesimpulan, kajian ini menunjukkan bahawa tahap pengetahuan dan sikap peserta kajian terhadap toxoplasmosis adalah tidak memuaskan, tetapi menjadi lebih baik setelah diberikan pendidikan tentang toxoplasmosis.

## ACKNOWLEDGEMENTS

First of all, I thank Allah for the Almighty on the great grace. Despite all the difficulties that it's faced me in my study, but his great mercy has enabled me to complete my study. I praise and thanks him a lot.

I would like to express my deep gratitude and appreciation for a lifetime to Dr. Titi Rahmawati Binti Hamedon for her wonderful advice, thoughtful guidance, continuous support and meaningful friendship throughout my Ph.D.'s degree. She encouraged me through the difficulties and inspired me to move to a higher level of the learning experience. Her positive attitude and support have helped me to become a better person.

I would also like to give my heartfelt thanks and deepest appreciation and gratitude to Dr. Anita Binti Abdul Rahman for her insightful suggestions and guidance, encouragement, patience, valuable advice that had helped me to carry on the study successfully.

I would like to express my special appreciation and very sincere gratitude to Dr. Suriani Binti Ismail. She gave her the time, effort, encouragement and valuable suggestions.

I would like to express my special appreciation and very sincere gratitude to Dr. Abdul Kareem Abdullah; he gave me the time, effort, encouragement and valuable suggestions.

They were exceptional role models in teaching, mentoring, and conducting research studies. Without their outstanding assistance and support, I would not have reached my goal.

Also, I would like to express a special thank you to Professor Dr. Bahaman Abu Sama for his exceptional advice, guidance, and assistance with the statistical aspects of my dissertation work.

I am so grateful to Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for allowing me to study.

I am so grateful to Ministry of higher education and scientific research, Foundation of Technical Education (Kufa Institute)- Iraq, to give me the opportunity to complete my Ph.D. study.

I also express appreciation for Al-Najaf Health Directorate\ Center of Training and Development of Staff to give us the permission to do this study.

I also express appreciation for volunteer women who have enrolled in the study and for the medical staff in the three hospitals in the study location.

I would like to extend a special thank you and deepest gratitude to my parents for their great kindness, devotion, encouragement and continuous support. I commend them efforts and them toleration. They will stay in my heart forever.

I would like to extend my deepest love and gratitude to my dearest my wife, and my beloved daughter Fatimah, for heartily encouragement and the unconditional support that they've offered through the long days stretching into months of my study.

I also wish to express my appreciation to all of my friends in Malaysia who've inspired me in all of these years. Thank you all.

*Atheer Kadhim Ibadi*



I certify that a Thesis Examination Committee has met on 10 October 2016 to conduct the final examination of Atheer Kadhim Ibadi on his thesis entitled "Effectiveness of Health Education in Improving Knowledge and Attitude Towards Toxoplasmosis among Pregnant Women in Al Najaf, Iraq" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Thesis Examination Committee were as follows:

**Sabrina binti Sukardi, PhD**

Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Chairman)

**Shamsul Bahri bin Hj. Mohd Tamrin, PhD**


Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Abdah binti Md Akim, PhD**

Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Sathaporn Jittapalapong, PhD**

Professor  
Kasetsart University  
Thailand  
(External Examiner)



---

**NOR AINI AB. SHUKOR, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 27 December 2016

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Titi Rahmawati Binti Hamedon, MD, M.**

Senior Lecturer (Medical)  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Chairman)

**Anita Binti Abdul Rahman, MD, M.**

Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Member)

**Suriani Binti Ismail, MD, M.**

Senior Lecturer (Medical)  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Member)

**Abdul Kareem Abdullah, PhD**

Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Member)

---

**ROBIAH BINTI YUNUS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:

## Declaration by graduate student

I hereby confirm that:

- This thesis is my original work;
- Quotations, illustrations and citations have been duly referenced;
- This thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- Intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- Written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before the thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- There is no plagiarism or data falsification/ fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism software.

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Name and Matric No.: Atheer Kadhim Ibadi, GS38376

## Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature:  
Name of  
Chairman of  
Supervisory  
Committee:

\_\_\_\_\_  
Titi Rahmawati Binti Hamedon, MD, M.

Signature:  
Name of  
Member of  
Supervisory  
Committee:

\_\_\_\_\_  
Anita Binti Abdul Rahman, MD, M.

Signature:  
Name of  
Member of  
Supervisory  
Committee:

\_\_\_\_\_  
Suriani Binti Ismail, MD, M.

Signature:  
Name of  
Member of  
Supervisory  
Committee:

\_\_\_\_\_  
Abdul Kareem Abdullah, PhD

## TABLE OF CONTENTS

	<b>Page</b>
<b>ABSTRACT</b>	i
<b>ABSTRAK</b>	ii
<b>ACKNOWLEDGEMENTS</b>	iii
<b>APPROVAL</b>	v
<b>DECLARATION</b>	vii
<b>LIST OF TABLES</b>	xiv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	
1.1 Background	1
1.2 Problem Statement	3
1.3 Study Justification	4
1.4 Objectives of the Study	5
1.4.1 General Objectives	5
1.4.2 Specific Objective	5
1.5 Study Hypothesis	5
1.6 Conceptual Framework	5
1.7 Definitions of Terms	7
<b>2 LITERATURE REVIEW</b>	
2.1 Introduction in Toxoplasmosis	
2.1.1 History of Toxoplasmosis	9
2.1.2 Classification of Toxoplasmosis	9
2.1.3 Mode Of Toxoplasmosis Transmission	9
2.1.4 T. Gondii Life Cycle	9
2.1.5 Toxoplasmosis in Pregnant Women	12
2.2 Epidemiology	13
2.2.1 Toxoplasmosis in The World	13
2.2.2 Toxoplasmosis in Arab Countries	12
2.2.3 Toxoplasmosis in Iraq	15
2.3 Pathogenesis	15
2.4 Congenital Toxoplasmosis	19
2.5 Diagnosis of Toxoplasmosis	20
2.5.1 Direct Microscope Examination Methods	21
2.5.1.1 Animal Inoculation and Cell Culture	21
2.5.1.2 Histological Diagnosis	21
2.5.1.3 Isolation of T. Gondii	21
2.5.2 Indirect Methods	21
2.5.2.1 Skin Test (Delayed Hypersensitivity Test)	21
2.5.2.2 Serological Tests	22
2.5.2.3 Sabin-Feldman Dye Test (DT)	22
2.5.2.4 Indirect Fluorescent Antibodies Test (IFAT)	22
2.5.2.5 Direct Agglutination Test (DAT)	22

2.5.2.6	Indirect Hemagglutination Test (IHAT)	22
2.5.2.7	Latex Agglutination Test (LAT)	22
2.5.2.8	Dipstick Dye Immunoassay (DDIA)	23
2.5.2.9	Enzyme-Linked Immunosorbent Assay (ELISA)	23
2.5.2.10	Enzyme-linked Fluorescent Assay (ELFA)	23
2.5.2.11	IgM-Immunosorbent Agglutination Assay (IgM-ISAGA)	23
2.5.2.12	IgG Avidity Test	23
2.5.2.13	Molecular Diagnosis	23
2.6	Treatment of Toxoplasmosis	24
2.7	Prevention of Toxoplasmosis	24
2.8	Risk Factors of Toxoplasmosis	26
2.8.1	Sociodemographic Characteristics	26
2.8.2	Geographical Distribution	26
2.8.3	Past Obstetric History	26
2.8.4	Past Medical History	27
2.8.5	Housing, Environmental And Location Conditions	27
2.8.6	History Immunization of Toxoplasmosis	27
2.9	Knowledge Toward Toxoplasmosis	27
2.10	Attitudes Toward Toxoplasmosis	29
2.11	Intervention Studies And Toxoplasmosis	30
<b>3</b>	<b>MATERIALS AND METHODS</b>	
3.1	Ethical Consideration	33
3.2	Study Location	33
3.3	Design Of Study	35
3.4	Sampling	35
3.4.1	Study Population	35
3.4.2	Sample of Study	35
3.4.3	Selection Criteria	35
3.4.3.1	Inclusion Criteria	35
3.4.3.2	Exclusion Criteria	35
3.4.4	Sampling Frame	36
3.4.5	Sample Selection	36
3.4.6	Randomization and Blinding Procedure	36
3.4.7	Sample Size	38
3.4.7.1	Calculation of Sample Size	38
3.5	Study Intervention	39
3.6	Instrument	40
3.6.1	Knowledge, Attitudes and Practices Related to Toxoplasmosis	43
3.6.1.1	Knowledge on Toxoplasmosis	43
3.6.1.2	Attitudes on Toxoplasmosis	43
3.7	Validity and Reliability of Questionnaire	43
3.7.1	Pre-Testing	43
3.7.1.1	Reliability Results for Pre-Test	43
3.7.2	Reliability Results for Current Study	44
3.7.3	Validity of Contents	44

3.8	Data Collection	44
3.9	Analysis of Data	46
3.10	Strengths and limitations of the study	46
<b>4</b>	<b>RESULTS</b>	
4.1	Response Rate	48
4.2	Results related to independent variables	48
4.2.1	Socio-Demographic Characteristics of Respondents	49
4.2.2	Housing, Environment, and Location Conditions	51
4.2.3	Past Obstetric History	52
4.2.4	Pet Ownership for Respondents	53
4.2.5	Cooking Preferences and Meat Consumption	54
4.3	Knowledge and attitudes score at baseline, first and second posttest	55
4.3.1	At Baseline	55
4.3.1.1	Baseline Scores of Knowledge on Toxoplasmosis Infection	55
4.3.1.2	Source of Information About Toxoplasmosis	57
4.3.1.3	Description of Attitudes Score of Respondents Toward Toxoplasmosis at Baseline	58
4.3.2	At First Post test Results	59
4.3.2.1	The Results of knowledge Toward Toxoplasmosis	59
4.3.2.2	Attitudes Scores of Respondents Toward Toxoplasmosis	62
4.3.3	At Second Post test	64
4.3.3.1	Knowledge Score of Respondents on The Toxoplasmosis Infection at Second Post test	64
4.3.3.2	Attitudes scores of respondents Toward Toxoplasmosis at Second Post test	67
4.4	Comparison of knowledge and attitudes score within three stages and between groups	68
4.4.1	Knowledge Scores Differences	69
4.4.2	Attitudes Score Differences on Toxoplasmosis Infection	70
4.5	Association between the score of knowledge and attitude for three stages for both groups with sociodemographic characteristics	72
4.5.1	Association of Score of Knowledge and Attitudes with Sociodemographic Characteristics	73
4.5.1.1	At Baseline	73
4.5.1.2	At the First Post test	75
4.5.1.3	At the Second Post test	77
4.5.1.4	Summary on Association of Knowledge and Attitudes with Sociodemographic Characteristics	79
4.6	Summarizing of the results	79

## 5 DISCUSSION

5.1	Sociodemographic Characteristics of Respondents	81
5.1.1	Age	81
5.1.2	Monthly income	82
5.1.3	Level of education	83
5.1.4	Occupation	83
5.1.5	Living location	84
5.2	Knowledge on Toxoplasmosis	85
5.2.1	Score of Knowledge at Baseline Level	85
5.2.1.1	Overall Knowledge Score	86
5.2.1.2	The Level of Knowledge Scores on Animals That Transmit of Toxoplasmosis	86
5.2.1.3	The Score of Knowledge on The Commonest Signs and Symptoms of Toxoplasmosis	86
5.2.1.4	The Score of Knowledge on Serious Complications of Toxoplasmosis	87
5.2.1.5	Knowledge on Toxoplasmosis Prevention	88
5.2.1.6	Source of Information	88
5.2.1.7	Knowledge on Source of Toxoplasmosis Infection	89
5.2.2	Scores of Knowledge at The First Post test	89
5.2.2.1	The Score of Knowledge About the Animals That Transmit of Toxoplasmosis	90
5.2.2.2	The Score of Knowledge on The Commonest Signs and Symptoms of Toxoplasmosis	90
5.2.2.3	According to The Serious Complications of Toxoplasmosis	91
5.2.2.4	The Score of Knowledge on The Methods of Toxoplasmosis Prevention	91
5.2.2.5	Knowledge on Source of Toxoplasmosis Infection	92
5.2.3	Score of Knowledge at The Second Post test	92
5.2.3.1	The Score of Knowledge About the Animals That Transmit of Toxoplasmosis	93
5.2.3.2	The Score of Knowledge on The Commonest Signs and Symptoms of Toxoplasmosis	93
5.2.3.3	According to The Serious Complications of Toxoplasmosis	94
5.2.3.4	The Score of Knowledge on The Methods of Toxoplasmosis Prevention	94
5.2.3.5	Knowledge on Source of Toxoplasmosis Infection	95
5.3	Differences in knowledge Among the Three Phases of Study	96
5.4	Difference of Attitudes and Practices Toward Toxoplasmosis Between Groups	99
5.4.1	At Baseline	99



5.4.2	At the First Post test	99
5.4.3	At the Second Post test	99
5.5	Differences in Attitudes and Practices Toward Toxoplasmosis Among the Three Phases of Study	100
5.6	The Relationship Between Knowledge, Attitudes, and Practices with The Sociodemographic Characteristics	101
<b>6</b>	<b>SUMMARY AND CONCLUSION</b>	
6.1	Acceptance and Rejection of the study hypothesis	106
6.2	Summary and Conclusion	106
6.3	Limitations and Strength of the study	107
6.4	Recommendations and Further studies	108
	<b>REFERENCES</b>	109
	<b>APPENDICES</b>	137
	<b>BIODATA OF STUDENT</b>	186
	<b>LIST OF PUBLICATIONS</b>	187

## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1.1	Prevalence of toxoplasmosis among women of childbearing age in selected continents	1
2.1	Studies on health education intervention about toxoplasma infection in pregnancy in selected countries	25
2.2	Different intervention studies in selected countries	32
3.1	Pre-testing reliability test result	41
3.2	Reliability test results for this study	44
4.1	Comparing the sociodemographic characteristics of respondent (Experiment and Control)	49
4.2	Comparison of the response of respondents according to housing conditions	51
4.3	Distribution of respondents who have a past obstetric history	52
4.4	Distribution of respondents according to their parity	52
4.5	Distribution of respondents according to animal's ownership	53
4.6	Distribution of respondents according to types of food and its preferences	54
4.7	Comparison the level of attitudes of respondents according to the types of food and its preferences	54
4.8	Knowledge score of respondents who had a correct answer about toxoplasmosis infection	56
4.9	Distribution of perception scores of respondents about toxoplasmosis infection	58
4.10	Comparing the levels of attitudes score of respondents on toxoplasmosis infection at baseline	59
4.11	Knowledge score of respondents who had a correct answer about toxoplasmosis infection at first post test	60
4.12	Comparing the score of knowledge on toxoplasmosis at first post test	61
4.13	Distribution of perception score of respondents about toxoplasmosis infection at first post test	63
4.14	Comparing the level attitudes score of respondents about toxoplasmosis infection at first post test	63

4.15	knowledge scores of respondents who had a correct answer about toxoplasmosis infection at second post test	65
4.16	Comparing the knowledge score of respondents on toxoplasmosis infection at second post test	66
4.17	Distribution of perception scores of respondents on toxoplasmosis infection at second post test	67
4.18	Comparing the levels of attitudes scores of respondents on toxoplasmosis infection at second post test	68
4.19	Comparison the knowledge score in baseline stage, after intervention and after three months from intervention within the Experiment and Control groups	69
4.20	Comparing the knowledge scores on toxoplasmosis between the Experiment and Control groups	70
4.21	Attitudes score of respondents for three stages of study on toxoplasmosis infection	71
4.22	Comparison the attitudes score in baseline stage, after intervention and after three months from intervention within the Experiment and control groups	72
4.23	Comparing the attitudes score on toxoplasmosis between the Experiment and Control groups	72
4.24	Summary the association of the score of knowledge and attitudes with sociodemographic characteristics	79
4.25	Correlation regression among overall knowledge and attitudes scores and sociodemographic characteristics between the Experiment and Control groups	79

## LIST OF FIGURES

Figure		Page
1.1	Conceptual framework of this study	6
2.1	Major routes of transmission of <i>T. gondii</i>	11
2.2	Life cycle of <i>T. gondii</i>	12
2.3	Prevalence of toxoplasmosis in Iraq from 2000 to 2012	16
2.4	Prevalence of toxoplasmosis in Al-Najaf according to the real time of work	17
2.5	Prevalence of toxoplasmosis according to the geographical distribution for years 2003-2012 from different studies	18
2.6	Number of toxoplasmosis cases in Iraq for years 1989-2001	19
3.1	Map of the study location	34
3.2	Study flow chart of Experiment and Control groups	37
3.3	Flow chart development of the questionnaire of this study	41
3.4	Flow chart of data collection process for the study of both groups	45
4.1	Distribution of respondents according to the source of respondent's information about toxoplasmosis	57
4.2	Scatter plot of correlation between the overall knowledge score and overall attitudes score of Experiment at baseline	73
4.3	Scatter plot of correlation between the overall knowledge score and overall attitudes score of Control group at baseline	74

## LIST OF ABBREVIATIONS

KAP	Knowledge, Attitude, and Practice
AIDS	Acquired immuno Deficiency Syndrome
LAT	Latex Agglutination Test
DLA	Direct Latex Agglutination
ELA	Enzyme Immuno Assay
DDIA	Dipstick Dye Immuno Assay
ELISA	Enzyme Linked Immuno Sorbent Assay
IgG	Immunoglobulin G
IgM	Immunoglobulin M
Kg	Kilogram
HIV	Human Immuno Viruses
CSF	Cerebral Spinal Fluid
AbS	Antibodies
DAT	Direct Agglutination Test
T. gondii	Toxoplasma gondii
IHAT	Indirect Haemagglutination Test
AF	Amniotic Fluid
CMI	Cell Mediated Immunity
IFAT	Indirect Fluorescent Antibodies
RH	Rimbunan Hijau Group
TSA	Tachyzoites soluble antigen
ELFA	Enzyme -Linked Fluorescent
ISAGA	Immuno Sorbent Agglutination Assay
PCR	Polymerase Chain Reaction
DNA	Deoxyribo Nucleic Acid
WHO	World Health Organization
UPM	University Putra Malaysia

US	United Nation of America
VIDAS	Routine batch or random access testing for serology, immunochemistry, antigen detection and immunohemostasis.
DT	Sabin-Feldman Dye Test
IFAT	Indirect Immuno-Fluorescent Antibody Test
n	Sample size
Ho	Null hypothesis
Ha	Alternative Hypothesis
ANOVA	Analysis of Variance
IQD	Iraqi Dinar
US	The United State of America
RM	Malaysian Ringet
Abs	Antibodies

# CHAPTER 1

## INTRODUCTION

### 1.1 Background

Toxoplasmosis is defined as a parasitic disease caused by an intracellular protozoan called *Toxoplasma gondii* (Jones et al., 2001). This parasite infect human and most of warm blooded animals genus, but cat are considered as the essential host. The dangerous implications of this disease can not only affect pregnant women but it can also have severe consequences on fetuses. The transmission rate of this disease to the fetus ranges from 10-15% at the first trimester of pregnancy and may reach to 68% in the third trimester of pregnancy (Remington, Thulliez, & Montoya, 2004).

Toxoplasmosis is the third leading cause of infectious disease in the US after salmonellosis and listeriosis (Dubey & Jones, 2008). The infection has a worldwide distribution. Onethird of all human beings have been exposed to this parasite. However, the seroprevalence of this disease varies considerably between countries, from less than 10% to more than 90% (Pawlowski et al., 2001). Toxoplasmosis infection occurs worldwide even though the rates of infection differs substantially geographically. A survey which was conducted on women of childbearing age from 44 countries and which included 99 studies found the areas with high prevalence of toxoplasmosis (Pappas, Roussos & Falagas, 2009). The findings are shown in Table 1 below.

**Table 1.1: Prevalence of toxoplasmosis among women of childbearing age in selected continents**

<i>Place</i>	<i>Prevalence (%)</i>
<i>Within Latin America</i>	50–80
<i>Parts of Eastern and Central Europe</i>	20–60
<i>The Middle East</i>	30-50
<i>Parts of Southeast Asia</i>	20–60
<i>Parts of Africa</i>	20–55

(Pappas, Roussos & Falagas, 2009)

In other studies, the prevalence of disease and risk factors of transmission of toxoplasmosis infection varies substantially between countries (Abu-Madi, Al-Molawi & Behnke, 2008).

In Iraq, many studies were done concerning that seroprevalence of toxoplasmosis by using different diagnostic techniques in various regions ( Al-Kalaby, 2008; Khalil, 2008; and Al- Mousawi, 2008). These studies found that the incidence rate of

toxoplasmosis among women who had abortion in Najaf, Baghdad, and Basrah were 31.9%, 25%, and 26.1% respectively (Taher, 2011). A study was conducted in 2007 in Al-Najaf to determine the prevalence of *Toxoplasma gondii* among female in the age group of 16-26 years old. This study showed that the highest prevalence of the infection was highest, 68%, among those respondents in the age group 25-26 years (Al-Nahi & Al-Abbas, 2007).

Toxoplasmosis can be avoided by giving health information about the source of infection to those who are at risk, especially pregnant women. This will encourage them to change their behavior, and thereby reducing the probability to acquire the infection during pregnancy (Hall, Ryan & Buxton, 2001). However, few studies have examined the effectiveness of health education despite numerous pleas in the published literature for a stronger focus on primary prevention of toxoplasma infection among pregnant women. Existing guidelines for such care are also lacking (Jones, et al, 2003; Conyn-van Spaendonck & Van Knapen, 1992; Baril, et al, 1999; Foulon, Naessens & Ho-Yen, 2000).

There are not many report available that address the effectiveness of health education in reducing toxoplasmosis. One exception is a 1994 survey of 196 health districts in the United Kingdom which demonstrated that health education was offered in approximately half of the health units surveyed. However there were serious deficiencies found in the monitoring to see whether information was given to all women (Newton and Hall., 1994). In France, primary prevention for toxoplasmosis was also recommended, but its practices were not assessed or evaluated (Ancelle, et al., 1996). One case control study of risk factors for toxoplasmosis seroconversion in pregnant women showed that controls were more likely to have received documentary advice on prevention than cases (Baril, et al., 1999).

Health care providers should make preconception, prenatal and natal investigations and health education to prevent toxoplasmosis as standard of care for pregnant women. Educational materials that contain messages on how to prevent pregnant women from becoming infected have resulted in reducing rates of seroconversion (Gollub et al., 2008). Effective prevention of congenital toxoplasmosis depends on avoidance of infection during pregnancy (Lebech, et al., 1999).

There is no much published work found on the information about the frequency of preventive practice behaviors on toxoplasmosis infection among pregnant women. While knowledge is a crucial determinant to establish behavioral change, precise knowledge may not lead to appropriate preventive behavior. Attitudes of pregnant women towards changing their behavior and their perception about the likelihood of contracting the infectious disease during their pregnancy may also be important contributors to establish behavioral change (Pereboom et al., 2013).



## 1.2 Problem statement

Toxoplasmosis is an important public health problem. This disease is responsible for substantial rate of neonatal morbidity and mortality, particularly in congenitally infected and immuno-compromised individuals (Tenter, Heckerth & Weiss, 2000; Luft, et al., 1993). In Iraq 2011, women with previous history of abortion and those with abnormal pregnancies had the highest prevalence of toxoplasmosis (57.1%), and those women are within the age range of 26 and 30 years old. One of the reasons for the high prevalence is related to female handling raw meat more frequently than male, and they spend more time cooking at home (Mohammed, Ahmed & Hussain, 2013). There were many studies conducted on toxoplasmosis among pregnant women in different regions in Iraq (Mohammed, 2011; Mohammad, Ahmed, & Hussain, 2013; Al- Mousawi, 2008). For example, in Tikrit, the prevalence of toxoplasmosis among pregnant women who attended gynecological clinics were about 49-95% (Al-Doori, 2010). In Thi-Qar, the figure was 50% (Hadi, 2011). Many studies were carried out to determine the seroprevalence of toxoplasmosis by using different diagnostic techniques among pregnant women in various regions of Iraq (Al-Kalaby, 2008; Khalil, 2008; Al- Mousawi, 2008). These studies found the incidence rate of toxoplasmosis among women with previous history of abortion in Najaf, Baghdad, and Basrah were 31.9%, 25%, and 26.1% respectively (Taher, 2011).

In 2007, the highest prevalence of *Toxoplasma gondii* infection detected using latex agglutination test (LAT) technique was recorded among female (pregnant and non-pregnant women) in the age group of 25-26 year (68 %) (Al-Nahi & Al-Abbas, 2007). Two years later in Al-Abbasiya Najaf, the same technique showed a prevalence of 43.7% positive cases of toxoplasmosis among respondent in the age group of 18-27 years old (Hussain, Yousif & Nassir, 2010). In Baghdad province in the same year, about 33.3% of women who had abortion found to have infected with toxoplasmosis (Al-Garawia, Al-Fartusie & Al-Bairmani, 2012). In 2013 in Kirkuk city there was 7.2% of pregnant women who had abortion twice were infected with toxoplasmosis (Salman, 2014). In 2011 in Al-Najaf province, 35% of newborns were infected with congenital toxoplasmosis (Al-haris, Saheb & Abdul-Sada, 2015). Most of the infection occurred among young pregnant women of 26-30-year-old group (Al-Nahi & Al-Abbas, 2007; Mohammed, Ahmed & Hussain, 2013). Also, in the same place in 2009, a study found that there was 48% of patient's children age less than six years (pre-school age) positive with IgG anti-toxoplasma antibodies (Taher, 2011). Although there were many studies conducted in Iraq generally and in Al-Najaf particularly to diagnose toxoplasmosis infection, there were none that looked into the knowledge and attitudes of pregnant women toward toxoplasmosis.

Globally, it is estimated that about one-third of the world's population is infected with toxoplasmosis (Pappas et al., 2009). In the Netherlands, the incidence rate of congenital toxoplasmosis is two children per 1000 live births, which is ten times higher than those in Denmark and twenty times higher than those in Ireland (Ross, Jones & Lynch, 2006). High prevalence of toxoplasmosis infection has been reported among pregnant women and women of reproductive age from different areas around the world including the in the Middle East (Pappas et al., 2009). The prevalence of toxoplasmosis among the pregnant women who attended to the gynecological clinic in

Saudi Arabia-Jazan was 24.1% in 2014 (Aqeely, et al., 2014), 31.6% in Jordan (Jumaian, 2005), 35.1% in Qatar, Abu-Madi et. al., (2010) and 55% in Lebanon Beirut Bouhamdan, et al., (2010). The prevalence of such infection was 34.6% in the United Arab Emirates (Abu-Zeid, 2002).

### 1.3 Study justification

Pregnant women are exposed to various health risks during pregnancy. The global maternal mortality from the year 1990 to 2015 dropped by about 44% (WHO, UNICEF & UNFPA, 2012). However, the maternal mortality is still higher in the poorer communities. The large gap in the number of maternal deaths in some regions in the world illustrates inequities of health care services and highlights the differences between the rich and poor. In developing countries, the maternal mortality rate is 239 for every 100 000 live births in 2015, whereas, in developed countries it is 12 per 100 000 live births (WHO, 2015).

Among the serious complications of toxoplasmosis during pregnancy are fetal death and stillbirths (Moncada & Montoya, 2012). Stillbirths that are caused by toxoplasmosis infections are more commonly occurring in developing than in developed countries (Goldenberg & Thompson, 2003). The knowledge about the epidemiology of *T. gondii* infection in women who had stillbirths, miscarriages and abnormal pregnancy are still poor (Adesiyun, et al., 2007).

In Iraq, health problems during with pregnancy and deaths associated with childbirth have increased. The estimation of maternal mortality for 2013 was 84 per 100 000 live births; the neonatal deaths rate was 19 per 100 000 live births (UNICEF, 2013). According to the Iraqi Ministry of Health in 2000, 24.3% of registered newborn babies had birth weight that is less than 2.5 kg (Wells et al., 2011). About 15.3% of women with history of abortion in Baghdad in 2013 were infected with toxoplasmosis (Hussan, 2013). The rate of toxoplasmosis infection among those with abnormal pregnancy was 2% in Kirkuk city in 2012 (Aljumaily & Alsamarai, 2013). In the same city in 2014, there was 26.7% of women with bad obstetric history infected with toxoplasmosis, 94.1% of them had a stillbirth and 74.1% of them had a miscarriage (Mohammad & Salman, 2014). In 2009, there were 55% of pregnant women who were infected with toxoplasmosis in Erbil Governorate and majority of them were illiterate (Hamad & Kadir, 2014). One study conducted in Kirkuk city in 2012 found that 21% of pregnant women were positive for IgG of toxoplasmosis antibodies, and most of them were not educated about toxoplasmosis (Aljumaily & Alsamarai, 2013). Many studies have been conducted to detect of toxoplasmosis in Iraq using different techniques. However, there were no studies that looked at the knowledge and attitudes related to toxoplasmosis.

Hence, establishing the database about the knowledge and attitudes related to toxoplasmosis is considered to be very crucial in order to plan for making prevention programs. The prevention programs will, hopefully reduce the maternity risks of developing toxoplasmosis and increase their quality life. Health education, if done regularly will increase their knowledge and awareness about the disease, and

eventually will reduce the incidence of getting toxoplasmosis.

## **1.4 Objectives of the study**

### **1.4.1 General Objectives**

The main purpose of this study is to determine the effect of health education intervention on the knowledge and attitudes related to toxoplasmosis among pregnant women with toxoplasmosis in Al- Najaf Al- Ashraf – Iraq.

### **1.4.2 Specific Objective**

- 1- To describe the socio-demographic characteristics of respondents.
- 2- To determine the knowledge and attitudes scores on toxoplasmosis among the respondents at different stages - at baseline stage, after there were given health education intervention and at three months after the intervention.
- 3- To compare the level of knowledge and attitudes on toxoplasmosis among the respondents within at different stages (as mentioned above).
- 4- To determine the association between the level of knowledge and attitudes with socio-demographic characteristics of the respondents.
- 5- To determine the correlation between the level of knowledge and attitudes on toxoplasmosis with sociodemographic characteristics of respondent.

## **1.5 Study Hypothesis**

- 1- There is a significant difference in the level of knowledge and attitudes towards toxoplasmosis between Experiment and Control groups.
- 2- The level of knowledge related to toxoplasmosis among pregnant women in Al-Najaf province is low.
- 3- The attitude related to toxoplasmosis among pregnant women in Al-Najaf province is negative.
- 4- There is a significant association between knowledge and attitudes related to toxoplasmosis among pregnant women in Al-Najaf province and socio-demographic characteristics.
- 5- There is a significant difference in the levels of knowledge and attitudes related to toxoplasmosis among three stages of data collection of the study.

## **1.6 Conceptual framework**

Toxoplasmosis has been known as a disease with various risk factors, which include past medical and obstetric history, socio-demographic characteristics, geographical distribution, housing conditions index, source of infection, pet ownership, toxoplasmosis immunization status, and knowledge, attitudes, and practices (KAP).




Figure 1.1: Conceptual framework of this study

## 1.7 Definitions of Terms

**Risk Factors:** Factors that increase a woman's chances of getting toxoplasmosis are called risk factors. Risk factors are not necessarily causes of toxoplasmosis but are associated with an increased chance of getting toxoplasmosis. Different risk factors can control (modifiable risk factors, e.g. diet and lifestyle) (Goldman & Hatch, 2000).

**Toxoplasmosis infection:** A parasitic disease caused by *Toxoplasma gondii*. This disease usually causes no symptoms in adult humans. Sometimes there may be a few weeks or months of mild flu-like illness such as muscle aches and tender lymph nodes. This parasite may invade tissues and damage the brain, especially of the fetus and newborn (CDC, 2012).

**Socio-demographic characteristics:** Define as a set of variables for any human such as a given population's age, ethnicity, and socioeconomic status, whether they reside in an urban or rural area (Fletcher & Hirdes, 2002).

**Housing conditions:** A group of elements of the quality for any house to be suitable for people who are living at that place such as availability of sufficient space in the dwelling, availability of basic sanitary facilities (such as a bath or shower or indoor flushing toilet), the wider residential area, availability of good ventilation and availability of natural and artificial lighting (Verma & Betti, 2006).

**Pet ownership:** A person who owns a pet such as dog or cat (Conlee, Stephens & Rowan, 2009).

**Cooking preferences:** The type of food that is liked, wanted and preferred more than another type. Cooking preferences can describe user-configured values for cooking parameters (Tasevska, et al., 2011).

**The source of infection:** Defined as the person, animal, object or substance have the ability to harboring the infectious agent and spread it to the host (Friis & Sellers, 2013).

**Past obstetric history:** Define as the previous information about gestational age, history of current pregnancy, antenatal history, previous pregnancies, menstrual history, sexual history, and gynecological conditions (O'Connor & Kovacs, 2003).

**Past medical history:** The patient's health status before the presenting problem, which is included the medical information about the past diseases and medical conditions (O'Connor & Kovacs, 2003).

**Immunization status:** Describes the current status of the vaccination event or a record of a vaccination as reported by a patient, a clinician's or another party (Shefer, et al., 2011).



## REFERENCES

- A'aiz, N. N. (2010). Genotyping analysis to determine the lineages types of *Toxoplasma Gondii* with study of autoantibodies production. Ph.D. Sc. Thesis. *College of Medicine. Kufa University, Iraq.*
- Abd El-Razek, H. D. (2012). *Toxoplasmosis related knowledge, perception of pregnant women in Menoufia, Egypt.* (Unpublished Ph.D.). Faculty of Nursing, Tanta University.
- Abu-Madi, M. A., Al-Molawi, N., & Behnke, J. M. (2008). Seroprevalence and epidemiological correlates of toxoplasma *Gondii* infections among patients referred for hospital-based serological testing in Doha, Qatar. *Parasites & Vectors, 1* (1), 39-3305-1-39. Available at <http://parasitesandvectors.biomedcentral.com/articles/10.1186/1756-3305-1-39> Accessed on January 4, 2016.
- Abu-Madi, M. A., Behnke, J. M., & Dabritz, H. A. (2010). *Toxoplasma gondii* seropositivity and co-infection with TORCH pathogens in high-risk patients from Qatar. *The American journal of tropical medicine and hygiene, 82*(4), 626-633. Available at <http://www.ajtmh.org/content/82/4/626.full> Accessed on February 15, 2016.
- Abu-Zeid, Y. (2002). Serological evidence for remarkably variable prevalence rates of *toxoplasma gondii* in children of major residential areas in United Arab Emirates. *Acta Tropica, 83*(1), 63-69. Available at <http://www.sciencedirect.com/science/article/pii/S0001706X0200061X> Accessed on March 15, 2016.
- Aday, L. A., & Cornelius, L. J. (2011). *Designing and conducting health surveys: A comprehensive guide* John Wiley & Sons. Available at [www.josseybass.com](http://www.josseybass.com) Accessed on January 4, 2016.
- Adesiyun, A. A., Gooding, R., Ganta, K., Seepersadsingh, N., & Ramsewak, S. (2007). Congenital toxoplasmosis in two health institutions in Trinidad. *West Indian medical journal, 56*(2), 166-170. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17910149>. Accessed on February 27, 2016.
- Ageel, N. F. (2003). *Serological and biochemical study of toxoplasmosis in Tikrit teaching hospital.* (Unpublished MSc. Thesis). College of Medicine, Tikrit University.
- Al Hamdani, M. M., & Mahdi, N. K. (1997). Toxoplasmosis among women with habitual abortion. Available at <http://apps.who.int/iris/handle/10665/117384> Accessed on February 6, 2016.
- Albarracín, D., & McNatt, P. S. (2005). Maintenance and decay of past behavior influences: Anchoring attitudes on beliefs following inconsistent actions. *Personality and Social Psychology Bulletin, 31*(6), 719-733. Available at <http://psp.sagepub.com/content/31/6/719.full.pdf> Accessed on January 4, 2016.

- Al-Doori, M. A. (2010). Epidemiological study of *Toxoplasma gondii* between couples in Tikrit city, and experimental trial about possibility of sexual transmission of infection in mice (Doctoral dissertation, M. Sc. Thesis, college of Education, University of Tikrit).
- Al-Garawia, Z.S, Al-Fartusiea, F. S & Al-Bairmani, H. K. (2012). Is hyperglycemia or toxoplasmosis the suspect of spontaneous abortion in neonates in Baghdad women? *J. Basic. Appl. Chem*, 2(6)29-34. ISSN 2090-424X. Available at <http://www.textroad.com/JBAC-%20June-August,%202012.html>. Accessed on February 5, 2016.
- Al-haris, F. M., Saheb, H. S., & Abdul-Sada, K. M. (2015). Investigation of Toxoplasmosis in Cord Blood of Newborns at Al-Najaf Province, Iraq by Searching for IgG and IgM Antibodies. *Int. J. Curr. Microbiol. App. Sci*, 4(2), 314-321. Available at <http://ijcmas.com/vol-4-2/Faris%20M.%20Al-haris,%20et%20al.pdf>. Accessed on February 5, 2016.
- Al-Hindi, A. I., & Lubbad, M. H. (2009). Seroprevalence of toxoplasmosis among Palestinian aborted women in Gaza. *Ann Alquds Med*, 5, 39-47.
- Ali, Z., Hossein, M. M., & Khadijeh, D. (2007). Toxoplasma chorioretinitis in primary school children in Tehran, Iran, 2003-2004. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*, 13(4), CR201-5. Available at <http://www.medscimonit.com/abstract/index/idArt/481106> Accessed on February 15, 2016.
- Aljumaily, Z., & Alsamarai, A. (2013). Risk factors for bad obstetric history in Kirkuk women, Iraq. *International Journal of Infection and Microbiology*, 2(3), 70-77. Available at <http://www.nepjol.info/index.php/IJIM/article/viewFile/8121/7276>. Accessed on February 27, 2016.
- Al-Kalaby, R. F. (2008). Sero-epidemiological study of toxoplasmosis among different groups of population in Najaf city (Doctoral dissertation, M. Sc. Thesis, Kufa University).
- Al-Mousawi, G. N. (2008): Some Immunological and sero-epidemiological aspects associated with toxoplasmosis in aborted and unmarried women in Basrah province. M. Sc. Thesis, University of Basrah.
- Al-Nahi, A. S., & Al-Abbas, S. K. A. (2007). Prevalence of *Toxoplasma Gondii* among females aged (16-26) years in Al-Najaf city. Available at <http://sci.uokufa.edu.iq/>. Accessed on February 17, 2016.
- Al-Nakib, W., Ibrahim, M. E. A., Hathout, H., Moussa, M. A. A., Deverajan, L. V., Thorburn, H., & Yousof, A. M. (1983). Seroepidemiology of viral and toxoplasmal infections during pregnancy among Arab women of child-bearing age in Kuwait. *International journal of epidemiology*, 12(2), 220-223. Available at <http://ije.oxfordjournals.org/content/12/2/220.short> Accessed on February 15, 2016.
- Al-Seadawy, M. A. H. (2010). Prevalence of Toxoplasmosis in pregnant women in Al Muthanna province/Iraq. *Kufa J Vet Med Sci*, 1, 166-73. Available at <http://uokufa.edu.iq/journals/index.php/kjvs/article/viewFile/2407/2051> Accessed on February 6, 2016.



- Al-Sheyab, N. A., Obaidat, M. M., Bani Salman, A. E., & Lafi, S. Q. (2015). Toxoplasmosis-Related Knowledge and Preventive Practices among Undergraduate Female Students in Jordan. *Journal of Food Protection*, 78(6), 1161-1166. Available at <http://dx.doi.org/10.4315/0362-028X.JFP-14-579> Accessed on January 4, 2016.
- Al-Shimmery, M. N., Al-Hilaly, H. A., & Al-Khafaji, A. (2011). Seroprevalence of cytomegalovirus and toxoplasmosis in cases of miscarriages women in Al-Diwaniyah province. *Al-Qadisiyah Med J*, 7, 160-168. Available at <http://www.iasj.net/iasj?func=fulltext&aId=14048> Accessed on February 6, 2016.
- Altintas, N., Kuman, H. A., Akisu, C., Aksoy, U., & Atambay, M. (1997). Toxoplasmosis in last four years in Aegean region, turkey. *Journal of the Egyptian Society of Parasitology*, 27(2), 439-443. Available at <http://europepmc.org/abstract/med/9257982> Accessed on February 15, 2016.
- AL-Waely, R. (1998). *The prevalence of seropositive toxoplasma gondii antibodies among Iraqi mothers and their new-born babies in Baghdad*. (Unpublished The diploma in primary health care). College of Medicine, Tikrit Univ. (Accessed December 10, 2015).
- Al-Warid, H. S., & Al-Qadhi, B. N. (2012). Evaluation of progesterone and estrogen hormonal levels in pregnant women with toxoplasmosis. *European Journal of Scientific Research*, 91 (4), 515-519. Available at <https://www.researchgate.net/publication/233993325> Accessed on February 4, 2016.
- Amin, T. T., Al Ali, M. N., Alrashid, A. A., Al-Agnam, A. A., & Al Sultan, A. A. (2013). Toxoplasmosis preventive behavior and related knowledge among Saudi pregnant women: an exploratory study. *Global journal of health science*, 5(5), 131. Available at <http://www.ccsenet.org/journal/html/gjhs/articles/27592.html> Accessed on January 4, 2016.
- Ancelle, T., Goulet, V., Tirard-Fleury, V., Baril, L., Du Mazaubrun, C., Thulliez, P. H., & Carme, B. (1996). La Toxoplasmose chez la femme enceinte en France en 1995. Resultats d'une enquete nationale perinatale. *Bull Epidemiol Hebd*, 51, 227-229. Available at <http://fulltext.bdsp.ehesp.fr/Invs/Beh/1996/51/51.pdf> Accessed on October 19, 2016.
- Andiappan, H., Nissapatorn, V., Sawangjaroen, N., Khaing, S. L., Salibay, C. C., Cheung, M. M. M., ... & Adenan, N. A. M. (2014). Knowledge and practice on Toxoplasma infection in pregnant women from Malaysia, Philippines, and Thailand. *Frontiers in microbiology*, 5, 291. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24966855> Accessed on January 4, 2016.
- Angel-Müller, E., Houghton, M. P., Eslava, C., Riaño, J., Rey, G. E., & Gómez-Marín, J. E. (2014). Gestational and congenital toxoplasmosis in two hospitals in Bogota, Colombia. *Revista Facultad De Medicina De La Universidad Nacional De Colombia*, 62(2), 179-185. Available at <http://dx.doi.org/10.15446/revfacmed.v62n2.45430> Accessed on February 15, 2016.
- Aqeely, H., El-Gayar, E. K., Perveen Khan, D., Najmi, A., Alvi, A., Bani, I., ... & Elhassan, I. M. (2014). Seroepidemiology of Toxoplasma gondii amongst

- pregnant women in Jazan province, Saudi Arabia. *Journal of tropical medicine*, 2014. Available at <http://downloads.hindawi.com/journals/jtm/2014/913950.pdf> Accessed on February 15, 2016.
- Aubert, D., Foudrinier, F., Villena, I., Pinon, J. M., Biava, M. F., & Renoult, E. (1996). PCR for diagnosis and follow-up of two cases of disseminated toxoplasmosis after kidney grafting. *Journal of clinical microbiology*, 34(5), 1347. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC229018/pdf/341347.pdf> Accessed on March 4, 2016.
- Avelino, M. M., Campos Júnior, D., Parada, J. B. d., & Castro, A. M. d. (2004). Risk factors for toxoplasma gondii infection in women of childbearing age. *Brazilian Journal of Infectious Diseases*, 8(2), 164-174. Available at <http://dx.doi.org/10.1590/S1413-86702004000200007> Accessed on May 9, 2016.
- Ayi, I., Edu, S., Apea-Kubi, K., Boamah, D., Bosompem, K., & Edoh, D. (2009). Sero-epidemiology of toxoplasmosis amongst pregnant women in the greater Accra region of Ghana. *Ghana Medical Journal*, 43(3). Available at <http://www.ajol.info/index.php/gmj/article/viewFile/55325/43789> Accessed on January 4, 2016.
- Azevedo, Kátia Martins Lopes de, Setúbal, S., Lopes, V. G. S., Camacho, L. A. B., & Oliveira, S. A. d. (2010). Congenital toxoplasmosis transmitted by human immunodeficiency-virus infected women. *Brazilian Journal of Infectious Diseases*, 14(2), 186-189. Available at [http://www.scielo.br/scielo.php?pid=S1413-86702010000200014&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S1413-86702010000200014&script=sci_arttext) Accessed on January 4, 2016.
- Aziz, F. M., & Druish, M. J. (2011). Toxoplasmosis: Serious disease during pregnancy. *Baghdad Sci J*, 8, 91-95. Available at <http://csw-journal.org/full/57.pdf> Accessed on January 28, 2016.
- Aziz, T. M., Krysiak, P., El-Gamel, A., Campbell, C., Rahman, A., Deiraniya, A., & Yonan, N., (1998). Bacteremia and endocarditis following endomyocardial biopsy. In *Transplantation proceedings* (Vol. 30, No. 5, pp. 2112-2113). Available at Elsevier. [http://dx.doi.org/10.1016/S0041-1345\(98\)00556-9](http://dx.doi.org/10.1016/S0041-1345(98)00556-9) Accessed on March 4, 2016.
- Babikian, T., Freier, M. C., Hopkins, G. L., DiClemente, R., McBride, D., & Riggs, M. (2004). An assessment of HIV/AIDS risk in higher education students in Yerevan, Armenia. *AIDS and Behavior*, 8(1), 47-61. Available at <http://link.springer.com/article/10.1023/B:AIBE.0000017525.92015.5d> Accessed on May 9, 2016.
- Baker, D. W., Parker, R. M., Williams, M. V., Clark, W. S., & Nurss, J. (1997). The relationship of patient reading ability to self-reported health and use of health services. *American Journal of Public Health*, 87(6), 1027-1030. Available at <http://ajph.aphapublications.org/doi/abs/10.2105/AJPH.87.6.1027> Accessed on May 9, 2016.
- Baril, L., Ancelle, T., Goulet, V., Thulliez, P., Tirard-Fleury, V., & Carme, B. (1999). Risk factors for Toxoplasma infection in pregnancy: a case-control study in France. *Scandinavian journal of infectious diseases*, 31(3), 305-309.

Available at <http://dx.doi.org/10.1080/00365549950163626> Accessed on October 19, 2016.

- Ben, A. R., Siala, E., Bouafsoun, A., Maatoug, R., Souissi, O., Aoun, K., & Bouratbine, A. (2013). [Toxoplasmosis mother-to-child screening: study of cases followed in the Pasteur Institute of Tunis (2007-2010)]. *Bulletin de la Societe de pathologie exotique (1990)*, 106(2), 108-112. Available at <http://europepmc.org/abstract/med/23576025> Accessed on March 15, 2016.
- Bonita, R., Beaglehole, R., & Kjellström, T. (2006). Basic epidemiology. *World Health Organization*, 212. Available at [http://colinmayfield.com/waterhealth/course2/content/Resources/WHO\\_Epidemiology\\_basics/9241547073\\_eng.pdf](http://colinmayfield.com/waterhealth/course2/content/Resources/WHO_Epidemiology_basics/9241547073_eng.pdf) Accessed on October 14, 2016.
- Bouhamdan, S. F., Bitar, L. K., Saghir, H. J., Bayan, A., & Araj, G. F. (2010). Seroprevalence of toxoplasma antibodies among individuals tested at hospitals and private laboratories in Beirut. *Le Journal Medical Libanais. the Lebanese Medical Journal*, 58(1), 8-11. Available at <http://www.fsedu.usj.edu.lb/unte/JML/articles/58-1/original2.pdf> Accessed on February 15, 2016.
- Boyer, C. B., Tschann, J. M., & Shafer, M. (1999). Predictors of risk for sexually transmitted diseases in ninth grade urban high school students. *Journal of Adolescent Research*, 14(4), 448-465. Available at <http://jar.sagepub.com/content/14/4/448.full.pdf> Accessed on May 9, 2016.
- Boyer, K. M., Holfels, E., Roizen, N., Swisher, C., Mack, D., Remington, J., \Toxoplasmosis Study Group. (2005). Risk factors for toxoplasma gondii infection in mothers of infants with congenital toxoplasmosis: Implications for prenatal management and screening. *American Journal of Obstetrics and Gynecology*, 192(2), 564-571. Available at <http://www.sciencedirect.com/science/article/pii/S0002937804007914> Accessed on January 4, 2016.
- Bradley, R. H., & Corwyn, R. F. (2002). Socioeconomic status and child development. *Annual Review of Psychology*, 53(1), 371-399. Available at <http://www.annualreviews.org/doi/pdf/10.1146/annurev.psych.53.100901.135233> Accessed on February 6, 2016.
- Branco, B. H. M., de Araújo, S. M., & Falavigna-Guilherme, A. L. (2012). Primary prevention of toxoplasmosis: Knowledge and attitudes of health professionals and pregnant women of public service of Maringa, Parana state, Brazil [abstract in English]. *Scientia Medica*, 22(4), 185-190. Available at <http://revistaseletronicas.pucrs.br/ojs/index.php/scientiamedica/article> Accessed on May 9, 2016.
- Breugelmans, M., Naessens, A., & Foulon, W. (2004). Prevention of toxoplasmosis during pregnancy—an epidemiologic survey over 22 consecutive years. *Journal of perinatal medicine*, 32(3), 211-214. Available at <https://www.degruyter.com/view/j/jpme.2004.32.issue3/jpm.2004.039/jpm.2004.039.xml> Accessed on October 14, 2016.
- Brown, C. R., Hunter, C. A., Estes, R. G., Beckmann, E., Forman, J., David, C., ... & McLeod, R. (1995). Definitive identification of a gene that confers resistance against Toxoplasma cyst burden and encephalitis. *Immunology*, 85(3), 419.

Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1383915/pdf/immunology00069-0075.pdf> Accessed on January 4, 2016.

- Brug J, Assema P, Lechner L. (Ed.). (2007). *In: Gezondheidsvoorlichting en gedragsverandering: Een planmatige* [In Health Education and behavior: a systematic approach.] (5th ed.). Open Universiteit Nederland: Heerlen: van Gorcum.
- Burg, J. L., Grover, C. M., Pouletty, P., & Boothroyd, J. C. (1989). Direct and sensitive detection of a pathogenic protozoan, *Toxoplasma gondii*, by polymerase chain reaction. *Journal of clinical microbiology*, 27(8), 1787-1792. Available at <http://jcm.asm.org/content/27/8/1787.full.pdf> Accessed on January 4, 2016.
- Camargo, M. E., Ferreira, A. W., Mineo, J. R., Takiguti, C. K., & Nakahara, O. S. (1978). Immunoglobulin G and immunoglobulin M enzyme-linked immunosorbent assays and defined toxoplasmosis serological patterns. *Infection and Immunity*, 21(1), 55-58. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2018360/pdf/amjpathol00734-0153.pdf> Accessed on March 4, 2016.
- Cannon, S., & Boswell, C. (2009). Filling gaps in knowledge: Educating nurses to provide appropriate patient materials. *Journal of Continuing Education in Nursing*, 40(4), 148-149. Available at <http://www.healio.com/nursing/journals/jcen/2009-4-40-4> Accessed on May 9, 2016.
- Cantos, G. A., Prando, M. D., & Siqueira, M. V. & Teixeira, R. M. (2000). Toxoplasmosis: Occurrence of antibodies ant *toxoplasma gondii* and diagnosis. *Rev. Assoc. Med. Bras. [Online]*, 46(4), pp.335-341. Available at <https://dx.doi.org/10.1590/S0104-4230200000400033> Accessed on February 15, 2016.
- Caprarelli, G., & Fletcher, S. (2014). A brief review of spatial analysis concepts and tools used for mapping, containment and risk modelling of infectious diseases and other illnesses. *Parasitology*, 141(05), 581-601. Available at <http://dx.doi.org/10.1017/S0031182013001972> Accessed on October 14, 2016.
- Carter, A. O., Gelmon, S. B., Wells, G. A., & Toepell, A. P. (1989). The effectiveness of a prenatal education programme for the prevention of congenital toxoplasmosis. *Epidemiology and infection*, 103(03), 539-545. Available at <http://dx.doi.org/10.1017/S0950268800030934> Accessed on October 14, 2016.
- Centers for Disease Control and Prevention (CDC). (2000). Preventing congenital toxoplasmosis, *recommendations and reports. Mmwr*, 49((RR02)), January 20 2016-57-75. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr4902a5.htm> Accessed on May 9, 2016.
- Centers for Disease Control and Prevention (CDC). (2012). Parasites-toxoplasmosis (toxoplasma infection). *Web Page, [Accessed 25 November 2012]*. Available at <http://www.Cdc.gov/parasites/toxoplasmosis> Accessed on January 4, 2016.

- Central Statistical Organization (CSO), (2013). Demographic statistics. Ministry of Planning, Republic of Iraq. Available at <http://cosit.gov.iq/en/population-manpower-statistics/life> Accessed on January 4, 2016.
- Chabbert, E., Lachaud, L., Crobu, L., & Bastien, P. (2004). Comparison of two widely used PCR primer systems for detection of toxoplasma in amniotic fluid, blood, and tissues. *Journal of Clinical Microbiology*, 42(4), 1719-1722. Available at <http://jcm.asm.org/content/42/4/1719.short> Accessed on January 4, 2016.
- Chacin-Bonilla, L., Sanchez-Chavez, Y., Monsalve, F., & Estevez, J. (2001). Seroepidemiology of toxoplasmosis in Amerindians from western Venezuela. *The American Journal of Tropical Medicine and Hygiene*, 65(2), 131-135. Available at <http://www.ajtmh.org/content/65/2/131.full.pdf> Accessed on August 19, 2015.
- Chapel, H., Haeney, M., Misbah, S., & Snowden, N. (Ed.). (2013). *Essentials of clinical immunology* (6 ed.) John Wiley & Sons. Available at <https://books.google.com.my/books> Accessed on January 15, 2016.
- Charest, H., Sedegah, M., Yap, G. S., Gazzinelli, R. T., Caspar, P., Hoffman, S. L., & Sher, A. (2000). Recombinant attenuated *Toxoplasma gondii* expressing the *Plasmodium yoelii* circumsporozoite protein provides highly effective priming for CD8+ T cell-dependent protective immunity against malaria. *The Journal of Immunology*, 165(4), 2084-2092. Available at <http://www.jimmunol.org/content/165/4/2084> Accessed on March 4, 2016.
- Chatterton, J. M., Evans, R., Ashburn, D., Joss, A. W. L., & Ho-Yen, D. O. (2002). *Toxoplasma gondii* in vitro culture for experimentation. *Journal of microbiological methods*, 51(3), 331-335. Available at <http://www.sciencedirect.com/science/article/pii/S016770120200101X> Accessed on March 4, 2016.
- Chow, S. C., Wang, H., & Shao, J. (2007). *Sample size calculations in clinical research*. CRC press. Taylor & Francis Group. 2<sup>nd</sup> edition. Pp 206.
- Conlee, K., Stephens, M., & Rowan, A. N. (2009). The Humane Society of the United States. Accessed on <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.683.6312> Accessed on October 10, 2016.
- Conyn-van Spaendonck, M. A. E., & Van Knapen, F. (1992). Choices in preventive strategies: experience with the prevention of congenital toxoplasmosis in The Netherlands. *Scandinavian Journal of Infectious Diseases Supplement*, 51-51. Available at <http://dx.doi.org/10.3109/inf.1992.24.suppl-84.01> Accessed on October 19, 2016.
- Cook, A. J. C., Holliman, R., Gilbert, R. E., Buffolano, W., Zufferey, J., Petersen, E., ... & Dunn, D. T. (2000). Sources of toxoplasma infection in pregnant women: European multicenter case-control study commentary: Congenital toxoplasmosis—further thought for food. *Bmj*, 321(7254), 142-147. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC27431/> Accessed on January 4, 2016.

- Cormier, C. M., & Kotrlik, J. W. (2009). Health literacy knowledge and experiences of senior baccalaureate nursing students. *The Journal of Nursing Education*, 48(5), 237-248. Available at <http://www.healio.com/nursing/nursing-education> Accessed on May 9, 2016.
- Costa, F. F., Gondim, A. P. S., de Lima, M. B., Braga, J. U., de Souza Vieira, L. J. E., & Araújo, M. A. L. (2012). Preventive behavior for toxoplasmosis in pregnant adolescents in the state of Ceara, Brazil. *BMC Public Health*, 12(1), 1. Available at <http://bmcpublikealth.biomedcentral.com/articles/10.1186/1471-2458-12-73> Accessed on January 4, 2016.
- Crano, W. D., & Prislin, R. (Eds.). (2011). *Attitudes and attitude change*. Psychology Press. Available at <https://books.google.com.my/books> Accessed on January 4, 2016.
- Davis, T. C., Williams, M. V., Marin, E., Parker, R. M., & Glass, J. (2002). Health literacy and cancer communication. *CA: A Cancer Journal for Clinicians*, 52(3), 134-149. Available at <http://onlinelibrary.wiley.com/doi/10.3322/canjclin.52.3.134/full> Accessed on May 9, 2016.
- De Sousa, S. R. (2009). Serotyping of toxoplasma gondii (Doctoral dissertation, University De Limoges).
- Desta, A. H. (2015). Knowledge, attitude and practice of community towards zoonotic importance of toxoplasma infection in central afar region, north east Ethiopia. *International Journal of Biomedical Science and Engineering*, 3(6), 74-81. Available at [http://article.sciencepublishinggroup.com/html/10.11648.j.ijbse.20150306.12.html](http://article.sciencepublishinggroup.com/html/10.11648/j.ijbse.20150306.12.html) Accessed on January 4, 2016.
- Di Mario, S., Basevi, V., Gagliotti, C., Spettoli, D., Gori, G., D'Amico, R., & Magrini, N. (2015). Prenatal education for congenital toxoplasmosis. *Cochrane Database Syst Rev*, 4. Pub. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26493047> Accessed on January 4, 2016.
- Di Mario, S., Basevi, V., Gagliotti, C., Spettoli, D., Gori, G., D'Amico, R., & Magrini, N. (2013). Prenatal education for congenital toxoplasmosis. *Cochrane Database Syst Rev*, 2. Available at [http://www.cochrane.org/CD006171/PREG\\_prenatal-education-congenital-toxoplasmosis](http://www.cochrane.org/CD006171/PREG_prenatal-education-congenital-toxoplasmosis) Accessed on May 9, 2016.
- Doehring, E., Reiter-Owona, I., Bauer, O., Kaisi, M., Hlobil, H., Quade, G., . . . Seitz, H. M. (1995). Toxoplasma gondii antibodies in pregnant women and their newborns in Dar es salaam, Tanzania. *The American Journal of Tropical Medicine and Hygiene*, 52(6), 546-548. Available at <http://europepmc.org/abstract/med/7611563> Accessed on February 15, 2016.
- Dubey, J. P. (2004). Toxoplasmosis—a waterborne zoonosis. *Veterinary parasitology*, 126(1), 57-72. Available at <http://dx.doi.org/10.1016/j.vetpar.2004.09.005> Accessed on January 15, 2016.
- Dubey, J. P. (2014). The history and life cycle of toxoplasma gondii. *Toxoplasma Gondii (Second Edition)*, 1-17. Available at <http://www.ars.usda.gov/pandp/people/people.htm?personid=1472> Accessed on January 28, 2016.

- Dubey, J. P., & Beattie, C. P. (Eds.). (1988). *Toxoplasmosis of animals and man*. (First ed.). Boca Raton, FL 33431, USA: CRC Press, Inc. pp 220.
- Dubey, J. P., & Jones, J. L. (2008). Toxoplasma gondii infection in humans and animals in the United States. *International journal for parasitology*, 38(11), 1257-1278. Available at <http://www.sciencedirect.com/science/article/pii/S0020751908001100> Accessed on February 4, 2016.
- Dubey, J. P., & Thulliez, P. H. (1993). Persistence of tissue cysts in edible tissues of cattle fed Toxoplasma gondii oocysts. *American journal of veterinary research*, 54(2), 270-273. Available at <http://europemc.org/abstract/med/8430937> Accessed on March 4, 2016.
- Dubey, J., Lago, E., Gennari, S., Su, C., & Jones, J. (2012). Toxoplasmosis in humans and animals in brazil: High prevalence, high burden of disease, and epidemiology. *Parasitology*, 139(11), 1375-1424. Available at <http://dx.doi.org/10.1017/S0031182012000765> Accessed on January 15, 2016.
- Dumas, P. N., Le Guenno, B., Digoutte, J. P., & Seguela, J. P. (1990). Toxoplasmosis in the republic of Senegal. sero-epidemiological survey. [Toxoplasmose en republique du Senegal. Sondage sero-epidemiologique] *Bulletin De La Societe De Pathologie Exotique* (1990), 83(2), 283-285. Available at <http://europemc.org/abstract/med/2208458> Accessed on February 15, 2016.
- Dunn, S. M., Beeney, L. J., Hoskins, P. L., & Turtle, J. R. (1990). Knowledge and attitude change as predictors of metabolic improvement in diabetes education. *Social Science & Medicine*, 31(10), 1135-1141. Available at <http://www.sciencedirect.com/science/article/pii/027795369090235K> Accessed on May 9, 2016.
- Durlach, R., Kaufer, F., Carral, L., Freuler, C., Ceriotto, M., Rodríguez, M., Corazza, R. (2008). Consenso argentino de toxoplasmosis congénita. *Medicina (Buenos Aires)*, 68(1), 75-87. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3826532/> Accessed on February 15, 2016.
- Ebbesen, P. (2000). Placenta physiology. In: Ambroise-Thomas, P.; Petersen, E. editors. *Congenital toxoplasmosis: Scientific background, clinical management and control*. Paris: Springer- Verlag, pp. 27-35. Available at [http://link.springer.com/chapter/10.1007/978-2-8178-0847-5\\_3](http://link.springer.com/chapter/10.1007/978-2-8178-0847-5_3) Accessed on March 4, 2016.
- Egbert, N., & Nanna, K. M. (2009). Health literacy: Challenges and strategies. *The Online Journal of Issues in Nursing*, 14(3). Available at <http://www.nursingworld.org> Accessed on October 14, 2016.
- El Awady, M. (2000). Comparison between toxoplasma gondii DNA and specific immunoglobulins during pregnancy. *EMHJ - Eastern Mediterranean Health Journal*, 6 (5-6), 888-897. Available at <http://www.who.int/iris/handle/10665/118943> Accessed on January 4, 2016.
- Elbez-Rubinstein, A., Ajzenberg, D., Dardé, M., Cohen, R., Dumètre, A., Yera, H., . . . Thulliez, P. (2009). Congenital toxoplasmosis and reinfection during pregnancy: Case report, strain characterization, experimental model of

- reinfection, and review. *Journal of Infectious Diseases*, 199(2), 280-285. Available at <http://jid.oxfordjournals.org/content/199/2/280.full> Accessed on January 4, 2016.
- Elliott, P., & Best, N. (1998). Geographic patterns of disease. *Encyclopedia of Biostatistics*. John Wiley & Sons, Inc.
- Elsheikha, H. M. (2008). Congenital toxoplasmosis: priorities for further health promotion action. *Public health*, 122(4), 335-353. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17964621> Accessed on May 9, 2016.
- Engstrom Jr., R. E., Holland, G. N., Nussenblatt, R. B., & Jabs, D. A. (1991). Current practices in the management of ocular toxoplasmosis. *American Journal of Ophthalmology*, 111(5), 601-610. Available at [http://ezproxy.upm.edu.my:2112/10.1016/S0002-9394\(14\)73706-7](http://ezproxy.upm.edu.my:2112/10.1016/S0002-9394(14)73706-7) Accessed on January 4, 2016.
- Eshete, H., Tessema, S., Abebe, S., & Abebe, A. (1994). Some notes on toxoplasmosis in pregnant women in Addis Ababa. *Ethiopian Medical Journal*, 32(2), 135-136. Available at <http://europepmc.org/abstract/med/8033880> Accessed on February 15, 2016.
- Evengard, B., Lilja, G., Capraru, T., Malm, G., Kussofsky, E., Öman, H., & Forsgren, M. (1999). A retrospective study of seroconversion against *Toxoplasma gondii* during 3,000 pregnancies in Stockholm. *Scandinavian journal of infectious diseases*, 31(2), 127-129. Available at <http://www.tandfonline.com/doi/abs/10.1080/003655499750006146> Accessed on March 4, 2016.
- Evering, T., & Weiss, L. M. (2006). The immunology of parasite infections in immunocompromised hosts. *Parasite immunology*, 28(11), 549-565. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3109637/> Accessed on February 4, 2016.
- Ferreira, M. R., Dolan, N. C., Fitzgibbon, M. L., Davis, T. C., Gorby, N., Ladewski, L., . . . Bennett, C. L. (2005). Health care provider-directed intervention to increase colorectal cancer screening among veterans: Results of a randomized controlled trial. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 23(7), 1548-1554. Available at <https://jco.ascopubs.org/content/23/7/1548.full> Accessed on May 9, 2016.
- Filisetti, D., & Candolfi, E. (2004). Immune response to *Toxoplasma gondii*. *Ann Ist Super Santa*, 40(1), 71-80. Available at <http://www.iss.it/binary/publ/publi/40171.1107854946.pdf> Accessed on January 4, 2016.
- Fischer, M. J., Rauch, J., & Levine, J. S. (2007, January). The antiphospholipid syndrome. In *Seminars in nephrology* (Vol. 27, No. 1, pp. 35-46). WB Saunders. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3440307/> Accessed on January 15, 2016.
- Flegr, J., Havlíček, J., Kodým, P., Malý, M., & Smahel, Z. (2002). Increased risk of traffic accidents in subjects with latent toxoplasmosis: a retrospective case-control study. *BMC infectious diseases*, 2(1), 11. Available at <http://bmcinfectdis.biomedcentral.com/articles/10.1186/1471-2334-2-11> Accessed on February 4, 2016.



- Flegr, J., Kodým, P., & Tolarová, V. (2000). Correlation of duration of latent *Toxoplasma gondii* infection with personality changes in women. *Biological psychology*, 53(1), 57-68. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10876065> Accessed on March 4, 2016.
- Flegr, J., Prandota, J., Sovičková, M., & Israili, Z. H. (2014). Toxoplasmosis—A global threat. correlation of latent toxoplasmosis with specific disease burden in a set of 88 countries. *PLoS One*, 9(3), e90203. Available at <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0090203> Accessed on February 4, 2016.
- Fletcher, P. C., & Hirdes, J. P. (2002). Risk factors for falling among community-based seniors using home care services. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57(8), M504-M510. Available at <http://biomedgerontology.oxfordjournals.org/content/57/8/M504.short> Accessed on October 10, 2016.
- Fonlon, W., Naessens, A., & Derde, M. (1994). Evaluation of the possibilities for preventing congenital toxoplasmosis. *Obstetrical & Gynecological Survey*, 49(9), 601-602. Available at <http://journals.lww.com/obgynsurvey/citation/> Accessed on May 9, 2016.
- Fonseca, A. L., Silva, R. A., Fux, B., Madureira, A. P., Sousa, F. F. d., & Margonari, C. (2012). Epidemiologic aspects of toxoplasmosis and evaluation of its seroprevalence in pregnant women. *Revista Da Sociedade Brasileira De Medicina Tropical*, 45(3), 357-364. Available at <http://dx.doi.org/10.1590/S0037-86822012000300015> Accessed on January 4, 2016.
- Fosu. A. K., (2015). Poverty and development. *Bulletin of the World Health Organization*, 85, 10. pp 733-810, October 2007. Available at <http://www.who.int/bulletin/volumes/85/10/07-045955/en/> Accessed on February 6, 2016.
- Foulon, W., Naessens, A., & Ho-Yen, D. (2000). Prevention of congenital toxoplasmosis. *Journal of perinatal medicine*, 28(5), 337-345. Available at <https://www.degruyter.com/view/j/jpme.2000.28.issue5/jpm.2000.043/jpm.2000.043.xml> Accessed on October 19, 2016.
- Frenkel, J. K., & Pfefferkorn, E. R. (1991). *U.S. Patent No. 5,045,313*. Washington, DC: U.S. Patent and Trademark Office. Available at <https://www.google.com/patents/US5045313> Accessed on February 4, 2016.
- Friis, R. H., & Sellers, T. (2013). *Epidemiology for public health practice*. Jones & Bartlett Publishers. Accessed on <http://samples.jbpub.com/9781449665494/frontmatter.pdf> Accessed on October 10, 2016.
- Furtado, J. M., Smith, J. R., Belfort, R., Jr, Gattey, D., & Winthrop, K. L. (2011). Toxoplasmosis: A global threat. *Journal of Global Infectious Diseases*, 3(3), 281-284. Available at <http://www.jgid.org/text.asp?2011/3/3/281/83536> Accessed on February 6, 2016.
- Gaheen, M. A. S. A. and Elkazeh, E. A. E. E. (2014). Knowledge and attitude of women regarding toxoplasmosis during pregnancy and measures to overcome it in slums areas. *International Journal of Current Research*, 6(4), 6356-6371.

Available at <http://www.journalcra.com/> Accessed on January 4, 2016.

- Garfield, R., & Waldman, R. (2003). Review of potential interventions to reduce child mortality in Iraq. USAID, BASICS 11. *Paper, November, 5*. Available at [http://pdf.usaid.gov/pdf\\_docs/PNACW617.pdf](http://pdf.usaid.gov/pdf_docs/PNACW617.pdf) Accessed on February 4, 2016.
- Gilot-Fromont, E., Mercier, A., Gotteland, C., Richomme, C., Aubert, D., Afonso, E., . . . Lélou, M. (2012). *The life cycle of toxoplasma gondii in the natural environment* INTECH Open Access Publisher. Available at <http://dx.doi.org/10.5772/48233> Accessed on January 15, 2016.
- Goldenberg, R. L., & Thompson, C. (2003). The infectious origins of stillbirth. *American journal of obstetrics and gynecology, 189*(3), 861-873. Available at <http://zoologia.biologia.uasnet.mx/protozoos/protozoa16.pdf>. Accessed on February 27, 2016.
- Goldman, M. B., & Hatch, M. C. (2000). Breast cancer epidemiology, treatment, and prevention. ed. G. Ursin. and D.V. Spicer, pp 871-883. New York: Academic Press.
- Golkar, M., Azadmanesh, K., Khalili, G., Khoshkholgh-Sima, B., Babaie, J., Mercier, C., ... & Cesbron-Delauw, M. F. (2008). Serodiagnosis of recently acquired *Toxoplasma gondii* infection in pregnant women using enzyme-linked immunosorbent assays with a recombinant dense granule GRA6 protein. *Diagnostic microbiology and infectious disease, 61*(1), 31-39. Available at <http://www.sciencedirect.com/science/article/pii/S0732889307003902> Accessed on January 4, 2016.
- Gollub, E. L., Leroy, V., Gilbert, R., Chêne, G., & Wallon, M. (2008). Effectiveness of health education on toxoplasma-related knowledge, behaviour, and risk of seroconversion in pregnancy. *European Journal of Obstetrics & Gynecology and Reproductive Biology, 136*(2), 137-145. Available at <http://dx.doi.org/10.1016/j.ejogrb.2007.09.010> Accessed on May 9, 2016.
- Gras, L., Wallon, M., Pollak, A., Cortina-Borja, M., Evengard, B., Hayde, M., ... & Gilbert, R. E. EMSCOT (2005). Association between prenatal treatment and clinical manifestations of congenital toxoplasmosis in infancy: a cohort study in 13 European centers. *Acta Paediatrica Scandinavia, 94*, 1-12. Available at <http://onlinelibrary.wiley.com/doi/10.1111/j.1651-2227.2005.tb01844.x/full> Accessed on March 4, 2016.
- Gutiérrez, J., Roldán, C., & Maroto, M. C. (1995). Seroprevalence of human toxoplasmosis. *Microbios, 85*(343), 73-75. Available at <http://europemc.org/abstract/med/8643032> Accessed on March 4, 2016.
- Hadi, N. J. (2011). Prevalence of antibodies to cytomegalovirus, rubella virus and toxoplasma Gondii among aborted women in Thi-Qar province. *J Educ. Coll, 1*, 3-9. Available at <http://utq.edu.iq/Final2/1.pdf> Accessed on February 4, 2016.
- Haines, T. P., Hill, A., Hill, K. D., McPhail, S., Oliver, D., Brauer, S., . . . Beer, C. (2011). Patient education to prevent falls among older hospital inpatients: A randomized controlled trial. *Archives of Internal Medicine, 171*(6), 516-524.

Available at <http://archinte.jamanetwork.com/article.aspx?articleID=226901> Accessed on May 9, 2016.

- Hall, S., Ryan, K. A., & Buxton, D. (2001). The epidemiological toxoplasmosis infection: in D H M Joynson and T G Wreghitt (ed). *Toxoplasmosis: A Comprehensive Clinical Guide*. Cambridge University press, Cambridge, pp 58-124.
- Hamad, N. R., & Kadir, M. A. (2014). Prevalence and comparison between the efficacy of different techniques for diagnosis of toxoplasma gondii among women in Erbil province-Iraqi kurdistan. *European Scientific Journal*, 9(21). Available at <http://eujournal.org/index.php/esj/article/viewFile/2537/2399>. Accessed on February 27, 2016.
- Hammoud, H. R. (2006). Illiteracy in the Arab world. *Adult Education and Development*, 66, 83. Available at [http://rasit.org/files/Illiteracy\\_in\\_Arab\\_World.pdf](http://rasit.org/files/Illiteracy_in_Arab_World.pdf) Accessed on February 6, 2016.
- Hasan, S. F. (2011). Seroprevalence toxoplasmosis among comers to marriage in Kerbela governorate. *Kerb J Pharm Sci*, 2, 97-102. Kerbela Journal of Pharmaceutical Sciences No.2. pp. 97-105. Available at <http://pharmacy.uokerbala.edu.iq/images/journal/second%20non/9.pdf> Accessed on February 4, 2016.
- Hazzard, A., Dabrow, S., Celano, M., McFadden - Garden, T., & Melhado, T. (2000). Training residents in pediatric literacy: Impact on knowledge, attitudes and practice. *Ambulatory Child Health*, 6(4), 237-246. Available at <http://onlinelibrary.wiley.com/doi/10.1046/j.1467-0658.2000.00085.x/abstract> Accessed on May 9, 2016.
- Hedman, K., Lappalainen, M., Seppäiä, I., & Mäkelä, O. (1989). Recent primary toxoplasma infection indicated by a low avidity of specific IgG. *Journal of infectious diseases*, 159(4), 736-740. Available at <http://jid.oxfordjournals.org/content/159/4/736.short> Accessed on January 4, 2016.
- Hess, J., & Whelan, J. S. (2009). Making health literacy real: Adult literacy and medical students teach each other. *Journal of the Medical Library Association: JMLA*, 97(3), 221-224. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2706439/> Accessed on May 9, 2016.
- Hill, D. E. & Dubey, J.P. (2014). *Toxoplasma gondii. Integrated approach to the detection and control of foodborne parasites and the impact on food safety* (2014th ed., pp. 1389:1-89). USDA: United States Department of Agriculture. Agricultural Research Service. Available at <http://www.ars.usda.gov/pandp/people/people.htm?personid=1472> Accessed on January 15, 2016.
- Hill, D., & Dubey, J. P. (2002). Toxoplasma gondii: transmission, diagnosis and prevention. *Clinical microbiology and infection*, 8(10), 634-640. Available at <http://www.sciencedirect.com/science/article/pii/S1198743X1462509X> Accessed on March 4, 2016.

- Hou, S. I. (2014). Health Education Theoretical Concepts, Effective Strategies and Core Competencies. *Health promotion practice*, 15(5), 619-621. Available at <http://hpp.sagepub.com/content/15/5/619.short> Accessed on October 14, 2016.
- Hughes, J. M., Colley, D. G., Lopez, A., Dietz, V. J., Wilson, M., Navin, T. R., & Jones, J. L. (2000). Preventing congenital toxoplasmosis. *Morbidity and Mortality Weekly Report: Recommendations and Reports*, 57-75. Available at <http://www.jstor.org/stable/42000708> Accessed on January 4, 2016.
- Hunter, C. A., & Sibley, L. D. (2012). Modulation of innate immunity by toxoplasma gondii virulence effectors. *Nature Reviews Microbiology*, 10(11), 766-778. Available at <http://www.nature.com/nrmicro/journal/v10/n11/pdf/nrmicro2858.pdf> Accessed on January 15, 2016.
- Hussain, Z., Yousif, J., & Nassir, K. M. (2010). The detection of toxoplasmosis infection among married women in Al-Abbasiya – Najaf region. - *Al-Kufa Journal for Biology*, Vol.2, No. (2). ISSN: 20738854 23116544 pp: 166-172. Available at <http://www.uokufa.edu.iq/journals/index.php/ajb/article/viewFile/686/597> Accessed on February 5, 2016.
- Hussan, B. M. (2013). Study the prevalence of ACL, APL, CMV, HSV, rubella and toxoplasma gondii in aborted women in Baghdad. *Med J Babylon*, 10, 455-464. Available at <http://www.medicaljb.com/article.aspx?jrid=161>. Accessed on February 27, 2016.
- Iraqi Ministry of Planning and Development Corporation. (24 May 2009, 31 December 2015). IRAQ: Over 20 percent of Iraqis live below the poverty line. *IRIN Middle East News*, pp. 1 Economy. Available at [http://www.cosit.gov.iq/en/press\\_poverty.php](http://www.cosit.gov.iq/en/press_poverty.php) Accessed on February 6, 2016.
- Israelski, D. M., & Remington, J. S. (1993). Toxoplasmosis in patients with cancer. *Clinical infectious diseases*, 17(Supplement 2), S423-S435. Available at [http://cid.oxfordjournals.org/content/17/Supplement\\_2/S423.short](http://cid.oxfordjournals.org/content/17/Supplement_2/S423.short) Accessed on February 4, 2016.
- Jamshidi Makiani, M., Davoodian, P., Golsha, R., & Dehghani, M. (2012). Seroepidemiology and risk factors of toxoplasmosis in the first trimester among pregnant women. *International Electronic Journal of Medicine*, 1(2). Available at [http://eprints.hums.ac.ir/1879/1/1%20\(39\).pdf](http://eprints.hums.ac.ir/1879/1/1%20(39).pdf) Accessed on December 15, 2015.
- Jara, M., Hsu, H. W., Eaton, R. B., & Demaria JR, A. (2001). Epidemiology of congenital toxoplasmosis identified by population-based newborn screening in Massachusetts. *The Pediatric Infectious Disease Journal*, 20(12), 1132-1135. Available at <http://journals.lww.com/pidj/pages/default.aspx> Accessed on May 9, 2016.
- Jekel, J. F., Katz, D. L., Elmore, J. G., & Wild, D. (2007). *Epidemiology, biostatistics and preventive medicine*. Elsevier Health Sciences. Saunders, an imprint of Elsevier Inc. 4<sup>th</sup> edition, Chap. 12, pp 157.
- Jin, S., Chang, Z. Y., Ming, X., Min, C. L., Wei, H., Sheng, L. Y., & Hong, G. X. (2005). Fast dipstick dye immunoassay for detection of immunoglobulin G

- (IgG) and IgM antibodies of human toxoplasmosis. *Clinical and diagnostic laboratory immunology*, 12(1), 198-201. Available at <http://cvi.asm.org/content/12/1/198.long> Accessed on March 4, 2016.
- Jones, J. E. F. F. R. E. Y., Lopez, A., & Wilson, M. (2003). Congenital toxoplasmosis. *American family physician*, 67(10), 2131-2146. Available at <http://www.foedeafdeling.hvidovrehospital.dk> Accessed on January 4, 2016.
- Jones, J. L., Kruszon-Moran, D., & Wilson, M. (2003). Toxoplasma gondii infection in the united states, 1999-2000. *Emerging Infectious Diseases*, 9(11), 1371-1374. Available at [http://wwwnc.cdc.gov/eid/article/9/11/03-0098\\_article](http://wwwnc.cdc.gov/eid/article/9/11/03-0098_article) Accessed on August 19, 2015.
- Jones, J. L., Kruszon-Moran, D., Wilson, M., McQuillan, G., Navin, T., & McAuley, J. B. (2001). Toxoplasma Gondii infection in the United States: Seroprevalence and risk factors. *American Journal of Epidemiology*, 154 (4), 357-365. Available at <https://aje.oxfordjournals.org/content/154/4/357.full> Accessed on February 4, 2016.
- Jukkala, A., Deupree, J. P., & Graham, S. (2009). Knowledge of limited health literacy at an academic health center. *The Journal of Continuing Education in Nursing*, 40(7), 298-302. Available at <http://www.healio.com/nursing/journals/jcen> Accessed on May 9, 2016.
- Jumaian, N. (2005). Seroprevalence and risk factors for toxoplasma infection in pregnant women in Jordan. *Eastern Mediterranean Health Journal*, 11(1/2), 45. Available at [http://apps.who.int/iris/bitstream/10665/116916/1/11\\_1-2\\_2005\\_45\\_51.pdf](http://apps.who.int/iris/bitstream/10665/116916/1/11_1-2_2005_45_51.pdf) Accessed on February 15, 2016.
- Kadir, M. A., Ghalib, A. K., Othman, N. F., & Ahmed, I. S. (2011). Seroprevalence of toxoplasma gondii among pregnant women in Kirkuk, Iraq. *J Kir Univ-Sci Studies*, 6, 1-11. Available at <http://en.kujss.com/files/pdf/vol6%20no.2/43107.pdf> Accessed on February 6, 2016.
- Kapperud, G., Jenum, P. A., Stray-Pedersen, B., Melby, K. K., Eskild, A., & Eng, J. (1997). Risk factors for Toxoplasma gondii infection in pregnancy: results of a prospective case-control study in Norway. *Obstetrical & gynecological survey*, 52(3), 158-159. Available at <http://aje.oxfordjournals.org/content/144/4/405.full.pdf> Accessed on January 4, 2016.
- Katzer, F., Burrells, A., & Opsteegh, M. (2014). Chapter twenty-one - toxoplasma gondii. In Gray, Steven L. Percival Marylynn V. Yates David W. Williams Rachel M. Chalmers Nicholas F. (Ed.), *Microbiology of waterborne diseases (second edition)* (pp. 417-440). London: Academic Press. Available at <http://dx.doi.org/10.1016/B978-0-12-415846-7.00021-4> Accessed on January 15, 2016.
- Khalil, H. I. (2008). Some aspects in Seroprevalence, diagnosis and influence of sex hormones on immunity during human toxoplasmosis. Ph.D. Thesis, College of Medicine. University of AL-Mustansiriyah. Pp: 173.

- Khandelwal, K. C., & Mahdi, S. S. (1988). *Biogas technology: a practical handbook*. Tata McGraw-Hill. V.1. publishing company limited. 12\4 Asaf Ali Road. New delhi. 110002.
- Kila, R. (1996). Dilemmas in AIDS care. *Papua and New Guinea medical journal*, 39(3), 218. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9795567> Accessed on January 4, 2016.
- Kimbita, E. N., Xuan, X., Huang, X., Miyazawa, T., Fukumoto, S., Mishima, M., ... & Suzuki, N. (2001). Serodiagnosis of *Toxoplasma gondii* infection in cats by enzyme-linked immunosorbent assay using recombinant SAG1. *Veterinary Parasitology*, 102(1), 35-44. Available at <http://www.sciencedirect.com/science/article/pii/S0304401701005222> Accessed on March 4, 2016.
- Kirsch, I. S. (1993). *Adult literacy in America: A first look at the results of the national adult literacy survey*. ERIC. Available at <http://eric.ed.gov/?id=ED358375> Accessed on May 9, 2016.
- Kravetz, J. D., & Federman, D. G. (2005). Prevention of toxoplasmosis in pregnancy: knowledge of risk factors. *Infectious diseases in obstetrics and gynecology*, 13(3), 161-165. Available at <http://www.hindawi.com/journals/idog/2005/753406/abs/> Accessed on January 4, 2016.
- Kravetz, J. D., & Federman, D. G. (2005). Toxoplasmosis in pregnancy. *The American journal of medicine*, 118(3), 212-216. Available at <http://www.migato.net/conocele/docs/Kravetz2005a.pdf> Accessed on January 4, 2016.
- Kutner, M., Greenburg, E., Jin, Y., & Paulsen, C. (2006). The health literacy of America's adults: Results from the 2003 national assessment of adult literacy. NCES 2006-483. *National Center for Education Statistics*. Available at <http://eric.ed.gov/?id=ED493284> Accessed on May 9, 2016.
- Lafferty, K. D. (2006). Can the common brain parasite, *Toxoplasma gondii*, influence human culture? *Proceedings. Biological Sciences / the Royal Society*, 273(1602), 2749-2755. Available at <http://rspb.royalsocietypublishing.org/content/273/1602/2749.short> Accessed on February 15, 2016.
- Lakhanpal, V., Schocket, S. S., & Nirankari, V. S. (1983). Clindamycin in the treatment of toxoplasmic retinochoroiditis. *American journal of ophthalmology*, 95(5), 605-613. Available at <http://ezproxy.upm.edu.my:2055/science/article/pii/0002939483903781> Accessed on January 4, 2016.
- Lebech, M., Andersen, O., Christensen, N. C., Hertel, J., Nielsen, H. E., Peitersen, B., ... & Danish Congenital Toxoplasmosis Study Group. (1999). Feasibility of neonatal screening for toxoplasma infection in the absence of prenatal treatment. *The Lancet*, 353(9167), 1834-1837. Available at <http://www.sciencedirect.com/science/article/pii/S0140673698112813> Accessed on October 19, 2016.

- Lehrer, E. W., Fredebaugh, S. L., Schooley, R. L., & Mateus-Pinilla, N. E. (2010). Prevalence of antibodies to *Toxoplasma gondii* in woodchucks across an urban-rural gradient. *Journal of Wildlife Diseases*, 46(3), 977-980. Available at <http://dx.doi.org/10.7589/0090-3558-46.3.977> Accessed on February 6, 2016.
- Lelong, B., Rahelimino, B., Candolfi, E., Ravelojaona, B. J., Villard, O., Rasamindrakotroka, A. J., & Kien, T. (1994). [Prevalence of toxoplasmosis in a population of pregnant women in Antananarivo (Madagascar)]. *Bulletin de la Societe de pathologie exotique (1990)*, 88(1), 46-49. Available at <http://europepmc.org/abstract/med/7787454> Accessed on February 15, 2016.
- Lélu, M., Langlais, M., Poulle, M. L., & Gilot-Fromont, E. (2010). Transmission dynamics of *Toxoplasma gondii* along an urban-rural gradient. *Theoretical population biology*, 78(2), 139-147. Available at <http://dx.doi.org/10.1016/j.tpb.2010.05.005> Accessed on February 6, 2016.
- Li, X., Wei, H., Zhang, H., Peng, H., & Lindsay, D. S. (2014). A meta-analysis on risks of adverse pregnancy outcomes in *Toxoplasma gondii* infection. *PloS One*, 9(5), e97775. Available at <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0097775> Accessed on January 15, 2016.
- Liesenfeld, O., Montoya, J. G., Kinney, S., Press, C., & Remington, J. S. (2001). Effect of testing for IgG avidity in the diagnosis of *Toxoplasma gondii* infection in pregnant women: experience in a US reference laboratory. *Journal of Infectious Diseases*, 183(8), 1248-1253. Available at <http://jid.oxfordjournals.org/content/183/8/1248.full> Accessed on March 4, 2016.
- Lindau, S. T., Tomori, C., McCarville, M., & Bennett, C. (2001). Improving rates of cervical cancer screening and pap smear follow-up for low-income women with limited health literacy. *Cancer Investigation*, 19(3), 316-323. Available at <http://www.tandfonline.com/doi/abs/10.1081/CNV-100102558> Accessed on May 9, 2016.
- Link, B. G. (2008). Epidemiological sociology and the social shaping of population health. *Journal of Health and Social Behavior*, 49(4), 367-384. Available at <http://hsb.sagepub.com/content/49/4/367.full.pdf> Accessed on February 6, 2016.
- Liu, Q., Wei, F., Gao, S., Jiang, L., Lian, H., Yuan, B., . . . Zhu, X. Q. (2009). *Toxoplasma gondii* infection in pregnant women in china. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 103(2), 162-166. Available at <http://trstmh.oxfordjournals.org/content/103/2/162.short> Accessed on May 9, 2016.
- Longstaffe, J. A. (1984). Helminths, arthropods and protozoa of domesticated animals (7th edition): E. J. L. Soulsby, 1982. London: Bailliere Tindall, 809 pp., illus. ISBN 0-7020-0820-6. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 78(3), 329. Available at [http://ezproxy.upm.edu.my:2112/10.1016/0035-9203\(84\)90110-X](http://ezproxy.upm.edu.my:2112/10.1016/0035-9203(84)90110-X) Accessed on March 4, 2016.

- Luft, B. J., Hafner, R., Korzun, A. H., Leport, C., Antoniskis, D., Bosler, E. M., . . . Jacobson, J. (1993). Toxoplasmic encephalitis in patients with the acquired immunodeficiency syndrome. *New England Journal of Medicine*, 329(14), 995-1000. Available at <http://www.nejm.org/doi/pdf/10.1056/NEJM199309303291403> Accessed on February 17, 2016.
- Macintyre, K., Rutenberg, N., Brown, L., & Karim, A. (2004). Understanding perceptions of HIV risk among adolescents in KwaZulu-Natal. *AIDS and Behavior*, 8(3), 237-250. Available at <http://link.springer.com/article/10.1023/B:AIBE.0000044072.71361.b3> Accessed on May 9, 2016.
- Mahler, H., Epp, J., Franklin, W., & Kickbusch, I. (1986). Ottawa charter for health promotion. *Health Promotion International*, 1(4), 405. Available at <http://heapro.oxfordjournals.org/content/1/4> Accessed on October 14, 2016.
- Mahmood, S. H., AL-Qadhi, B. N., & Zghair, K. H. (2013). Prevalence of Toxoplasmosis of Males Blood Donors in Baghdad. *Iraqi Journal of Science*, Vol 54, No.4, pp:832-841. Available at <http://ijs.scbaghdad.edu.iq/issues/Vol54/No4/Vol54Y2013No4P832-841.pdf> Accessed on February 4, 2016.
- Martinez Sanchez, R., Bacallao Gordo, R., Alberti Amador, E., & Alfonso Berrio, L. (1994). Prevalence of toxoplasmosis in pregnant women of the province of la Habana. *Revista do Instituto De Medicina Tropical De Sao Paulo*, 36(5), 445-450. Available at <http://europepmc.org/abstract/med/7569612> Accessed on February 15, 2016.
- Meerburg, B. G., & Kijlstra, A. (2009). Changing climate—changing pathogens: *Toxoplasma gondii* in North-Western Europe. *Parasitology research*, 105(1), 17-24. Available at <http://link.springer.com/article/10.1007/s00436-009-1447-4> Accessed on October 16, 2016.
- Meroni, V., & Genco, F. (2008). Toxoplasmosis in pregnancy: evaluation of diagnostic methods. *Parassitologia*, 50(1/2), 51. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18693557> Accessed on March 4, 2016.
- Millar, P. R., Moura, F. L. d., Bastos, O. M. P., Mattos, Danuza Pinheiro Bastos Garcia de, Fonseca, A. B. M., Sudré, A. P., . . . Amendoeira, M. R. R. (2014). Toxoplasmosis-related knowledge among pregnant and postpartum women attended in public health units in Niterói, Rio de Janeiro, Brazil. *Revista do Instituto De Medicina Tropical De São Paulo*, 56(5), 433-438. Available at <http://dx.doi.org/10.1590/S0036-46652014000500011> Accessed on May 9, 2016.
- Mishra, G., Thakur, S., Singhal, P., Ghosh, S. N., Chauhan, D., & Jayam, C. (2016). Assessment of child behavior in dental operatory in relation to sociodemographic factors, general anxiety, body mass index and role of multi media distraction. *Journal of Indian Society of Pedodontics and Preventive Dentistry*, 34(2), 159. Available at <http://www.jisppd.com/article.asp?issn=09704388;year=2016;volume=34;issue=2;page=159;epage=164;aulast=Mishra> Accessed on February 6, 2016.
- Mohamed, N. A., & Ibrahim, H. D. F. (2012). Evaluate the health education program for Pregnant women about toxoplasma infection at qena university



- hospital. *Journal of American Science*, 8(12), 1306-1315. Available at <http://www.jofamericanscience.org/> Accessed on January 4, 2016.
- Mohammad, E. A. K., & Salman, Y. J. (2014). Study of TORCH infections in women with bad obstetric history (BOH) in Kirkuk city. *Int.J.Curr.Microbiol.App.Sci*, 3(10), 700-709. Available at <http://www.ijcmas.com/>. Accessed on February 27, 2016.
- Mohammad, H. A., Amin, T., Balaha, M., & Moghannum, M. A. (2010). Toxoplasmosis among the pregnant women attending a Saudi maternity hospital: Seroprevalence and possible risk factors. *Annals of Tropical Medicine and Parasitology*, 104(6), 493-504. Available at <http://www.tandfonline.com/doi/abs/10.1179/136485910X12786389891443> Accessed on February 6, 2016.
- Mohammad, M., Ahmed, S., & Hussain, A. (2013). Seroprevalence of toxoplasma gondii between couples in Ramadi city using enzyme linked Immunosorbent Assay (ELISA). *International Journal*, 5(6), 295-299. Available at <http://www.academicjournals.org/journal/IJMMS/article-full-text-pdf/2007C1338>. Accessed on February 17, 2016.
- Mohammed, T. K. (2011). Seroprevalence of Toxoplasma gondii among pregnant women in Baghdad city. *Iraq Acad Sci J*, 24(24), 21-28. Available at <http://www.iasj.net/iasj?func=fulltext&aId=29334> Accessed on January 4, 2016.
- Moncada, P. A., & Montoya, J. G. (2012). Toxoplasmosis in the fetus and newborn: an update on prevalence, diagnosis and treatment. *Expert review of anti-infective therapy*, 10(7), 815-828. Available at <http://www.tandfonline.com/doi/abs/10.1586/eri.12.58>. Accessed on February 27, 2016.
- Montoya, J. G., & Remington, J. S. (2000). "Toxoplasma gondii". In: Mandell G., Dolin, A., and Bennett, J., (eds). Principles and Practice of Infectious Diseases. Fifth Edition. Pp 2858-88. Available at <http://www.pamf.org/serology/bibliography.html> Accessed on February 4, 2016.
- Morampudi, V., & Braun, M. (2010). Study of the modulation of innate immune responses in intestinal epithelial cells by toxoplasma gondii and its correlation with parasite virulence (Master Thesis). Interuniversity Library of the French Community of Belgium- Free University of Bruxelles. Available at [http://theses.ulb.ac.be/ETD\\_db/collection/available/ULBetd-05252010-151346](http://theses.ulb.ac.be/ETD_db/collection/available/ULBetd-05252010-151346) Accessed on January 28, 2016.
- Moura, L. D., Oliveira, Lilian Maria Garcia Bahia, Wada, M. Y., Jones, J. L., Tuboi, S. H., Carmo, E. H., . . . Graça, R. M. (2006). Waterborne toxoplasmosis, brazil, from field to gene. Available at <http://dx.doi.org/10.1590/0074-02760150262> Accessed on February 15, 2016.
- Nabias, R., Ngouamizokou, A., Migot-Nabias, F., Mbou-Moutsimbi, R. A., & Lansoud-Soukate, J. (1997). [Serological investigation of toxoplasmosis in patients of the MIP center of Franceville (Gabon)]. *Bulletin de la Societe de*

- pathologie exotique* (1990), 91(4), 318-320. Available at <http://europepmc.org/abstract/med/9846226> Accessed on February 15, 2016.
- Nascimento, F. S., Suzuki, L. A., & Rossi, C. L. (2008). Assessment of the value of detecting specific IgA antibodies for the diagnosis of a recently acquired primary Toxoplasma infection. *Prenatal diagnosis*, 28(8), 749-752. Available at <http://onlinelibrary.wiley.com/doi/10.1002/pd.2052/abstract> Accessed on January 4, 2016.
- Nasir, I. A., Aderinsayo, A. H., Mele, H. U., & Aliyu, M. M. (2015). Prevalence and associated risk factors of toxoplasma gondii antibodies among pregnant women attending Maiduguri teaching hospital, Nigeria. *Journal of Medical Sciences*, 15(3), 147. Available at <http://scialert.net/abstract/?doi=jms.2015.147.154> Accessed on February 15, 2016.
- Ndumbe, P. M., Andela, A., Nkemnkeng-Asong, J., Watonsi, E., & Nyambi, P. (1992). Prevalence of infections affecting the child among pregnant women in Yaoundé, Cameroon. *Medical Microbiology and Immunology*, 181(3), 127-130. Available at <http://link.springer.com/article/10.1007/BF00202052> Accessed on February 15, 2016.
- Newton, L. H., & Hall, S. M. (1994). Survey of local policies for prevention of congenital toxoplasmosis. *Communicable disease report. CDR review*, 4(10), R121-4. Available at <http://europepmc.org/abstract/med/7527277> Accessed on October 19, 2016.
- Newton, L. H., & Hall, S. M. (1995). A survey of health education material for the primary prevention of congenital toxoplasmosis. *Communicable Disease Report. CDR Review*, 5(2), R21-7. Available at <http://europepmc.org/abstract/med/7532519> Accessed on May 9, 2016.
- Novotna, M., Hanusova, J., Klose, J., Preiss, M., Havlicek, J., Roubalova, K., & Flegr, J. (2005). Probable neuroimmunological link between toxoplasma and cytomegalovirus infections and personality changes in the human host. *BMC Infectious Diseases*, 5, 54. Available at <http://www.biomedcentral.com/1471-2334/5/54> Accessed on February 4, 2016.
- Ocak, S., Duran, N., Eskiocak, A. F., & Aytac, H. (2005). Anti-toxoplasma gondii antibodies in hemodialysis patients receiving long-term hemodialysis therapy in turkey. *Saudi Medical Journal*, 26(9), 1378-1382. Available at <http://smj.psmmc.med.sa/index.php/smj/article/view/5493> Accessed on January 4, 2016.
- O'Connor, V., & Kovacs, G. (2003). *Obstetrics, gynaecology and women's health*. Cambridge University Press.
- Ogunmodede, F., Scheftel, J., Jones, J. L., & Lynfield, R. (2005). Toxoplasmosis prevention knowledge among pregnant women in Minnesota. *Minnesota medicine*, 88(2), 32-34. Available at <http://europepmc.org/abstract/med/17886796> Accessed on January 4, 2016.
- Olariu, T. R., Remington, J. S., McLeod, R., Alam, A., & Montoya, J. G. (2011). Severe congenital toxoplasmosis in the United States: clinical and serologic

- findings in untreated infants. *The Pediatric infectious disease journal*, 30(12), 1056-1061. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21956696> Accessed on March 4, 2016.
- Pappas, G., Roussos, N., & Falagas, M. E. (2009). Toxoplasmosis snapshots: Global status of toxoplasma Gondii seroprevalence and implications for pregnancy and congenital toxoplasmosis. *International Journal for Parasitology*, 39 (12), 1385-1394. Available at <http://dx.doi.org/10.1016/j.ijpara.2009.04.003>. Accessed on February 4, 2016.
- Paquet, C., Yudin, M. H., Allen, V. M., Bouchard, C., Boucher, M., Caddy, S., . . . Ogilvie, G. (2013). Toxoplasmosis in pregnancy: Prevention, screening, and treatment. *Journal of Obstetrics and Gynaecology Canada*, 35(1), 78-79. Available at <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.661.6294&rep=rep1&type=pdf> Accessed on December 15, 2015.
- Patz, J. A., Graczyk, T. K., Geller, N., & Vittor, A. Y. (2000). Effects of environmental change on emerging parasitic diseases. *International journal for parasitology*, 30(12), 1395-1405. Available at <http://www.sciencedirect.com/science/article/pii/S0020751900001417> Accessed on October 16, 2016.
- Pawlowski, Z. S., Gromadcka-Sutkiewicz, M., Skommer, J., Paul, M., Rokossowski, H., Suchocka, E., & Schantz, P. M. (2001). Impact of health education on knowledge and prevention behavior for congenital toxoplasmosis: the experience in Poznan, Poland. *Health Education Research*, 16(4), 493-502. Available at <http://her.oxfordjournals.org/content/16/4/493.long> Accessed on January 4, 2016.
- Peadon, E., Payne, J., Henley, N., D'Antoine, H., Bartu, A., O'Leary, C., ... & Elliott, E. J. (2011). Attitudes and behaviour predict women's intention to drink alcohol during pregnancy: the challenge for health professionals. *BMC Public Health*, 11(1), 1. Available at <http://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-584> Accessed on January 4, 2016.
- Pelloux, H., Brun, E., Vernet, G., Marcillat, S., Jolivet, M., Guergour, D., ... & Ambroise-Thomas, P. (1998). Determination of anti-Toxoplasma gondii immunoglobulin G avidity: adaptation to the Vidas system (bioMérieux). *Diagnostic microbiology and infectious disease*, 32(2), 69-73. Available at <http://www.sciencedirect.com/science/article/pii/S0732889398000777> Accessed on March 4, 2016.
- Penney, D. P., Powers, J. M., Frank, M., Willis, C., & Churukian, C. (2002). Analysis and testing of biological stains--the biological stain commission procedures. *Biotechnic & histochemistry*, 77(5-6), 237-275. Available at <http://www.tandfonline.com/doi/abs/10.1080/bih.77.5-6.237.275> Accessed on March 4, 2016.
- Pereboom, M. T., Manniën, J., Spelten, E. R., Schellevis, F. G., & Hutton, E. K. (2013). Observational study to assess pregnant women's knowledge and behaviour to prevent toxoplasmosis, listeriosis and cytomegalovirus. *BMC pregnancy and childbirth*, 13(1), 1. Available at

<http://www.biomedcentral.com/1471-2393/13/98> Accessed on January 4, 2016.

- Petersson, K., Stray-Pedersen, B., Malm, G., Forsgren, M., & Evengård, B. (2000). Seroprevalence of toxoplasma gondii among pregnant women in Sweden. *Acta Obstetrica Et Gynecologica Scandinavica*, 79(10), 824-829. Available at <http://www.tandfonline.com/doi/abs/10.1080/00016340009169209> Accessed on August 19, 2015.
- Pinon, J. M., Dumon, H., Chemla, C., Franck, J., Petersen, E., Lebech, M., ... & Johnson, J. (2001). Strategy for diagnosis of congenital toxoplasmosis: evaluation of methods comparing mothers and newborns and standard methods for postnatal detection of immunoglobulin G, M, and A antibody. *Journal of clinical microbiology*, 39(6), 2267-2271. Available at <http://jcm.asm.org/content/39/6/2267.full> Accessed on March 4, 2016.
- Powers, B. J., Trinh, J. V., & Bosworth, H. B. (2010). Can this patient read and understand written health information? *Jama*, 304(1), 76-84. Available at <http://jama.jamanetwork.com/article.aspx?articleid=186175> Accessed on May 9, 2016.
- Prabhakar, P., Bailey, A., Smikle, M. F., & Ashley, D. (1992). Seroprevalence of cytomegalovirus infection in a selected population in Jamaica. *The West Indian Medical Journal*, 41(4), 133-135. Available at <http://europepmc.org/abstract/med/1337805> Accessed on February 15, 2016.
- Pratkanis, A. R., Breckler, S. J., & Greenwald, A. G. (2014). *Attitude structure and function*. Psychology Press. Available at <https://books.google.com.my/books> Accessed on January 4, 2016.
- Prislin, R. A. D. M. I. L. A., & Crano, W. D. (2008). Attitudes and attitude change. *Attitudes and attitude change*, 3-15. Available at <https://books.google.com.my/books> Accessed on January 4, 2016.
- Pullan, R. L., Sturrock, H. J., Magalhaes, R. J. S., Clements, A. C., & Brooker, S. J. (2012). Spatial parasite ecology and epidemiology: a review of methods and applications. *Parasitology*, 139(14), 1870-1887. Available at <https://www.cambridge.org/core/services/aop-cambridge-core/content/view/S0031182012000698> Accessed on October 14, 2016.
- Remington J. S., McLeod, R., Thulliez, P., & Desmonts, G., (Ed.) (2001). *Toxoplasmosis. in: Remington JS, Klein J, eds. infectious diseases of the fetus and newborn infant* (5th ed.). Philadelphia: W.B. Saunders. pp 205–346.
- Remington JS, Desmonts G. (1990). Toxoplasmosis. In Remington JS, Klein JO (ed), *Infectious diseases of the fetus and newborn infants*. WB Saunders, Philadelphia. PA. pp 89-195. Available at <https://books.google.com.my/books> Accessed on March 4, 2016.
- Remington, J. S., Thulliez, P., & Montoya, J. G. (2004). Recent developments for diagnosis of toxoplasmosis. *Journal of Clinical Microbiology*, 42 (3), 941-

945. Available at <http://jcm.asm.org/content/42/3/941.short> Accessed on February 4, 2016.
- Robert-Gangneux, F., Aubert, D., & Villena, I. (2015). Toxoplasmosis: A widespread zoonosis diversely affecting humans and animals. *Zoonoses-infections affecting humans and animals* (pp. 355-376) Springer. Available at [http://link.springer.com/chapter/10.1007/978-94-017-9457-2\\_14](http://link.springer.com/chapter/10.1007/978-94-017-9457-2_14) Accessed on January 15, 2016.
- Rodier, M., Berthonneau, J., Bourgoin, A., Giraudeau, G., Agius, G., Burucoa, C., . . . Jacquemin, J. (1995). Seroprevalences of toxoplasma, malaria, rubella, cytomegalovirus, HIV and treponemal infections among pregnant women in Cotonou, republic of Benin. *Acta Tropica*, *59*(4), 271-277. Available at <http://www.sciencedirect.com/science/article/pii/0001706X9500087U> Accessed on February 15, 2016.
- Roelke, M. E., Forrester, D. J., Jacobson, E. R., Kollias, G. V., Scott, F. W., Barr, M. C., ... & Pirtle, E. C. (1993). Seroprevalence of infectious disease agents in free-ranging Florida panthers (*Felis concolor coryi*). *Journal of Wildlife Diseases*, *29*(1), 36-49. Available at <http://www.bioone.org/doi/abs/10.7589/0090-3558-29.1.36> Accessed on March 4, 2016.
- Roitt, I., Brostoff, J., & Male, D. (2001). Immunology. 6th eds. USA, Mosby, 323-383.
- Rorman, E., Zamir, C. S., Rilkis, I., & Ben-David, H. (2006). Congenital toxoplasmosis—prenatal aspects of toxoplasma gondii infection. *Reproductive Toxicology*, *21*(4), 458-472. Available at <http://dx.doi.org/10.1016/j.reprotox.2005.10.006> Accessed on January 4, 2016.
- Rosenthal, M. S., Werner, M. J., & Dubin, N. H. (2004). The effect of a literacy training program on family medicine residents. *Family Medicine-Kansas City*, *36*(8), 582-587. Available at <https://www.stfm.org/fmhub/fm2004/September/Marjorie582.pdf> Accessed on May 9, 2016.
- Ross, D. S., Jones, J. L., & Lynch, M. F. (2006). Toxoplasmosis, cytomegalovirus, listeriosis, and preconception care. *Maternal and Child Health Journal*, *10*(1), 189-193. Available at <http://link.springer.com/article/10.1007/s10995-006-0092-0> Accessed on May 9, 2016.
- Ross, D. S., Victor, M., Sumartojo, E., & Cannon, M. J. (2008). Women's knowledge of congenital cytomegalovirus: results from the 2005 HealthStyles™ survey. *Journal of women's health*, *17*(5), 849-858. Available at <http://online.liebertpub.com/doi/abs/10.1089/jwh.2007.0523> Accessed on January 4, 2016.
- Sackett, D. L., Straus, S. E., Richardson, W. S., Rosenberg, W., & Haynes, R. B. (2000). How to practice and teach EBM. *Edinburgh: Churchill Livingstone*. Available at <http://ktclearinghouse.ca/cebml/products> Accessed on January 4, 2016.

- Sahm, D. F., & Weissfeld, A. S. (2007). *Study Guide for Bailey & Scott's Diagnostic Microbiology*. Mosby. Available at <http://www.store.elsevierhealth.com/asia/> Accessed on March 4, 2016.
- Saker, L., Lee, K., Cannito, B., Gilmore, A., & Campbell-Lendrum, D. H. (2004). Globalization and infectious diseases: A review of the linkages. Available at <http://apps.who.int/iris/handle/10665/68726> Accessed on February 6, 2016.
- Salih, H. (2010). Prevalence of toxoplasmosis among pregnant women in Najaf city. *Kufa J Vet Med Sci*, 1, 101-108. Available at <http://uokufa.edu.iq/journals/index.php/kjvs/article/view/2399> Accessed on February 4, 2016.
- Salman, Y. J. (2014). Role of toxoplasma gondii and human herpes simplex virus type-2 in women with abortions and congenital abnormalities in Kirkuk city. *International Journal of Current Research in Biosciences and Plant Biology*, 1(2), pp. 1-8. doi:ISSN: 2349-8080. Available at <http://www.ijcrbp.com/vol-1-2/Yahya%20Jirjees%20Salman.pdf>. Accessed on February 27, 2016.
- Samaka, H. M., Alhatami, A. O., & Mohammed, K. G. (2013). Evaluation of the toxocell latex agglutination test as screening test and distribution of toxoplasma antibodies among sexually active people in Najaf city. *Kufa Journal for Nursing Sciences*, 2(1). Available at [http://www.uokufa.edu.iq/journals/index.php/kjns/article/view/139/pdf\\_89](http://www.uokufa.edu.iq/journals/index.php/kjns/article/view/139/pdf_89) Accessed on February 4, 2016.
- Savva, M., Duda, E., & Huang, L. (1999). A genetically modified recombinant tumor necrosis factor- $\alpha$  conjugated to the distal terminals of liposomal surface grafted polyethyleneglycol chains. *International journal of pharmaceuticals*, 184(1), 45-51. Available at <http://www.sciencedirect.com/science/article/pii/S0378517399000927> Accessed on January 4, 2016.
- Schlichting, J. A., Quinn, M. T., Heuer, L. J., Schaefer, C. T., Drum, M. L., & Chin, M. H. (2007). Provider perceptions of limited health literacy in community health centers. *Patient Education and Counseling*, 69(1), 114-120. Available at <http://www.sciencedirect.com/science/article/pii/S0738399107003096> Accessed on May 9, 2016.
- Sclar, D. A., Chin, A., Skaer, T. L., Okamoto, M. P., Nakahiro, R. K., & Gill, M. A. (1990). Effect of health education in promoting prescription refill compliance among patients with hypertension. *Clinical therapeutics*, 13(4), 489-495. Available at <http://europemc.org/abstract/med/1934001> Accessed on May 9, 2016.
- Setta, A. A., & Yamani, R. (2008). Prevalence of toxoplasmosis in non-pregnant women in Tripoli, Libya. *The Egyptian J Oumal of Hospital Medicine*, 31, 198-202. Available at [http://egyptianjournal.net78.net/31\\_4.pdf](http://egyptianjournal.net78.net/31_4.pdf) Accessed on March 15, 2016.
- Shefer, A., Atkinson, W., Friedman, C., Kuhar, D. T., Mootrey, G., Bialek, S. R., & Lorick, S. A. (2011). Immunization of health-care personnel: recommendations of the Advisory Committee on Immunization Practices

- (ACIP). *MMWR Recomm Rep*, 60, 1-45. Accessed on [http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6007a1.htm?s\\_cid%3Drr6007a1\\_x](http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6007a1.htm?s_cid%3Drr6007a1_x) Accessed on October 10, 2016.
- Shin, D., Cha, D., Hua, Q. J., Cha, G., & Lee, Y. (2009). Seroprevalence of toxoplasma gondii infection and characteristics of seropositive patients in general hospitals in Daejeon, Korea. *The Korean Journal of Parasitology*, 47(2), 125-130. Available at <http://synapse.koreamed.org/> Accessed on February 15, 2016.
- Sibley, L. D., Khan, A., Ajioka, J. W., & Rosenthal, B. M. (2009). Genetic diversity of toxoplasma gondii in animals and humans. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 364(1530), 2749-2761. Available at <http://rstb.royalsocietypublishing.org/content/364/1530/2749.short> Accessed on January 15, 2016.
- Slomko, Z. and Szczapa, J. (1997). How to avoid infectious diseases in pregnancy. in pawlowski, Z. (ed.), przewodnik dla przyszłych matek (information for mothers-to-be) [in polish]. polskie towarzystwo osóby zdrowotnej, Poznań, Poland, *health education research*, 16(4), 493-502. pp. 1–15.
- Snelling, W. J., Lin, Q., Moore, J. E., Millar, B. C., Tosini, F., Pozio, E., ... & Lowery, C. J. (2007). Proteomics analysis and protein expression during sporozoite excystation of *Cryptosporidium parvum* (Coccidia, Apicomplexa). *Molecular & cellular proteomics*, 6(2), 346-355. Available at <http://www.mcponline.org/content/6/2/346.full> Accessed on January 4, 2016.
- Sood, N., Soni, S., Vegad, M., & Gupta, P. (2009). Seroprevalence of Toxoplasma gondii in women with bad obstetric history in Ahmedabad. *Gujarat Med J*, 64(64), 35-37. Available at <http://medind.nic.in/gaa/t09/i2/gaat09i2p35.pdf> Accessed on January 4, 2016.
- Sroka, S., Bartelheimer, N., Winter, A., Heukelbach, J., Ariza, L., Ribeiro, H., . . . Liesenfeld, O. (2010). Prevalence and risk factors of toxoplasmosis among pregnant women in Fortaleza, northeastern Brazil. *The American Journal of Tropical Medicine and Hygiene*, 83(3), 528-533. Available at <http://www.ajtmh.org/content/83/3/528.short> Accessed on February 6, 2016.
- Stanley, K. (2007). Design of randomized controlled trials. *Circulation*, 115(9), 1164-1169. Available at <http://circ.ahajournals.org/content/115/9/1164.full> Accessed on January 4, 2016.
- Sultan, B.A., Al-Klaby, R.F., Obaid, R.F. (2010). Sero-diagnosis of toxoplasmosis among pregnant women and immunocompromised patients. *Kufa med. Journal*. Vol.13.No.2. Available at [https://www.researchgate.net/profile/Baqur\\_Sultan/publication/274316525](https://www.researchgate.net/profile/Baqur_Sultan/publication/274316525) Accessed on February 4, 2016.
- Suzuki, Y. (2002). Host resistance in the brain against *Toxoplasma gondii*. *Journal of Infectious Diseases*, 185(Supplement 1), S58-S65. Available at [http://jid.oxfordjournals.org/content/185/Supplement\\_1/S58.short](http://jid.oxfordjournals.org/content/185/Supplement_1/S58.short) Accessed on February 4, 2016.

- Taher, J. H. (2011). Seroepidemiological aspects of toxoplasmosis among pre-school children in Najaf province. *Al-Kufa Journal for Biology*, 3 (1). Available at <http://www.uokufa.edu.iq/journals/index.php/ajb/article/viewFile/714/623> Accessed on January 22, 2016.
- Tamma P., & Serwint J.R. (2007). Toxoplasmosis. *Pediatrics in Review*, 28(12), 470-471. doi:10.1542/pir.28-12-470.
- Tasevska, N., Cross, A. J., Dodd, K. W., Ziegler, R. G., Caporaso, N. E., & Sinha, R. (2011). No effect of meat, meat cooking preferences, meat mutagens or heme iron on lung cancer risk in the prostate, lung, colorectal and ovarian cancer screening trial. *International journal of cancer*, 128(2), 402-411. Accessed on <http://onlinelibrary.wiley.com/doi/10.1002/ijc.25327/full> Accessed on October 10, 2016.
- Taylor, M. R., Lennon, B., Holland, C. V., & Cafferkey, M. (1997). Community study of Toxoplasma antibodies in urban and rural schoolchildren aged 4 to 18 years. *Archives of disease in childhood*, 77(5), 406-409. Available at <http://adc.bmj.com/content/77/5/406.long> Accessed on February 4, 2016.
- Tenter, A. M., Heckeroth, A. R., & Weiss, L. M. (2000). Toxoplasma gondii: From animals to humans. *International Journal for Parasitology*, 30(12-13), 1217-1258. Available at [http://dx.doi.org/10.1016/S0020-7519\(00\)00124-7](http://dx.doi.org/10.1016/S0020-7519(00)00124-7) Accessed on February 17, 2016.
- The World Bank. IBRD. IDA. (2015). Overview. Working for a world free of poverty, Iraq. *Countries / Iraq*, Available at <http://www.worldbank.org/en/country/iraq> Accessed on February 6, 2016.
- Torrey, E. F., & Yolken, R. H. (2003). Toxoplasma gondii and schizophrenia. *Emerging infectious diseases*, 9(11), 1375. Available at <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3035534/> Accessed on February 4, 2016.
- Torrey, E. F., Bartko, J. J., & Yolken, R. H. (2012). Toxoplasma gondii and other risk factors for schizophrenia: An update. *Schizophrenia Bulletin*, 38(3), 642-647. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22446566> Accessed on December 15, 2015.
- Tung, W., Ding, K., & Farmer, S. (2008). Knowledge, attitudes, and behaviors related to toxoplasmosis among college students in Taiwan. *Journal of the Association of Nurses in Care*, 19(5), 397-408. Available at <http://www.sciencedirect.com/science/article/pii/S1055328008001416> Accessed on May 9, 2016.
- Turner, T., Cull, W. L., Bayldon, B., Klass, P., Sanders, L. M., Frintner, M. P., . . . Dreyer, B. (2009). Pediatricians and health literacy: Descriptive results from a national survey. *Pediatrics*, 124 Suppl 3, S299-305. Available at <https://kuscholarworks.ku.edu/handle/1808/14499> Accessed on May 9, 2016.
- Un, D. E. S. A. (2015). World Urbanization Prospects: The 2014 Revision. Available at <http://www.un.org/en/development/desa/publications/2014-revision-world-urbanization-prospects.html> Accessed on February 6, 2016.



- UNICEF, (2008). *Progress for children: A report card on maternal mortality* (7th ed.). United Nations Children's Defense Fund. New York: UNICEF.
- UNICEF, (2013). Iraq, Statistics. Available at [http://www.unicef.org/infobycountry/iraq\\_statistics.html#0](http://www.unicef.org/infobycountry/iraq_statistics.html#0) Accessed on February 27, 2016.
- United Nations Population Fund (UNFPA) and Family Care International (FCI). (2007). *Living testimony: Obstetric fistula and inequities in maternal health*. In UNFPA & FCI. New York (Ed.), *Progress for children: A report card on maternal mortality* (7th ed., pp. 1). New York: UNICEF. Available at [http://www.unicef.org/publications/index\\_45454.html](http://www.unicef.org/publications/index_45454.html) Accessed on February 6, 2016.
- Verma, V., & Betti, G. (2006). EU Statistics on Income and Living Conditions (EU-SILC): Choosing the survey structure and sample design. *Statistics in Transition*, 7(5), 935-970. Accessed on <http://ec.europa.eu/eurostat/statistics-explained/index.php/Glossary:Eurostat> Accessed on October 10, 2016.
- Wallon, M., Peyron, F., & Chêne, G. (2006, April). Impact of health education for the primary prevention of Toxoplasma infection in pregnancy: lessons from the ERIS study. In *16th European Congress of Clinical Microbiology and Infectious Diseases (ECCMID)* (pp. 1-4). Available at <http://congrex.com/> Accessed on October 14, 2016.
- Watts, A. G., Lukasik, V. M., Fortin, M. J., & Alexander, S. M. (2015). Urbanization, grassland, and diet influence coyote (*Canis latrans*) parasitism structure. *EcoHealth*, 12(4), 645-659. Available at <http://link.springer.com/article/10.1007/s10393-015-1040-5> Accessed on February 6, 2016.
- Wells, T. S., Miller, S. C., Adler, A. B., Engel, C. C., Smith, T. C., & Fairbank, J. A. (2011). Mental health impact of the Iraq and Afghanistan conflicts; a review of US research, service provision, and programmatic responses. *International review of psychiatry*, 23(2), 144-152. Available at <http://www.tandfonline.com/doi/abs/10.3109/09540261.2011.558833> Accessed on February 5, 2016.
- Wickrama, K. A., Conger, R. D., Lorenz, F. O., & Jung, T. (2008). Family antecedents and consequences of trajectories of depressive symptoms from adolescence to young adulthood: A life course investigation. *Journal of Health and Social Behavior*, 49(4), 468-483. Available at <http://hsb.sagepub.com/content/49/4/468.short> Accessed on February 6, 2016.
- Williams, M. V., Baker, D. W., Parker, R. M., & Nurss, J. R. (1998). Relationship of functional health literacy to patients' knowledge of their chronic disease: a study of patients with hypertension and diabetes. *Archives of internal medicine*, 158(2), 166-172. Available at <http://archinte.jamanetwork.com/article.aspx?articleid=191123&resultclick=1> Accessed on May 9, 2016.
- Wilson, M., Jones, J. L., McAuley, J. B., Murray, P. R., Baron, E. J., Jorgensen, J. H., ... & Pfaller, M. A. (2006). Toxoplasma. *Manual of clinical microbiology: Volume 2*, (Ed. 9), 2070-2081. Available at

<http://www.cabdirect.org/abstracts/20073141651.html> Accessed on March 4, 2016.

World Health Organization (WHO). 2015. Maternal mortality in Iraq. (n.d.). Retrieved December 25, 2015, Available at <http://www.who.int/mediacentre/factsheets/fs348/en/> Accessed on February 5, 2016.

World Health Organization, UNICEF, & United Nations Fund for Population Activities. (2012). *Trends in maternal mortality: 1990 to 2010: WHO, UNICEF, UNFPA, and the world bank estimates* World Health Organization. Available at [https://www.unfpa.org/sites/default/files/pubpdf/Trends\\_in\\_Maternal\\_Mortality\\_1990-2015\\_eng.pdf](https://www.unfpa.org/sites/default/files/pubpdf/Trends_in_Maternal_Mortality_1990-2015_eng.pdf) . Accessed on February 27, 2016.

World Health Organization. Regional Office for Africa (Ed.). (2006). *The health of the people: The African regional health report*. Geneva: WHO. p.19.

Yan, C., Liang, L. J., Zheng, K. Y., & Zhu, X. Q. (2016). The impact of environmental factors on the emergence, transmission and distribution of *Toxoplasma gondii*. *Parasites & vectors*, 9(1), 1. Available at <https://parasitesandvectors.biomedcentral.com/articles/10.1186/s13071-016-1432-6> Accessed on October 16, 2016.

## BIODATA OF STUDENT

The student was born on 16<sup>th</sup> October 1977, in Al-Najaf, Iraq. He passed Elementary, Secondary and High school education in 1994 in Al-Najaf. In 1995 he pursued two years Professional Diploma of Public Health in Medical Institute of Al-Qadisiyah province. In 1997 he pursued 3 (three) years Bachelor of Public Health in Faculty of Medical and Health Technology\ Foundation of Technical Education-Baghdad. He joined as a Bachelor Lecturer in Faculty of Medical and Health Technology\ Foundation of Technical Education-Baghdad. In 2004, he has enrolled as a Master student in the field of Community Health at the Department of Community Health, Faculty of Medical and Health Technology\ Foundation of Technical Education-Baghdad. In 2013, he has enrolled as a Ph.D. student in the field of Community Health at the Department of Community Health, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia under the supervision of Dr. Titi Rahmawati Hamedon.

## LIST OF PUBLICATIONS

- Ibadi, A. K., & Hamedon, T. R. (2015). The Sociodemographic Characteristics Of Pregnant Women With Toxoplasmosis In Al-Qadisiyah-Iraq. *International Journal Of Public Health And Clinical Sciences*, 2(6), 59-67.
- Ibadi<sup>12</sup>, A. K., & Hamedon, T. R. Sociodemographic Characteristics Of Pregnant Women With Toxoplasmosis In Al-Najaf-Iraq. *Malaysian Journal Of Public Health Medicine* 2016, Vol. 16 (3):





**UNIVERSITI PUTRA MALAYSIA**

**STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT**

**ACADEMIC SESSION :** \_\_\_\_\_

**TITLE OF THESIS / PROJECT REPORT :**

EFFECTIVENESS OF HEALTH EDUCATION IN IMPROVING KNOWLEDGE AND ATTITUDE  
TOWARDS TOXOPLASMOSIS AMONG PREGNANT WOMEN IN AL NAJAF, IRAQ

**NAME OF STUDENT:** ATHEER KADHIM IBADI

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

\*Please tick (v )

**CONFIDENTIAL**

(Contain confidential information under Official Secret Act 1972).

**RESTRICTED**

(Contains restricted information as specified by the organization/institution where research was done).

**OPEN ACCESS**

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

**PATENT**

Embargo from \_\_\_\_\_ until \_\_\_\_\_  
(date) (date)

**Approved by:**

\_\_\_\_\_  
(Signature of Student)  
New IC No/ Passport No.:

Date :

\_\_\_\_\_  
(Signature of Chairman of Supervisory Committee)  
Name:

Date :

**[Note : If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted. ]**