

UNIVERSITI PUTRA MALAYSIA

ANTINOCICEPTIVE ACTIVITIES OF SENDUDUK (MELASTOMA MALABATHRICUM L.) LEAVES METHANOLIC EXTRACT AND ITS PETROLEUM ETHER FRACTIONS

ERMAN SHAH JAIOS

FPSK(m) 2016 53



ANTINOCICEPTIVE ACTIVITIES OF SENDUDUK (Melastoma malabathricum L.) LEAVES METHANOLIC EXTRACT AND ITS PETROLEUM ETHER FRACTIONS



Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of the Requirements for the Degree of Master of Science

November 2016

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

ANTINOCICEPTIVE ACTIVITIES OF SENDUDUK (Melastoma malabathricum L.) LEAVES METHANOLIC EXTRACT AND ITS PETROLEUM ETHER FRACTION

By

ERMAN SHAH JAIOS

November 2016

Chairperson: Associate Professor Zainul Amiruddin Zakaria, PhD

Faculty : Medicine and Health Sciences

Natural products that obtained from the extraction process of medicinal plants are being studied scientifically and endeavor to discover new potential therapeutic agents with less, or no side effect. Melastoma malabathricum L. is one of the medicinally important plants belonging to the family Melastomaceae, commonly known as "Senduduk" in Malay culture. Traditionally, leaves are claimed to relieve diverse pain-related ailments. Therefore, the objective of the present study was to examine the antinociceptive activities of M. malabathricum L. leaves methanolic extract (MEMM) and its petroleum ether (PEMM) fraction by using the in vivo models of nociception in both thermal- and chemical-induced pain tests. The dose of extracts (100, 250, and 500 mg/kg) was administered via orally 60 minutes (min) prior to subjection of the respective test in the volume of 10 mL/kg. Throughout this study, rats and mice (n=6) were pre-treated with the drugs or extract per group. The study was designed as a preventive method and the potential of MEMM and PEMM against nociception has never been reported. In the first stage, we were attempted to evaluate the extract antinociceptive activities, the *in vivo* thermal (hot plate test; HT), chemicals (acetic acid-induced abdominal constriction; ACT and formalin-induced paw licking test; FT) models of nociception were used. In order to elucidate the mechanisms of action involved, the role of opioid, vanilloid receptors (capsaicin), glutamate system (glutamatergic) and nitric-oxide/cyclic guanosine phosphate (NO/cGMP) pathway in modulation of the extract antinociceptive activities were determined. In the second stage, MEMM was partitioned into three fractions: petroleum ether (PEMM), ethyl acetate (EAMM), and aqueous (AQMM). Nevertheless, our objective in this second stage was to investigate the most potent fraction among the three extracts. Therefore, the experiment ED₅₀ (effective dose producing a 50% reduction in relative to control value) and its 95% confidence intervals (CI) values were conducted to determine the most potent fraction and the ACT was used to screen the antinociceptive effect. From the calculation, PEMM is the most effective fraction was further used to assess the antinociceptive properties

using the *in vivo* models of nociception. Moreover, all the extracts (MEMM, PEMM, EAMM and AQMM) underwent the phytochemical screening such as Flavanoids Test, Saponins Test, Tannins and Polyphenolic Compounds Test, Steroids / Triterpenes Test, and Alkaloids Test were recorded. Analysis and identification of phytochemical constituents with the aid of High-Performance Chromatography (HPLC) and Gas Chromatography (GC-MS) technique were performed. In the first stage, MEMM significantly (P < 0.05) was exhibited antinociceptive activities in all the chemically- and thermally-induced nociception models. Naloxone (5 mg/kg), a non-selective opioid antagonist, significantly (P < 0.05) was failed to affect the antinociceptive activity of MEMM. Moreover, MEMM antinociception significantly (P < 0.05) was reversed the capsaicin- and glutamateinduced paw licking test. Whereas, L-arginine (a nitric oxide precursor), L-NAME (an inhibitor of NO synthase), methylene blue, MB (an inhibitor of cGMP), or their combination significantly (P < 0.05) was failed to change the intensity of MEMM antinociception. In the second stage, it was shown the verified screening of the antinociceptive effect of PEMM, EAMM and AQMM fractions assessed by ACT. Likewise, the PEMM and EAMM had similar efficacy to produce antinociceptive effect [max. inhibitions of 24.17 ± 1.33 (70.94%) and 18.83 ± 0.91 (77.36%)] at the dose 500 mg/kg, respectively. As a result, the PEMM was more effective than the EAMM with the calculation of ED₅₀ values [with 95% confidence interval (C.I)] of 119.5 mg/kg (97.03 – 147.1 mg/kg) and 125.9 mg/kg (109.9 – 144.1 mg/kg), respectively. PEMM significantly (P < 0.05) was exhibited antinociceptive activity in all the chemically- and thermally-induced nociception models. Naloxone (5 mg/kg), a non-selective opioid antagonist, significantly (P < 0.05) was failed to reverse the antinociceptive effect of PEMM assessed using the HT and FT. PEMM antinociception significantly (P < 0.05) was reversed the capsaicin- and glutamateinduced paw licking test. Furthermore, L-arginine, L-NAME, MB, or their combination significantly (P < 0.05) was also failed to interfere the PEMM antinociception effect. The phytochemical analysis was screened for all the extracts, and presence of flavonoids, tannins, saponins, triterpenes and steroids, but no alkaloids. In addition, the HPLC analysis of MEMM and PEMM were demonstrated the presence of flavonoids as its major constituents. In the GC-MS analysis, the phytoconstituents were screened and majority of these identified compounds are palmitic acid, terpene, diterpene, α-Linolenic acid and fatty acid ester. Together, these results indicate that the MEMM produced dose-dependent antinociception in the in vivo nociception models of chemical and thermal with the aids of the phytoconstituents, whereby, the PEMM was considered to have the best activity of antinociceptive activities among the fractions, which warrants further investigation.

AKTIVITI ANTINOSISEPTIF OLEH EKSTRAK METANOL DARI DAUN SENDUDUK (*Melastoma malabathricum* L.) DAN PECAHAN PETROLEUM ETER

Oleh

ERMAN SHAH JAIOS

November 2016

Pengerusi : Professor Madya Zainul Amiruddin Zakaria, PhD

Fakuti : Perubatan dan Sains Kesihatan

Produk semulajadi yang diperolehi daripada tumbuhan perubatan melalui proses pengekstrakan sedang giat dikaji secara saintifik dan berusaha untuk menemui agen terapeutik baru yang berpontensi mempunyai kurang atau tidak kesan sampingan. Melastoma malabathricum L. merupakan salah satu tumbuhan perubatan yang berpontesi dan penting di dalam keluarga Melastomaceae dan kebiasannya dikenali sebagai Senduduk dikalangan kaum Melayu. Secara tradisional, daunnya banyak digunakan sebagai rawatan untuk menghilangkan masalah kesihatan atau penyakit berkaitan kesakitan. Objektif kajian ini adalah untuk menilai aktiviti antinosiseptif ekstrak metanol dari daun M. malabathricum L. (MEMM) dan pecahan petroleum eter (PEMM) menggunakan ujian model in vivo nosiseptif secara haba- dan bahan kimia-penghambatan kesakitan. Dos ekstrak (100, 250, and 500 mg/kg) dimasukkan melalui mulut 60 minit awal sebelum dikenakan ujian dengan menggunakan kiraan isipadu 10mL/kg. Sepanjang kajian ini, bilangan (n = 6) bagi tikus dan mencit telah digunakan untuk rawatan bagi ekstrak/dadah per satu kumpulan rawatan. Kajian ini menggunakan kaedah pencengahan dan tiada sebarang laporan berkaitan MEMM dan PEMM melawan nosiseptif direkodkan. Pada peringkat yang pertama, penilaian akitiviti MEMM antinosiseptif menggunakan model ujian nosiseptif teknik in vivo secara haba (ujian plat panas, UPP) dan kimia (ujian asid asetik-penghambatan pengeliatan perut; UAA dan ujian formalin-penghambatan penjilatan tapak kaki; UF). Penjelasan lebih lanjut mengenai tindakan mekanisma yang terlibat telah dilakukan dengan menilai fungsi reseptor opiat, vaniloid (capsaicin) sistem glutamat (glutamatergik) dan laluan nitrik-oksida/siklik-guanosin fosfat (NO/cGMP) dalam modulasi aktiviti ekstrak antinosiseptif. Pada peringkat yang kedua, MEMM telah dipecahkan kepada tiga pecahan iaitu petroleum eter (PEMM), etil esetat (EAMM) dan akueus (AQMM). Namun begitu, objektif kami pada peringkat kedua ini adalah untuk menyiasat pecahan yang paling poten diantara tiga pecahan tersebut. Oleh itu, ujian ED₅₀ (dos efektif menghasilkan pengurangan 50% relatif mengawal nilai kawalan) dan 95% selang keyakinan (CI) telah digunapakai untuk menentukan dos yang paling efektif diantara tiga pecahan, dan ujian UAA telah digunapakai bagi

saringan aktiviti antinosiseptif tersebut. Daripada pengiraan tersebut, pecahan PEMM adalah yang paling berkesan dan susulan ujian telah dilakukan untuk mendapatkan profil antinosiseptif menggunakan model teknik in vivo nosiseptif. Tambahan lagi, semua ekstrak (MEMM, PEMM, EAMM dan AQMM) telah menjalani ujian saringan fitokimia seperti penentuan flavonoid, saponins, tannins dan sebatian polifenol, steroid/tritepen, dan alkaloid. Analisis dan identifikasi juzuk juzuk fitokimia telah dilakukan dengan penggunaan teknik kromatografi cecair berprestasi tinggi (HPLC) serta kromatografi gas (GC-MS). Di peringkat pertama, MEMM secara signifikasi (P < 0.05) mempamerkan aktiviti antinosiseptif dalam semua in vivo model nosiseptif melibatkan ujian secara kimia- dan habapenghambatan. Naloxon (5 mg/kg), antagonis opiad-tidak terpilih, secara signifikasi (P < 0.05) telah gagal memberi kesan kepada aktiviti MEMM antinosiseptif. Selain itu, MEMM antinosiseptif secara signifikasi (P < 0.05) membalikkan aktiviti capsaicin dan glutamat di dalam ujian penghambatan penjilatan tapak kaki, UF. Manakala, L-argina (pencetus nitrik oksida, NO), L-NAME (perencat NO sintase), metilena biru, MB (perencat cGMP), atau gabungannya secara signifikasi (P < 0.05) telah gagal mengubah intensiti MEMM antinosiseptif. Pada peringkat kedua pula, penilaian menggunakan UAA telah mengesahkan bahawa terdapat kesan antinosiseptif daripada semua pecahan (PEMM, EAMM dan AQMM) tersebut. Walaubagaimanapun, pecahan PEMM dan EAMM mempunyai tahap keberkesanan yang serupa untuk menghasilkan kesan antinosiseptif mak. kekangan daripada 24.17±1.33 (70.94%) dan 18.83±0.91 (77.36%)] pada setiap dos 500 mg/kg. Keputusannya, secara respektif, pecahan PEMM adalah lebih berkesan atau poten daripada pecahan EAMM dengan nilai kiraan ED₅₀ (paras keyakinan 95%) iaitu 119.5 mg/kg (97.03 – 147.1 mg/kg) dan 125.9 mg/kg (109.9 – 144.1 mg/kg). PEMM secara signifikasi (P < 0.05) mempamerkan aktiviti antinosiseptif di dalam semua model ujian nosiseptif teknik in vivo secara kimia- dan haba-penghambatan. Penilaian menggunakan antagonis opiat-tidak terpilih (naloxon (5 mg/kg), secara signifikasi (P < 0.05) gagal untuk membalikkan kesan antinosiseptif daripada PEMM melalui ujian UPP dan UF. PEMM antinosiseptif secara signifikasi (P < 0.05) membalikkan aktiviti capsaicin dan glutamat di dalam ujian penghambatan penjilatan kaki. Tambahan lagi, L-arginina, L-NAME, MB, atau gabungannya juga secara signifikasi (P < 0.05) telah gagal untuk mengganggu kesan PEMM antinosiseptif. Penilaian saringan analisis fitokimia bagi semua ekstrak mendapati, kehadiran flavonoid, tannin, saponin, triterpen and steroid, kecuali alkaloid. Tamabahan lagi, analisis HPLC bagi MEMM dan PEMM telah merekodkan dan menunjukkan kehadiran flavonoid sebagai juzuk utamanya dan di dalam analisis GC-MS pula, saringan fitokomponen telah dilakukan dan majoriti sebatian yang dikenalpasti adalah sebatian asid palmitik, terpene, diterpene, asid α-Linolenik dan asid lemak ester. Sebagai kesimpulan, keputusan ini menunjukkan bahawa MEMM menghasilkan dos-hubungan antinosiseptif di dalam model nosiseptif ujian teknik in vivo melalui secara kimia dan haba, dan mencadangkan PEMM menunjukkan aktiviti antinosiseptif yang terbaik di antara pecahan - pecahan serta memerlukan siasatan atau kajian lebih lanjut.

ACKNOWLEDGEMENT

ALHAMDULILLAH, Praise to Allah Subhanahu Wa Ta'ala for granting me grace and strength to preserve throughout my master study and to overcome all the challenges that I had gone through in the study.

First of all, I wish to extend my deepest appreciation to my principle supervisor as chairman of supervisory committee, Associate Professor Dr. Zainul Amiruddin Zakaria for his unrelenting guidance, valuable advice, patience, support, continually and convincingly supervision throughout the adventure in regards of this research project. I truly thank him for giving me opportunity to be his postgraduate student without any hesitation.

I would like to express my gratitude and appreciation to my co-supervisor, Associate Professor Dr. Suzanah Abd Rahman, and Y. Bhg. Professor Dr. Muhammad Nazrul Hakim Abdullah as my supervisory committee for their kindness to provide the invaluable advice and motivation in order to complete this project. Without their guidance and persistent help, this dissertation would not have been possible.

My sincere appreciation dedicates to the Faculty of Medicine and Health Sciences (FPSK), Universiti Putra Malaysia (UPM) and Kulliyyah of Allied Health Sciences (KAHS), International Islamic University Malaysia (IIUM) for giving me the permission and opportunity to carry out this project.

I take this opportunity to record our sincere thanks to all the Faculty/Kulliyyah members of the Department of Basic Medical Sciences (BMS), KAHS, IIUM, for their constant companion, assistant and encouragement. I would like to thank all my fellow friends from UPM and IIUM for their friendship and care towards me, inspiration, and cooperation in completing this project successfully. I would also place on record, my sense of gratitude to one and all who, directly or indirectly, have lent their helping hand in this project.

Words are not enough to express to my beloved wife, Maizatul Akmar Haini for her personal and great patience at all the time, and family, especially my dearest parents, Jaios Hj. Kassim and Hamidah Hj. Abdullah for their unwavering love, encouragement and advice that truly motivated me to accomplish my master study successfully. I owe a depth gratitude to them which I can never be repaid. May Allah SWT shower His countless blessings upon them.

Thank you so much

I certify that a Thesis Examination Committee has met on 10 November 2016 to conduct the final examination of Erman Shah bin Jaios on his thesis entitled "Antinociceptive Activities of Senduduk (*Melastoma malabathricum* L.) Leaves Methanolic Extract and its Petroleum Ether Fractions" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Sabrina binti Sukardi, PhD

Associate Professor Faculty of Medicine and Health Science Universiti Putra Malaysia (Chairman)

Mohamad Taufik Hidayat bin Baharuldin, PhD

Associate Professor Faculty of Medicine and Health Science Universiti Putra Malaysia (Internal Examiner)

Muhammad Taher bin Bakhtiar, PhD

Associate Professor International Islamic University Malaysia Malaysia (External Examiner)

NOR AINI AB. SHUKOR, PhD

Professor and Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date: 28 February 2017

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Zainul Amiruddin Zakaria, PhD

Associate Professor Faculty of Medicine and Health Sciences Universiti Putra Malaysia (Chairman)

Muhammad Nazrul Hakim Abdullah, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Suzanah Abdul Rahman, PhD

Associate Professor Kulliyyah of Allied and Health Sciences International Islamic University Malaysia (Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean School of Graduates Studies Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustration and citation have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institution;
- intellectual property from the thesis and copyright of the thesis are fully-owned by Universiti Putra Malaysia (Research Rules 2012);
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, poster, report, lecture, notes, learning modules or any other materials stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature:			Date:	

Name and Matric No: Erman Shah Jaios, GS32659

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of the thesis was under my supervision;
- supervision and responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012 2013) are adhered to

Signature: Name of Chairman of Supervisory Committee:	Signature: Name of Member of Supervisory Committee:
Signature: Name of Member of Supervisory Committee:	

TABLE OF CONTENTS

ABS ACE APP DEC LIST LIST	ROVAI LARA FOF TA FOF FI	TION	Page i iii v vi viii xv xviii xxv
CHA	PTER		
1.	INT	RODUCTION	1
•	T TTT	ERATURE REVIEW	
2.	2.1	Drug Discovery and Development from medicinal plants	6
	2.1	to natural products	O
	2.2	Synthesis and Function of Plant Primary & Secondary	7
		metabolites	
	2.3	Extraction, Isolation and Identification of bioactive	8
	2.4	compounds in medicinal plant.	0
	2.4	Nature and Classification of Pain 2.4.1 Acute	9 10
		2.4.1 Acute 2.4.2 Chronic	10
		2.4.3 Nociceptive	11
		2.4.4 Neuropathic	11
	2.5	Pain Pathway	12
	2.6	Antinociceptive-related to natural products	14
	2.7	Involvement of Various Systems in Pain	
		2.7.1 Involvement of Opioid System	15
		2.7.2 Response to heat (thermal)	16
		2.7.3 Abdominal constriction response (chemical)	17
		2.7.4 Formalin-induced pain	18
		2.7.5 Nociception induced by Glutamate (glutamatergic system)	19
		2.7.6 Nociception induced by Capsaicin (Vanilloid receptors)	20
		2.7.7 Role of nitric oxide/cylic-guanosine monophosphate (NO/cGMP) pathway on mechanisms of pain.	21
	2.8	Melastoma Malabathricum Linn.	23
		2.8.1 Plant Morphology	23
		2.8.2 Southeast Asian Genus <i>Melastoma</i> (<i>Melastomataceae</i>)	24
		2.8.3 Classification and Nomenclature	25
		2.8.4 Ethno-Botanical Description and Natural Habitat	26

	2.9	2.8.5 Ethno-Medicinal Uses of <i>M. malabathricum</i> L. Phytochemical and bioactivity studies of <i>M</i> .	26 30
	2.9	malabathricum L.	30
	2.10		32
	2.11	ϵ	38
	2.11	1 hytochemical and 7 metoxidant 1 toperties	30
3.	MET MAL	ALUATION OF ANTINOCICEPTIVE ACTIVITY OF THANOLIC EXTRACT OF MELASTOMA LABATHRICUM L. (MEMM) LEAVES AND ITS SIBLE MECHANISM OF ACTION	
	3.1	Introduction	40
	3.2		
		3.2.1 Plant collection	41
		3.2.2 Preparation of MEMM	41
		3.2.3 Preparation of Chemicals and Drugs	41
		3.2.4 Experimental animals	42
		3.2.5 Acute toxicity study of MEMM	42
		3.2.6 Antinociceptive activity of MEMM	
		3.2.6.1 Acetic Acid-Induced Abdominal	43
		Constriction Test (ACT)	
		3.2.6.2 Hot Plate Test (HPT)	43
		3.2.6.3 Formalin-Induced Paw Licking Test (FT)	44
		3.2.7 Investigation of the mechanisms of	
		antinociceptive activity of MEMM	
		3.2.7.1 Capsaicin-Induced Paw Licking Test	44
		3.2.7.2 Glutamate-Induced Paw Licking Test	44
		3.2.7.3 Involvement of the opioid receptor	45
		3.2.7.4 Involvement of the nitric oxide/cyclic-	45
		guanosine monophosphate (NO/cGMP)	
		pathway	
	3.3	Statistical analysis	45
	3.4	Results	
		3.4.1 Acute toxicity effect of MEMM	46
		3.4.2 Acetic acid-induced abdominal constriction Test	46
		3.4.3 Hot plate Test	46
		3.4.4 Formalin-induced paw licking Test	47
		3.4.5 Possible Mechanisms of action MEMM	
		3.4.5.1 Capsaicin-induced paw licking test	54
		3.4.5.2 Glutamate-induced paw licking test	54
		3.4.5.3 Involvement of opioid receptors	54
		3.4.5.4 Involvement of NO/cGMP pathway	57
	3.5	Discussion	60

4.	EVALUATION OF ANTINOCICEPTIVE ACTIVITIES OF FRACTIONS DERIVED FROM METHANOLIC CRUDE				
				MELASTOMA MALABATHRICUM L.	
				AND ITS POSSIBLE MECHANISMS	
	•	CTION	AVES	AND THE TOSSIBLE MECHANISMS	
	4.1	Introdu	ection		67
	4.2				07
	1.2	4.2.1	Prepar	ation of fractions: petroleum ether	68
		1.2.1		M), ethyl acetate (EAMM) and aqueous	00
			(AQM	· · · · · · · · · · · · · · · · · · ·	
		4.2.4		ation of Chemicals and Drugs	70
			Anima	<u> </u>	70
				ociceptive assays	
				Acetic acid-induced abdominal	70
				constriction test	
			4.2.4.2	Hot plate test	71
				Formalin-induced paw licking test	73
		4.2.5		le mechanisms of antinociceptive action of	
			PEMN	I fraction	
			4.2.5.1	Investigation on the role of vanilloid	73
				receptors using the capsaicin-induced paw	
				licking test	
			4.2.5.2	Investigation on the role of glutamatergic	73
				system using the glutamate-induced paw	
				licking test	
			4.2.5.3	Investigation on the involvement of opioid	74
			, <	receptors system using the hot plate and	
			1	formalin-induced paw licking tests	
			4.2.5.4	Investigation on the involvement of nitric	74
				oxide/cyclic-guanosine monophosphate	
				(NO/cGMP) pathway using the abdominal	
	4.2	Q		constriction test	7.4
	4.3	Statisti			74
		4.4	Results		75
			4.4.1	Screening on antinociceptive activity of	75
				fractions assessed using the abdominal constriction test	
			4.4.2		75
			4.4.2	Antinociceptive profile of PEMM fraction assessed using the hot-plate test	73
			4.4.3	Antinociceptive profile of PEMM fraction	75
			4.4.5	assessed using the formalin-induced paw	13
				licking test	
			4.4.4	Mechanisms of antinociceptive activities	
				of PEMM fraction	
				4.4.4.1 Involvement of vanilloid receptors	76
				in the antinociceptive activity of	. 0
				PEMM fraction	
				4.4.4.2 Involvement of glutamatergic	76
				system in the antinociceptive	
				·	

	activity of PEMM fraction	
	4.4.4.3 Involvement of opioid receptors in	76
	the antinociceptive activity of	
	PEMM fraction	
	4.4.4.4 Involvement of NO/cGMP	77
	pathway in the antinociceptive	
	activity of PEMM fraction	96
	4.5 Discussion	86
	ALUATION ON PHYTOCHEMICAL SCREENING, GH PERFORMANCE LIQUID CHROMATOGRAPHY	
	PLC) AND GAS CHROMATOGRAPHY (GC-MS)	
	ALYSIS OF THE METHANOLIC EXTRACT M.	
	labathricum L. (MEMM) LEAVES AND ITS	
	ACTIONS (PEMM, EAMM & AQMM).	
5.1		92
5.2	Methodology	
	5.2.1 Phytochemical screening of the MEMM, PEMM,	
	EAMM and AQMM	
	5.2.1.1 Test for alkaloids	93
	5.2.1.2 Test for flavonoids	93
	5.2.1.3 Test for steroids and triterpenes	94
	5.2.1.4 Test for tannins	94
	5.2.1.5 Test for saponins	94
	5.2.2 HPLC profiling of MEMM	95
	5.2.2 Identification of flavonoids in MEMM via HPLC analysis	95
	5.2.3 HPLC analysis of the most effective fraction	95
	(PEMM) at various wavelengths	,,,
	5.2.4 GC-MS analysis of MEMM and PEMM	95
5.3	Results	, ,
	5.3.1 Phytochemical screening	96
	5.3.2 HPLC profile of MEMM	97
	5.3.3 Identification of flavonoids in MEMM via HPLC	98
	analysis	, 0
	5.3.4 HPLC profiling of PEMM at different wavelength	99
	5.3.5 GC-MS profile of MEMM and PEMM fraction	100
5.4	Discussion	106

6.	GENERAL DISCUSION	109
7.	SUMMARY, CONCLUSION AND RECOMMENDATION FOR FURTHER STUDY	114
	FERENCES	116
APP	PENDICES	147
BIO	DATA OF STUDENT	158
LIS	Γ OF PUBLICATIONS	159



LIST OF TABLES

Table 1	The species of <i>M. malabathricum</i> L. (subspecies <i>malabathricum</i> and <i>normale</i>), found in Malaysia and the Morphological Variability (Meyer, 2001)	Page 24
2	The vernacular name of <i>M. malabathricum</i> L. depending on tribes/language and country/origin (S Mohd Joffry <i>et al</i> , 2012).	25
3	The ethno-Medicinal uses of <i>M. malabathricum</i> L. according to its part	27
4	The medicinal uses of <i>M. malabathricum</i> L. according to several communities/tribes	29
5	The phytochemical compound(s) of <i>M. malabathricum</i> L. leaves and its extracts	30
6	Pharmacology properties of M. malabathricum L. leaves	33
7a	Percentage changes in body weight of rats in acute oral toxicity study of MEMM	50
7b	Biochemical analysis of rats in acute oral toxicity study of MEMM	50
8	Antinociceptive profile of MEMM assessed using the hotplate test in mice. * $P < 0.05$ when compared with the control group at same the respective interval. Data are the mean \pm SEM; n = 6 mice per group.	52
9	Antinociceptive profile of PEMM assessed using the hotplate test in mice. Statistical significant was determined by two-way ANOVA followed by Bonferroni's post hoc test. $*P < 0.05$ when compared with the control group at same the respective interval. Data are the mean \pm SEM; $n = 6$ mice per group.	78

10	Effect of PEMM in formalin-induced paw licking test. Data represents the mean \pm SEM of 6 rats. The rats were pretreated with vehicle (10% DMSO), PEMM (100, 250, and 500 mg/kg, p.o.), acetylsalicylic acid (ASA, 100 mg.kg, p.o.), or morphine (5 mg/kg, p.o.), 60 min before <i>i.pl</i> injection of formalin. The asterisks denote the significance levels as compared to control. Data analyzed by one-way ANOVA followed by Dunnett's post hoc test.	79
11	Phytochemical screening of <i>M. malabathricum</i> L. leaves in Powder of MEMM, PEMM, EAMM and AQMM	96
12	GC-MS analysis of the volatile compounds from the leaves of methanolic extract <i>M. malabathricum</i> L. (MEMM).	101
13	GC-MS analysis of the volatile compounds from of semi-purified petroleum ether (PEMM) of methanolic extract <i>M. malabathricum</i> L. (MEMM) Leaves.	102
14	Some of the nature chemical compounds present in MEMM and PEMM and its medical properties. Therapeutic activity source: Dr. Duke's Phytochemical and Ethnobotanical databases	104

LIST OF FIGURES

Figure		Page
1	Schematic representation of a typical medicinal plant drug discovery process and development (Balunas and Kinghorn, 2005).	7
2	Main pathways leading to secondary metabolites (Taiz and Zeiger, 2006).	8
3	A simplified representation of nociceptive process. 1st order neuron-a noxious stimulus at the periphery activates a primary afferent fibre and transmits the information to the dorsal horn of the spinal cord; 2nd order neuron- ascending in a spinal tract to the level of the thalamus intervenes; 3rd order neuron-transmits the modified noxious stimulus to higher brain centres (cerebral cortex) for perception (Lamount LA <i>et al.</i> , 2000).	12
4	Melastoma malabathricum L. tree, leaf, scrub	23
5	Histological sections in acute toxicity study of MEMM	48
6	Antinociceptive activity of MEMM assessed by the acetic acid-induced abdominal constriction test in mice. Acetic acid administrated by intraperitoneally 60 min before pre-treated with DMSO as vehicle (control), acetylsalicylic acid (ASA), or MEMM (100, 250 and 500 mg/kg, $p.o.$). All treatments administrated via oral route. The asterisks denote the significant level as compared to control, *** P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test. ***Data differed significantly (P < 0.001) when compared to 10% DMSO-treated group.	51
6a	Effect of MEMM on Formalin-induced paw licking test in rats (early / neurogenic phase, $0-5$ min). Each column represents the mean \pm SEM of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control, MEMM (100, 250, and 500 mg/kg, $p.o.$), acetylsalicylic acid (ASA, $p.o.$), or morphine (5 mg/kg, $p.o.$), 60 min before i.pl injection of formalin. The asterisks denote the significance levels as compared to control, *** $P < 0.001$ by one-way ANOVA followed by Dunnett's post hoc test. *** Data differed significantly ($P < 0.001$) when compared to the 10% DMSO-treated group.	53

6b Effect of MEMM on Formalin-induced paw licking test in rats (late / inflammatory phase, 15-30 min). Each column represents the mean \pm SEM of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control, MEMM (100, 250, and 500 mg/kg, p.o.), acetylsalicylic acid (ASA, p.o.), or morphine (5 mg/kg, p.o.), 60 min before i.pl injection of formalin. The asterisks denote the significance levels as compared to control, *** P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test. *** Data differed significantly (P < 0.001) when compared to the 10% DMSO-treated group.

53

- Antinociceptive activity of MEMM assessed using the capsaicin-induced paw licking test in rats. Each column represents the mean ± S.E.M. of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control or MEMM (100, 250 and 500 mg/kg, p.o) 60 min before of capsaicin (1.6μg/paw, 20μL, *i.pl*.).The asterisks denote the significance levels as compared to control, ***P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test.
- Antinociceptive activity of MEMM assessed using the glutamate-induced paw licking test in rats. Each column represents the mean ± S.E.M. of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control or MEMM (100, 250 and 500 mg/kg, *p.o*) 60 min before of glutamate (10 μmol/paw, 20μL, *i.pl.*). The asterisks denote the significance levels as compared to control, ****P* < 0.001 by one-way ANOVA followed by Dunnett's post hoc test.
- 9 Schematic representation of the standard solvent partitioning methods described by Sowndhararajan and Kang, 2013 with modification, for preparation of semi- purified extracts from the crude methanolic extract *M. malabathricum* L. (MEMM) leaves.
- Involvement of NO/cGMP pathway in the modulation of antinociceptive activity of MEMM. Effect of L-arginine, L-NAME and their combination on MEMM antinociception as assessed by acetic acid-induced abdominal constriction test. The asterisks denote the significance levels as compared to control, *** P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test.
- 9b Involvement of NO/cGMP pathway in the modulation of antinociceptive activity of MEMM. Effect of L-arginine, MB and their combination on MEMM antinociception as assessed by acetic acid-induced abdominal constriction test. The asterisks denote the significance levels as compared to

control, *** $P < 0.001$	by one-way	ANOVA	followed	by
Dunnett's post hoc test.				

72

- Schematic representation of the standard solvent partitioning, screening antinociceptive activity of partitions, calculation for determination of most effective fraction and PEMM-antinociceptive profiles.
- Screening on antinociceptive activity of fractions assessed by the acetic acid-induced abdominal constriction test in mice
- 12a Effect of PEMM extract on formalin-induced paw licking test in rats (early / neurogenic phase, 0 5 min). Each column represents the mean ± SEM of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control, PEMM (100, 250, and 500 mg/kg, p.o.), acetylsalicylic acid (ASA, p.o.), or morphine (5 mg/kg, p.o.), 60 min before i.pl injection of formalin. The asterisks denote the significance levels as compared to control, *** P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test. *** Data differed significantly (P < 0.001) when compared to the 10% DMSO-treated group.
- 12b Effect of PEMM extract on formalin-induced paw licking test in rats (late / inflammatory phase, 15 30 min). Each column represents the mean ± SEM of 6 rats. The rats were pretreated with vehicle (10% DMSO) as control, PEMM (100, 250, and 500 mg/kg, p.o.), acetylsalicylic acid (ASA, p.o.), or morphine (5 mg/kg, p.o.), 60 min before i.pl injection of formalin. The asterisks denote the significance levels as compared to control, *** P < 0.001 by one-way ANOVA followed by Dunnett's post hoc test. *** Data differed significantly (P < 0.001) when compared to the 10% DMSO-treated group.
- 13 Effect of PEMM on capsaicin-induced paw licking test in rats. Each column represents the mean ± SEM of 6 rats. The rats were pre-treated with vehicle (control, 10% DMSO), Capsazepine (0.17 mmol/kg) or PEMM (100, 250, and 500 mg/kg, *p.o.*) 60 min before injection of capsaicin (1.6 μg/paw, 50 μl, *i.pl.*). The asterisks denote the significance levels as compared to control, *** *P* <0.001 by one-way ANOVA followed by Dunnett's post hoc test.
- 14 Effect of PEMM on glutamate-induced paw licking test in rats. Each column represents the mean ± SEM of 6 rats. The rats were pre-treated with vehicle (control, 10% DMSO), ASA (100mg/kg) or PEMM (100, 250, and 500 mg/kg, *p.o.*) 60 min before injection of glutamate (10 μmol/paw, 50 μl,

	compared to control, *** $P < 0.001$ by one-way ANOVA followed by Dunnett's post hoc test.	
15a	Effect of L-arginine, L-NAME and their combination with PEMM extract assessed by acetic acid-induced abdominal constriction test. The asterisks denote the significance levels as compared to control, *** $P < 0.001$ by one-way ANOVA followed by Dunnett's post hoc test.	84
15b	Effect of L-arginine, MB and their combination with PEMM extract assessed by acetic acid-induced abdominal constriction test. The asterisks denote the significance levels as compared to control, *** $P < 0.001$ by one-way ANOVA followed by Dunnett's post hoc test.	85
16	The HPLC analysis of MEMM. A) The HPLC profile of MEMM at the wavelength of 366 nm. B) The UV spectra analysis of MEMM at 366 nm. The chromatogram demonstrated the presence of several peaks, with the only one major peak detected at the retention time (R_T) of 28.584 min. This peak was observed at the λ_{max} within the region of 284.9 – 349.4 nm, suggesting, in part, the presence of flavonoid-based compounds.	97
16C	Comparison of the HPLC chromatograms of quercitrin alone or in combination with MEMM against the chromatogram of MEMM alone focusing on the retention time (R _T) and UV spectra (λ_{max}) of the major peak, Peak 1. The HPLC analysis was performed at 366 nm. The R _T for Peak 1 in MEMM, quercitrin and their combination as 'MEMM + Quercitrin' was 28.585, 28.870 and 28.922 min, respectively. The λ_{max} for the respective peak falls in the range of 204.9–349.4, 204.9–349.4 and 206.1–347.0 nm, respectively.	98
17	A), The HPLC of PEMM at 254 nm and 366nm, B), The HPLC profile of PEMM demonstrated the presence of rutin.	99
18	A typical gas chromatogram of the chemical constituents of methanol crude extract (MEMM) from the leaves of <i>M. malabathricum</i> L.	104
19	A typical gas chromatogram of the chemical constituents of PEMM fraction.	105
20	Schematic diagram of the proposed mechanisms of antinociceptive action demonstrated by MEMM and PEMM exerts antinociceptive activity at central and peripheral levels	113

i.pl.). The asterisks denote the significance levels as

via modulation of several receptors.

LIST OF ABBREVIATIONS

MEMM Methanolic crude extract *Melastoma malabathricum* L. leaves PEMM Petroleum ether semi-purified extract of *M. malabathricum* L.

leaves

EAMM Ethyl acetate semi-purified extract of *M. malabathricum* L. leaves AQMM Aqueous semi-purified extract of *M. malabathricum* L. leaves

MeOH Methanol
PE Petroleum ether
EA Ethyl acetate
AQ Aqueous
dH₂O Distilled water
NaCL Normal saline

p.o Orally

i.p Intraperintoneally
 i.pl Intraplantarly
 s.c Subcutaneously
 HPT Hot-Plate Test

ACT Acetic Acid-Abdominal Constriction Test

FT Formalin Test NO Nitric oxide

cGMP Cyclic Guanosine Monophospate

DMSO Dimethyl sulfoxide ASA Acetyl salicylic acid

NLX Naloxone Capz Capsazepine L-arg L-arginine

L-NAME N^G-nitro-L-arginine methyl esters

MB methylene blue
mL Milliliter
L Liter
kg Kilogram
gm Gram

°C Degree Celsius

Minute min Second(s) sec μL Microliter Micromole µmol Milimoles mM cm centimeter μM Micrometer h Hour(s)

ID₅₀ Effective dose producing a 50% reduction in relative to control

value

CI Confidence Interval
ANOVA Analysis of variance
S.E.M Standard error mean
USA United States of America

WHO World Health Organization UPM Universiti Putra Malaysia

IIUM International Islamic University Malaysia
IASP International Association for the Study of Pain

BC Before Century

p P-value

NSAIDs Non-steroidal anti-inflammatory drugs

% Percent

GIT Gastrointestinal tract
COX Cyclooxygenase
LOX Lipooxygenase

TRPA1 Transient receptor potentially A1

TRPV1 Transient receptor potential cation channel subfamily V member 1

CNS Central nervous system
PNS Peripheral nervous system

AMPA α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid

NMDA N-methyl-D-aspartic acid receptor

PGE₂ Prostaglandin E2

sGC Soluble guanylate cyclase

GC Guanylate cyclase

HPLC High Performance Liquid Chromatography
GC-MS Gas Chromatography-Mass Spectrometry
MPLC Medium-Pressure Liquid Chromatography
LC-MS Liquid chromatography—mass spectrometry

SFE Supercritical Fluid Extraction

IR Infrared

NMR Nuclear Magnetic Resonance

RP Reversed-phase

TPC Total phenolic content

DPPH 2, 2-diphenyl-1- picrylhydrazyl
SOD Superoxide anion radical scavenging
ORAC Oxygen radical absorbance capacity

PMS-NADH Phenazine methosulphate - nicotinamide adenine dinucleotide

NBT Nitroblue tetrazolium

AAPH 2,2'-Azobis(2-amidinopropane) dihydrochloride

UV-Vis Ultra Violet-Visible Spectrophometer

NSIT National Institute Standard and Technology

R_T Retention time Nanometer

 $\begin{array}{lll} TE & Trolox \ Equivalent \\ GAE & Gallic \ Acid \ Equivalent \\ PKC & Protein \ Kinace \ C \\ K^+ & Potassium \ ion \\ Ca2^+ & Calcium \ ion \\ \end{array}$

Ca₂⁺ Calcium ion
Na⁺ Sodium ion
CO₂ Carbon dioxide

ADME Absorption, Distribution, Metabolism and Excretion

EAA Excitatory amino acids

5HT Serotonin



CHAPTER 1

INTRODUCTION

Pain is the most common symptom of ailments, which refers to an individual experience that accompanies us from childhood. It is a protective mechanism or warning signal to which the body responds to harmful stimulus (Swieboda P *et al.*, 2013). According to the International Association for the Study of Pain (IASP), pain is an unpleasant sensory and emotional experience associated with potential or actual tissue damaged and/or reaction of the body to harmful stimuli or indicates a protective early signal/warning to the body system (MH Mohd Sani *et al.*, 2012). However, the experience of pain depends on the individual strength of the stimulus, tendency and resistance to pain. Nevertheless, the way of pain perception is differ from time to time in the same individual, and also depending on the several factors such as arousal, attention, distraction and expectations (Swieboda P *et al.*, 2013). Therefore, no one patient with pain experience can be treated with exactly the same methods or medications as another patient.

Subsequently, several reports have claimed that, pain experience is the most common reason for any individual to seek for health medication (Hui Ming Ong et al., 2010). For example, in year 2008, the Ministry of Health Malaysia (MOH) recognized Pain as the fifth vital sign among the other major ailments, which giving emphasis and serious observation in their strategy to improve pain management in the hospitals (MOH Malaysia, 2013). Furthermore, according to the World Health Organization (WHO) estimated that approximately 80% of the world population has either no or insufficient access for treatment of pain (Jaganath IB and LT Ng, 2002). In addition, every year tens of millions of people around the world are suffering from pain without treatment (MOH Malaysia, 2013). Therefore, relieve from pain is desirable when the duration and intensity of pain alters the ability of a subject to function efficiently. In such situation, analgesics drugs are useful due to these agents could relieve pain without producing a loss of consciousness. Commonly, the treatment of pain is using nonopioid analgesic drugs such as non-steroidal anti-inflammatory drugs (NSAIDs) that act to reduce the generation process of the pain mediators at the site tissue damaged, despite several of the opioid analgesic drugs are also have some effects within the central nervous system (CNS). Meaning that, the opioid analgesic drugs are unique, which not only block the incoming nociceptive signals to the brain, however, they are also capable to control the affective components of the pain, which act at higher brain centres (Sanda P Welch and Billy R Martin, 1997).

Considering, various ailments have been detected recently, use and practice alternative medicinal treatment, which approach the uses of medicinal plants due to minimum or no adverse side effect. Moreover, the use of plant-related natural products relatively, has indicates lower incidence of adverse side effect reaction in comparing to modern or conventional pharmaceutical products, as well as linked with their reduced cost and encouraging effort from both public consumer and health

care, which highlighted medicinal plant as an alternatives approach to replace the synthetic drugs. Therefore, the provision and preparation for a range of ailments, that using medicinal plants for the treatment are commonly experienced by contemporary community of people such as pain and inflammation (SK Raghav *et al.*, 2006; HP Rang *et al.*, 2011). Such that approaches, one of the medicinal plants, which possess medicinal values, and being used by the various communities is called *Melastoma malabathricum* L. known as 'Senduduk' of the family Melastomataceae. This plant has attribute valuable healing properties to relieve fever (antipyretics), pain reducer (analgesics), treating vaginal discharge (leucorrhoea), reduction of inflammation, excessive menstrual bleeding, and treating burns or bleeding, inflammation of the walls of blood vessels with blood clots in the vessels (Dalimartha, 2000). It also has been used extensively in traditional medicine both locally and abroad.

Generally, the M. malabathricum L. can be described as a small shrub, which commonly can be found in waste places, previously cleared land and along roadside throughout the Southeast Asian countries including Malaysia (JLCH Van Valkenberg and N Bunyapraphatsara, 2001). In Malaysia, the plant commonly growth in the lowland and slope mountain forests, especially in open places. It is also native to tropical and temperate in Asia and Pacific Islands (KH Ling et al., 2009). Interestingly, it has different vernacular names depending on the location/countries (e.g., Malaysia, Indonesia, China and India), and the communities or tribes (e.g., Malay, Chinese and Indian), which traditionally used for medicinal purposes (FA Abdul Majid and LY Ting, 2011). Moreover, parts of the M. malabathricum L. plant have also been widely used in traditional remedies. Commonly, every part of the plant including roots can be used as the biomaterial resources for the preparation of traditional medicine, and as natural food colourants due to the presence of anthocyanins (Janna et al., 2006). Furthermore, Jaganath IB and LT Ng, 2002 has also reported that, puerperal disease and infectious diarrhea could be treated by eating the raw 'Senduduk' leaves. It was also proven to have anti-inflammation and antinociceptive effect on mice (Sulaiman et al., 2004; ZA Zakaria et al., 2006; Zakaria ZA et al., 2008). On the other hand, various parts of the M. malabathricum L. have been claimed to possess medicinal values, which is supported particularly by the Malay and Indian traditional uses of the plants in the treatment of a number of ailments as described earlier. Therefore, a systematic scientific research of the M. malabathricum L. was prepared as extracts using different types of solvents and tested using a range of in vitro and in vivo test models, which was demonstrated various pharmacological findings that required in-depth studies. Such that plant, regardless of the parts used has been shown to exert anti-bacterial, anti-viral, antiantioxidant, cytotoxicity, anti-coagulant, platelet-activating inhibitory, wound healing, anti-ulcer, anti-diarrheal, anti-venom, anti-inflammatory, antinociceptive and anti-pyretic activities at different doses or concentrations (Sulaiman et al., 2004; ZA Zakaria et al., 2006; Zakaria ZA et al., 2008; S Mohd Joffry et al, 2012).

Problem statement

Generally, response of pain could be the first treatment with the non-opioid analgesics drugs (NSAIDs) such as acetylsalicylic acid (ASA), which commonly useful for treatment of pain, fever, and inflammation but less effective than the opioids to relieve pain (mild-to-moderate). Impressively, opioid analgesic drugs such as morphine known as narcotic analgesics could relieve severe pain (moderate-tosevere) by selective acting on CNS to reduce the pain reaction, at the same time do not dissipate the function of peripheral nerves, which means it is capable of inhibiting pain of any origin (Sanda P Welch and Billy, 1997). However, consumption of these drugs associated with their prolong used to treat pain has several adverse side effects and most commonly are nausea, vomiting, dry mouth, constipation, urinary retention, and bring to mental confusion (Sanda P Welch and Billy R Martin, 1997). Moreover, the most serious adverse side effect that associated with chronic use of opioid analgesic drugs, which cause physical dependency and development of patience (Henry Hitners and Barbara Nagle, 1999). Based on the report from Ministry of Health Malaysia 2014, Malaysian statistics on medicine 2009 & 2010 indicates the total opioid consumption in Malaysia that used for pain control, which was recorded 0.3643 DDD/1000 population/day in year 2010 and in year 2009 was 0.3174 DDD/1000 population/day, respectively showing an increase of 15% in year 2010 compared to year 2009, and morphine remains as the most commonly used strong opioid, with an increasing trend over the years approximately 73.2% of all strong opioids in 2009 and 86.8% in 2010, as compared to 65% in 2008. Moreover, the report is also mentioned the increase in expenditure on medicines every year is an indication of the increasing burden of diseases whereby the commitment and responsibility of the healthcare industries in the country to treat the population and to fight against the emerging diseases is essential. Hence, there is need to find alternative agents with less or possibly no side effects, lower cost and medicinal plant is one of source of these agents.

Justification for studying the antinociceptive potential of *M. malabahricum* L. leaves

It is time demanding to explore and develop potential new drugs from natural resources as many ailments are continuously arising. Furthermore, due to unpleasant side effects, limitation on dosage consumption as well as high cost of available drugs, many synthetic drugs are withdrawn years after their introduction into the market. Therefore, the bioactive compounds, which discovered in medicinal plants as an alternative medicine for treating of ailment related to pain as a substitute for available or current drugs that have less or no side effect, and considered cheaper as well as widely available. In addition, the number of researchers, which studied the potential plant extracts, produce the antinociceptive and anti-inflammatory agents have been increased, and the interest has been enhanced lately (Ferguson *et al.*, 2003; Bighetti *et al.*, 2005; Orhan *et al.*, 2007), thus, in this study, we aimed to discover the potential antinociceptive activities of the *M. malabathricum* leaves that might add to

another candidates to the list, which also support with the various evidences on the traditional application or ethno-medicinal uses of M. malabathricum leaves to treat pain-related ailments (Sharma et al., 2001; Zakaria ZA et al., 2008; Umali-Stuart and Stiuart-Santiago, 2010). Subsequently, scientific studies, reported that M. malabathricum leaves possess antinociceptive activities (Sulaiman et al., 2004; ZA Zakaria et al., 2006; Zakaria ZA et al., 2008). However, the three researches have reported the use of ethanol, aqueous and chloroform extracts as their source of antinociceptive study, and was administered systemically either by intraperitoneal or subcutaneous routes. In contrast to those reports, our present study used methanolic extract that was administered orally, which represent the traditional way of consuming the plant's extract. Furthermore, this plant is considered one of the most common weeds that grow wildly and available in open places such as lowland and mountain forests especially in the moist areas (S Mohd Joffry et al., 2012). In consideration of the ethno-medicinal and scientific reports together with the identification of phytoconstituents through the phytochemical screening and analysis via chromatography, therefore, this study is expected to discover the capacity of M. malabathricum leaves for antinociceptive activity.

Hypothesis

Methanolic extract of *M. malabathricum* L. (MEMM) leaves possess the antinociceptive activity in thermal- and chemicals-induced nociception assays, and petroleum ether (PEMM) fraction is expected to exert antinociceptive effects induced by thermal and chemicals in animal models.

General objectives:

To determine the antinociceptive activities of methanolic extract of *Melastoma* malabathricum leaves and its petroleum ether fractions in animal models.

Specific objectives:

- 1. To determine safety of methanolic extract of *M. malabathricum* leaves (MEMM) using the single high-dose acute toxicity model.
- 2. To determine the antinociceptive profile of MEMM using various animal models.
- 3. To elucidate the possible mechanisms of action that takes part in the antinociceptive of MEMM.
- 4. To determine the most effective fraction from MEMM using acetic acidinduced abdominal constriction test.
- 5. To elucidate the possible mechanisms of action that takes part in the antinociceptive of petroleum ether (PEMM) fraction.
- 6. To screen and identify the possible bioactive compounds that present in the MEMM and its fractions triggered antinociception using the phytochemical screening test, HPLC and GC-MS analysis.

REFERENCES

- Abacioglu, N., Tun, Tan, B., Akbulut, E., Cakici, I. (2000). Participation of the components of L-arginine/nitric oxide/cGMP cascade by chemically-induced abdominal constriction in the mouse. *Life Sci.* 67: 1127 1137.
- Abbott, R. C., Chomel, B. B., Kasten, R. W., Floyd-Hawkins, K. A., Kikuchi, Y. *et al.*, (1997). Experimental and natural infection with Bartonella henselae in domestic cats. *Comp Immunol Microbiol Infect Dis.* 20: 41 51.
- A, Beirith., A. R. S, Santos., and J. B. Calixto. (2002). Mechanisms underlying the nociception and paw oedema caused by injection of glutamate into the mouse paw. *Brain Research*. 924 (2): 219 228.
- A. B. Malmberg., T. L. Yakash. (1995). Cyclooxygenase inhibition and the spinal release of prostaglandin E₂ and amino acid evoked by paw formalin injection: a microdialysis study in unanesthetized rats. *J. Neurosci.* 14: 2768 2776.
- Adeyanju, O., Olutayo, O. O., Michael, A., and Arifalo, K. M. (2011). Preliminary phytochemical, antimicrobial screening of the leaf extract *Pilostigma reticulatum* (dc) *Hochst. African Journal of Pure and Applied Chemistry*. 5(3): 43 46.
- A. Dickenson. (1995). Central acute pain mechanisms. *Ann. Med.* 27: 223 227.
- Arif Ullah H. M., Zaman, S., Juhara, F., Akter, L., Tareq, S.M., Masum, E. H., Bhattacharjee R. (2014). Evaluation of antinociceptive, *in-vitro* anti-inflammatory activity of ethanolic extract of *Curcuma zedoaria* rhizome. *BMC Complement Alternate Med.* doi; 10.1186/1472-6882-14-346.
- Ahmad, A., Alkarkhi, A. A., Hena, S., Lim, H. K. (2009). Extraction, Separation and Identification of Chemical Ingredients of *Elephantopus Scaber* (L) using Factorial Design of Experiment. *Int J Chem.* 1:36 49.
- Ahmed, F., Hossain, M.H., Rahman, A.A., Shahid, I. Z. (2006). Antinociceptive and sedative effects of the bark of *Cerbera odollam* Gaertn. *OPEM*. 6:344 48.
- Akinmoladun, A. C., Ibukun, E. O., Afor, E., Obuotor, E. M., Farombi, E. O. (2007).
- Alamgeer., Taseer, Ahmad., Iradat, Hussain., Muhammad, Naveed, Mushtaq., Nadeem, Irshad., Muhammad, Wasim., Sultan, Uallah., Maria, Mukkaram. (2013). The use of plant extract in a mouse model of depression: A Review. *J App Pharm.* 5(4): 838 842.
- Alan, Fein. (2012). Nociceptors and the Perception of Pain. *Revised February 2012*. Chapter 8:133 145.

- Al-Hasani, R., and Bruchas, MR. (2011). Review: Molecular mechanisms of opioid receptor-dependent signaling and behavior. *Anesthesiology*. 115(6):1363 81.
- Ali, D. M. H., Wong, K. C., Boey, L. P. (2010). Triterpenoids, glycolipids and flavonoids of *Melastoma malabathricum*: Isolation, spectrometric characterization and antibacterial activity. *Germany: VDM Verlag.* 1 228.
- Allison, M. C., Howatson, A. G., Torrance, C. J., Lee, F. D., Russell, R. I. (1992). Gastrointestinal damage associated with the use of non-steroidal anti-inflammatory drugs. *N Engl J Med.* 327: 749 54.
- Almeida, R. N., Navarro, D. S., and Barbosa-Filho, J. M. (2001). Plants with central analgesic activity. *Phytomed*. 8: 310 322.
- Ames, B. N., Shigenaga, M. K., Hagen, T. M. (1993). Oxidants, antioxidants, and the degenerative diseases of aging. *Proc. Natl. Acad. Sci.* 90: 7915 7922.
- Amresh, G., Reddy, G. D., Rao, C., Singh, P. N. (2007). Evaluation of antiinflammatory activity of *Cissampelos pareira* root in rats. *J Ethnopharmacol*. 110: 526 – 531.
- Amresh, G., Singh, P. N., Rao, C. V. (2007). Antinociceptive and antiarthritic activity of *Cissampelos pareira* roots. *J. Ethnopharmacol.* 111: 531 536.
- Anbar, M., and Gratt, B. M. (1997). Role of nitric oxide in the physiopathology of pain. *J Pain Symptom Manage*. 14: 225 254.
- Anjali, Soni., and Sheetal, Sosa. (2013). Phytochemical Analysis and Free Radical Scavenging Potential of Herbal and Medicinal Plant Extracts. *Journal of Pharmacognosy and Phytochemistry*. 2 (4): 22 29.
- Angela, Caunii., George, Pribac., Ioana, Grozea., Dorin, Gaitin., Ionel, Samfira. (2012). Design of optimal solvent for extraction of bio–active ingredients from six varieties of *Medicago sativa*. *J Chem Ctrl*. 6:123.doi: 10.1186/1752-153X-6-123.
- Anonymous. (2007a). Nature cure senduduk. (Online) available from URL :http://www.forestexplorers.com/naturecures/senduduk.shtml. (Accessed on 13/11/2007).
- Anonymous. (2007b). Database on important medicinal and aromatic plants. (Online) available from http://www.ics.triesta.it/MedicinalPlant/_MedicinalPlant_EthnobotanicalInfo.a sp?id=86 (accessed on 14/11/2007).
- Ardenghi, J. V., Kanegusuku, M., Niero, R., Filho, V. C., Monache, F. D., Yunes R. A., De, Souza, M. M. (2006). Analysis of the mechanism of antinociceptive action of niga-ichigoside F1 obtained from *Rubus imperialis* (Rosaceae). *J Pharm Pharmacol*. 12: 1669 75.

- Aruoma, O. J. (1998). Journal of American Oil Chemical Society. 75: 199 212.
- Audette, J. F., and Bailey, A. (2008). Contemporary Pain Medicine: Integrative Pain Medicine: The Science and Practice of Complementary and Alternative Medicine in Pain Management. Humana Press, Totowa, NJ. 18 33.
- Ayuvedaherbs. (2005). Importance of medicinal plants. ayuvedaherbs.wordpress.com.
- Backonja, M. M., Malan, T. P., Vanhove, G. F., Tobias, J. K. (2010). NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomized, double-blind, controlled study with an open-label extension. *Pain Med.* 11: 600 608.
- Balick, M. J., and Cox, P.A. (1997). Plants People and Culture: the science of ethnobotany. Scientific American Library, New York, NY.
- Balunas, M. J and Kinghorn, D. A. (2005). Drug discovery from medicinal plants. Review article. *Life Sci.*, 78 (5): 431 441.
- Balandrin, M. F., Klocke, J. A, Wurtele, E. S., Bollinger, W. H. (1996). Natural plant chemicals: sources of industrial and medicinal materials. *Science*. 228: 1154 1160.
- Basbaum, A. L., and Woolf, C. J. (1999). *Pain. Curr Biol.* 9, R429 431.
- Beirith A, Santos A.R, Rodrigues, A.L., Creczynski-Pasa, T.B., Calixto, J.B. (1998). Spinal and supraspinal antinociceptive action of dipyrone in formalin, capsaicin and glutamate tests. Study of the mechanism of action. *Eur J. Pharmacol.* 345: 233 45.
- Beirith, A., Santos, A.R., Calixto, J.B. (2002). "Mechanisms underlying the nociception and paw oedema caused by injection of glutamate into the mouse paw". *Brain Research*. 924: 219 228.
- Beirith, A., Santos, A.R, Calixto J.B., Hess, S.C., Messana, I., Ferrari, F., Yunes, R.A. (1999). Study of the antinociceptive action of the ethanolic extract and the triterpene 24-hydroxytormentic acid isolated from the stem bark of *Ocotea suaveolens*. Planta Med. 65:50–5.
- Beissert, S., and Schwarz, T. (2002). Role of immunomodulation of diseases responsive to phototherapy. *Methods*. 28 (1): 138 144.
- Bentley, G.A., Newton, S.H., Starr, J. (1981). Evidence for an action of morphine and the enkephalins on sensory nerve endings in the mouse peritoneum. *Br. J. Pharmacol.* 73: 325 32.

- Bentley, G.A., Newton, S.H., Starr, J. (1983). Studies on the antinociceptive action of 'a-agonist drugs and their interaction with opioid mechanisms. *British Journal of Pharmacology*, 79: 125 134.
- Birt, D.A. (2006). *Journal of the American Dietetic Association*. 106: 20 24.
- Block, L.H. (1964). DMSO: Medicinal and Pharmaceutical Aspects. *Drug and Cosmetic Industry*. 95: 342 465.
- Bors, W., Saran, M. (1987). Radical scavenging by flavonoid antioxidants. *Free Radic Res Commun.* 2: 289 294.
- Braggio, M.M., Lima, M.E.L., Veasey, E.A., Haraguchi, M. (2002). Atividades farmacológicas das folhas da *Sesbania virgata* (Cav.) *Pers. Arq. Inst. Biol.* 69 (4): 49 53.
- Burns, J.P.T., Gardner, D., Mathew, G.G., Duthie, M., Lean, E., Crozier, A. (2001). Journal of Agriculture and Food Chemistry. 49: 5797 – 5808.
- Burkill, I.H. (1966) A Dictionary of the Economic Products of the Malay Peninsular Ministry of Agriculture and Cooperatives, Kuala Lumpur, Malaysia.. 2: 1463 1465.
- Butler, M.S. (2004). The role of natural product chemistry in the drug discovery. J Nat. Prod. 67 (12): 2141 2153.
- Caterina, M.J., Schumacher, M.A., Tominaga, M., Rosen, T.A., Levine, J.D., Julius, D. (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature*. 389: 816 24.
- Caterina, M.J., Julius, D. (2001). A molecular gateway to the pain pathway. *Ann. Rev. Neurosci.* 24 (1): 487 517.
- C.A. Giglio., H.L.A. Defino., C.A. da-Silva., A.S. de-Souza., and E.A. del, Bel. (2006). Behavioral and physiological methods for early quantitative assessment of spinal cord injury and prognosis in rats. *Brazilian Journal of Medical and Biological Research*. 39 (12): 1613 1623.
- Cesare, P., and McNaughton, P. (1997). Peripheral pain mechanisms. *Curr Opin Neurobiol*. 7: 493 499.
- Chas, Bountra., Rajesh, Munglani., & William, K., Schmidt. (2005). Published in The Taylor & Francis e-Library by Marcel Dekker, Inc. (ISBN: 0-8247-8865-6); *PAIN*: Current Understanding, Emerging Therapies, and Novel Approaches to Drug Discovery.
- Chen, Y.F., Tsai, H.Y., Wu, T.S. (1995). Antiinflammatory and analgesic activity from roots of *Angelica pubescens*. *Planta Medica*. 61: 2 8.

- Chhetri HP, Yogo NS, Sherchan J, Anupa KC, Mansoor S and Thapa P (2008). Phytochemical and Antimicrobial Evaluations of some Medicinal Plants of Nepal, Kathmandu University. *Journal of Science, Engineering and Technology*. 1(5): 49 54.
- Choi J, Jung HJ, Lee KT, Park HJ (2005). Antinociceptive and anti-inflammatory effects of the saponin and sapogenins obtained from the stem of *Akebia quinata*. *J Med Food*. 1: 78 85.
- Chopra A (2000). Ayurvedic medicine and arthritis. *Rheum. Disease Clin. N. AM*. 26(1): 133 144.
- C Kokate, AP Purohit, SB Gokhale (2010). Pharmacognosy. *Nirali Prakashan India*. 10.28 10.29.
- Clark SJ, Follenfant RL, Smith TW (1988). Evaluation of opioid-induced antinociceptive effects in anaesthetized and conscious animals. *Br J Pharmacol*. 1:275 283.
- CN Serhan and JZ Haeggstrom (2010). Lipid mediators in acute inflammation and resolution: eicosanoids, PAF, resolvins and proteins. *In fundamentals of Inflammation*, C. N. Serhan, P. A. Ward *et al.*, Eds. 153–174, Cambridge University press, Cambridge, UK.
- CS Patil, NK Jain, A Singh and SK Kulkarni (2003). Modulatory effect of cyclooxygenase inhibitors on sildenafilinduced antinociception. *Pharmacology.* 69 (4): 183 189.
- CS Kalaivani, S Sahaya Sathish, N Janakiraman, M Jonhson (2012). GC-MS studies on *Andrographis paniculata* (Burm. F) Wall. Ex Nees medicinally important plant. *Int. J. Med. Arom. Plants*. 2 (1): 69 74.
- Corner EJH (1951). Wayside Trees of Malaya. Malayan Nature Society: Kuala Lumpur. 1: 445 453.
- Cortright DN, Szallasi A (2004). Biochemical pharmacology of the vanilloid receptor TRPV1. An update. *Eur J Biochem.* 271: 1814 1819.
- Cottele N (2001). Role of flavonoids in oxidative stress. *Curr. Top. Med. Chem.* 1: 569 590.
- Cui M, Honore P, Zhong C, Gauvin D, Mikusa J, Hernandez G, Faltynek CR (2006). TRPV1 receptors in the CNS play a key role in broad-spectrum analgesia of TRPV1 antagonists. *J Neuro*. 37: 9385 9393.
- Cunha TM, Roman-Campos D, Lotufo CM, Duarte HL, Souza GR, Verri WA Jr, Funez MI, Dias QM, Schivo IR, Domingues AC, Sachs D, Chiavegatto S, Teixeira MM, Hothersall JS, Cruz JS, Cunha FQ, Ferreira SH (2010). Morphine peripheral analgesia depends on activation of the

- PI3Kγ/AKT/nNOS/NO/K_{ATP} signaling pathway. *Proc Nat Acad Sci USA*. 107: 4442 4447.
- Cury Y, Picolo G, Gutierrez VP, Ferreira SH (2011). Pain and analgesia: the dual effect of nitric oxide in the nociceptive system. *Nitric Oxide*. 3: 243 254.
- Dai J and Mumper RJ (2010). Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 15: 7313 7352.
- Dalimartha S (2000). Atlas tumbuhan Obat Indonesia. Jilid I. Jakarta: Tubus Agriwidya. 130 132.
- Damas J, Liegeois JF (1999). The inflammatory reaction induced by formalin in the rat paw. *Naunyn Schmiedebergs Arch Pharmacol.* 359: 220 227.
- Daniel P Alford, Peggy Compton and Jeffrey H. Samet (2006). Acute Pain Management for Patients Receiving Maintenance Methadone or Buprenorphine Therapy. Ann Intern Med. 144(2): 127–134
- Dawson, Taylor and Reide (2002). Pharmacology, 2nd Edition, Elsevier Science Limited. 3 18.
- D Begun and SC Nath (2000). Ethnobotanical review of medicinal plants used for skin diseases and related problems in Northeastern India. *Journal of Herbs*, Species and Medicinal Plants. 7: 55 93.
- DC Pope and WT Oliver (1966). Dimethyl Sulfoxide (DMSO). Can. J. Med. Vet. Sci. Vol. 30: 3 8.
- Déciga-Campos M, Montiel-Ruiz RM, Navarrete-Vázquez G, López-Muñoz FJ (2007).. Palmitic acid analogues exhibiting antinociceptive activity in mice. *Proc West Pharmacol Soc.* 50: 75 7.
- de Oliveira AM de Araújo AF, Lyra Lemos RP, Conserva LM, de Souza Ferro JN, Barreto E (2015). Antinociceptive and anti-inflammatory activity of the siaresinolic acid, a triterpene isolated from the leaves of *Sabicea grisea Cham. & Schltdl. var. grisea. J Nat Med.* 69:232–40.
- Derardt R, Jougney S, Delevalcese F, Falhout M (1980). Release of prostaglandins E and F in an algogenic reaction and it inhibition. *Eur J Pharmacol*. 51: 17 24.
- Dewan SMR, Amin MN, Adnan T, Uddin SMN, Shahid-Ud-Daula AFM, Sarwar G, Hossain MS (2013). Investigation of analgesic potential and in vitro antioxidant activity of two plants of Asteraceae family growing in Bangladesh. *J Pharm Res.* 6: 599 603.
- Dickson M and Gagnon JP (2004). Key factor in the rising cost of new drug discovery and development. *Nat. Rev. Drug Discovery*. 3 (5): 417 429.

- Dickenson AH, Sullivan AF (1987). Evidence for a role of the NMDA receptor in the frequency dependent potentiation of deep rat dorsal horn nociceptive neurones following C fibre stimulation. *Neuropharmacol.* 26: 1235 1238.
- DL Jackson, CB Craff, JD Richardson, KM Hargreaves (1995). Glutamate participates in the peripheral modulation of thermal hyperalgesia in rats. *Eur. J. Pharmacol.* 284: 661 664.
- Downey MO, Rochfort S (2008). Simultaneous separation by reversed-phase high performance liquid chromatography and mass spectral identification of anthocyanins and flavonols in Shiraz grape skin. *J. Chromatogr.* 1201: 43 47.
- Du J, Yu Y, Ke Y, Wang C, Zhu L, Qian ZM (2007). Ligustilide attenuates pain behaviour induced by acetic acid or formalin. *J. Ethnopharmacol*. 112: 211 214.
- Duarte I, Lorenzetti B, Ferreira S (1990). Peripheral analgesia and activation of the nitric oxide-cyclic GMP pathway. *Eur J Pharmacol.* 186:289 293.
- Duarte IDG, Nakamura M, Ferreira SH (1988). Participation of the sympathetic system in acetic acid-induced writhing in mice. *Braz J Med Biol Res.* 21: 341–43.
- Ehab S, Desoky EL, Ihab A, Fouad (2005). Pharmacological evidence for the role of nitric oxide cGMP in antinociception. *J Appl Res.* 5: 451 9.
- EK Perimal, MN Akhtar, AS Mohamad, MH Khalid, OH Ming, S Khalid, LM Tatt, MN Kamaldin, ZA Zakaria, DA Israf, N Lajis, MR Sulaiman (2011). Zerumbone-induced antinociception: involvement of the L-arginine-nitric oxide-cGMP-PKC-K+ ATP channel pathways. *Basic Clin. Pharmacol. Toxicol.* 108: 155 162.
- Ellof JN (1998). Which extractant should be used for the screening and isolation of antimicrobial components from plants?. *J Ethnopharmacol*. 60: 1 6.
- Emily O Dumas and Gary M Pollack (2008). Opioid Tolerance Development: A Pharmacokinetic/Pharmacodynamic Perspective. AAPS J. 10(4): 537.
- ES Jaios, Abdul Rahman S, Ching SM, Arifah AK, Desa MN, Zakaria ZA (2016). Possible mechanisms of antinociception of methanol extract of *Melastoma malabathricum* Leaves. *Braz J Pharmacognocy*. doi.org/10.1016/j.bjp.2016.01.011.
- FA Abdul Majid and LY Ting (2011). "Basic in vitro studies of human skin culture: *Melastoma malabathricum* extract on fibroblasts growth," February 2011, http://eprints.utm.my/2850/1/75127.pdf.

- Fasihuddin B. Ahmad and Ghazally Ismail (2003). *Medicinal Plants Used by Kadazandusun Communities around Crocker Range*. ASEAN Review of Biodiversity and Environmental Conservation (ARBEC).
- Farnsworth NR and Soejarto DD (1985). Potential consequence of plant extinction in the United States on the current and future availability of prescription drugs. *Economic Botany.* 39: 231 240.
- Ferdous M, Rouf R, Shilpi JA, Uddin SJ (2008), Antinociceptive activity of the ethanolic extract of *Ficus racemosa* Linn. (Moraceae). *OPEM*. 8: 93 96.
- Ferreira SH, Duarte IDG and Lorenzetti BB (1991). The Molecular Mechanism of Action of Peripheral Morphine Analgesia: Stimulation of the cGMP System via Nitric Oxide Release. *European Journal of Pharmacology*. 201: 121 122.
- Ferreira J, da Silve GL, Pizzolatti MG, Kassuya CAL, Calixto JB and Santos ARS (2006). Contribution of vanilloid receptors to the overt nociception induced by B2 kinin receptor activation in mice. *British journal of pharmacology*. 141: 787 794.
- Fine PG and Ashburn MA (1998). Functional Neuroanatomy and Nocieption. In Ashburn MA, Rice LJ (editors). The Management of Pain. Churchill Livingstone, Philadelphia. 1 16.
- F Yahya, SS Mamat, MFF Kamarolzaman, AA Sayedan, KF Jakius, ND Mahmood, MS Shahril, Z Suhaili, N Mohtaruddin, D Susanti, MN Somchit, LK Teh, MZ Salleh and ZA Zakaria. (2013). Hepatoprotective Activity of Methanolic Extract of *Bauhinia purpurea* Leaves against Paracetamol-Induced Hepatic Damage in Rats. *Evidence-Based Complimentary and Altenative Medicine*. 636580: 1 10.
- Fezai M, Senovilla L, Jemaà M, Ben-Attia M (2013). Analgesic, anti-inflammatory and anticancer activities of extra virgin olive oil. *J Lipids*. 2013: 129736.
- Fundytus ME (2001). Glutamate receptors and nociception: implications for the drug treatment of pain. CNS Drug. 1: 29 58.
- Ghani A (2003). "Medicinal plants of Bangladesh with chemical constituents and users", 2nd edn.: Asiatic Society of Bangladesh, Dhaka, Rama. 42.
- GA Bentley, SH Newton and J Starr (1981). Evidence for an action of morphine and the enkephalins on sensory nerve endings in the mouse peritoneum. *British Journal of Pharmacology*. 73(2): 325 332.
- Gaurav K, Karthik L and Bhaskara RKV (2011). A Review on Pharmacological and Phytochemical Properties of *Zingiber officinale* Roscoe (Zingiberaceae). *Journal of Pharmacy Research*. 4(9): 2963 2966.

- Garthwaite J and Boulton CL (1995). Nitric Oxide Signaling in the Central Nervous System. *Annual Review of Physiology*. 57: 683 706.
- G Bhave, F Karim, SM Carlton and RW Gereau IV (2001). "Peripheral group I metabotropic glutamate receptors modulate nociception in mice," *Nature Neuroscience*. 4(4): 417 423.
- GB Martini-Bettolo (1980). Present aspects of the use of plants in traditional medicine. *Journal of Ethnopharmacology*. 2(1): 5 7.
- Geysen, HM Schoenen, F Wagner, D Wagner R (2003). Combinatorial compound libraries for drug discovery: an ongoing challenge. *Nat. Rev. Drug Discovery*, 2 (3): 222 230.
- George JA and Joseph K (2012). Antinociceptive and anti-inflammatory activities of leaf methanol extract of *Cotyledon orbiculata* L. (Crassulaceae). *Advances in Pharmacological Sciences*. 862625: 1 10.
- George NJ, Obot IB, Ikot AN, Akpan AE & Obi-Egbed NO (2010). Phytochemical and Antimicrobial Properties of Leaves of *Alchonea Cordifolia*. *EJournal of Chemistry*. 7(3): 1071 1079.
- GG Vivancos, CA Parada and SH Ferreira (2003). "Opposite nociceptive effects of the arginine/NO/cGMP pathway stimulation in dermal and subcutaneous tissues," *British Journal of Pharmacology*. 138(7): 1351 1357.
- Gibuła-Bruzda E (2010). Animal models of nociceptive pain. *Pharmacy and Medical Sciences*, XXII (4): 75 85.
- Giglio CA, Defino HLA, Da-Silva CA, De-Souza AS, Del Bel EA (2006). Behavioral and physiological methods for early quantitative assessment of spinal cord injury and prognosis in rats. *Braz J Med Bio Res.* 39: 1613 1623.
- Gloria A Ayoola, Solomon S Ipav, Margaret O Sofidiya, Aderonke A Adepoju-Bello, Herbert AB Coker, Tolu O Odugbemi (2008). Phytochemical Screening and Free Radical Scavenging Activities of the Fruits and Leaves of *Allanblackia floribunda* Oliv (Guttiferae). *International Journal of Health Research*. 1(2): 87 93.
- Goldberg MR, Tung CS, Robertson D (1983). Influence of alphamethyldopa treatment on adrenergic receptor binding in rat forebrain. *Clin Exp Hypertens*. A 5: 1589 1596.
- Goncalves CE, Araldi D, Panatieri RB, Rocha JB, Zeni G and Nogueira CW (2005). Antinociceptive properties of acetylenic thiophene and furan derivatives: evidence for the mechanism of action. *Life Sciences*. 76: 2221 2234.

- Gonzalez-Trujano ME, Pena EI, Martinez AL, Moreno J, Guevara-Fefer P, Deciga-Campos M *et al.*, (2007). Evaluation of the antinociceptive effect of *Rosmarinus officinalis* L. using three different experimental models in rodents. *J Ethnopharmacol*. 111: 476 482.
- Goudet C, Chapuy E, Alloui A, Acher F, Pin JP, Eschalier A (2008). Group III metabotropic glutamate receptors inhibit hyperalgesia in animal models of inflammation and neuropathic pain. *Pain*. 137: 112 124.
- Gureje O, Von Korff M, Simon GE, Gater R. (1998). Persistent pain and well-being: a World Health Organization Study in Primary Care. *JAMA*. 280: 147 51.
- Gurib-Fakim A (2006). Medicinal plants: Tradition of yesterday and drugs of tomorrow. *Review article. Mol. Aspects Med.* 27 (1):1 93.
- Cragg GM and Newman DJ (2005). Biodiversity: A continuing source of novel drug leads. *Pure App Chem.* 77 (1): 7 24.
- Granados-Soto V, Rufino MDO, Gomes Lopes LD, Ferreira SH (1997). Evidence for the involvement of the nitric oxide-cGMP pathway in the antinociception of morphine in the formalin tests. *Eur J Pharmacol*. 2 3: 177 180.
- Graul Al (2000). The year's new drugs. *Drug new Perspect.* 14 (1):12 31.
- GS Kelly (1999). *Altern. Med. Rev.* 4: 29 36.
- Hagiwara K, Nojima H and Kuraishi Y (1999). Serotonin-induced biting of the hind paw is itch-related response in mice. *Neuroscience Letter*. 37: 17 22.
- H Hosseinzadeh and HM Younesi (2002). Antinociceptive and anti-inflammatory effects of *Crocus sativus* L. stigma and petal extracts in mice. *BMC Pharmacology*, 2: 7.
- Halliwell B, Gutteridge JMC, Cross CE (1992). Free radicals, antioxidants, and human disease: where are we now? *J Lab Clin Med*. 119: 598 620.
- Halliwell B (2008). Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and in vivo studies? Arch. Biochem. Biophys. 476: 107 112.
- Han T, Zhang QY, Zhang H, Wen J, Wang Y, Huang BK, Rahman K, Zheng HC and Qin LP (2009). Authentication and quantitative analysis on the chemical profile of Xanthium fruit (Cang-Er-Zi) by high-performance liquid chromatography-diodearray detection tandem mass spectrometry method. *Analytica Chimica Acta*. ISSN 0003-2670. 634 (2): 272 278.
- Harborne JB, Williams CA (2000). Advances in flavonoids research since 1992, *Phytochemistry*. 55: 481 504.

- Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH, Alam MA, Faruque A, Rana S, Rahman S (2010). Analgesic activity of the different fractions of the aerial parts of *Commelina benghalensis* Linn, *International J Pharmacol*. 6: 63 67.
- Hasler G (2004). Discovering endophenotypes for major depression. Neuropsychopharmacology. 29: 1765 – 1781.
- Hassan Azaizeh, Bashar Saad, Edwin Cooper and Omar Said. (2010). Review: Traditional Arabic and Islamic Medicine, a Re-Emerging Health Aid. Evidence-Based Complementary and Alternative Medicine. 7(4): 419 424.
- Helnrich M, Barnes J, Gibbons S, Williamson EM (2004). Fundamentals of Pharmacognosy and Phytotherapy. Churchill Livingston, Elsevier Science Ltd., UK.
- Heldt H-W, Heldt F (2005). Secondary metabolites fulfil specific ecological functions in plants. *Plant Biochemistry. Academic Press, Burlington, USA*. 3: 403 412.
- Hemani T, Panihar MS (1998) Reactive oxygen species and oxidative property of *Nardostachys jatamanasi. Indian J. Expt. Biol.* 34: 1150 1151.
- Henry Hitner and Barbara T Nagle (1999). Basic Pharmaclogy, 4th Edition (Glencoe MacGraw-Hill).
- Herdegen T, Rudiger S, Mayer B, Bravo R and Zimmermann M (1994). Expression of Nitric Oxide Synthase and Colocalisation with Jun, Fos and Krox Transcription Factors in Spinal Cord Neurons Following Noxious Stimulation of the Rat Hindpaw. *Molecular Brain Research*. 22: 245 258.
- HG Vogel and WH Vogel (1997). Pharmacological assays in Drug Discovery and Evaluation. J. A. Majors Company, Lewisville, Texas, USA. 360 418
- HO Collier, LC Dinneen, CA Johnson and C Schneider (1968). The abdominal constriction response and its suppression by analgesic drugs in the mouse. *British Journal of Pharmacology*. 32 (2): 295 310.
- Hosseinzadeh H, Younesi HM (2002). Antinociceptive and anti-inflammatory effects of *Crocus sativus* L. stigma and petal extracts in mice, *BMC Pharmacol*. 2: 18.
- HP Rang, MM Dale, JM Ritter, RJ Flower and G Henderson. (2011). *Pharmacology*, Elsevier Churchill Livingstone, Edinburgh, UK, 7th edition.
- Hui Ming Ong, Azam Shah Mohamad, Noor Adilah Makhtar, Mohamed Hanief Khalid, Syamimi Khalid, Enoch Kumar Perimal, Siti Nurulhuda Mastuki, Zainul Amiruddin Zakaria, Nordin Lajis, Daud Ahmad Israf, Mohd Roslan Sulaiman (2010). Antinociceptive activity of methanolic extract of *Acmella uliginosa* (Sw.) Cass. *Journal of Ethnopharmacology*. 227 233.

- Hunskaar S, Hole K (1987). The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. *Pain*. 30: 103 114.
- Hurley RW, Grabow TS, Tallarida RJ and Hammond DL (1999). Interaction between medullary and spinal delta 1 and delta 2 opioid receptors in the production of antinociception in the rat. *J. Pharmacol Exp Ther.* 289: 993 999.
- Ibironke GF, Ajiboye KI (2007). Studies on the anti–inflammatory and analgesic properties of Chenopodium ambrosioides leaf extract in rats. *Int. J. Pharmacol.* 3: 111 115.
- I Cock, WN Setzer, KD Ruebhart, OA El Dahshan, M Tomczyk (2009). An antidiabetic and hypolipidemic effects from *Azadirachta Indica* leaves, *African Journal of Biotechnology*. 8 (13): 3084 – 3091.
- IDG Duarte, BB Lorenzetti and SH Ferreira (1990). Peripheral analgesia and activation of the nitric oxide-cyclic GMP pathway. *European Journal of Pharmacology*, vol. 186, no. 2-3: 289 293.
- IDG Duarte and SH Ferreira (2000). L-NAME cause antinociception by stimulation of the arginine-NO-cGMP pathway. *Mediators of Inflammation*. 9: 25 30.
- IH Burkill (1966). "A Dictionary of Economic Products of Malay Peninsular," Ministry of Agriculture and Cooperates Malaysia, Kuala Lumpur. 1463 1465.
- Ikhiri K, Boureima D and Dan-Kouloudo D (1992). Chemical screening of medicinal plants used in the traditional pharmacopoeia of Niger. *Int J Pharmacog*. 30:251–62.
- Ioannides C (2002). Pharmcokinetic interactions between herbal remedies and medicinal drugs. *Xenobiotica*. 32: 451 478.
- Jaganath IB and LT Ng (2002). Herbs: The Green Pharmacy of Malaysia. Published by Vinpress Sdn. Bhd. in collaboration with Malaysia Agricultural Research and Development Institute (MARDI).
- Jain NK, Patil CS, Singh A, Kulkarni SK (2003). Sildenafil-induced peripheral analgesia and activation of the nitric oxide-cyclic GMP pathway. *Brain Res*. 909: 170 8.
- Jaijoy K, Soonthornchareonnon N, Lertprasertsuke N, Panthong A Sireeratawong S (2000) Acute and Chronic oral toxicity of standardized water extract from fruit of Phyllanthus emblica Linn. *International Journal of Applied and Basic Medical Research*, 3: 45 58.

- James Hamuel Doughari (2012). Phytochemicals: Extraction Methods, Basic Structures and Mode of Action as Potential Chemotherapeutic Agents, Phytochemicals A Global Perspective of Their Role in Nutrition and Health, Dr Venketeshwer Rao (Ed.), ISBN: 978-953-51-0296-0.
- Janna OA, A Khairul, M Maziah and Y Mohd (2006). Flower pigment analysis of *Melastoma malabathricum. Afr. J. Biotechnol.* 5: 170 174.
- Jayaprakasha GK and Rao J (2000) Phenolic constituents from lichen Parmontrema stuppeum. Hale and antioxidant activity. *Zeitschrift Für Naturforscung*. 55: 1018 1022.
- Jayanthi P, Lalitha P & Sripathi SK (2011). Phytochemical investigation of the extracts of *Eichhornia crassipes* and its solvent fractionates. *Journal of Pharmacy Research*. 4(5): 1405 1406.
- Jean-Pierre Nowicki and Bernard Scatton (2008). *The Practice of Medicinal Chemistry*, 3rd Edition. 82 83.
- Jesse CR, Rocha JB, Nogueira CW, Savegnago L (2009). Further analysis of the antinociceptive action caused by p-methoxyldiphenyl diselenide in mice. *Pharmacol Biochem Behav.* 91: 573–80.
- J Ferreira, ARS Santos, JB Calixto (1999). The role of systemic, spinal and supraspinal L-arginine-nitric oxide-cGMP pathway in thermal hyperalgesia caused by intrathecal injection of glutamate in mice. *Neuropharmacology*. 38: 835 842.
- J Garthwaite and CL Boulton (1995). Nitric oxide signaling in the central nervous system. *Annual Review of Physiology*. 57: 683 706.
- Jhaveri MD, Elmes SJR, Kendall DA, Chapman V (2005). Inhibition of peripheral vanilloid TRPV1 receptors reduces noxious heat-evoked responses of dorsal horn neurons in naïve, carrageenan-inflamed and neuropathic rats. *Eur J Neuro*. 2: 361 370.
- Jin Dai and Russell J Mumper (2010). A Review: Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties. *Molecules*.;doi:10.3390/molecules15107313. 15: 7313 7352.
- JLCH van Valkenberg and N Bunyapraphatsara. (2001). "*Melastoma malabathricum* L.," in Plant Resources of South-East Asia No. 12(2): Medicinal and Poisonous Plants 2, JLCH van Valkenburg and N Bunyapraphatsara, Eds., Backhuys, Leiden, The Netherlands. 365 366.
- Jimuty Roy, MD Ruhul Kuddus, Bilkis Begum and Choudhury M Hasan (2012). Evaluation of analgesic, cytotoxic and antioxidant activities of *Blumea Membranacea* DC. *Int J Pharm Bio Sci.* 3 (4): 566 572

- J Moa, DD Price, RL Hayes, J Lu, DJ Mayer (1992). Differential roles of NMDA and non-NMDA receptor activation in induction and maintenance of thermal hyperalgesia in rats with painful peripheral mononeuropathy. *Brain Res.* 598: 271 278.
- Joa°o B Calixto, Alessandra Beirith, Juliano Ferreira, Adair R S Santos, Valdir Cechinel Filho and Rosendo A Yunes (2000). Review Article: Naturally Occurring Antinociceptive Substances from Plants. *Phytother. Res.* 14: 401 418.
- Joy PP, Thomas J, Mathew S, Skaria BP (1998). Medicinal Plants. Kerala Agricultural University, Aromatic and Medicinal Plants Research Station, Kerala, India, PIN: 683 549: 3.
- Jungi T, Adler H, Thöny M, Krampe M and Peterhans E (1996). Inducible Nitric Oxide Synthase of Macrophages: Present Knowledge and Evidence for Species-specific Regulation. *Veterinary Immunology and Immunopathology*. 54: 323 330.
- Jun Zheng Goh, Sook Nai Tang, Hoe Siong Chiong, Yoke Keong Yong, Ahmad Zuraini, Muhammad Nazrul Hakim (2015). Evaluation of antinociceptive activity of nanoliposome-encapsulated and free-form diclofenac in rats and mice. International Journal of Nanomedicine. 10: 297–303.
- Kamisan FH, Yahya FA, Mamat SS, Kamarolzaman MFF, Suhaili Z, Mohtarrudin N, et al., (2014). Effect of Methanol Extract of Dicranopteris linearis Against Carbon Tetrachloride-induced Acute Liver injury in Rats. BMC Compliment Alt Med. 14: 123.
- Kamba AS & Hassan LG (2010). Phytochemical Screening and Antimicrobial Activities of *Euphorbia balasamifera* Leaves Stems and Root against Some Pathogenic Microorganisms. *African Journal of Pharmaceutical Sciences and Pharmacy*. 4(9): 645 652.
- Kalayou S, M. Haileselassie, G. Gebre-Egziabher S. Sahle and H. Taddele *et al.* (2012). *In-vitro* antimicrobial activity screening of some ethnoveterinary medicinal plants traditionally used against mastitis, wound and gastrointestinal tract complication in Tigray Region, Ethiopia. *Asian Pacific J. Tropical Biomed.*, 2: 512-522. DOI: 10.1016/S2221-1691(12)60088 4.
- Kapoor LD (1990). CRC Handbook of Ayurvedic medicinal plants. CRC Press, Boca Raton, USA.
- Kara Capriotti MD and Joseph A capriotti MD (2012). Dimethyl Sulfoxide History, Chemistry, and Clinical Utility in Dermatology. *The Journal of Clinical and Aesthetic Dermatology*. Vol. 5(9): 1 3.

- Kathryn Bayne (2000). Assessing Pain and Distress: A Veterinary Behaviourist's Perspective. Definition of Pain and Distress and Reporting Requirements for Laboratory Animals (Proceedings of the Workshop Held June 22, 2000).
- Kawabata A, Umeda N, Takagi H (1993). L-Arginine Exerts a Dual Role in Nociceptive Processing in the Brain: Involvement of the Kyotorphinmet-enkephalin Pathway and NO-cyclic GMP Pathway. *Br J Pharmacol*.109: 73 79.
- Kawabata A, Manabe S, Manabe Y and Takagi H (1994). Effect of Topical Administration of L-arginine on Formalin-induced Nociception in the Mouse: A Dual Role of Peripherally Formed Nitric Oxide in Pain Modulation. *British Journal of Pharmacology*. 112: 547 550.
- K Meyer (2001). "Revision of the Southeast Asian genus Melastoma (Melastomataceae)," Blumea: Journal of Plant Taxonomy and Plant Geography. 46(2): 351 398.
- Khairatkar-Joshi N, Szallasi A (2009). TRPV1 antagonists: the challenges for therapeutic targeting. *Trends in Mol Med*. 1: 14 22.
- KH Ling, CT Kian and TC Hoon (2009). A Guide to Medicinal Plants. An Illustrated, Scientific and Medicinal Approach, World Scientific, Singapore.
- Klejdus B, Mikelova R, Adam V, Zehnalek J, Vacek J, Kizek R (2004). Liquid chromatographic-mass spectrophotometric determination of genistin and daidzin in soybean food samples after accelerated solvent extraction with modified content of extraction cell. *Anal Chimica Acta*. 517: 1 11.
- Klejdus B, Mikelova R, Petrolova J, Potesil D, Adam V, Stiborova M (2005). Evaluation of isoflavone aglycon and glycoside distribution in soy plants and soybeans by fast column high-performance liquid chromatography coupled with a diode-array detector. *J Agr and Food Chem.* 53: 5848 5852.
- Kumar S, SA Hassan, S Dwivedi, AK Kukreja, A Sharma, AK Singh, S Sharma and R Tewari (eds). (2000). Proceedings of the National Seminar on the Frontiers of Research and Development in Medicinal Plants, 16 18 September 2000. *Journal Medicinal and Aromatic Plant Sciences*. Vol. 22/4A and Vol. 23/1A. Central Institute of Medicinal and Aromatic Plants (CIMAP). Lucknow, India.
- Kumar VK, Kumar PS, Rajan M, Kumar AV, Boppana R, Reddy PS & Alzeber HFH (2011). Qualitative Phytochemical analysis of *Bauhinia tomentosa* Linn flower by HPTLC. *Journal of Pharmacy Research*. 4(9): 2868 2880.
- Koehn FE and Carter GT (2005). The evolving role of natural products in drug discovery. *Nat. Rev. Drug Discovery.* 4 (3): 206 220.

- K Omote, T Kawamata, M Kawamata, A Namiki (1998). Formalin-induced release of excitotary amino acids in the skin of the rat hindpaw. *Brain Res.* 787: 161 164.
- Koay SS (2008). Establishment of cell suspension culture of *Melastoma* malabathricum L. for the production of anthocyanin, PhD. Thesis. Pulau Pinang, Malaysia: *Universiti Sains Malaysia*.
- Kou J, Ni Y, Li N, Wang J, Liu L, Jiang Z.H (2005). Analgesic and antiinflammatory activities of total extract and individual fractions of Chinese medicinal plant Polyrhachis lamellidens. *Biol. Pharm. Bull.* 28: 176 – 180.
- Lamont LA, Tranquilli WJ and Grimm KR (2000). Physiology of pain. Veterninary Clinics of North America: Small Animal Practice. Vol. 3, no. 3.
- Larson AA, Koyacs KJ, Cooper JC, Kitto KF (2000). Transient changes in the synthesis of nitric oxide result in long-term as well as short term changes in acetic acid-induced writhing in mice. *Pain*. 86: 103 11.
- Le Bars D, Gozariu M, Cadden SW (2001). Animal models of nociception. *Pharmacol. Rev.* 53: 597 652.
- Levine JD (1998). New directions in pain research: molecules to maladies. *Neuron*. 20: 649 54.
- Ley SV and Baxendale IR (2002). New tool and concepts for modern organic synthesis. *Nat. Rev. Drug Discovery.* 1(8): 573 586.
- Li SZ (2003). Compendium of materia medica (Bencao Gangmu) (1st edition), Foreign Language Press, ISBN 9787119032603, Beijing.
- Lombardino JG and Lowe III JA (2004). The role of medicinal chemist in drug discovery-then and now. *Nat. Rev. Drug Discovery*. 3(10): 853 862.
- Longhi-Balbinot DT, Martins DF, Lanznaster D, SilvaMD, Facundo VA, Santos AR (2011). Further analyses of mechanisms underlying the antinociceptive effect of the triterpene 3β, 6β, 16β-trihydroxylup-20(29)-ene in mice. *Eur J Pharmacol.* 653:32–40.
- Loeser JD, Melzack R (1999). Pain: an overview. *Lancet*. 353: 1607 1609.
- Loeser JD and Treede RD (2008). The Kyoto protocol of IASP basic pain terminology. *Pain*. 137: 473 477.
- Lopes-Lutz D, Alviano SD, Alviano SC, Kolodziejczyk PP (2008). *Phytochemistry*. 69: 1732 1738.

- Machelska H, Labuz D, Przewlocki R and Przewlocka B (1997). Inhibition of Nitric Oxide Synthase Enhances Antinociception Mediated by *Mu*, *Delta* and *Kappa* Opioid Receptors in Acute and Prolonged Pain in the Rat Spinal Cord. *Journal of Pharmacology and Experimental Therapeutics*. 282: 977 984.
- Mamat SS, Kamarolzaman MFF, Yahya F, Mahmood ND, Shahril MS, Jakius KF, Mohtarrudin N, Ching SM, Deny S, Taher M, Zakaria ZA (2013). Methanol extract of *Melastoma malabathricum* (L) leaves exerted antioxidant and liver protective activity in rats. *BMC Complement Alternate Med.* 13:326.
- Manicam C, Abdullah JO, Mohd Tohit ER, Seman Z, Sieo CC, Hamid M (2010). In vitro anticoagulant activities of *Melastoma malabathricum* (L) aqueous leaf extract: a preliminary novel finding. *J Med Plants Res.* 4:1464–72.
- M Déciga-Campos, FJ López-Muñoz (2004). Participation of the L-arginine-nitric oxide-cyclic GMP-ATP-sensitive K+ channel cascade in the antinociceptive effect of rofecoxib. *Eur. J. Pharmacol.* 484: 193 199.
- Md Shihab Hasan, Md Iqbal Ahmed, Sukla Mondal, Shaikh Jamal Uddin, Mohammad Mehedi Masud, Samir Kumar Sadhu and Masami Ishibashi (2006). Antioxidant, antinociceptive activity and general toxicity study of *Dendrophthoe falcata* and isolation of quercitrin as the major component. *Oriental Pharmacy and Experimental Medicine*. 6(4): 355 360
- Mahadevan N, Shivali, Kamboj P (2009). *Natural Product Radiance*. 8 (1): 77 83.
- Magner LN (1992). A history of medicine. Marcel Dekker Inc: New York.
- Marchand F, Perretti M, McMahon SB (2005). Role of the immune system in chronic pain. *Nat Rev Neurosci.* 6: 521 532.
- Maria de Lourdes Reyes-Escogido, Edith G Gonzalez-Mondragon and Erika Vazquez-Tzompantzi (2011). A Review: Chemical and Pharmacological Aspect of capcaisin. *Molecules*. 16: 1253 1270.
- Marchioro M, Blank MFA, Mourão RHV, Antoniolli AR (2005). Antinociceptive activity of the aqueous extract of *Erythrina velutina* leaves. *Fitoterapia*. 76: 637 642.
- Martínez AL, González-Trujano ME, Chávez M, Pellicer F (2012). Antinociceptive effectiveness of triterpenes from rosemary in visceral nociception. *J Ethnopharmacol*. 1: 28 34.
- Max MB, Payne R and Edwards WT (1999). Principle of Analgesic Use in the Treatment of Acute Pain and Cancer Pain. 4th ed. Glenview, IL: *American Pain Society*.
- McQuay H, Moore A, Justins D (1997). Fortnightly review: treating acute pain in hospital. *BMJ*. 314: 1531.

- McCurdy CR, Scully SS (2005). Analgesic substances derived from natural products (natureceuticals). *Life Sci.* 78: 476 484.
- McNamara CR, Mandel-Brehm J, Bautista DM, Siemens J, Deranian KL, Zhao M (2007). TRPA1 mediates formalin-induced pain. *Proc Natl Acad Sci.* 104: 13525 30.
- Md Abdul Muhit, Syed Mohammed Tareq, Apurba sarker Apu, Debasish Basak and Mohammad S. Islam (2010). Isolation and Identification of Compounds from the Leaf Extract of Dillenia indica Linn. *Bangladesh Pharmaceutical Journal*. ISSN no.: 0301-4606. Vol. 13: 1.
- Megalah Thevi Rajenderan (2010). Ethno Medicinal Uses and Antimicrobial Properties of *Melastoma malabathricum*. ISSN 1985-5672. 3(2): 34 44.
- Merskey H and Bugduk N (1994). Classification of Chronic Pain. Descriptions of Chronic Pain Syndromes and Definitions of Pain Terms.
- Merken HM, Beecher GR (2000). Measurement of food flavonoids by high-performance liquid chromatography: A review. J. Agric. Food Chem. 48: 577 599.
- Millan MJ (1999). The induction of pain: an integrative review. *Prog. Nuerobiol.* 57: 1 164.
- Misha-Miroslav Backonja MD (2002) Use of anticonvulsants for treatment of neuropathic pain. *Neurology*. 5: 14 17.
- MJ Hudspith (1997). Glutamate: a role in normal brain function, anaesthesia, analgeisic and CNS injury. *Br. J. Anaesth.* 79: 731 747.
- Mohamed Hanief Khalid, Muhammad Nadeem Akhtar, Azam Shah Mohamad, Enoch Kumar Perimal, Ahmad Akira, Daud Ahmad Israf, Nordin Lajis, Mohd Roslan Sulaiman (2011). Antinociceptive effect of the essential oil of *Zingiber zerumbet* in mice: Possible mechanisms. *Journal of Ethnopharmacology*. 137: 345 351.
- Mohd Sani MH, Zakaria ZA, Balan T, Teh LK and Salleh MZ (2012). *Antinociceptive Activity of Methanol Extract of Muntingia calabura* Leaves and the Mechanisms of Action Involved. *Evidence-Based Complementary and Alternative Medicine*. 890361: 1 10.
- Mohd Yusuf, M Teh L, Salleh M and Zakaria ZA (2011). Antinociceptive activity of *Muntingia calabura* leaves. *Planta. Med.* 77 PM38.

- Mohamad Yusof MI, Salleh MZ, Teh LK, Ahmat N, Nik Azmin NF, Zakaria ZA (2013). Activity-guided isolation of bioactive constituents with antinociceptive activity from *Muntingia calabura* L. leaves using the formalin test. *Evidence Based Complementary Alternative Medicine.*, 715074, http://dx.doi.org/10.1155/2013/715074
- Momin A, McNaughton PA (2009). Regulation of firing frequency in nociceptive neurons by pro-inflammatory mediators. *Exp Brain Res.* 196 (1): 45 52
- Moncada S, Palmer RMJ and Higgs EA (1989). Biosynthesis of Nitric Oxide from Larginine: A Pathway for the Regulation of Cell Function and Communication. *Biochemical Pharmacology* 38: 1709 1715.
- Moreira J, Klein-Júnior LC, Cechinel Filho V, de Campos Buzzi F (2013). Antihyperalgesic activity of corilagin, a tannin isolated from *Phyllanthus niruri* L. (Euphorbiaceae). *J Ethnopharmacol*. 1: 318 23.
- Morioka N, Inoue A, Hanada T, Kumagai K, Takeda K, Ikoma K, Hide I, Tamura Y, Shiomi H, Dohi T, Nakata Y (2002). Nitric oxide synergistically potentiates interleukin-1β-induced increase of cyclooxygenase-2 mRNA levels, resulting in the facilitation of substance P release fromprimary afferent neurons: involvement of cGMP-independent mechanisms. *Neuropharmacol.* 5: 868 876.
- MT Rajenderan (2010). "Ethno medicinal uses and antimicrobial properties of *Melastoma malabathricum*," SEGi Review. 3: 34 44.
- Mungole A, Chaturvedi A (2011). Journal of Pharmaceutical Sciences Rewiev and Research. 6(1): 83 87.
- M Zimmermann (1983). "Ethical guidelines for investigations of experimental pain in conscious animals," *Pain*. 16(2): 109 110.
- N Abaclolu, B Tun Tan, E Akbulut and I Akici (2000). Participation of the components of L-arginine/nitric oxide/ cGMP cascade by chemically-induced abdominal constriction in the mouse. *Life Sciences*. 67(10): 1127 1137.
- Naczk M, Shahidi F (2004). Extraction and analysis of phenolics in food. *J. Chromatogr. A.* 1054: 95 111.
- Naczk M, Shahidi F (2006). Phenolics in cereals, fruits and vegetables: occurrence, extraction and analysis. *J. Pharm. Biomed. Anal.* 41, 1523 1542.
- Nemirovsky A, Chen L, Zelman V and Jurna I (2001). The antinociceptive effect of the combination of spinal morphine with systemic morphine or buprenorphine. *Anesthesia and Analgesia*. 93: 197 2003.
- Neugebauer V (2001a). Metabotropic glutamate receptors: novel targets for pain relief. *Expert Review of Neurotherapeutics*. 1: 207 224.

- Neugebauer V (2001b). Peripheral metabotropic glutamate receptors: fight the pain where it hurts. *Trends in Neurosciences*. 24: 550 552.
- Neugebauer V (2002). Metabotropic glutamate receptors—important modulators of nociception and pain behavior. *Pain.* 98(1-2): 1-8.
- Neuwinger H.D. (2000). Afrin traditional. MedPharm Scientific Publishers, Stuttgart, Germany.
- Newman H. J, Cragg G.M and Snader K.M (2000). The influence of natural products upon drug discovery, *Nat. Prod. Rep.* 17 (3): 215 234.
- Newman H. J, Cragg G.M and Snader K.M (2003). Natural products as source of new drugs over the period 1981-2002. *J. Nat. Prod. Rep.* 66 (7): 1022 1037.
- Nicholas D. Moore (2009). In serach of an ideal analgesic for common acute pain. *Acute Pain.* 11: 129 137.
- Niv D, Kreitler S (2001). Pain and quality of life. *Pain Pract*. 1: 150 61.
- Ni Q, Xu G, Lu G, Gao Q, Zhou C, Zhang Y (2012). African Journal of Biotechnology. 11 (14): 3379 3387.
- Negus SS, Vanderah TW, Brandt MR, Bilsky EJ, Becerra L, Borsook D (2006). Preclinical assessment of candidate analgesic drugs: recent advances and future challenges. *J Pharmacol Exp Ther.* 319: 507 514.
- Numazaki M, Tominaga M (2004). Nociception and TRP channels. Curr Drug Targets CNS Neurol Disord 3: 479 485.
- Nunes PH, Cavalcanti PM, Galvao SM, Martins MC (2009). Antiulcerogenic activity of *Combretum leprosum*. *Pharmazie*. 64: 58 62.
- Nur Izzati Ismail, Lee Ming-Tatt, Nordin Lajis, Muhammad Nadeem Akhtar, Ahmad Akira, Enoch Kumar Perimal, Daud Ahmad Israf and Mohd Roslan Sulaiman (2016). Antinociceptive Effect of 3-(2,3-Dimethoxyphenyl)-1-(5-methylfuran-2-yl)prop-2-en-1-one in Mice Models of Induced Nociception. Molecules, 21, 1077; doi:10.3390/molecules21081077.
- Nuresti S, Baek SH, Asari A (2003). Chemical components of *Melastoma malabathricum*. ACGC Chem Res Comm. 16:28–33.
- Notman R, Anwar J, O'Malley B (2006). Molecular Basis for Dimethyl sulfoxide (DMSO) Action on Lipid Membranes. *Journal of the American Chemical Society*. 128 (43): 13892 13893.
- Oboh U (2006). European Food Research Technology. 224: 61 65.

- Obouayeba Abba Pacome, DJYH Nazaire Bernard, DIABATE Sekou, DJAMAN Allico Joseph, N'GUESSAN Jean David, KONE Mongomake and KOUAKAU Tanoh Hilaire (2014). Phytochemical and Antioxidant Activity of Roselle (Hibiscus Sabdariffa L.) Petal Extracts. Research Journal of Pharmaceutical, Biological and Chemical Sciences. 5 (2): 1453 1465.
- Oh YS, Lee JH, Yoon SH, Oh CH, Choi DS, Choe E, Jung MY (2008). Characterization and quantification of anthocyanins in grape juices obtained from the grapes cultivated in Korea by HPLC/DAD, HPLC/MS, and HPLC/MS/MS. *J. Food Sci.* 73: C378 389.
- Oh YT, Lee JY, Lee JH, Kim JE, Ha J, Kang I (2010). Oleamide suppresses lipopolysaccharide-induced expression of iNOS and COX-2 through inhibition of NF-kappa B activation in BV2 murine microglial cells. *Neurosci Lett.* 474: 148 153.
- Ooi KK, Chear YL, Rodzi R, Othman F, Mohtarrudin N, Suhaili Z, Zakaria ZA (2014). Anti-carcinogenic activity of methanol extract of *Melastoma malabathricum* leaves in DMBA/croton oil-induced mouse skin carcinogenesis. *Africa J Complement Trad Alt Med*.11: 66–70.
- Ogundipe O T and Ayodele J (1999). *Bioscience Research Communication*. 12: 17 21.
- Okokon JE, Udoh AE, Frank SG, Amazu LU (2012). Anti-inflammatory and analgesic activities of *Melanthera scandens*. Asian Pac J Trop Biomed. 2: 144 148.
- OP Tandon, V Malhotra, S. Tandon, I D'Silva (2003). NEUROPHYSIOLOGY OF PAIN: INSIGHT TO OROFACIAL PAIN. *Indian J Physiol Pharmacol*. 47 (3): 247 269.
- Otuki MF, Lima FV, Malheiros A, Cechinel-Filho V, Monache FD, Yunes RA, Calixto JB (2001) Evaluation of the antinociceptive action caused by ether fraction and a triterpene isolated from resin of *Protium kleinii*. *Life Sci*. 69 (19): 2225 2236.
- Otuki MF, Ferreira J, Lima FV, Meyre-Silva C, Malheiros A, Muller LA, Cani GS, Santos ARS, Yunes RA, Calixto JB (2005). Antinociceptive properties of mixture of α-amyrin and β-amyrin triterpenes: evidence for participation of protein kinase C and protein kinase a pathways. *J Pharmacol Exp Ther*. 313:310–8
- Ossipov MH, Lai J, King T, Vanderah TW, Malan TP Jr, Hruby VJ, Porreca F (2004). Antinociceptive and nociceptive actions of opioids. *J Neurobiol*. 1: 126 48.

- Owen RW, Haubner R, Mier W, Giacosa A, Hull WE, Spiegelhalder B, Bartsch H (2003). Isolation, structure elucidation and antioxidant potential of the major phenolic and flavonoid compounds in brined olive drupes. *Food Chem Toxicol*. 41: 703 17.
- Padua de L.S, Bunyapraphatsara N and Lemmens R.H.M.J. (1999). Plant resources of South-East Asia, No 12(1). Medicinal and Poisonous Plant 1. Backhuys Publishers, Leiden, The Netherlands.
- Park JH, Son KH, Kim SW, Chang HW, Bae K Kang SS, Kim HP (2004).
- Parida N.K, Sahu M.R, Debata P.C. and Panda P.K. (2010). Antinociceptive and Anti-inflammatory effects of methanolic extract of *Benincasa hispida* Fruit Peel in rodents. *Asian journal of chemistry*. 22(10):7573 7579.
- Parada CA, Tambeli CH, Cunha FQ, Ferreira SH (2001). The major role of peripheral release of histamine and 5-hydroxytryptamine in formalin-induced nociception. *Neurosci.* 102: 937 44.
- Pasero C (2004). Pathophysiology of neuropathic pain. *Pain Manag. Nurs.* 5:3 8.
- Pasero C, Paice JA and McCaffery M (1999a). Basic mechanisms underlying the causes and effects of pain. In: McCaffery M, Pasero C, eds. *Pain Clinical Manual*. 2nd ed. St. Louis MO: Mosby Inc. 15 34.
- Pasero C, Portenoy RK and McCaffery M (1999b). Opioid analgesic. In: McCaffery M, Pasero C, eds. Pain Clinical Manual 2nd ed. St. Louis, MO: Mosby Inc; 15 34.
- Patricia HB, Covington EC, Dahl JL, Katz JK and Miaskowski C (2001). Background and Significant. Pain: Current Understanding of Assessment, Management, and Treatments. *American Pain Society*. 2 18.
- Pasternak GW (1993). Review: Pharmacological mechanisms of opioid analgesics. Clin Neuropharmacol. 16(1):1 – 18
- Patwarhan B, Warude D, Pushpangadan P and Bhatt N (2005). Ayurveda and traditional Chinese medicine: A comparative overview. e.C.A.M.2 (4): 465 473.
- Pèrez MB, Calderon NL and Croci CA (2007). Radiation-induced enhancement of antioxidant oxidation activity in extracts of rosemary *Rosmarinus officinalis* L. *Food Chemistry*. 104: 585 592.
- Peter Winstanley and Tom Walley (1996). Analgesic: Pharmacology. *Churchill's Mastery of Medicine*. 74 78.

- Portenoy Russell K, Brennan Michael J (1994). "Chronic Pain Management". In Good, David C, Couch James R. Handbook of Neurorehabilitation. *Informa Healthcare*. ISBN 0-8247-8822-2.
- Portenoy RK and Kanner RM (1996). Definition and Assessment of Pain. In Portenoy RK, Kanner RM, eds. Pain Management: Theory and Practice. Philadelphia: FD Davis. 3 –18.
- Prior RL, Lazarus SA, Cao G, Muccitelli H, Hammerstone JF (2001). Identification of procyanidins and anthocyanins in blueberries and cranberries (Vaccinium spp.) using highperformance liquid chromatography/mass spectrometry. *J. Agric. Food Chem.* 49: 1270 1276.
- Pieroni A. (2000). Medicinal plants and food medicines in the folk traditions of the Lucca Province, Italy. *J. Ethnopharmacol*. 70: 235 273.
- Purushoth Prabhu T, Panneerselvam P, Suresh R, Clement Atlee W, Balasubramanian S (2013). GC-MS analysis of ethanolic extract of *Canthium parviflorum* Lamk Leaf. *Journal of Applied Pharmaceutical Science*. DOI: 10.7324/JAPS.2013.30229. Vol. 3(2): 166 168.
- Ramaswamy S, Langford RM (2016). Antinociceptive and immunosuppressive effect of 3 opioids in an acute postoperative setting: a review. *BJA Education*. doi: 10.1093/030.
- Ram RL (2001). Advances in Plant Sciences. 14: 525 530.
- Ramzi AA Mothana (2011). Anti-inflammatory, antinociceptive and antioxidant activities of the endemic Soqotraen *Boswellia elongate* Balf. f. and *Jatropha unicostata* Balf. f. in different experimental models. *Food and Clinical Toxicology*. 49: 2594 2599.
- Rauf A, Uddin G, Siddiqui BS, Khan H, Shah SU, Hadda TB, Mabkhot YN, Farooq U, Khan A (2016). Antinociceptive and anti-inflammatory activities of flavonoids isolated from *Pistacia integerrima* galls. *Complement Ther Med*. 25: 132 8.
- Reginaldo Vicente Ribeiro, Regilane Matos da Silva, Joaquim Corsino da Silva Lima, Domingos Tabajara de Oliveira Martins (2010). Antiinflammatory, antinociceptive and antipyretic effects of hydroethanolic extract from *Macrosiphonia velame* (A.St.-Hil) M. Arg. in animal models. *Brazilian Journal of Pharmaceutical Sci.* 46: 1 9.
- Reichert J.M. (2003). Trends in development and approval times for new therapeutics in the United States. *Nat. Rev. Drug. Discovery* 2 (9): 695 702.
- Reisine T and Pasternak G (1996). Opioid analgesic and antagonist. In: Hardman JG, Li,bard LE, eds. Goodman & Gillman's: Pharmacologic Basic of Therapeutics 9th ed. New York: MacGraw-Hill. 531 555.

- Ribeiro RA, Vale ML, Thomazzi SM, Paschoalato AB, Poole S, Ferreira SH et al. (2000). Involvement of resident macrophages and mast cells in the writhing nociceptive response induced by zymosan and acetic acid in mice. *Eur J Pharmacol.* 387: 111–8.
- Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radic. Res.* 22: 375 383.
- Rice-Evans CA, Miller NJ, Paganga G (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.* 20, 933 956.
- Rivot JP, Montagne-Clavel J, Besson JM (2002). Subcutaneous formalin and intraplantar carrageenan increase nitric oxide release as measured by in vivo voltammetry in the spinal cord. *Eur J Pain*. 6: 25 34.
- R Negrete, A Hervera, S Leánez, JM Martín-Campos, O Pol (2011). The antinociceptive effects of JWH-015 in chronic inflammatory pain are produced by nitric oxide cGMP-PKG-KATP pathway activation mediated by opioids. *PLoS One*, 6: e26688.
- RNS Yadav and Munin Agarwala (2011). Phytochemical analysis of some medicinal plants. *Journal of Phytology*. ISSN: 2075-62403(12): 10 14.
- Rustoen T, Stubhaug A, Eidsmo I, Westheim A, Paul SM, Miaskowski C. (2008). Pain and quality of life in hospitalized patients with heart failure. *J Pain Symptom Manage*. 36: 497–504.
- Robards K (2003). Strategies for the determination of bioactive phenols in plants, fruit and vegetables. *J. Chromatogr. A.* 1000: 657 691.
- Robinson M. M. & Zhang X. (2011), *The World Medicine situation 2011: Traditional medicine:Global situation, issues and challenges*, WHO Press, Geneva, Retrived from http://www.who.int/medicines/areas/policy/world_medicines_situation/WMS_ch18_wTraditionalMed.pdf.
- Robbins RJ (2003). Phenolic acids in foods: an overview of analytical methodology. *J. Agric. Food Chem.* 51: 2866 2887.
- R Seabra, P Valentao, F Ferresres and P Andrade (2002). Phenolic profile in the definitions of natural products authenticity. *Natural Products in the New Millennium: Prospects and Industrial Application Proceedings of the Phytochemical Society of Europe.* 47: 183 194.
- Sabina E, Chandel S, Rasool M.K (2009). Evaluation of analgesic, antipyretic and ulcerogenic effect of Withaferin A. *Int. J. Integ. Biol.* 6(2): 52 56.

- Sakakibara H, Honda Y, Nakagawa S, Ashida H, Kanazawa K (2003). Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *J. Agric. Food Chem.* 51: 571 581.
- Sakurada, T Matsumara, T Moriyama, T Sakurada, C Ueno S and Sakurada S. Differential effects of intraplantar capsazepine and ruthenium red on capsaicin-induced desensitization in mice. *Pharmacol Biochem Behav.* 75: 115 121.
- Samuelsson G (2004). Drugs of natural origin: A text book pharmacology 5th Swedish Pharmaceutical Press, Stockholm.
- Sanda P Welch and Billy R Martin (1997). Chapter 26: Opioid & Non-opioid Analgesic, Modern Pharmacology with Clinical Applications by Craig & Charles R, 6th Edition, ISBN 13: 9780316159340.
- Sawynok J (2005). Topical analgesics in neuropathic pain. *Curr. Pharm. Des.*, 11, 2995 3004.
- Saravanan Ramalingan, Arthi Mohan, Balakrishnan Sethuramali, Saravanan Ramalingam (2015). GC-MS Analysis of Methanolic Extract of *Phyllanthus Amarus* leaves Collected from Salem Region. *Asian Journal of Pharmacology and Toxicology*. 03 (10): 54 59.
- Siddall PJ and Cousins MJ (1997). Neurobiology of pain. *Int Anesthesiol Clin*. 35: 1 26.
- Siddiqui S, Verma A, Rather AA, Jabeen F & Meghvansi MK (2009). Preliminary Phytochemicals Analysis of Some Important Medicinal and Aromatic Plants. *Advances in Biological Research*. 3: 188 195.
- Schmidtko A, Tegeder I, Geisslinger G (2009). A Review: No NO, no pain? The role of nitric oxide and cGMP in spinal pain processing. *Trends Neurosci*. 32(6): 339 46.
- Singleton VL, Orthofer R, Lamuela-Raventos RM (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.* 299: 152 178.
- Singleton VL, Rossi JA Jr (1965). Colorimetry of total phenolics with phosphomolybdicphosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16: 144 158.
- Shanmugasundaram P and Venkataraman S (2005). Antinociceptive activity of *Hygrophila auriculata* (Schum) Heine. *Afr J Trad* CAM. 2(1): 62 69.
- Sharma A, VK Patel, S Rawat, P Ramteke and R Verma (2010). Identification of the antibacterial component of some Indian medicinal plants against *Klebsiella pneumonia*. *Int. J. Pharmacy Pharmaceutical Sci.*, 2: 123 127.

- Shibata M, Ohkubo T, Takahashi H, Inoki R. (1989). Modified formalin test; characteristic biphasic pain response. *Pain*. 38: 347 352.
- S Bharadwaj and SK Gakhar (2005). Ethnomedicinal plants used by tribals of Mizorum to use cuts and wounds. *Indian Journal of Traditional Knowledge*. 4: 75 80.
- SK Raghav, B Gupta, C Agrawal, K Goswami and HR Das (2006). "Antiinflammatory effect of *Ruta graveolens* L. Inmurine macrophage cells," *Journal of Ethnopharmacology*. 104(1-2): 234-239.
- SM Carlton, GL Hargett, RE Coggeshall (1995). Localization and activation of glutamate receptors in unmyelinated axons of rats glabrous skin. *Neurosci*. *Lett*. 197: 25 28.
- S Mohd Joffry, NJ Yob, MS Rofiee, MMR Meor Mohd Affandi, Z Suhaili, E Othman, A Md Akim, MNM Desa and ZA Zakaria (2012). *Melatoma malabathricum* (L.) Smith Ethnomedicinal Uses, Chemical Constituents, and Pharmacological Properties: A Review. *Evidence-Based Complimetary and Alternative Medicine*, doi:10.1155/2012/258434: 1 48.
- Smith TW, Buchan P, Parsons DN, Wilkinson S (1982). Peripheral antinociceptive effects of N-methyl morphine. *Life Sci.* 31:1205 1208.
- Soares de Moura R, Rios AAS, Santos EJA, Amorim Nascimento AB, de Castro Resende ff, Lemos Neto M *et al.* (2004). Role of the NOcGMP pathway in the systemic antinociceptive effect of clonidine in rats and mice. *Pharmacol Biochem Behav.* 78: 247 53.
- Sowndhararajan K and Kang SC (2013). Protective effect of ethyl acetate fraction of *Acacia ferruginea* DC. Against ethanol-induced gastric ulcer in rats. *Journal of Ethnopharmacology*. 148: 175 181.
- Staniszewka I, Krolicka A, Malinski E, Lojkowska E and Szafranek J (2003). Elicitation of secondary metabolites in *in vitro* cultures of *Ammi majus* L. Enzymes Microb. Technol. 33: 565 568.
- Suh H, Song D, Huh S, Son K, Kim Y (2000). Antinociceptive mechanisms of dipsacus saponin C administered intrathecally in mice. *J Ethnopharmacol*. 1-2: 211-8.
- Suganuma T, Suzuki T, Oshimi M, Hanano M. (1998). Change of beta-endorphin concentration in rat brain after administration of indomethacin or carrageenin. *Biol Pharm Bull.* 21: 756 760.
- Susanti D. and Rasadah M.A. (2007). Anti-inflammatory action of components from *Melastoma malabathricum. Pharmaceutical biology*. 45(5): 372 375.

- Sulaiman MR, MN Somchit, DA Israf, Z Ahmad and S Moin (2004). Antinociceptive effect of *Melastoma malabathricum* ethanolic extract in mice. Fitoterapia, 75: 667 672.
- Sulaiman MR, Perimal EK, Zakaria ZA, Mokhtar F, Akhtar MN, Lajis NH and Israf DA (2009). Preliminary analysis of the antinociceptive activity of zerumbone. *Fitoterapia*.
- Sousa OV, Del-Vechio-Vieira G, Kaplan MAC (2007). Propriedades analgésica e antiinflamatória do extrato metanólico de folhas de *Annona coriacea* Mart. (Annonaceae). *Lat. Am. J. Pharm.* 26: 872 877.
- Sousa OV, Del-Vechio-Vieira G, Amaral MPH, Pinho JJRG, Yamamoto CH, Alves MS (2008). Efeitos antinociceptivo e antiinflamatório do extrato etanólico das folhas de *Duguetia lanceolata* St. Hil. (Annonaceae). *Lat. Am. J. Pharm.* 27: 398 402.
- South African Medicines Formulary (SAMF). (2010). Health and Medical Publishing Group of the South African Medical Association, Cape Town, South Africa, 9th edition.
- SS Koay (2008). Establishment of cell suspension culture of *Melastoma* malabathricum L. for the production of anthocyanin. *PhD Thesis*, *Universiti* Sains Malaysia, *Pulau Pinang*, Malaysia.
- Stalikas CD (2007). Extraction, separation, and detection methods for phenolic acids and flavonoids. *J. Sep. Sci.* 30: 3268 3295.
- Stanley W Jacob and Robert Herschler (1986). Pharmacology of DMSO. *Cyrobiology*. Vol. 23(1): 14 27.
- ST Meller, GF Gebhart (1993). Nitric oxide (NO) and nociceptive processing in the spinal cord. *Pain*. 52: 127 136.
- ST Meller, C Dykstra, GF Gebhart (1996). Acute thermal hyperalgesia in the rats in produced by activation of *N*-methyl-D-aspartate receptors and protein kinase C and production of nitric oxide. *Neuroscience*. 71: 327 335.
- S Talarek and S Fidecka (2002). Role of nitric oxide in benzodiazepines-induced antinociception in mice," *Polish Journal of Pharmacology*, vol. 54, no. 1, pp. 27 34.
- Stoclet J, Muller B, Gyorgy K, Andriantsiothaina R and Kleschyov A (1999). The Inducible Nitric Oxide Synthase In Vascular and Cardiac Tissue. *European Journal of Pharmacology* 375: 139 155.
- Svensson CI, Yaksh TL (2002). The spinal phospholipase-cyclooxygenase-prostanoid cascade in nociceptive processing. *Annu Rev Pharmacol Toxicol.*, 42: 553 83.

- Świeboda P, Filip R, Prystupa A, Drozd M (2013). Assessment of pain: types, mechanism and treatment. Ann Agric Environ Med. 2013; Special Issue 1: 2–7.
- S Zhou, L. Bonasera, S.M. Carlton (1996). Peripheral administration of NMDA, AMPA or KA results in pain behaviours in rats. *Neuro report*. 7: 895 900.
- Szolcsányi J (1993). Actions of capsaicin on sensory receptors. In *Capsaicin in the study of pain*, ed Wood J; Academic, London. 1 26.
- Taiz L & Zeiger E (2006). Plant physiology, 4th edn.. Sinauer Associates, Inc. Publishers, Massachusetts. 13: 315 344.
- Tang L, Chen Y, Chen Z, Blumberg PM, Kozikowski AP, Wang ZJ (2007). Antinociceptive pharmacology of N-(4-chlorobenzyl)-N'-(4-hydroxy-3-iodo-5-methoxybenzyl) thiourea, a high-affinity competitive antagonist of the transient receptor potential vanilloid 1 receptor. *J Pharm Exp Ther*. 2: 791 798.
- Tapas AR, Sakarkar DM and Kakde RB (2008). Flavonoids as nutraceuticals: a review. *Tropical Journal of Pharmaceutical Research*. 7: 1089 1099.
- Tesfaye S (2009). Advances in the management of diabetic peripheral neuropathy. *Curr. Opin. Support. Palliat. Care.* 3: 136 143.
- Tiwari P, Kumar B, Kaur M, Kaur G & Kaur H. (2011). Phytochemical screening and Extraction: A Review. *Internationale Pharmaceutica Sciencia*. 1(1): 98 106.
- T L Yaksh (1997). Acta Anaesthesiol. Scand. 41, 94.
- TM Cunha, D Roman-Campos, CM Lotufo *et al.*, (2010). Morphine peripheral analgesia depends on activation of the PI3Kγ/AKT/nNOS/NO/KATP signaling pathway. *Proceedings of the National Academy of Sciences of the United States of America*. 107(9): 4442 4447.
- Toda N, Kishioka S, Hatano Y and Toda H (2009). Modulation of opioid actions by nitric oxide signaling. *Anesthesiol*. 1: 166 181.
- TRL Romero, IDG Duarte (2009). a2 Adrenoceptor agonist xylazine induces peripheral antinociceptive effect by activation of the L-arginine/nitric oxide/cyclic GMP pathway in rat. *Eur. J. Pharmacol.* 613: 64 67.
- Troullos E, Hargreaves KM, Dionne RA (1997). Ibuprofen elevates immunoreactive beta-endorphin levels in humans during surgical stress. *Clin Pharmacol Ther*. 62: 74 81.
- Tsimogiannis D, Samiotaki M, Panayotou G, Oreopoulou V (2007). Characterization of flavonoid subgroups and hydroxyl substitution by HPLC-MS/MS. *Molecules*. 12: 593 606.

- TW Hugh Tan and CK Yeo (2009). The Potential of Native Woody Plants for Enhancing the Urban Waterways and Waterbodies Environment in Singapore. Raffles Museum of Biodiversity Research and Singapore-Delft Water Alliance, National University Singapore. P 14.
- Usoh IF, Akpan EJ, Etim EO, Farombi EO (2005). *Pakistan Journal of Nutrition*. 4 (3): 135 141.
- Van Agtmael MA, Eggelte TA and van Boxtel CJ (1999). Artemisinin drugs in the treatment of malaria: from medicinal herb to registered medication. *Trends Pharmacol. Sci.* 20 (5): 199 205.
- Vane JR, Botting RM (1990). The mode of action of anti-inflammatory drugs. *J. Postgrad. Med.* 66: S2 S17.
- Vaz ZR, Cechinel V, Yunes RA, Calixto JB (1996). Antinociceptive action of 2-(4-bromobenzoyl)-3-methyl-4-6-dimethoxy bezofuran, a novel xanthoxyline derivative of chemical and thermal models of nociception in mice. *J Pharm Exp Ther*. 278: 304 312.
- V Neugebauer (2002). "Metabotropic glutamate receptors-important modulators of nociception and pain behavior," *Pain.* 98(1-2): 1 8.
- Verma PR, Joharapurkar AA, Chatpalliwar VA, Asnani A (2005). Antinociceptive activity of alcoholic extract of *Hemidesmus indicus* Rbr. in mice. *J Ethnopharmacol.* 102: 298 301.
- Verpoorte R (1998). Exploration of nature's chemodiversity: the role of secondary metabolites as leads in drug development. *Drug Discov. Today.* 3: 232 238.
- Vicker A and Zollman C (1999). ABC of complementary medicine: herbal medicine. *B.M.J.* 319 (7222): 1422.
- Vivancos GG, Parada CA, Ferreira SH (2003). Opposite nociceptive effects of the Larginine/NO/cGMP pathway stimulation in dermal and subcutaneous tissues. *Brit J Pharmacol.* 138: 1351 1357.
- Vongtau HO, Abbah J, Ngazal IE, Kunle OF, Chindo BA, Otsapa PB and Gamaniel KS (2004). Antinociceptive and anti-inflammatory activities of the methanolic extract of *Parinari polyandra* stem bark in rats and mice. *J. Ethnopharmacol*. 90(1): 115 121.
- Vyklicky L (1979). Techniques for the study of pain in animals. In Bonica, J.J., Liebeskind, J.C, Albe-Fessard, D.G (eds.). Advances in Pain Research and Therapy. Raven Press, New York.
- Walsh TD (1990). Prevention of opioid side effects. *J Pain Symptom Manage*. 5: 362 7.

- Weiher, E, van, der, Werf., A. Thompson., K. Roderick., M. Garnier, E., and Ericksson, O. (1999). Challenging Theophrastus: A common core list of plant traits for functional ecology. *J. Veg. Sci.* 10 (5): 609-620.
- Weisburger, J.H. (2002). Lifestyle, health and disease prevention: the underlying mechanisms. *Eur. J. Cancer Prev.* 11: S1 7.
- Willow, J.H. Liu. (2011). Traditional Herbal Medicines, Ch 1: 1-25.
- Williams, M., Kowaluk, E.A., Arneric, S.P. (1999). Emerging molecular approaches to pain therapy. *J Med Chem.* 42: 1481 500.
- Wink, M. (1999). Biochemistry of plant secondary metabolism. *Annual plant reviews*. Sheffield Academic Press. 2(1): 1 6.
- World Health Organization (2002). Traditional medicine strategy launched. *Buletin of the World Health Organization*. 80:7
- Y.F. Chen., H.Y. Tsai., and T.S. Wu. (1995). Anti-inflammatory and analgesic activities from roots of *Angelica pubescens*. *Planta Medica*. 61: 2 8.
- Yakash, T.L. (1999). Central pharmacology of nociceptive transmission. In: Wall PD, Melzack R. eds, Textbook of Pain, 4th edition. Churchill Levingston, Endiburgh. 253 308.
- Yang, S.Z. (2005). The divine famer's materia; medica: a translation of the shen nong ben cao jing (1st edition)., Blue Poppy Press, ISBN 0585105464, Boulder, Colorado.
- Yang, M., Sun, J., Lu, Z., Chen, G., Guan, S., Liu, X., Jiang, B., Ye, M., and Guo, D.A. (2009). Phytochemical analysis of traditional Chinese medicine using liquid chromatography coupled with mass spectrometry. *Journal of Chromatography A.* ISSN 0021-9673. Vol. 1216 (11): 2045 2062.
- Yanagida, A., Shoji, T., Kanda, T. (2002). Characterization of polymerized polyphenols by sizeexclusion HPLC. *Biosci. Biotechnol. Biochem.* 66: 1972 1975.
- Zabidi, Z., Wan, Zainulddin, W.N., Mamat, S.S., Shamsahal, Din, S., Kamisan, F.H., Yahya, F., Ismail, N.A., Rodzi, R., Hussain, H., Mohtarrudin, N., Somchit, M.N., Zakaria, Z.A. (2012). Antiulcer activity of methanol extract of *Melastoma malabathricum* leaves in rats. *Med Princ Pract*. 21:501–503.
- Zainul, Amiruddin, Zakaria., Mohd, Roslan, Sulaiman., Muhammad, Nazrul., Somchit., Abdul, Manan, Mat, Jais., Daud, Israf, Ali. (2005). The effects of larginine, d-arginine, l-name and methylene blue on *channa striatus*-induced peripheral antinociception in mice. *J. Pharm Pharmaceutical Sci.* 8(2): 199 206.

- Z.A, Zakaria., R.N.S, Raden, Mohd, Nor., Z.D.F, Abdul, Ghani., G, Hanan, Kumar., M.R, Sulaiman., and C.A. Fatimah. (2006). Antinociceptive and Anti-inflammatory Properties of *Melastoma malabathricum* Leaves Chloroform Extract in Experimental Animals. *Journal of Pharmacology and Toxicology*. 1(4): 337 345.
- Zakaria, Z.A., Raden, Mohd, R.N., Nor, S., Hanan, G., Kumar, Z., Abdul, Ghani, D.F. *et al.*, (2008). Antinociceptive, anti-inflammatory and anti-pyretic properties of *Melastoma malabathricum* leaves aqueous extract in experimental animals. *Canadian Journal of Physiology and Pharmacology*. 84(12): 1291 1299.
- Zakaria, Z.A., Nelendran, M., Pubalan, S., Sulaiman, M.R., and Abdullah, F.C. (2006). Effect of opoid and non-opoid antagonists, pH and enzyme on *Corchorus olitorius* antinociceptive in mice. *Oriental Pharmacy and Experimental Medicine*. 6 (3): 186 195.
- Zakaria, Z.A., Abdul, Rahman., N.I. Loo. Y.W., Abdul, Ayub, A.H., Sulaiman, M.R., Mat, Jais, A.M., Gopalan, H.K., and Fatimah, C.A. (2009). Antinociceptive and anti-inflammatory activities of the chloroform extract of *Bauhinia purpurea* L. (Leguminosae) leaves in animal models. *International Journal of Tropical Medicine*. 4(4): 140 145.
- Zakaria, Z.A., Somchit, M.N., Mat, Jais, A.M., Teh, L.K., Salleh, M.Z., and Long, K. (2011). *in vivo* antinociceptive and anti-inflammatory activities of dried and fermented processed virgin coconut oil. *Med. Princ. Pract.* 20 (3): 231 236.
- Zakaria, Z.A., Rofiee, M.S., Mohamed, A.M., Teh, L.K., Salleh, M.Z. (2011). In vitro antiproliferative and antioxidant activities and total phenolic contents of the extracts of *Melastoma malabathricum* leaves. *J Acupunct Meridian Stud*. 4:248–256.
- Zakaria, Z.A., Balan, T., Mamat, S.S., Mohtarrudin, N., Teh, L.K., Salleh, M.Z. (2015). Mechanisms of gastroprotection of methanol extract of *Melastoma malabathricum* leaves. *BMC Complement Alt Med* 2015, 15: 135.
- Zhou, S., Koh, H.L., Gao, Y., Gong, Z., Lee, E.J.D. (2004). Herbal bioactivation: The good, the bad and the ugly. *Life sci*. 74: 935 968.

BIODATA OF STUDENT

Erman Shah Bin Jaios was born in Muar, Johor on September 14th, 1979. He received his early primary education at Sekolah Rendah Kebangsaan Ismail 1 & 2, Muar, Johor (1986 – 1991). Then he continued his secondary education at Sekolah Menengah Dato' Sri Amar Diraja, Muar, Johor (1992 – 1994) from Form 1 – 3, and Sekolah Menengah Teknik Melaka, Bukit Piatu, Melaka (1995 – 1996) from Form 4 – 5. He then enrolled into UPM Matriculation, PTPL, Ampang, Selangor (May 1997 – 1998) as preparation to enter UPM.

In 1998, He was accepted to pursue his first Bachelor Degree in Biomedical Sciences at Universiti Putra Malaysia (1998 – 2002), one of the Research Universities. During four years of study period, his passion, curiosity and interest towards has been developed. After graduation, he was working at Laboratory Scientific & Medical Supplier as Sales Executive (2002 – 2003) to enhanced the knowledge on the type, technical & application on laboratory equipment & instrumentation as well as apparatus & chemicals while main responsibilities on sales and technical scopes.

In 2003, He was hired as Science Officer serving in the Department of Basic Medical Sciences, Kulliyyah of Pharmacy, IIUM, Kuantan (April 2003 – Dec 2011). He was responsible on the purchasing, arrangement, coordination on the laboratory & teaching materials for undergraduate laboratory practical, to ensure good safety and health practice is implemented by laboratory staff, student and other parties, and safe environment. In addition, he used to work on the laboratory animal management, which involved in the animal laboratory practical such as pharmacology & physiology, at the same time assist the researchers either postgraduate or lecturer on their research activities.

Based on the expertise and experiences on the laboratory management, he decided and was accepted to further his studies in Master of Science (Pharmacology & Toxicology), admission in second semester 2011/2012, at Faculty of Medicine and Health Sciences, Universiti Putra Malaysia.

His thesis entitled Antinociceptive Activities of *Melastoma malabathricum* L. (Senduduk) Leaves Methanolic Extract and Its Petroleum Ether Fractions.

LIST OF PUBLICATIONS

- E.S. Jaios., Abdul, Rahman, S., Ching, S.M., Arifah, A.K., Desa, M.N., Zakaria, Z.A. (2016). Possible mechanisms of antinociception of methanol extract of *Melastoma malabathricum* Leaves. *Braz J Pharmacognocy*. doi.org/10.1016/j.bjp.2016.01.011.
- Z.A. Zakaria., E.S. Jaios., M.H. Omar., S. Abd. Rahman., S.S.A. Hamid., S.M. Ching., L.K. Teh., M.Z. Salleh., S. Deny., and M. Taher. (2016). Antinociception of petroleum ether fraction derived from crude methanol extract of *Melastoma malabathricum* leaves and its possible mechanisms of action in animal models. *BMC Complementary and Alternative Medicine*. doi. 10.1186/s12906-016-1478-1.

Presentations

Poster Presentation; 1st Allied Health Scientific Colloquium (AHSC) in conjunction with 2nd KAHS Research Week (KRW) at Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, Kuantan, Pahang., dated: 21st – 25th November 2016.



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION: <u>SECOND SEMESTER 2016/2017</u>

TITLE OF THESIS / PROJECT REPORT:

ANTINOCICEPTIVE ACTIVITIES OF SENDUDUK (Melastoma malabathricum L.)
LEAVES METHANOLIC EXTRACT AND ITS PETROLEUM ETHER FRACTIONS

NAME OF STUDENT: ERMAN SHAH JAIOS

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

- 1. This thesis/project report is the property of Universiti Putra Malaysia.
- 2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
- 3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as:

*Please tick (V)	
CONFIDENTIAL	(Contain confidential information under Official Secret Act 1972).
RESTRICTED	(Contains restricted information as specified by the organization/institution where research was done).
OPEN ACCESS	I agree that my thesis/project report to be published as hard copy or online open access.
This thesis is submitted for :	
PATENT	Embargo from until (date) (date)
	Approved by:
(Signature of Student) New IC No/ Passport No.:	(Signature of Chairman of Supervisory Committee) Name:
Date :	Date :

[Note: If the thesis is CONFIDENTIAL or RESTRICTED, please attach with the letter from the organization/institution with period and reasons for confidentially or restricted.]