



UNIVERSITI PUTRA MALAYSIA

***ANTHROPOGENIC ACTIVITIES IMPACT ON WATER QUALITY, FISH
COMMUNITY AND BACTERIA PRESENCE IN DISTURBED AND
UNDISTURBED NORTH SELANGOR PEAT SWAMP FOREST,
MALAYSIA***

SULE HUSSEIN ALIU

FS 2016 78



**ANTHROPOGENIC ACTIVITIES IMPACT ON WATER QUALITY, FISH
COMMUNITY AND BACTERIA PRESENCE IN DISTURBED AND
UNDISTURBED NORTH SELANGOR PEAT SWAMP FOREST, MALAYSIA**

By

SULE HUSSEIN ALIU

**Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in fulfilment of the requirement for the degree of Master of Science**

December 2016

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



*To my loving parents, for their immense support
To my beloved wife Rabiati Aliu, for her love, patience and wholehearted support
&
To my beautiful daughters Aqeelah Aliu & Hafeezah Aliu, for making everything
worthwhile*



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**ANTHROPOGENIC ACTIVITIES IMPACT ON WATER QUALITY, FISH
COMMUNITY AND BACTERIA PRESENCE IN DISTURBED AND
UNDISTURBED NORTH SELANGOR PEAT SWAMP FOREST**

By

SULE HUSSEIN ALIU

December 2016

Chairman: Ahmad Ismail, PhD
Faculty: Science

This study is concerned with the impact of anthropogenic activities on the water quality, the fish community and the bacteria presence in disturbed and undisturbed north Selangor peat swamp forest (NSPSF). Field sampling and analyses were conducted during the months of June and October 2015, and January 2016. Environmental conditions, and water physicochemical data were measured in peat swamp, paddy field and oil palm plantation. Fish was collected and bacterial isolation and identification from fish, water and sediment using standard microbiological techniques. Statistical analyses employed include ANOVA and Turkey LSD, Kruskal-Wallis and Mann-Whitney tests, Principal Component Analysis (PCA) and Canonical Correspondence Analysis (CCA). In addition, fish community structure was calculated using Shannon-Weaver diversity, Pielou's evenness and Margalef's richness indexes.

Peat swamp recorded a significantly ($p < 0.05$) higher phosphate ($1.937 \pm 0.196 \text{ mg L}^{-1}$), sulphate ($28.917 \pm 20.398 \text{ mg L}^{-1}$) and chlorine ($0.915 \pm 0.361 \text{ mg L}^{-1}$), while low for pH (3.684 ± 0.251), dissolved oxygen ($0.590 \pm 0.169 \text{ mg L}^{-1}$) and ammonia-nitrogen ($0.346 \pm 0.081 \text{ mg L}^{-1}$) when compared to paddy field and oil palm plantation areas. However, there was no significant ($p > 0.05$) difference observed for water electrical conductivity, total dissolved solids, salinity and nitrite for at least two of the sampling habitats, and for all of the sampling habitats for water temperature. The phosphate and chlorine levels of all sites, the pH of peat swamp (3.684 ± 0.251) and paddy field (4.962 ± 0.462), and the dissolved oxygen level in peat swamps were above the standard level for support of aquatic life and fall into class V of the National Water Quality Standard (NWQS) of Malaysia.

Paddy field recorded a significantly ($p < 0.05$) lower score of accessibility (1.50 ± 0.548) and deforestation level (1.00 ± 0.000), while oil palm recorded the lowest score of pollution level (2.00 ± 0.000). There was no significant ($p > 0.05$) difference on level of human development, water depth and pollution in at least two of the sampling habitats, and for all of the sampling habitats in term of the distance from human settlement. With the exception of water depth, all of the measured environmental conditions showed their importance in determining the water quality of the NSPSF.

Statistical analyses revealed that pollution level is the most important environmental factor determining water quality in the NSPSF.

A total of 1382 individual fish belonging to 20 species were collected during this study. Of this, only 15 species were collected from peat swamp, compared to 19 and 20 species from paddy field and oil palm plantation respectively. In the same vein, peat swamp recorded the lowest species diversity (2.4243) and richness (2.4768), compared to paddy field and oil palm plantation. Family Cyprinidae has the highest number of representative species. A notable concern however is the absence of several endangered species such as *Betta livida* and *Parosphromenus harveyi*, which were previously recorded from the NSPSF, highlighting the need for a reassessment of their conservation status. Multivariate statistics reveal that the water physicochemical parameters such as water temperature, conductivity, ammonia, sulphate, chlorine and dissolved oxygen influence fish community.

A total of 3,421 isolates from 39 species of bacteria were obtained during this study. The most dominant bacterial species were *Escherichia coli*, *Salmonella* spp., and *Streptococcus agalactiae*. Paddy field and oil palm plantation had the highest number of isolates and species, due to higher anthropogenic inputs in these areas. Among the species isolated were several pathogens such as *Aeromonas hydrophila*, *Edwardsiella tarda*, *Staphylococcus* spp., and *S. agalactiae*. Water physicochemical parameters such as water temperature, conductivity, ammonia, sulphate, chlorine and dissolved oxygen were important in influencing bacterial composition.

This study identifies peat land conversion to agriculture, anthropogenic inputs such as domestic and industrial effluents discharge as the greatest threats to the resident biological communities in the NSPSF. Among others, the importance of inclusion of water quality improvement strategies in peatland conservation was highlighted.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PERKAITAN ANTARA AKTIVITI ANTROPOGENIK, KUALITI AIR,
STRUKTUR KOMUNITI IKAN DAN KEHADIRAN BAKTERIA DI HUTAN
PAYA GAMBUT UTARA SELANGOR.**

Oleh

SULE HUSSEIN ALIU

Disember 2016

Pengerusi: Ahmad Ismail, PhD
Fakulti: Sains

Kajian ini menitik beratkan perkaitan antara aktiviti antropogenik, kualiti air, dan struktur komuniti ikan serta kehadiran bakteria di kawasan hutan paya gambut Selangor utara (NSPSF). Persampelan di lapangan dan analisis telah dilaksanakan pada bulan Jun dan Oktober 2015, serta Januari 2016. Keadaan persekitaran dan fizikokimia air telah diukur di dalam kawasan paya gambut, sawah padi, dan ladang kelapa sawit. Pelbagai spesies ikan dikutip serta pengasingan dan pengenpastian bakteria daripada ikan, air, dan juga sedimen menggunakan teknik mikrobiologi piawai. Analisis statistik juga telah dijalankan menggunakan ANOVA dan Turkey LSD, Kruskal-Wallis dan ujian Mann-Whitney, Analisis Komponen Utama (PCA) dan Analisis Menyurut Canonical (CCA). Tambahan pula, struktur komuniti ikan telah dikira menggunakan kepelbagaian Shannon-Weaver, kesamarataan Pielou dan indeks kekayaan Margalef.

Paya gambut merekodkan jumlah fosfat yang signifikan ($1.937 \pm 0.196 \text{ mg L}^{-1}$), sulfat ($28.917 \pm 20.398 \text{ mg L}^{-1}$) dan klorin ($0.915 \pm 0.361 \text{ mg L}^{-1}$), manakala bagi nilai pH yang paling rendah (3.684 ± 0.251), oksigen terlarut ($0.590 \pm 0.169 \text{ mg L}^{-1}$) dan amonia-nitrogen ($0.346 \pm 0.081 \text{ mg L}^{-1}$) apabila dibandingkan dengan sawah padi dan juga ladang kelapa sawit. Walau bagaimanapun, tiada perbezaan yang signifikan diperhatikan bagi konduktiviti elektrik dalam air, jumlah pepejal terlarut, saliniti, nitrit bagi sekurang-kurangnya dua habitat persampelan, serta bagi semua kawasan persampelan untuk suhu air. Tahap klorin dan fosfat pada setiap kawasan, pH bagi kawasan paya gambut (3.684 ± 0.251) serta sawah padi (4.962 ± 0.462) dan juga tahap oksigen terlarut di dalam paya gambut adalah melebihi tahap piawai untuk menampung hidupan akuatik dan jatuh kedalam kelas V bagi Pawaian Kualiti Air Nasional (NWQS) Malaysia.

Sawah padi merekodkan jumlah signifikan skor yang lebih rendah bagi kebolehcapaian (1.50 ± 0.548) dan tahap pembasmian hutan (1.00 ± 0.000), manakala ladang kelapa sawit mencatatkan skor paling rendah bagi tahap pencemaran (2.00 ± 0.000). Terdapat tiada perbezaan yang signifikan bagi tahap pembangunan manusia, kedalaman air, serta pencemaran sekurang-kurangnya pada dua kawasan habitat persampelan, serta bagi semua habitat persampelan dalam erti kata lebih jauh daripada kawasan kediaman manusia. Dengan keberkecualian bagi kedalaman air, kesemua keadaan yang telah

diukur menunjukkan kepentingan dalam mengenalpasti tahap kualiti air bagi NSPSF. Analisis statistik memaparkan tahap pencemaran adalah faktor alam sekitar yang terpenting untuk mengenalpasti kualiti air didalam NSPSF.

Sejumlah 1382 individu ikan tergolong kepada 20 spesies telah dikutip sepanjang kajian ini. Hanya 15 spesies dikutip di kawasan paya gambut, berbanding dengan 19 dan 20 spesies daripada kawasan sawah padi serta ladang kelapa sawit. Pada masa yang sama, kawasan paya gambut, merekodkan jumlah kepelbagaian spesies paling rendah (2.4243) dan kekayaan (2.4768), berbanding sawah padi dan ladang kelapa sawit. Keluarga Cyprinidae mempunyai nombor tertinggi dalam perwakilan spesies. Pemerhatian perlu diambil bagi beberapa spesies yang menghadapi kepupusan sebelum daripada catatan NSPSF. Sebagai contoh, spesies *Betta livida* dan *Parosphromenus harveyi* telah disenarai sebagai menghadapi kepupusan oleh Senarai Merah Spesies Terancam IUCN tidak terdapat didalam kajian ini dan perlu dinilai semula status konservasinya. Statistik pelbagai menunjukkan parameter fizikokimia air seperti suhu air, konduktiviti, amonia, sulfat, klorin serta keterlarutan oksigen mempengaruhi kehadiran ikan.

Sejumlah 3421 pengasingan daripada 39 spesies bakteria diasingkan daripada kajian ini. Spesies paling dominan ialah *Escherichia coli*, *Salmonella* spp., dan *Streptococcus agalactiae*. Sawah padi dan ladang kelapa sawit mencatatkan pengasingan dan juga jumlah spesies tertinggi, berikutan terdapat input antropogenik yang tinggi dikawasan ini. Terdapat beberapa spesies patogen antaranya ialah *Aeromonas hydrophila*, *Edwardsiella tarda*, *Staphylococcus* spp., dan *Streptococcus agalactiae*. Parameter fizikokimia air seperti suhu air, konduktiviti, amonia, sulfat, klorin dan oksigen terlarut adalah penting dalam mempengaruhi komposisi bakteria.

Kajian ini mengenalpasti kawasan paya bertukar kepada pertanian, input antropogenik seperti baja, domestik serta pembuangan kumbahan industri sebagai ancaman terbesar kepada komuniti biologi didalam NSPSF. Selain itu, kajian ini juga mengangkat kepentingan penglibatan strategi pembaikpulih kualiti air dalam memelihara paya gambut.

ACKNOWLEDGEMENTS

My gratitude to almighty Allah for His blessings, protection, strength, good health and abilities He showered on me to enable me get to this point in my life. My heart deep appreciation goes to my supervisory committee Prof. Dr. Ahmad Ismail, Dr. Mohammad Noor Amal Azmai and Dr. Syaizwan Zahmir Zulkifli, who have given me individual help and support. I thank them for their invaluable guidance, ideas, patience, and encouragement.

My heartfelt gratitude to the management of Kogi State College of Education Ankpa, Nigeria for giving me the opportunity for this program. Special thanks to the Tertiary Education Trust Fund (TETFund), Abuja, Nigeria for the grant to undergo this program. In a very special way, I thank Dr. Drisu J. Okai, the Registrar, Kogi State College of Education Ankpa, Nigeria, for his immeasurable and unrivalled support throughout my life. Thank you for being there for me, Dad.

My deepest appreciation to my beautiful family, my loving and caring wife Mrs Rabiat Aliu, my dearest children Aqeelah Aliu and Hafeezah Aliu who has been my inspirations. I cannot quantify your patience, sacrifices and support throughout my studies. Your understanding with daddy for leaving you for this long is deeply appreciated. I also thank my parents, Mr. Hussein F. Sule and Mrs. Haishat A. Sule for their prayers and love. Not forgetting my siblings, I thank all of you for your prayers.

To my friends and colleagues at University Putra Malaysia, thank you for your support and time. Finally, I thank my colleagues in Kogi State College of Education who have assisted me in one way or the other.

This thesis submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Ahmad Ismail, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Mohammad Noor Amal Azmai, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

Syaizwan Zahmir Zulkifli, PhD

Senior Lecturer
Faculty of Science
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by Members of Supervisory committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) were adhered to.

Signature: _____
Name of
Chairman of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

Signature: _____
Name of
Member of
Supervisory
Committee: _____

TABLE OF CONTENTS

ABSTRACT	Page i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
CHAPTER	
1 INTRODUCTION	1
1.1 Overview	1
1.2 Objectives of the study	3
1.3 Approach of study	3
2 LITERATURE REVIEW	5
2.1 Peat swamp forest	5
2.2 North Selangor peat swamp forest	5
2.3 Fish diversity of north Selangor peat swamp forest	7
2.4 Bacterial communities in north Selangor peat swamp forest	14
2.5 Bacterial flora of fish	14
2.6 Degradation of peat swamp forests (PSFs)	16
2.7 Importance of peat swamp forests conservation	19
3 MATERIALS AND METHODS	22
3.1 Study area	22
3.2 Sampling sites	22
3.2.1 Peat swamp sites	25
3.2.2 Paddy field sites	25
3.2.3 Oil palm plantation sites	26
3.3 Environmental conditions measurement	27
3.4 Water physicochemical analyses	29
3.5 Fish collection and identification	29
3.6 Bacteriological examination	29
3.6.1 Water and sediment samples for bacteriological analyses	29
3.6.2 Fish samples for bacteriological analyses	30
3.6.3 Bacterial isolation from water and sediment	30
3.6.4 Bacterial isolation from fish	30
3.6.5 Bacterial identification from water, sediment and fish	31
3.7 Statistical analyses	31

4	RESULTS	32
4.1	Environmental factors	32
4.2.	Physicochemical water parameters	34
4.3	Principal component analysis of water physicochemical parameters	36
4.4	Principal component analysis of environmental factors	37
4.5	Canonical correspondence analysis of environmental factors	37
4.6	Effects of environmental factors on water physicochemical parameters	38
4.7	Taxonomic outline of fish collected (Appendix 1)	39
4.8	Fish community structure	47
4.9	Canonical correspondence analysis of water physicochemical parameters contribution to the CCA ordination axes influencing fish occurrence	47
4.10	Relationship between water physicochemical parameters and fish occurrence	49
4.11	Bacteria count from water, sediment and fish	52
4.12	Taxonomic composition of bacteria	52
4.13	Canonical correspondence analysis of water physicochemical parameters contribution to the CCA ordination axes influencing fish occurrence	60
4.14	Relationship between water physicochemical parameters and bacterial composition	62
5	DISCUSSION	65
5.1	Influence of anthropogenic activities on the water quality in NSPSF	65
5.2	Fish diversity and the influence of water quality on their occurrence in the NSPSF	70
5.3	Bacteria composition and the influence of water quality on their presence in the NSPSF	73
6	SUMMARY, CONCLUSION AND RECOMMENDATIONS	76
	BIBLIOGRAPHY	78
	APPENDICES	94
	BIODATA OF STUDENT	129
	LIST OF PUBLICATIONS	130

LIST OF TABLES

Table	Page
2.1 Forest Reserves and area in the NSPSF	6
2.2 Fish species recorded in north Selangor Peat Swamp Forest	8
3.1 Aspects of environmental factors and human activities	27
4.1 Comparison of environmental factors score between the three habitats investigated	33
4.2 Comparison of water quality parameters between the three habitats investigated	35
4.3 Principal component loadings from principal component analysis of water quality parameters from peat swamp, paddy field and oil palm of north Selangor peat swamp forest	36
4.4 Eigenvalues, percentage variance explained and component loadings from principal component analysis of environmental variables	37
4.5 Canonical correspondence analysis summary statistics for environmental variables	38
4.6 List of family, fish species, IUCN status, number and percentages of collected from peat swamp, paddy field, and oil palm plantation	40
4.7 Fish community structure in peat swamp, paddy field and oil palm expressed as diversity, evenness and richness indexes	47
4.8 Canonical correspondence analysis summary statistics of physicochemical parameters of water	48
4.9 Bacterial abundance (mean $\pm$ standard deviation of 10^8 cfu g ⁻¹ , cfu mL ⁻¹ or cfu cm ⁻²) in sediment, water, gill, surface and intestine of fish	52
4.10 List of bacteria, abbreviation used, number and percentage of isolate from peat swamp, paddy field and oil palm of north Selangor peat swamp forest	53
4.11 List of bacteria, abbreviation used, number and percentage of isolate in sediment, water, gill, surface and intestine in peat swamp	55
4.12 List of bacteria, abbreviation, number and percentage of isolate in sediment, water, gill, surface and intestine in paddy field	56
4.13 List of bacteria, abbreviation, number and percentage of isolate from sediment, water, gill, surface and intestine in oil palm plantation	58
4.14 Canonical correspondence analysis summary statistics of physicochemical parameters of water	61
5.1 Comparison of water quality parameters in peats swamps from studies conducted between 1989 to current observation	66
A2 Normality tests, Kruskal-Wallis and Wilcoxon-Mann-Whitney U-test of environmental factors	97
A3 Normality tests, ANOVA and LSD pairwise comparison of water physicochemical parameters	101
A4 Test of normality, ANOVA and pairwise comparison of bacterial colony forming unit	106
A5 Shannon-Weaver, Pielou and Margalef indexes	110
A6 Principal Component Analysis (PCA) of environmental factors	113

A7	PCA of water physicochemical parameters	115
A8	Canonical Component Analyses (CCA)	118



LIST OF FIGURES

Figure	Page
1.1	4
2.1	17
2.2	17
2.3	17
2.4	17
2.5	18
2.6.	18
2.7	18
2.8	18
2.9	19
2.10	19
2.11	19
3.1	23
3.2	24
3.3	24
3.4	24
3.5	24
3.6	25
3.7	25
3.8	25
3.9	25
4.1	39
4.2	49
4.3	50
4.4	51
4.5	62
4.6.	63
4.7	64
A1	94

CHAPTER 1

INTRODUCTION

1.1 Overview

Peat swamps are biotopes of great ecological importance. Like most brackish water habitats, there have a unique characteristic of being associated with a particular ecological community, with high biological diversity and trophic migrations, which could be in peril without appropriate management (Cognetti and Maltagliati 2000). Over the last decades, peat swamps in particular and aquatic systems in general have been globally impacted by a wide array of human factors (Allard et al. 1991; Dynesius and Nilsson 1994; Rahel 2000). Anthropogenic activities modify the physical, chemical and biological processes of water resources, which results in modification of the resident biological community. This modification could be in the form of reduced richness, diversity and extinction, or a replacement of the community with more resilient ones (Giam et al. 2012; Kuklina et al. 2013). As a result of the imperative biological response of communities to environmental changes, they serve as precise indicators of the status of the aquatic system as they are subject to the full range of chemical and physical influences, with additive and synergistic effects included (Adams et al. 1999; Schiemer 2000; Wolska et al. 2007).

In a delicate semi-closed ecosystem such as peat swamps, environmental and ecological factors cloud organisms' response to environmental stressors (Adams et al. 1989; Adams et al. 1999). Ellis (1999) mentioned that such stressors predisposes organisms, especially fish to chronic immune suppression, thus stimulating bacterial invasion. Generally, fish are continuously exposed to the microorganisms present in water and in sediment (El-Shafai et al. 2004). These organisms change the microbial flora on external surfaces of the fish, including the gills, with high accumulation at sites of damage, such as missing scales or abrasions (Austin and Austin 2012). Similarly, food and water populated with microorganisms are delivered to the digestive tract via ingestion (Olafsen 2001), and they eventually colonise the digestive tract (Austin and Austin 2012).

Fisheries represent an important sector in a country's national income structure. According to Huq et al. (1983), swamps serve as natural water bodies for various fish species of differing sizes, and an important source of income for small scale farmers. Swamp fish provides the cheapest protein, and involves small investment and little or no monitoring as the fish can grow easily without care. Peat swamps with its unique characteristics does not just provide fish for consumption or ornamental values but also serve as a reservoir of bio-indicator organisms of great research importance (United Nations Development Programme (UNDP) 2006; Yule 2010). Therefore, unregulated alteration to natural swamps, including peat swamps, can have its toll on a nation's economy.

The biological responses of animals and plants to environmental stressors have been well researched (Pounds et al. 2006; Whitfield et al. 2007; Dong et al. 2010). Also the bacterial community of several cultured and wild fishes (both freshwater and marine) have been well studied and documented (Austin and Austin 1999; Olafsen 2001;

Moustafa et al. 2010; Khan and Ghosh 2012; Saad 2013). But, only few studies have been conducted on the response of swamp fishes to human induced environmental stress, and on the bacterial communities of swamps. Such studies on peat swamp forest is completely fragmented, incomplete and mostly without a major focus on the response of biological communities (Mishra et al. 2014). In Malaysia especially, such studies on peat swamps and associated habitats are completely lacking.

The north Selangor Peat Swamp Forest (NSPSF), located on a plain coast in the northern part of the state of Selangor is home to diverse fish species, some of which are stenotopic to the peat swamp of the forest (Ng et al. 1994; Beamish et al. 2003). The desire for agricultural development and expansion in recent times, had resulted in destruction of a large part of the forest and its subsequent conversion to paddy fields and oil palm plantations (Giam et al. 2012). Consequently, the agricultural parts of the NSPSF is now characterized with intense anthropogenic activities (Posa et al. 2011). This has led to further deterioration of the environment, a scenario that can make it inhabitable to native fish species especially (Stoeckl et al. 2015), hence reducing their chance of survival, their richness and diversity.

As a result of the low pH and other unique water attributes, fish species that can survive and breed in peat swamps are limited (Ng et al. 1994). Many species found in peat swamps are not found in any other habitat. Thus fish resources in the NSPSF are exploited for the aquarium trade and consumption (Zakaria et al., 1999). A review of fish surveys reveal that 114 fish species belonging to 23 families have been recorded from the NSPSF (Davies and Abdullah 1989; IPT-AWB 1993; Ng et al. 1994; Beamish et al. 2003; Giam et al. 2012; Ahmad et al. 2013; Ismail et al. 2013; Siow et al. 2013). Some recent surveys had reported a decline in the fish community structure resulting from their displacement following peatland conversions in NSPSF (Beamish et al. 2003; Giam et al. 2012). Some of the fish species listed as endangered by the IUCN Red List of Threatened Species have been rarely sighted during recent surveys (Giam et al. 2012; Ahmad et al. 2013; Ismail et al. 2013; Siow et al. 2013). This trend of peatland conversion which has continued despite the associated risks could result in the extinction of already endangered species in the near future.

Water from the peat swamps is used in irrigating the paddy fields and oil palm plantations, and occasional overflow occurs especially during high rainfall. Hence, fish is suspected to migrate from the adjacent paddy fields and oil palm plantations into the peat swamps of the NSPSF and vice versa, especially during seasons of high rainfall (Irvine et al. 2013). By implication therefore, there is the possibility of bacteria being introduced from the highly impacted anthropogenic agricultural areas into the peat swamps, especially through trophic migrations of fish. This could increase the incidence of bacterial infections, which are one of the causative agents of fish diseases, and the primary cause of mortality in wild and cultured fishes (Posa et al. 2011; Coll et al. 2016). It could also lead to or increase the risk of infection of humans by human pathogenic bacteria in and around fish. Such scenario of intense human activities, deterioration of the environment and migrations of fish, could pose a risk to the brackish water environment without appropriate management and risk assessment.

1.2 Objectives of the study

This study has three principal objectives, summarized as follows:

1. To determine the influence of anthropogenic activities on the water physicochemical parameters in peat swamp, paddy field and oil palm plantation of the NSPSF.
2. To determine the fish community structure in peat swamp, paddy field and oil palm plantations of NSPSF, and their relationship to the water physicochemical parameters.
3. To determine the bacterial presence in fishes, water and sediments of peat swamps, paddy fields and oil palm plantations of NSPSF, and their relationship to the water physicochemical parameters.

1.3 Approach of study

The approach in implementing this study involved a focused preliminary observation, identification of human activities and resultant environmental conditions, study sites selection, physicochemical analyses of water, fish sampling for fish identification and community structure, bacteriological examination of water, sediment and fish samples (Fig. 1.1). The methods adopted in selection of sampling sites, the duration and period of sampling, collection of fish, water and sediment, techniques in isolation and identification of bacteria, and the statistical methods in analysing data obtained are based on the insights and experiences derived from review of literatures and a careful scrutiny of the strengths and weaknesses of various available techniques. In addition, a preliminary study was conducted in the study area prior to the actual research to identify habitat characteristics, inhomogeneity in space or time, and possible complications to aid in stratification of sampling sites, and selection of best-fit sampling techniques.




Figure 1.1: Flowchart of the study

BIBLIOGRAPHY

- Adams SM, Bevelhimer MS, Greeley MS, et al (1999) Ecological risk assessment in a large river-reservoir: 6. Bioindicators of fish population health. *Environ Toxicol Chem* 18:628–640.
- Adams SM, Shepard KL, Greeley MS, et al (1989) The use of bioindicators for assessing the effects of pollutant stress on fish. *Mar Environ Res* 28:459–464.
- Ahl E (1922) Einige neue Süßwasserfische des Indo-Malayischen Archipels. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin* 1922:30–36.
- Ahmad A, Mazlan AG, Syed Ahmad Rizal TN, et al (2005) Freshwater fishes of Sungai Bebar and adjacent areas. In: Summary Findings PSF Technical Series No. 4. Peat Swamp Forest Project. UNDP, GEF, Pahang Forestry Department & Universiti Kebangsaan, pp 145–148
- Ahmad AB, Ahmad MF, Nek SART (2013) Freshwater fishes recorded from North Selangor Peat Swamp Forest, Selangor. In: Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest. Selangor, pp 72–76
- Ahmad MF, Samat A (2015) Species composition and abundance of peat swamp fishes in selected areas of Selangor, Malaysia. *Malaysian Appl Biol* 44:139–145.
- Al-Badaii F, Othman MS, Gasim MB (2013) Water quality assessment of the Seminyih river, Selangor, Malaysia. *J Chem* 2013:10. doi: 10.1155/2013/871056
- Al-Harbi AH (2003) Faecal coliforms in pond water, sediments and hybrid tilapia *Oreochromis niloticus* × *Oreochromis aureus* in Saudi Arabia. *Aquac Res* 34:517–524.
- Al-Harbi AH, Uddin N (2003) Quantitative and qualitative studies on bacterial flora of hybrid tilapia (*Oreochromis niloticus* × *O. aureus*) cultured in earthen ponds in Saudi Arabia. *Aquac Res* 34:43–48.
- Al-Harbi AH, Uddin N (2005) Bacterial diversity of tilapia (*Oreochromis niloticus*) cultured in brackish water in Saudi Arabia. *Aquaculture* 250:566–449.
- Alberto WD, María del Pilar D, María Valeria A, et al (2001) Pattern recognition techniques for the evaluation of spatial and temporal variations in water quality. A case study: Suquia River basin (Cordoba-Argentina). *Water Res* 35:2881–2894. doi: 10.1016/S0043-1354(00)00592-3
- Allard B, Craun GF, Oude NT, et al (1991) *Water Pollution*. Springer Berlin Heidelberg, Berlin, Heidelberg
- Allen DA, Austin B, Colwell RR (1983) Numerical Taxonomy of Bacterial Isolates Associated with a Freshwater Fishery. *Microbiology* 129:2043–2062.
- Allen DJ (2012) *Barbonymus schwanenfeldii*. In: IUCN Red List Threat. Species 2012 e.T181160A1705016. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T181160A1705016.en>. Accessed 20 Mar 2016
- Amal MNA, Zamri-Saad M, Siti-Zahrah A, et al (2010a) Prevalence of *Streptococcus agalactiae* in tilapia kept in different water bodies. *Online J Vet Res* 11:153–162.
- Amal MNA, Zamri-Saad M, Siti-Zahrah S, Zulkafli AR (2015) Water quality influences the presence of *Streptococcus agalactiae* in cage cultured red hybrid tilapia, *Oreochromis niloticus* × *Oreochromis mossambicus*. *Aquac Res* 46:313–323.
- Amal MNA, Zamri-Saad M, Zulkafli AR, et al (2010b) Water thermocline confirms

- susceptibility of tilapia cultured in lakes to *Streptococcus agalactiae*. J Anim Vet Adv 9:2811–2817.
- Ambak MA, Isa MM, Zakaria MZ, Ghaffar MA (2010) Fishes of Malaysia. Pernebit UMT, Universiti Malaysia Terengganu, Terengganu
- Anderson JAR (1983) The tropical Peat Swamps of Western Malesia. In: Gore AJP (ed) Systems of the World 4B. Mires: Swamp, Bog, Fen and Moor, 4B edn. Elsevier Sci. Publ., Amsterdam, pp 181–199
- Apun K, Yusof AM, Jugang K (1999) Distribution of bacteria in tropical freshwater fish and ponds. Int J Environ Health Res 9:285–292.
- Arce RG, de la Cruz CR (1977) Design/layout considerations in a freshwater rice-fish farming in the Philippines. Joint SCSP/SEAFDEC Workshop on Aquaculture Engineering (with emphasis on small-scale aquaculture projects). Volume 2 - Technical report. 335–346.
- Astel A, Biziuk M, Przyjazny A, Namieśnik J (2006) Chemometrics in monitoring spatial and temporal variations in drinking water quality. Water Res 40:1706–1716. doi: 10.1016/j.watres.2006.02.018
- Austin B (2006) The bacterial microflora of fish, revised. Sci World J 6:931–45. doi: 10.1100/tsw.2006.181
- Austin B (1982) Taxonomy of bacteria isolated from a coastal, marine fish-rearing unit. J Appl Bacteriol 53:253–268.
- Austin B, Austin DA (2012) Bacterial Fish Pathogens: Disease of Farmed and Wild Fish, Fifth Ed. Springer Science & Business Media
- Austin B, Austin DA (1999) Bacterial Fish Pathogens: Disease of Farmed and Wild Fish, Third ed. Springer Science & Business Media
- Azad IS, Al-Marzouk A, James CM, et al (2012) Outbreak of natural Streptococcosis in hatchery produced silver pomfret (*Pampus argenteus* Euphrasen) larvae in Kuwait. Aquaculture 330–333:15–20.
- Bailey RA, Clark HM, Ferris JP, et al (2002) Chemistry of the Environment. Elsevier
- Beamish FWH, Beamish RB, Lim SL (2003) Fish assemblages and habitat in a Malaysian blackwater peat swamp. Environ Biol Fishes 68:1–13.
- Beamish FWH, Saardrit P, Tongnunui S (2006) Habitat characteristics of the cyprinidae in small rivers in Central Thailand. Environ Biol Fishes 76:237–253.
- Bell L (2014) What is peat swamp, and why should I care? In: Mongabay, Peatlands. <http://news.mongabay.com/2014/07/what-is-peat-swamp-and-why-should-i-care/>. Accessed 11 Jan 2016
- Bertin L, Estève R (1948) Catalogue des types de poissons du Muséum national d'Histoire naturelle. 4e partie. Ostariophysaires (Cypriniformes). Muséum National d'Histoire Naturelle, Paris
- Biodic (2013) Threatened Species. ww.biodic.go.jp. Accessed 1 Mar 2015
- Bleeker P (1851a) Nieuwe bijdrage tot de kennis der ichthyologische fauna van Celebes. Natuurkundig Tijdschr voor Ned Indië 2:209–224.
- Bleeker P (1846a) Siluroideorum bataviensium conspectus diagnosticus [Overzicht der Siluroïden, welke te Batavia voorkomen]. Typis Societatis Artium et Scientiarum Bataviensis, Batavia
- Bleeker P (1846b) Overzicht der Siluroïden, welke te Batavia voorkomen. Natuur-en Geneeskd Arch voor Neêrland's Indië 3:135–184.
- Bleeker P (1851b) Vijfde bijdrage tot de kennis der ichthyologische fauna van Borneo met beschrijving van eenige nieuwe soorten van zoetwatervisschen.

- Natuurkundig Tijdschr voor Ned Indië 2:415–442.
- Bleeker P (1853) Nalezingen op de ichthyologische fauna van het eiland Banka. Natuurkundig Tijdschr voor Ned Indië 5:175–194.
- Bleeker P (1857) Descriptiones specierum piscium javanensium novarum vel minus cognitarum diagnosticae. Natuurkundig Tijdschr voor Ned Indië 13:323–368.
- Bleeker P (1849) Bijdrage tot de kennis der ichthyologische fauna van Midden- en Oost-Java, met beschrijving van eenige nieuwe species. Verh van het Bataviaasch Genoot van Kunsten en Wet 23:1–23.
- Bleeker P (1855) Verslag van eenige verzamelingen van visschen van Oost-Java. Natuurkundig Tijdschr voor Ned Indië 9:391–414.
- Bleeker P (1854a) Nieuwe tientallen diagnostische beschrijvingen van nieuwe of weinig bekende vischsoorten van Sumatra. Natuurkundig Tijdschr voor Ned Indië 5:495–534.
- Bleeker P (1860) Achtste bijdrage tot de kennis der vischfauna van Sumatra. Visschen van Benkoelen, Priaman, Tandjong, Palembang en Djambi. Acta Soc Regiae Sci Indo-Neêrlandicae 8:1–88.
- Bleeker P (1854b) Zevende bijdrage tot de kennis der ichthyologische fauna van Borneo. Zoetwatervisschen van Sambas, Pontianak en Pangaron. Natuurkundig Tijdschr voor Ned Indië 5:427–462.
- Bleeker P (1852a) Zesde bijdrage tot de kennis der ichthyologische fauna van Borneo. Visschen van Pamangkat, Bandjermassing, Praboeakarta en Sampit. Natuurkundig Tijdschr voor Ned Indië 3:407–442.
- Bleeker P (1850) Bijdrage tot de kennis der ichthyologische fauna van Borneo, met beschrijving van 16 nieuwe soorten van zoetwatervisschen. Natuurkundig Tijdschr voor Ned Indië 1:1–16.
- Bleeker P (1852b) Bijdrage tot de kennis der ichthyologische fauna van Blitong (Billiton), met beschrijving van eenige nieuwe soorten van zoetwatervisschen. Natuurkundig Tijdschr voor Ned Indië 3:87–100.
- Bloch ME (1792) Naturgeschichte der Ausländischen Fische. Sechster Theil, Morino, Berlin
- Borodin NA (1930) Scientific results of the yacht “Ara” expedition during the years 1926 to 1930, while in command of William K. Vanderbilt. Fishes (collected in 1929). Bull Vanderbilt Mar Museum 1:39–64, pls. 1–2.
- Brett JR (1972) The metabolic demand for oxygen in fish, particularly salmonids and a comparison with other vertebrates. Respir Physiol 14:151–170.
- Britz R, Kumar K, Baby F (2012) *Pristolepis rubripinnis*, a new species of fish from southern India (Teleostei: Percomorpha: Pristolepididae). Zootaxa 3345:59–68.
- Chapin M (2004) A Challenge to Conservationists. World Watch 17–31.
- Chernow BA, Vallasi GA (2003) Columbia Electronic Encyclopedia, 6th edn. Columbia University Press
- Chong VC, Lee PKY, Lau CM (2010) Diversity, extinction risk and conservation of Malaysian fishes. J Fish Biol 76:2009–2066. doi: 10.1111/j.1095-8649.2010.02685.x
- Christensen PJ (1977) The history, biology, and taxonomy of the Cytophaga group. Can J Microbiol 23:1599–653.
- Cognetti G, Maltagliati F (2000) Biodiversity and Adaptive Mechanisms in Brackish Water Fauna. Mar Pollut Bull 40:7–14.
- Coll M, Shannon LJ, Kleisner KM, et al (2016) Ecological indicators to capture the

- effects of fishing on biodiversity and conservation status of marine ecosystems. *Ecol Indic* 60:947–962. doi: 10.1016/j.ecolind.2015.08.048
- Colwell RR (1962) The bacterial flora of puget sound fish. *J Appl Bacteriol* 25:147–158.
- Corwin DL, Loague K, Ellsworth TR (1999) Advanced information technologies for assessing nonpoint source pollution in the vadose zone: Conference overview. *J Environ Qual* 28:357–365.
- Craft C, Krull K, Graham S (2007) Ecological indicators of nutrient enrichment, freshwater wetlands, Midwestern United States (U.S.). *Ecol Indic* 7:733–750. doi: 10.1016/j.ecolind.2006.08.004
- Cruz EM, de la Cruz CR (1991) Production of common carp (*Cyprinus carpio*) with supplemental feeding in rice fields in north Sumatra, Indonesia. *Asian Fish Sci* 4:31–39.
- Cuvier G (1829) Le règne animal distribué d'après sa organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Tome II. Déterville, Paris
- Cuvier G, Valenciennes A (1831) Histoire naturelle des poissons. Tome septième. Levrault, Paris
- Cuvier G, Valenciennes A (1840) Histoire naturelle des poissons. Tome quatorzième. Pitois-Levrault, Paris
- Cuvier G, Valenciennes A (1842) Histoire naturelle des poissons. Tome seizième. Bertrand, Paris
- Daldorff DC de. (1797) Natural history of *Perca splendens*. *Trans Linn Soc London* 3:62–63.
- Damodhar U, Reddy MV (2013) Impact of pharmaceutical industry treated effluents on the water quality of river Uppanar, South east coast of India: A case study. *Appl Water Sci* 3:501–514. doi: 10.1007/s13201-013-0098-x
- Davies J, Abdullah AR (1989) Freshwater Fish Survey of the North Selangor Peat Swamp Forest. Asian Wetland Bureau Publication No:46, IPT-AWB/WWF, Kuala Lumpur
- DePaola A, Capers GM, Alexander D (1994) Densities of *Vibrio vulnificus* in the intestines of fish from the U.S. Gulf Coast. *Appl Environ Microbiol* 60:984–988.
- Department of Environment (DOE) (2006) Water Quality Index. Ministry of Science, Technology and Environment Malaysia, Kuala Lumpur
- Diler Ö, Altun S, Çalikuşu F, Diler A (2000) A study on qualitative and quantitative bacterial flora of the Rainbow trout (*Oncorhynchus mykiss*) living in different fish farms. *Turkish J Vet Anim Sci* 24:251–260.
- Diler Ö, Diler A (1998) Quantitative and qualitative changes of the gastrointestinal microflora of Pike-perch (*Stizostedion Lucioperca* L. 1758) in Egirdir Lake. *Turk J Vet Anim Sci* 22:325–328.
- Dong H, Jiang H, Yu B, Liu X (2010) Impacts of environmental change and human activity on microbial ecosystems on the Tibetan Plateau, NW China. *GSA Today* 20:4–10.
- Dooley SR, Treseder KK (2012) The effect of fire on microbial biomass: a meta-analysis of field studies. *Biogeochemistry* 109:49–61.
- Dubey VK, Sarkar UK, Pandey A, et al (2012) The influence of habitat on the spatial variation in fish assemblage composition in an unimpacted tropical River of Ganga basin, India. *Aquat Ecol* 46:165–174.

- Froose R, Pauly D (2016g) *Osteochilus vittatus* (Valenciennes, 1842). In: FishBase, World Wide Web Electron. Publ. <http://fishbase.org/summary/Osteochilus-vittatus.html>. Accessed 20 Mar 2016
- Froose R, Pauly D (2016h) *Parachela oxygastroides* (Bleeker, 1852). In: FishBase, World Wide Web Electron. Publ. <http://www.fishbase.org/summary/4947>. Accessed 20 Mar 2016
- Froose R, Pauly D (2016i) *Helostoma temminckii* Cuvier, 1829. In: FishBase, World Wide Web Electron. Publ. <http://www.fishbase.org/summary/Helostoma-temminckii.html>. Accessed 20 Mar 2016
- Froose R, Pauly D (2016j) *Notopterus notopterus* (Pallas, 1769). In: FishBase, World Wide Web Electron. Publ. <http://www.fishbase.org/summary/Notopterus-notopterus.html>. Accessed 20 Mar 2016
- Froose R, Pauly D (2016k) *Trichopodus pectoralis* Regan, 1910. In: FishBase, World Wide Web Electron. Publ. <http://fishbase.org/summary/Trichopodus-pectoralis.html>. Accessed 21 Mar 2016
- Froose R, Pauly D (2016l) *Trichopodus trichopterus* (Pallas, 1770). In: FishBase, World Wide Web Electron. Publ. <http://www.fishbase.org/summary/Trichopodus-trichopterus.html>. Accessed 21 Mar 2016
- Froose R, Pauly D (2016m) *Wallago leerii* Bleeker, 1851. In: FishBase, World Wide Web Electron. Publ. fishbase.se/summary/Wallago-leerii.html. Accessed 21 Mar 2016
- Gasim MB, Azmin WN, Yaziz MI (2012) Land use change and their impact on water quality in the Semenyih River, Selangor, Malaysia. *J Technol Miner* 2:103–111.
- Giam X, Koh LP, Tan HH, et al (2012) Global extinctions of peat swamp fishes follow peatland conversion in Sundaland. *Front Ecol Environ* 10:465–470.
- Global Environment Centre (2014) Integrated Management Plan for North Selangor Peat Swamp Forest 2014-2023 for Selangor State Forestry Department. 183.
- González CJ, Encinas JP, García-López ML, Otero A (2000a) Characterization and identification of lactic acid bacteria from freshwater fishes. *Food Microbiol* 17:383–391.
- González CJ, López-Díaz TM, García-López ML, et al (1999) Bacterial microflora of wild brown trout (*Salmo trutta*), wild pike (*Esox lucius*), and aquacultured rainbow trout (*Oncorhynchus mykiss*). *J Food Prot* 62:1270–7.
- González CJ, Santos JA, García-López ML, et al (2001) Mesophilic aeromonads in wild and aquacultured freshwater fish. *J Food Prot* 64:687–91.
- González CJ, Santos JA, García-López ML, Otero A (2000b) Psychrobacters and related bacteria in freshwater fish. *J Food Prot* 63:315–21.
- Gorlach-Lira K, Pacheco C, Carvalho LCT, et al (2013) The influence of fish culture in floating net cages on microbial indicators of water quality. *Brazilian J Biol* 73:457–463.
- Grimes DJAY, Jacobs DAN, Swartz DG, et al (1993) Numerical Taxonomy of Gram-Negative , Oxidase-Positive Rods from Carcharhinid Sharks. *Int J Syst Bacteriol* 43:88–98.
- Grisez L, Reyniers J, Verdonck L, et al (1997) Dominant intestinal microflora of sea bream and sea bass larvae, from two hatcheries, during larval development. *Aquaculture* 155:387–399.
- Grover JH (1979) Rice-fish culture and the green revolution. *Advances in Aquaculture*.

- Food and Agriculture Organization of the United Nations. 223–224.
- Günther A (1876) Contributions to our knowledge of the fish-fauna of the Tertiary deposits of the highlands of Padang, Sumatra. Geol Mag New Ser Series 2,:433–440, pls. 16–19.
- Günther A (1861) Catalogue of the acanthopterygian fishes in the collection of the British Museum. Vol. 3. British Museum, London
- Hagi T, Tanaka D, Iwamura Y, Hoshino T (2004) Diversity and seasonal changes in lactic acid bacteria in the intestinal tract of cultured freshwater fish. Aquaculture 234:335–346.
- Hansen GH, Strøm E, Olafsen JA (1992) Effect of different holding regimens on the intestinal microflora of herring (*Clupea harengus*) larvae. Appl Environ Microbiol 58:461–470.
- Harris N, Minnemeyer S, Sizer N, et al (2015) With latest fire crisis, Indonesia surpasses Russia as world's fourth largest emitter. World Resour. Inst.
- Hem JD (1985) Study and interpretation of the chemical characteristics of natural water, Volume 2254 of US Geological Survey Water-Supply Paper, Third Edit. U.S. Department of the Interior, U.S. Geological Survey
- Hettiarachchi DC, Cheong CH (1994) Some characteristics of *Aeromonas hydrophila* and *Vibrio* species isolated from bacterial disease outbreaks in ornamental fish culture in Sri Lanka. J Natl Sci Counc Sri Lanka 22:261–269.
- Holben WE, Williams P, Gilbert MA, et al (2002) Phylogenetic analysis of intestinal microflora indicates a novel *Mycoplasma* phylotype in farmed and wild salmon. Microb Ecol 44:175–185.
- Holt JG, Krieg NR, Sneath PHA, Williams ST (1994) Bergey's Manual of Determinative Bacteriology, 9th ed. Williams and Wilkins, Baltimore
- Hora SL, Gupta JC (1941) Notes on Malayan fishes in the collection of the Raffles Museum, Singapore. I. Catfishes of the families Siluridae, Bagridae, Amblycepidae, Akysidae, Sisoridae, Chacidae, Schilbeidae and Clariidae. Bull Raffles Museum 17:12–43, pls. 2–4.
- Huat KK, Tan ESP (1980) Review of rice-fish culture in Southeast Asia. In: Pullin RS V., Shehadeh ZH (eds) Proceedings of the ICLARM Conference on integrated agriculture-aquaculture farming systems. International Centre for Living Aquatic Resources Management, Manila and the Southeast Asian Center for Graduate Study and Research in Agriculture, Los Banos, Laguna, Philippines, p 4
- Huq A, Small EB, West PA, et al (1983) Ecological relationships between *Vibrio cholerae* and planktonic crustacean copepods. Appl Environ Microbiol 45:275–83.
- IPT-AWB (1993) Fish Sampling Demonstration in North Selangor Peat Swamp Forest. 23.
- Irvine K, Vermette S, Mustafa FB (2013) The “black waters” of Malaysia: tracking water quality from the peat swamp forest to the sea. Sains Malaysiana 42:1539–1548.
- Ismail A, Amal MNA, Abdullah T, et al (2013) Fish diversity in north Selangor peat swamp forest. In: Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest. Shah Alam, Selangor, pp 57–71
- IUCN (2015a) IUCN Red List of Threatened Species. www.iucnredlist.org. Accessed 22 Feb 2015
- IUCN (2015b) The IUCN Red List of Threatened Species. In: Version 2015-4.

- <http://www.iucnredlist.org/>. Accessed 19 Mar 2016
- Jackson CR, Liew KC, Yule CM (2009) Structural and functional changes with depth in microbial communities in a tropical Malaysian peat swamp forest. *Microb Ecol* 57:402–12.
- Jenkins A, Kullander FF, Tan HH (2009a) *Parachela oxygastroides*. In: IUCN Red List Threat. Species 2009 e.T169547A6645702. <http://dx.doi.org/10.2305/IUCN.UK.2009-2.RLTS.T169547A6645702.en>. Accessed 20 Mar 2016
- Jenkins A, Kullander FF, Tan HH (2009b) *Rasbora tornieri*. In: IUCN Red List Threat. Species 2009 e.T169513A6640735. <http://dx.doi.org/10.2305/IUCN.UK.2009-2.RLTS.T169513A6640735.en>. Accessed 20 Mar 2016
- Johnson DS (1967a) Distributional patterns of Malayan freshwater fish. *Ecol Soc Am* 48:722–730.
- Johnson DS (1967b) On the chemistry of freshwaters in southern Malaya and Singapore. *Arch fur Hydrobiol* 63:477–496.
- Johnson DS (1968) Malayan blackwaters. In: Misra R, Gopal B (eds) Proceedings of the Symposium on recent advances in tropical ecology, Varanasi. International Society for Tropical Ecology, Varanasi, pp 303–310
- Jufri K (2013) Peatland forest clearance in Sumatra. In: Greenpeace / A For. Conserv. Policy. <http://www.greenpeace.org/australia/en/photosandvideos/photos/slideshows/Forest-Beauty--Destruction/Peatland-Forest-Clearance-in-Sumatra/>. Accessed 27 Jan 2016
- Kamei Y, Sakata T, Kakimoto D (1985) Microflora in the alimentary tract of tilapia: Characterization and distribution of anaerobic bacteria. *J Gen Appl Microbiol* 31:115–124.
- Kaup J (1860) *Anabas trifolius* n. sp. *Arch für Naturgeschichte*, Berlin 26:124–128.
- Khan A, Ghosh K (2012) Characterization and identification of gut-associated phytase-producing bacteria in some fresh water fish cultured in ponds. *Acta Ichthyol Piscat* 42:37–45.
- Kim DH, Brunt J, Austin B (2007) Microbial diversity of intestinal contents and mucus in rainbow trout (*Oncorhynchus mykiss*). *J Appl Microbiol* 102:1654–64.
- Kottelat M (2013) The Fishes of the Inland Waters of Southeast Asia: A Catalogue and Core Bibliography of the Fishes Known to Occur in Freshwaters, Mangroves and Estuaries. *Raffles Bull Zool Supplement* No. 27: 1-663.
- Kottelat M (1996a) *Betta livida*. In: IUCN Red List Threat. Species 1996 e.T2775A9479874. <http://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T2775A9479874.en>. Accessed 27 Mar 2016
- Kottelat M (1996b) *Parosphromenus harveyi*. In: IUCN Red List Threat. Species 1996 e.T16264A5594174. <http://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T16264A5594174.en>. Accessed 27 Mar 2016
- Kottelat M (1996c) *Betta hipposideros*. In: IUCN Red List Threat. Species 1996 e.T2774A9479841. <http://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T2774A9479841.en>. Accessed 27 Mar 2016
- Kottelat M, Whitten AJ, Kartikasari SN, Wirjoatmodjo S (1993) Freshwater fishes of

- Western Indonesia and Sulawesi. Periplus Editions, Hong Kong
- Kottelat M, Whitten T (1996) Freshwater biodiversity in Asia with special reference to fish.
- Kottelat M, Widjanarti E (2005) The fishes of Danau Sentarum National park and the Kapuas Lakes area, Kalimantan Barat, Indonesia. *Raffles Bull Zool* 13:13–173.
- Kuklina I, Kouba A, Kozák P (2013) Real-time monitoring of water quality using fish and crayfish as bio-indicators: a review. *Environ Monit Assess* 185:5043–53.
- La Cépède E (1800) Histoire naturelle des poissons. Tome second. Plassan, Paris
- Ladiges W (1934) Tropische Zierfische. Aquarium Hamburg, Hamburg
- Langner A, Miettinen J, Siegert F (2007) Land cover change 2002–2005 in Borneo and the role of fire derived from MODIS imagery. *Glob Chang Biol* 13:2329–2340.
- Langner A, Siegert F (2009) Spatiotemporal fire occurrence in Borneo over a period of 10 years. *Glob Chang Biol* 15:48–62.
- LeaMaster BR, Walsh WA, Brock JA, Fujioka RS (1997) Cold stress-induced changes in the aerobic heterotrophic gastrointestinal tract bacterial flora of red hybrid tilapia. *J Fish Biol* 50:770–780.
- Lee S, Lee Y (1995) Identification of Intestinal Microflora in Rainbow Trout. *Jour Microbiol* 33:273–277.
- Lee S, Najiah M, Wendy W, Nadirah M (2010) Antibigram and heavy metal resistance of pathogenic bacteria isolated from moribund cage cultured silver catfish (*Pangasius sutchi*) and red hybrid tilapia (*Tilapia* sp.). *Front Agric China* 4:116–120.
- Lim P, Lek S, Touch ST, et al (1999) Diversity and spatial distribution of freshwater fish in Great Lake and Tonle Sap River (Cambodia, Southeast Asia). *Aquat Living Resour* 12:379–386.
- Liston J (1957) The occurrence and distribution of bacterial types on flatfish. *J Gen Microbiol* 16:205–216.
- Liu C-W, Lin K-H, Kuo Y-M (2003) Application of factor analysis in the assessment of groundwater quality in a blackfoot disease area in Taiwan. *Sci Total Environ* 313:77–89. doi: 10.1016/S0048-9697(02)00683-6
- Lo J, Parish F (2013) Peatlands and Climate Change in Southeast Asia. *ASEAN Peatl. For. Proj. Sustain. Manag. Peatl. For. Proj.* 26.
- MacDonald NL, Stark JR, Austin B (1986) Bacterial microflora in the gastro-intestinal tract of Dover sole (*Solea solea* L.), with emphasis on the possible role of bacteria in the nutrition of the host. *FEMS Microbiol Lett* 35:107–111.
- Machan B (1931) Neue fische aus Java (gesammelt von Regierungsrat Dr. V. Pietschmann, Wien im August 1929). *Anzeiger, Akad der Wissenschaften Wien, Math Naturwissenschaftliche Klasse* 68:221–222.
- Mahapatra SS, Sahu M, Patel RK, Panda BN (2012) Prediction of water quality using Principal Component Analysis. *Water Qual Expo Heal* 4:93–104. doi: 10.1007/s12403-012-0068-9
- Mallya P (2015) Illustration showing degradation of tropical peat swamp forest. http://mongabay-images.s3.amazonaws.com/graphics/peatlands_art_full.jpg. Accessed 27 Jan 2016
- Mandal SN, Choudhury JP, Chaudhuri SRB (2012) Neuro-PCA-Factor Analysis in prediction of time series data. *Am J Intell Syst* 2:45–52.
- Margalef R (1958) Information theory in ecology. *Gen Syst Yearb* 3:36–71.
- Mishra S, Lee WA, Hooijer A, et al (2014) Microbial and metabolic profiling reveal

- strong influence of water table and land-use patterns on classification of degraded tropical peatlands. *Biogeosciences* 11:1727–1741.
- Mohsin AKM, Ambak MA (1983) Freshwater fishes of Peninsular Malaysia. Penerbit Universiti Pertanian Malaysia, Malaysia
- Momblanch A, Paredes-Arquiola J, Munné A, et al (2015) Managing water quality under drought conditions in the Llobregat River Basin. *Sci Total Environ* 503–504:300–18. doi: 10.1016/j.scitotenv.2014.06.069
- Montes M, Pérez MJ, Nieto TP (1999) Numerical taxonomy of gram-negative, facultative anaerobic bacteria isolated from skin of turbot (*Scophthalmus maximus*) and surrounding water. *Syst Appl Microbiol* 22:604–18.
- Mostafaei A (2014) Application of multivariate statistical methods and water-quality index to evaluation of water quality in the Kashkan River. *Environ Manage* 53:865–81. doi: 10.1007/s00267-014-0238-6
- Moustafa M, Mohamed LA, Mahmoud MA, et al (2010) Bacterial infections affecting marine fishes in Egypt. *J Am Sci* 6:603–612.
- Mullally G, Dunphy N (2015) State of play review of Environmental Policy Integration literature, a report for the National Economic and Social Council. Cork
- Munro PD, Moffat CF, Couper L, et al (1997) Degradation of synthetic mud based fluids in a solid-phase test system, Fisheries Research Services No. 1/97. 45.
- Neal C, House WA, Leeks GJL, et al (2000) Conclusions to the special issue of Science of the Total Environment concerning “The water quality of UK rivers entering the North Sea.” *Sci Total Environ* 251–252:557–573. doi: 10.1016/S0048-9697(00)00395-8
- Nedoluha PC, Westhoff D (1995) Microbiological Analysis of Striped Bass (*Marone saxatilis*) Grown in Flow-Through Tanks. *J Food Prot* 58:1363–1368.
- Ng HH (2002) The identity of *Mystus nigriceps* (Valenciennes in Cuvier & Valenciennes, 1840), with the description of a new bagrid catfish (Teleostei: Siluriformes) from southeast Asia. *Raffles Bull Zool* 50:161–168.
- Ng HH (2010) *Notopterus notopterus*. In: IUCN Red List Threat. Species 2010 e.T166433A6208173. <http://dx.doi.org/10.2305/IUCN.UK.2010-4.RLTS.T166433A6208173.en>. Accessed 20 Mar 2016
- Ng HH (2012) *Hemibagrus nemurus*. In: IUCN Red List Threat. Species 2012 e.T180954A1681839. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T180954A1681839.en>. Accessed 19 Mar 2016
- Ng HH, Kottelat M (2013) The identity of the cyprinid fishes *Rasbora dusonensis* and *R. tornieri* (Teleostei: Cyprinidae). *Zootaxa* 3635:62–70.
- Ng PKL, Ng HH (1995) *Hemibagrus gracilis*, a new species of large riverine catfish (Teleostei: Bagridae) from Peninsular Malaysia. *Raffles Bull Zool* 43:133–142.
- Ng PKL, Tay JB, Lim KKP (1994) Diversity and conservation of blackwater fishes in Peninsular Malaysia, particularly in the North Selangor peat swamp forest. *Hydrobiologia* 285:203–218. doi: 10.1007/BF00005667
- Ng TP, Shamsudin I (2001) Common trees in peat swamp forests of Peninsular Malaysia. 97.
- Nieto TP, Toranzo AE, Barja JL (1984) Comparison between the bacterial flora associated with fingerling rainbow trout cultured in two different hatcheries in the North-West of Spain. *Aquaculture* 42:193–206.
- Novis J (2010) United Nations warned that corruption is undermining grants to stop logging. In: Obs. <http://www.theguardian.com/environment/2010/jul/04/united->

- nations-corruption-logging. Accessed 27 Jan 2016
- Novotny L, Dvorska L, Lorencova A, et al (2004) Fish : a potential source of bacterial pathogens for human beings. *Vet Med* 49:343–358.
- Nowell LH, Norman JE, Moran PW, et al (2014) Pesticide Toxicity Index—A tool for assessing potential toxicity of pesticide mixtures to freshwater aquatic organisms. *Sci Total Environ* 476–477:144–157. doi: 10.1016/j.scitotenv.2013.12.088
- O'Brien R, Castelloe J (2012) Exploiting the Link Between the Wilcoxon-Mann-Whitney Test and a Simple Odds Statistic. In: *JSM Proceedings, Survey Research Methods Section*. American Statistical Association, Alexandria, VA, pp 2192–198
- Ochsenreiter T, Selezi D, Quaiser A, et al (2003) Diversity and abundance of Crenarchaeota in terrestrial habitats studied by 16S RNA surveys and real time PCR. *Environ Microbiol* 5:787–97.
- Olafsen JA (2001) Interactions between fish larvae and bacteria in marine aquaculture. *Aquaculture* 200:223–247.
- Onarheim AM, Wiik R, Burghardt J, Stackebrandt E (1994) Characterization and Identification of Two *Vibrio* Species Indigenous to the Intestine of Fish in Cold Sea Water; Description of *Vibrio iliopiscarius* sp. nov. *Syst Appl Microbiol* 17:370–379.
- Paepke H-J (2009) The nomenclature of *Trichopodus pectoralis* Regan, 1910; *Trichopus cantor* Sauvage, 1884 and *Osphronemus saigonensis* Borodin, 1930 (Teleostei: Perciformes: Osphronemidae). *Vertebr Zool* 59:53–60.
- Page SE, Hoschilo A, Langner A, et al (2009) Tropical peatland fires in Southeast Asia. In: *Cochrane MA (ed) Tropical Fire Ecology: Climate Change, Land Use and Ecosystem Dynamics*. Springer, pp 263–287
- Page SE, Rieley JO, Banks CJ (2011) Global and regional importance of the tropical peatland carbon pool. *Glob Chang Biol* 17:798–818.
- Pal M, Chaudhry S (2010) *Anabas testudineus*. In: *IUCN Red List Threat. Species 2010* e.T166543A6232945. <http://dx.doi.org/10.2305/IUCN.UK.2010-4.RLTS.T166543A6232945.en>. Accessed 19 Mar 2016
- Pallas PS (1769) *Spicilegia zoologica quibus novae imprimis et obscurae animalium species iconibus, descriptionibus atque commentariis illustrantur. Fasciculus septimus*. Lange, Berlin
- Pallas PS (1770) *Spicilegia zoologica quibus novae imprimis et obscurae animalium species iconibus, descriptionibus atque commentariis illustrantur. Fasciculus octavus*. Lange, Berlin
- Parish F (2002) Overview on peat, biodiversity, climate change and fire. In: *Parish F, Padimanabhan E, Lee CL, Thang HC (eds) Prevention and control of fire in peatlands, 19-21 March 2002*. Global Environmental Centre & Forestry Department, Peninsular Malaysia, Cetaktama, Kuala Lumpur, Malaysia, pp 51–56
- Parker SP (2003) *McGraw-Hill Dictionary of Scientific and Technical Terms*, 6th edn. McGraw-Hill Education
- Pawłowski D, Okupny D, Włodarski W, Zieliński T (2014) Spatial variability of selected physicochemical parameters within peat deposits in small valley mire: a geostatistical approach. *Geologos* 20:269–288.
- Pielou EC (1969) *An introduction to mathematical ecology*. Wiley, New York
- Popta CML (1904) *Descriptions préliminaires des nouvelles espèces de poissons*

- recueillies au Bornéo central par M. le Dr. A. W. Nieuwenhuis en 1898 et en 1900. Notes from Leyden Museum 24:179–202.
- Popta CML (1906) Résultats ichthyologique des voyages scientifiques de M. le professeur Dr. A. W. Nieuwenhuis dans le centre de Bornéo (1898 et 1900). Notes from Leyden Museum 27:1–304, pls. 1–10.
- Posa MRC, Wijedasa LS, Corlett RT (2011) Biodiversity and conservation of tropical peat swamp forests. Bioscience 61:49–57.
- Pounds JA, Bustamante MR, Coloma LA, et al (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. Nature 439:161–167.
- Prentice C, Parish D (1992) Conservation of peat swamp forest: A forgotten ecosystem. Malayan Nat J 45:128–144.
- Prokhorov AM (1982) Great Soviet encyclopedia, Volume 30. Macmillan
- Putro B, Kjeldsen TR, Hutchins MG, Miller J (2016) An empirical investigation of climate and land-use effects on water quantity and quality in two urbanising catchments in the southern United Kingdom. Sci Total Environ 548–549:164–172. doi: 10.1016/j.scitotenv.2015.12.132
- Rahel FJ (2000) Homogenization of Fish Faunas Across the United States. Science (80-) 288:854–856.
- Rainboth WJ (1996) Fishes of the Cambodian Mekong. FAO species identification field guide for fishery purpose. FAO, Rome
- Rainboth WJ (1994) Inland fishes of India and adjacent countries. Rev Fish Biol Fish 4:135–136. doi: 10.1007/BF00043269
- Ranganathan K, Jeyapaul S, Sharma DC (2007) Assessment of water pollution in different bleaching based paper manufacturing and textile dyeing industries in India. Environ Monit Assess 134:363–72. doi: 10.1007/s10661-007-9628-z
- Rasul MG, Islam MS, Yahaya f. M, et al (2015) Effects of Anthropogenic Impact on Water Quality in Bertam Catchment, Cameron Highlands, Malaysia. Int. J. Ecol. Environ. Sci. 41:75–86.
- Razmkhah H, Abrishamchi A, Torkian A (2010) Evaluation of spatial and temporal variation in water quality by pattern recognition techniques: A case study on Jajrood River (Tehran, Iran). J Environ Manage 91:852–60. doi: 10.1016/j.jenvman.2009.11.001
- Regan CT (1910) The Asiatic fishes of the family Anabantidae. Proc Gen Meet Sci Bus Zool Soc London 1909:767–787, pls. 77–79.
- Rheinheimer G (1985) Aquatic Microbiology, 3rd ed. John Wiley and Sons, Chichester
- Ringø E, Bendiksen HR, Gausen SJ, et al (1998) The effect of dietary fatty acids on lactic acid bacteria associated with the epithelial mucosa and from faecalia of Arctic charr, *Salvelinus alpinus* (L.). J Appl Microbiol 85:855–64.
- Ringø E, Strøm E, Tabachek JA (1995) Intestinal microflora of salmonids: a review. Aquac Res 26:773–789.
- Ringø E, Wesmajervi MS, Bendiksen HR, et al (2001) Identification and characterization of Carnobacteria isolated from fish intestine. Syst Appl Microbiol 24:183–91.
- Roberts TR (1989) The Freshwater fishes of Western Borneo (Kalimantan Barat, Indonesia). Mem Calif Acad Sci 14:1–210.
- Roberts TR (1993) The freshwater fishes of Java, as observed by Kuhl and van Hasselt in 1820–23. Zool Verh 285:1–94.

- Roberts TR (1992) Systematic revision of the old world freshwater fish family Notopteridae. *Ichthyol Explor Freshwaters* 2:361–383.
- Romano N, Zeng C (2007) Effects of potassium on nitrate mediated alterations of osmoregulation in marine crabs. *Aquat Toxicol* 85:202–8. doi: 10.1016/j.aquatox.2007.09.004
- Rosli N, Gandaseca S, Ismail J, Jailan MI (2010) Comparative Study of Water Quality at Different Peat Swamp Forest of Batang Igan, Sibu Sarawak. *Am J Environ Sci* 6:416–421. doi: 10.3844/ajessp.2010.416.421
- Saad TT (2013) Some Study on Heavy Metal and Bacterial infection Relationship in Some Cultured Marine Fish. *J Arab Aquac Soc* 8:143–162.
- Scheili A, Rodriguez MJ, Sadiq R (2015) Development, application, and sensitivity analysis of a water quality index for drinking water management in small systems. *Environ Monit Assess* 187:685. doi: 10.1007/s10661-015-4908-5
- Schiemer F (2000) Fish as indicators for the assessment of the ecological integrity of large rivers. *Hydrobiologia* 422–423:271–278.
- Schulz R, Bundschuh M, Gergs R, et al (2015) Review on environmental alterations propagating from aquatic to terrestrial ecosystems. *Sci Total Environ* 538:246–261. doi: 10.1016/j.scitotenv.2015.08.038
- Seale A (1910) New species of Philippine fishes. *Philipp J Sci Sect. A*, 4:491–543, pls. 1–13.
- Selvam S, Manimaran G, Sivasubramanian P, et al (2013) GIS-based Evaluation of Water Quality Index of groundwater resources around Tuticorin coastal city, south India. *Environ Earth Sci* 71:2847–2867. doi: 10.1007/s12665-013-2662-y
- Seppola M, Olsen RE, Sandaker E, et al (2006) Random amplification of polymorphic DNA (RAPD) typing of carnobacteria isolated from hindgut chamber and large intestine of Atlantic cod (*Gadus morhua* L.). *Syst Appl Microbiol* 29:131–7.
- Shah ASRM, Zarul HH, Chan KY, et al (2006) A recent survey of freshwater fishes of the Paya Beriah Peat Swamp Forest, North Perak, Malaysia. *J Biosains* 17:51–64.
- Shannon CE, Weaver W (1963) The mathematical theory of communication. Univeristy of Illinois Press, Urbana
- Shuhaimi-Othman M, Lim EC, Mushrifah I (2007) Water quality changes in Chini Lake, Pahang, West Malaysia. *Environ Monit Assess* 131:279–92. doi: 10.1007/s10661-006-9475-3
- Siow R, Ramli MN, Shakori MHM, Asmuni M (2013) Fish fauna assemblages and their distribution patterns in the North Selangor Peat Swamp Forest. In: *Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest, 28th November 2013, Shah Alam, Selangor, Malaysia*. Shah Alam, Selangor, pp 77–82
- Steindachner F (1870) Ichthyologische Notizen (X). (Schluss). *Sitzungsberichte der Kais Akad der Wissenschaften Wien, Math Cl Abt. 1*, 61:623–642, 5 pls.
- Stoeckl N, Chaiechi T, Farr M, et al (2015) Co-benefits and trade-offs between agriculture and conservation: A case study in Northern Australia. *Biol Conserv* 191:478–494. doi: 10.1016/j.biocon.2015.07.032
- Suanyuk N, Kong F, Ko D, et al (2008) Occurrence of rare genotypes of *Streptococcus agalactiae* in cultured red tilapia *Oreochromis* sp. and Nile tilapia *O. niloticus* in Thailand - Relationship to human isolates? *Aquaculture* 284:35–40.
- Sugita H, Miyajima C, Deguchi Y (1991) The vitamin B12-producing ability of the

- intestinal microflora of freshwater fish. *Aquaculture* 92:267–276. doi: 6
- Sugita H, Nakamura T, Tanaka K, Deguchi Y (1994) Identification of *Aeromonas* species isolated from freshwater fish with the microplate hybridization method. *Appl Environ Microbiol* 60:3036–8.
- Sugita H, Shibuya K, Hanada H, Deguchi Y (1997) Antibacterial Abilities of Intestinal Microflora of the River Fish. *Fish Sci* 63:378–383.
- Sullan KE, Essingen SD, Lozupone A, et al (2012) Environmental and ecological factors that shape the gut bacterial communities of fish: a meta-analysis. *Mol Ecol* 21:3363–3378.
- Tan HH, Kottelat M (2009) The fishes of Batang Hari drainage, Sumatra, with description of six new species. *Ichthyol Explor Freshwaters* 20:13–69.
- Tekile A, Kim I, Kim J (2015) Mini-review on river eutrophication and bottom improvement techniques, with special emphasis on the Nakdong River. *J Environ Sci* 30:113–121. doi: 10.1016/j.jes.2014.10.014
- Thinh D V., Van NS, Nguyen THT (2012) *Barbonymus gonionotus*. In: IUCN Red List Threat. Species 2012 e.T166914A1151554. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T166914A1151554.en>. Accessed 20 Mar 2016
- Toranzo AE, Novoa B, Romalde JL, et al (1993) Microflora associated with healthy and diseased turbot (*Scophthalmus maximus*) from three farms in northwest Spain. *Aquaculture* 114:189–202.
- tree-nation (2012) Highlight a fact about peat swamp forest. <http://www.tree-nation.com/forum/2092/highlight-a-fact-about-peat-swamp-forest>. Accessed 27 Jan 2016
- Trust TJ (1975) Bacteria Associated with the Gills of Salmonid Fishes in Freshwater. *J Appl Bacteriol* 38:225–233.
- Trust TJ, Sparrow RA (1974) The bacterial flora in the alimentary tract of freshwater salmonid fishes. *Can J Microbiol* 20:1219–28.
- United Nations Development Programme (UNDP) (2006) Malaysia's Peat Swamp Forests: Conservation and Sustainable Use. United Nations Development Programme, Kuala Lumpur, Malaysia
- Vaillant L (1902) Résultats zoologiques de l'expédition scientifique néerlandaise au Bornéo central. Notes from Leyden Museum 24:1–166, pls. 1–2.
- Varol M, Sen B (2009) Assessment of surface water quality using multivariate statistical techniques: a case study of Behrimaz Stream, Turkey. *Environ Monit Assess* 159:543–53. doi: 10.1007/s10661-008-0650-6
- Vickers AJ (2005) Parametric versus non-parametric statistics in the analysis of randomized trials with non-normally distributed data. *BMC Med Res Methodol* 5:1–12. doi: 10.1186/1471-2288-5-35
- Vidthayanon C (2012a) *Channa melasoma*. In: IUCN Red List Threat. Species 2012 e.T180782A1662338. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T180782A1662338.en>. Accessed 20 Mar 2016
- Vidthayanon C (2012b) *Cyclocheilichthys apogon*. In: IUCN Red List Threat. Species 2012 e.T181284A1717065. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T181284A1717065.en>. Accessed 20 Mar 2016
- Vidthayanon C (2012c) *Cyclocheilichthys heteronema*. In: IUCN Red List Threat. Species 2012 e.T180974A1684051. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T180974A1684051.en>.

- Vidthayanon C (2012d) *Osteochilus vittatus*. In: IUCN Red List Threat. Species 2012 e.T180750A1658850. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T180750A1658850.en>.
- Vidthayanon C (2012e) *Helostoma temminkii*. In: IUCN Red List Threat. Species 2012 e.T181326A1722150. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T181326A1722150.en>. Accessed 20 Mar 2016
- Vidthayanon C (2012f) *Trichopodus pectoralis*. In: IUCN Red List Threat. Species 2012 e.T188087A1852593. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T188087A1852593.en>. Accessed 20 Mar 2016
- Vidthayanon C (2012g) *Trichopodus trichopterus*. In: IUCN Red List Threat. Species 2012 e.T187981A1841751. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T187981A1841751.en>. Accessed 20 Mar 2016
- Vidthayanon C (2012h) *Trichopodus leerii*. In: IUCN Red List Threat. Species 2012 e.T187958A1839297. <http://dx.doi.org/10.2305/IUCN.UK.2012-1.RLTS.T187958A1839297.en>. Accessed 27 Mar 2016
- von Herbing IH (2002) Effects of temperature on larval fish swimming performance: the importance of physics to physiology. *J Fish Biol* 61:865–876.
- Wantzen KM, Yule CM, Mathooko JM, Pringle CM (2011) Organic Matter processing In Tropical Streams. In: *Tropical Stream Ecology*. Academic Press, pp 44–60
- Wardle DA, Bardgett RD, Klironomos JN, et al (2004) Ecological linkages between above-ground and below-ground biota. *Science* (80-) 304:1629–1633.
- Wetlands International (2014) Tropical peat swamp forests. In: *Peatlands*. <http://www.wetlands.org/Whatarewetlands/Peatlands/Tropicalpeatswampforests/tabid/2739/Default.aspx>. Accessed 11 Jan 2016
- Whitfield SM, Bell KE, Philippi T, et al (2007) Amphibian and reptile declines over 35 years at La Selva, Costa Rica. In: *Proceedings of the National Academy of Sciences*. Washington, D.C., pp 8352–8356
- Whitmore TC (1984) *Tropical Rain Forests of the Far East*, Second. Clarendon Press, Oxford
- Whitmore TC (1988) Forest Types and Forest Zonation. In: Cranbrook E of (ed) *Key Environments: Malaysia*. Elsevier, pp 20–30
- Whitten T, Damanik SJ, Anwar J, Hisyam N (2000) *The ecology of Sumatra*. Periplus Editions (HK) Ltd, Singapore
- WHO (1998) *Guidelines for Drinking Water Quality*, Second. WHO, Geneva, Switzerland
- Wilbers G-J, Becker M, Nga LT, et al (2014) Spatial and temporal variability of surface water pollution in the Mekong Delta, Vietnam. *Sci Total Environ* 485–486:653–65. doi: 10.1016/j.scitotenv.2014.03.049
- Wilkie MP, Wood C (1991) Nitrogenous waste excretion, acid base regulation and ionoregulation in rainbow trout (*Oncorhynchus mykiss*) exposed to extremely alkaline water. *Physiol Zool* 64:1069–1086.
- Wolska L, Sagajdakow A, Kuczyńska A, Namieśnik J (2007) Application of ecotoxicological studies in integrated environmental monitoring: Possibilities and problems. *Trends Anal Chem* 26:332–344. doi: 10.1016/j.trac.2006.11.012
- Wyatt-Smith J (1959) Peat Swamp Forests in Malaya. *Forester* 22:5–31.
- Wyatt-Smith J (1964) A preliminary vegetation map of Malaya with description of the vegetation types. *J Trop Geogr* 200–213.
- Xu X, Ma Z, Man K (2015) *The Chlorine Pollution Mechanism of Groundwater in*

- Jiaozuo Area. In: Bian F, Xie Y (eds) Geo-Informatics in Resource Management and Sustainable Ecosystem. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 332–337
- Yerel S, Ankara H (2012) Erratum to: Application of multivariate statistical techniques in the assessment of water quality in Sakarya River, Turkey. J Geol Soc India 79:115–115. doi: 10.1007/s12594-012-0019-x
- Yoshimizu M, Kimura T (1976) Study on Intestinal Microflora of Salmonids. Fish Pathol 10:243–259.
- Yule CM (2010) Loss of biodiversity and ecosystem functioning in Indo-Malayan peat swamp forests. Biodivers Conserv 19:393–409.
- Yule CM (2015) Stop the Haze: Protect our peat swamp forests. In: Monash Univ. Res. say. <http://www.monash.edu.my/research/researchers-say/stop-the-haze-protect-our-peat-swamp-forests>. Accessed 11 Jan 2016
- Yule CM, Gomez LN (2008) Leaf litter decomposition in a tropical peat swamp forest in Peninsular Malaysia. Wetl Ecol Manag 17:231–241.
- Zakaria R (2015) Peat swamp forests, a real lifesaver. In: New Straits Times Online. <http://www.nst.com.my/news/2015/12/115585/peat-swamp-forests-real-lifesaver>. Accessed 11 Jan 2016
- Zakaria R, Mansor M, Ali AB (1999) Swamp-riverine tropical fish population: a comparative study of two spatially isolated freshwater ecosystems in Peninsular Malaysia. Wetl Ecol Manag 6:261–268.
- Zhang Q, Li Z, Zeng G, et al (2009) Assessment of surface water quality using multivariate statistical techniques in red soil hilly region: a case study of Xiangjiang watershed, China. Environ Monit Assess 152:123–31. doi: 10.1007/s10661-008-0301-y
- Zhang X, Wang Q, Liu Y, et al (2011) Application of multivariate statistical techniques in the assessment of water quality in the Southwest New Territories and Kowloon, Hong Kong. Environ Monit Assess 173:17–27. doi: 10.1007/s10661-010-1366-y
- Zhao J, Shi B, Qing-ru J, Ke CH (2012) Changes in gut-associated flora and bacterial digestive enzymes during the development stages of abalone (*Haliotis diversicolor*). Aquaculture 341:147–153.
- Zhong W, Gu T, Wang W, et al (2010) The effects of mineral fertilizer and organic manure on soil microbial community and diversity. Plant Soil 326:511–522.
- Zimmerman DW (2012) A note on consistency of non-parametric rank tests and related rank transformations. Br J Math Stat Psychol 65:122–44. doi: 10.1111/j.2044-8317.2011.02017.x
- Zmysłowska I, Lewandowska D, Nowakowski T, Kozłowski J (2001) Occurrence of Bacteria in Water and in Vendace (*Coregonus albula*) during Rearing in Tanks. Polish J Environ Stud 10:51–56.

LIST OF PUBLICATIONS

Journal

Sule HA, Ismail A, Amal MNA (2016) A review of the ichthyofauna of Malaysian peat swamp forests. *Pertanika J Trop Agric Sci.* 39(4):421-458

Sule HA, Ismail A, Amal MNA, Zulkifli SZ, Mohd Fauzul AMR (2016) Influence of anthropogenic activities on water physicochemical parameters in peat swamp, paddy field and oil palm plantation of north Selangor peat swamp forest. *Environ Monit Assess*, *Submitted*.

Sule HA, Ismail A, Amal MNA, Zulkifli SZ (2016) Threats and conservation of tropical peat swap forest, *Pertanika J Sch Res Rev*, *Submitted*.

Sule HA, Go R., Ismail A, Amal, MNA (2016) Addressing biodiversity loss: A review. *Pak J Bot*, *Submitted*.

Ismail A, Amal MNA, Talib A, Sepet J, Aziz A, Hassan NI, Hassan MA, **Sule HA** (2016) A survey on fish diversity in North Selangor Peat Swamp Forest. *Check List*, *Submitted*.

Sule HA, Ismail A, Amal MNA, Zulkifli SZ, Mohd Fauzul AMR (2016) The influence of water physicochemical parameters on fish community structure in peat swamp, paddy field and oil palm plantation in north Selangor peat swamp forest. *PeerJ*, *Submitted*.

Sule HA, Ismail A, Amal MNA, Zulkifli SZ, Mastor SS (2016) The influence of water physicochemical parameters on bacterial composition and abundance in peat swamp, paddy field and oil palm plantation in north Selangor peat swamp forest, *Draft*.

Proceeding

Sule HA, Ismail A, Amal MNA, Syaizwan ZZ and Aidil MF (2016) The influence of water physicochemical parameters on fish occurrence in peat swamp, paddy field and oil palm plantation of north Selangor peat swamp forest, Selangor, Malaysia. Abstract presented at Malaysia International Biology Symposium '16 (i-SIMBIOMAS), held in Putrajaya International Convention Centre, Putrajaya, Malaysia, 26–27th October 2016.

Sule HA, Ismail A, Amal MNA, Syaizwan ZZ, Siti-Suhaiba M (2016) The influence of water physicochemical parameters on bacterial composition in peat swamps, paddy field and oil palm plantations in vicinity of north Selangor peat swamp forest, Selangor, Malaysia. Abstract presented at Fundamental Science Congress '16 held in Faculty of Science, Universiti Putra Malaysia, Selangor, Malaysia, 9–10th August 2016

Sule HA, Ismail A, Amal MNA, Zulkifli SZ (2015) Human activities increase bacterial composition of North Selangor Peat Swamp Forest. In: Ismail A, Zulkifli SZ, Hanapiah M, Aziz FZA (eds) Proceedings of Seminar Ekologi Malaysia '15 held in Putrajaya International Convention Centre, Putrajaya, Malaysia, 19th October 2015. Department of Biology, Universiti Putra Malaysia, Serdang, Selangor Darul Ehsan, pp 101–103

Ismail A, Amal MNA, Talib A, Sepet J, Aziz A, **Sule HA** (2015). Fish diversity of North Selangor Peat Swamp Forest, Selangor Malaysia. In: Ismail A, Zulkifli SZ, Hanapiah M, Aziz FZA (eds) Proceedings of Seminar Ekologi Malaysia '15 held in Putrajaya International Convention Centre, Putrajaya, Malaysia, 19th October 2015. Department of Biology, Universiti Putra Malaysia, Serdang, Selangor Darul Ehsan, pp 115–118

Sule HA, Ismail A, Amal MNA, Zulkifli, SZ, Rahman F (2015) Assessment of water quality in North Selangor Peat Swamp Forest. In: Ismail A, Zulkifli SZ, Hanapiah M, Aziz FZA (eds) Proceedings of Seminar Ekologi Malaysia '15 held in Putrajaya International Convention Centre, Putrajaya, Malaysia, 19th October 2015. Department of Biology, Universiti Putra Malaysia, Serdang, Selangor Darul Ehsan, pp 189–192



UNIVERSITI PUTRA MALAYSIA

STATUS CONFIRMATION FOR THESIS / PROJECT REPORT AND COPYRIGHT

ACADEMIC SESSION : _____

TITLE OF THESIS / PROJECT REPORT :

ANTHROPOGENIC ACTIVITIES IMPACT ON WATER QUALITY, FISH COMMUNITY AND
BACTERIA PRESENCE IN DISTURBED AND UNDISTURBED NORTH SELANGOR PEAT
SWAMP FOREST, MALAYSIA

NAME OF STUDENT: SULE HUSSEIN ALIU

I acknowledge that the copyright and other intellectual property in the thesis/project report belonged to Universiti Putra Malaysia and I agree to allow this thesis/project report to be placed at the library under the following terms:

1. This thesis/project report is the property of Universiti Putra Malaysia.
2. The library of Universiti Putra Malaysia has the right to make copies for educational purposes only.
3. The library of Universiti Putra Malaysia is allowed to make copies of this thesis for academic exchange.

I declare that this thesis is classified as :

*Please tick (✓)

☐

CONFIDENTIAL

(Contain confidential information under Official Secret Act 1972).

☐

RESTRICTED

(Contains restricted information as specified by the organization/institution where research was done).

☐

OPEN ACCESS

I agree that my thesis/project report to be published as hard copy or online open access.

This thesis is submitted for :

☐

PATENT

Embargo from _____ until _____
(date) (date)

Approved by:

(Signature of Student)
New IC No/ Passport No.:

Date :

(Signature of Chairman of Supervisory Committee)
Name:

Date :

[Note : If the thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach with the letter from the organization/institution with period and reasons for confidentiality or restricted.]